

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The role of histamine in neurogenic inflammation

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/136398> since

Published version:

DOI:10.1111/bph.12266

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Histamine in the neurogenic inflammation

Rosa A.C.¹ and Fantozzi R.¹

¹Dipartimento di Scienza e Tecnologia del Farmaco, University of Turin, Italy

Correspondence: Arianna Carolina Rosa, PhD,

Dipartimento di Scienza e Tecnologia del Farmaco, University of Turin, Italy,

Via P. Giuria 9, 10125, Turin, Italy

Phone: +390116707955

Fax: +390116707688

e-mail: ariannacarolina.rosa@unito.it

Running Title: Histamine in the neurogenic inflammation



UNIVERSITÀ DEGLI STUDI DI TORINO

This is the accepted version of the following article:

Rosa AC, Fantozzi R. The role of histamine in neurogenic inflammation. *Br J Pharmacol.* 2013 Sep;170(1):38-45. doi: 10.1111/bph.12266.

which has been published in final form at
<http://onlinelibrary.wiley.com/doi/10.1111/bph.12266/abstract>

Summary

The term “neurogenic inflammation” has been adopted to describe the local release of inflammatory mediators from neurons such as substance P and calcitonin gene related peptide. Once released, these neuropeptides cause histamine release from adjacent mast cells. In turn, histamine evokes the substance P and calcitonin gene related peptide release, thus a bidirectional link between histamine and neuropeptides in neurogenic inflammation is established. The aim of this review is to summarize the most recent findings on the role of histamine in neurogenic inflammation with particular regards to nociceptive pain as well as neurogenic inflammation in the skin, in the airways and in the bladder.

Keywords histamine, neurogenic inflammation, histamine receptors, neuropathic pain, skin, airways, bladder

Abbreviations calcitonin gene related peptide, CGRP; cell adhesion molecule, CADM; gastro-oesophageal reflux disease, GERD; histidine decarboxylase, HDC; interstitial cystitis, IC; oestrogen receptors, ERs; painful bladder syndrome, PBS; platelet-activating factor, PAF; substance P, SP; vasoactive intestinal peptide, VIP

Introduction

The term “neurogenic inflammation” describes the local release of inflammatory mediators from afferent neurons such as substance P (SP) and calcitonin gene related peptide (CGRP). Since the first evidences of a neurogenic vasodilatation in response to noxious stimuli in the skin of humans and other mammals (Schmelz M *et al.*, 2001), nowadays it is recognised that neurogenic inflammation occurs also in visceral organs. The inflammatory response evoked by the activation of the sensory nerve fibres, including local vasodilatation, plasma extravasation, leukocyte and platelet adhesion, and mast cell degranulation, is brought about by neuropeptides released from peripheral endings of sensory neurons upon stimulation of the primary sensory terminals. Both SP in neurons and histamine in mast cells have a dual mediator role in neurogenic inflammation (Figure 1). In fact, as described by Foreman and Jordan (1984), injury causes activation of sensory nerve endings either directly or through the release of histamine from adjacent mast cells. The action potential generated travels orthodromically to the dorsal horn of the spinal cord, and spreads to other branches of the same neurons which will then release SP from their terminals or varicosities. The released SP, besides contributing by itself to local vasodilatation, causes histamine release from adjacent mast cells which produce flare and further activates other sensory nerve endings (Foreman JC *et al.*, 1984).

Neutrophils, whose responsiveness to SP has been repeatedly demonstrated, contribute to the inflammatory soup following neurogenic inflammation. In fact, neutrophils express the SP receptors NK₁, NK₂ and NK₃, and their challenge with SP induces COX-2 expression and PGE₂ release in the nanomolar range (Gallicchio M *et al.*, 2008; Gallicchio M *et al.*, 2009). Furthermore, SP at micromolar concentrations primes neutrophils, thus enhancing O₂⁻ production evoked by platelet-activating factor (PAF) (Brunelleschi S *et al.*, 1991). In the picomolar range SP still induces neutrophil adhesion to human umbilical vein endothelial cells (Dianzani C *et al.*, 2003).

Together with histamine, other mast cell-derived mediators (such as, serotonin, **renin**, adenosine, heparin, tryptase, chymase, elastase, carboxypeptidase A and B, cathepsin, b-galactosidase, b-glucuronidase, matrix metalloproteinase, chemotactic factors for eosinophils and neutrophils, platelet derived factor, prostaglandin D₂, leukotriene B₄, C₄ and D₄, thromboxane A₂ and B₂, nitric oxide, TNF- α and other cytokines, neuropeptides) contribute to the functional picture of neurogenic inflammation (Tore F *et al.*, 2009). Experimental evidences support their role demonstrating their neuronal receptors.

Histamine and neurogenic inflammation: experimental evidences

The evidences for a neurogenic mechanism in the involvement of SP in the flare response stems from data showing that local anaesthetic injection into the human skin reduced the spread of the flare response although without affecting the development of the wheal response (Foreman JC *et al.*, 1983). Otherwise the role of histamine has been investigated mostly by specific receptor-oriented approaches. The effects of SP on flare response in human skin are mimicked only by the neuropeptide fragments able to degranulate **rat** mast cells *in vitro* (Fewtrell CM *et al.*, 1982). The H₁ histamine antagonist chlorpheniramine (20 mg i.v.) has been reported to prevent the spread of the flare response to SP **in the human skin** (Foreman JC *et al.*, 1983). These first observations of the role of histamine in the neurogenic inflammation have been further elucidated thanks to immunohistochemical studies demonstrating the proximity of mast cells and nerve fibres containing neuropeptides, with first studies in dura mater and gut, while larger body of evidence is available for the skin (Nakanishi M *et al.*, 2008). Moreover, Janiszewski *et al.* (1990) have demonstrated in a co-culture system that the activation of peritoneal mast cells induce depolarization and decrease membrane resistance in sympathetic neurons (Janiszewski J *et al.*, 1990). A bidirectional communication between nerves and mast cells has been supported by the presence of receptors for many neuropeptides (such as vasoactive intestinal peptide, neuropeptide Y, SP, CGRP, proopiomelanocortin, galanin, neuromedin U, pituitary adenylate cyclase-activating polypeptide,

neurotensin and corticotrophin-releasing factor) on mast cells surface. Thus these mediators can exert both paracrine and autocrine effects on mast cells (Figure 1). Similarly, also histamine exerts autocrine effects on mast cells as they express all the histamine receptors H₁₋₄ (Tore F *et al.*, 2009).

Notably, the interactions between mast cells and adjacent nerves is also accomplished by adhesion molecules such as N-cadherin or cell adhesion molecule (CADM)-1, a calcium-dependent adhesion protein that can be cleaved by the metalloproteinase MT5-MMP of the peptidergic receptors in the dorsal root ganglion (Guillot X *et al.*, 2012; Nakanishi M *et al.*, 2008).

Histamine, neurogenic inflammation and nociceptive pain

While the contribution of neurogenic inflammation to nociception has been previously elucidated (Julius D *et al.*, 2001), more recently detailed analysis of the role of histamine receptors has been investigated.

Mechanoinsensitive C-fibres are known to be activated by histamine and to be responsible for the neuropeptide release, for example in the skin inducing the axon reflex flare (Groetzner P *et al.*, 2010). Histamine-immunoreactive nerve fibres have been found in the superficial laminae of the dorsal horn, an important site for nociceptive transmission. The mRNA of histamine H₁ receptor genes has been detected in many SP and CGRP immunoreactive neurons following the peripheral nerve injuries (Kashiba H *et al.*, 1999), and histamine has been reported to mediate the release of SP and glutamate in inflammatory conditions (Riedel W *et al.*, 2001). Moreover, a bidirectional relationship between CGRP and histamine can be proposed according to the data showing that: CGRP induces histamine release from mast cells and potentiates histamine effects in the rat (Mobarakeh JI *et al.*, 2006); intraperitoneally injection of histamine induces CGRP release into the cerebrospinal fluid (Bileviciute I *et al.*, 1994); histamine administered to the nasal mucosa causes CGRP release from peripheral terminals of trigeminal ganglion in the guinea pig (Tani E *et al.*, 1990). Further support of a cooperation between histamine and CGRP is afforded by the

observation of a colocalization of both the histamine H₃ receptor and CGRP on A δ fibres; both mediators contributing to an high threshold mechanical nociceptive effect (Cannon KE *et al.*, 2007).

More recently, it has been demonstrated that both the H₁ receptor antagonist pyrilamine and the H₂ receptor antagonist ranitidine produce a dose-dependent antinociceptive response in the formalin test, and that a histamine intrathecally-induced hyperalgesia has been blunted by the GCRP antagonist CGRP 8-37 (Mobarakeh JI *et al.*, 2011).

Studies with histamine H₁ receptor knockout mice have demonstrated that both the receptor and its natural ligand are necessary to facilitating pain transmission at both peripheral and central levels (Mobarakeh JI *et al.*, 2002; Mobarakeh JI *et al.*, 2000). By using histidine decarboxylase (HDC) gene knockout mice it has been shown that the NK₁ receptors in the spinal cord mediate the histamine-induced hyperalgesic responses (Yoshida A *et al.*, 2005). Several *in vivo* studies have evaluated the antinociceptive efficacy of antihistamines. The administration of H₁ and H₂ receptor antagonists, chlorpheniramine and cimetidine respectively, was found to inhibit the development of both thermal and mechanical hyperalgesia, although chlorpheniramine was more potent (Zuo Y *et al.*, 2003). Even the most recently discovered histamine receptors, the H₃ and H₄ receptors, have been shown to be involved in mediating nociception. The pre-treatment with H₃/H₄ receptors dual antagonist thioperamide attenuated c48/80-induced thermal pain 30 min after challenge without causing analgesia (Chatterjea D *et al.*, 2012). Moreover, it was reported that H₃ receptor antagonists, such as GSK-189254 and ABT-239, are effective in reducing allodynia and hyperalgesia in models of neuropathic, and inflammatory pain (Hsieh GC *et al.*, 2010b; Medhurst AD *et al.*, 2007; Medhurst SJ *et al.*, 2008).

Focusing on H₄ receptors, both the experimental antagonists JNJ7777120 (10 and 30 mg kg⁻¹, s.c.) and VUF6002 (10 mg kg⁻¹, s.c.) significantly reduced the paw oedema and hyperalgesia provoked by subplantar injection of carrageenan (Coruzzi G *et al.*, 2007). According to these data, JNJ7777120 reversed hyperalgesia in both acute and chronic pain models (Hsieh GC *et al.*, 2010a).

Notably, the same study has reported no effect for H₁ receptor antagonist diphenhydramine, H₂ receptor antagonists ranitidine, or H₃ receptor antagonist ABT-239, thus **suggesting** a dominant role of the H₄ receptors in animal models of inflammatory and neuropathic pain.

All together, these data from animal models may be regarded as indicative of a potential efficacy of antihistamines in clinical setting. However, clinical evidence is still lacking and previously trials **were negative** (Raffa RB, 2001).

Histamine and neurogenic inflammation in the skin

The involvement of histamine in cutaneous neurogenic inflammation stems from the observation that this amine triggers the so called “triple response”. Nowadays, local erythema results from the axon reflex and antidromic sensory nerve stimulation-induced release of different vasoactive mediators, not only histamine, but also SP, histamine, purines and CGRP. Itch is mediated via dedicated afferent nonmyelinated C-type nerve fibres different from the polymodal C-fibres being unresponsive to mechanical stimulation. The so called ‘pruriceptors’ are differentiated in histamine-sensitive and histamine-independent itch-specific C-fibres. In particular, histamine-sensitive C-fibres, about the 5% of afferent C-fibres in human cutaneous nerves, are characterized by slow conduction velocities and extensive terminal branching (Raap U *et al.*, 2011; Shim WS *et al.*, 2008). When activated by histamine, they transmit electrical signals to the superficial layer of the dorsal horn of the spinal cord. These signals then ascend to the thalamus through contralateral spinothalamic tracts and are eventually conducted to the somatosensory and cingulate cortex (**Figure 2**). All the histamine receptors are involved in mediating histamine signal on neurons, but with a different functional weight. **The failure of cimetidine to suppress histamine-induced itch in BalbC mice (Bell JK *et al.*, 2004) do not support an involvement of H₂ receptor in itch. The H₁ receptor is the most extensively studied. Indeed, H₁ receptor blockers have an established and valued place in the treatment of itching of allergic and non-allergic origin (Skidgel RA *et al.*, 2011).**

Intriguing data on the H₃ receptor seem to contradict the aforementioned histamine-induced itch pathway (Summey BT *et al.*, 2005), as **antagonists** such as thioperamide or AQ0145 were found to increase significantly the incidence of scratching behaviour in mice (Sugimoto Y *et al.*, 2004). **Histamine H₃ receptor antagonists might block the modulatory role afforded by the presynaptic H₃ receptor, thus favouring the release of neuropeptides from sensory endings (Cannon KE *et al.*, 2007).** More recently, data obtained with clobenoprit, an H₃ receptor antagonists and H₄ receptor agonist, suggested that H₄ receptor also is involved in itch. Clobenoprit caused scratching responses in the mouse that were attenuated by pretreating animals with the H₄ receptor antagonist, JNJ7777120 (Dunford PJ *et al.*, 2007). Moreover, it has been reported that H₄ receptor agonists are able to induce itch through a direct action on peripheral nerves (Bell JK *et al.*, 2004), and that the H₄ antagonist inhibit SP-induced itch, which has been reported to be resistant to H₁ receptor antagonists (Yamaura K *et al.*, 2009). More recently, Suwa *et al.* (2011) have demonstrated that JNJ7777120 (10 and 30 mg kg⁻¹ die), and not the histamine H₁ receptor antagonist fexofenadine (30 mg kg⁻¹ die), reduces the scratching behaviour and ameliorates the skin lesions induced by repeated challenge with 2,4,6-trinitrochlorobenzene in HR-1 mice, in a dose-dependent manner (Suwa E *et al.*, 2011).

Interestingly, in the mice blocking both the H₁ receptors and the H₄ receptors, by a dual antagonist or a combination therapy gave the maximum response, almost completely **abolish** the itch response (Dunford PJ *et al.*, 2007).

Besides, other published data indicate that **H₁ receptors and H₄ receptors** share a similar pathway in the neuron excitation by increasing intracellular Ca²⁺ levels (Shim WS *et al.*, 2008). Histamine elevates calcium levels through H₁ receptor in rat cultured sensory neurons, and this effect is blocked by the PLC inhibitor U73122 (Nicolson TA *et al.*, 2002).

Moreover, in mouse sensory neurons stimulated by capsaicin, histamine induces inward currents and calcium influx in a TRPV1-dependent manner, as demonstrated by the blocking effect of

TRPV1 antagonists; in keeping with these results histamine-induced scratching is significantly lower in TRPV1-deficient mice (Kajihara Y *et al.*, 2010; Shim WS *et al.*, 2007).

All these evidences explain at least in part the antipruritic effects of both mirtazapine and doxepin, two antidepressant that share also additional antihistaminic effects (Raap U *et al.*, 2011). Moreover, the well recognised risk-benefit profile of H₁ antihistamines, especially the non-sedative second generation ones, lead the European Academy of Allergy and Clinical Immunology to recommend in its guidelines an up dosing of non-sedating antihistamines up to four-fold in patients with chronic urticaria. Clinical studies on H₄ receptor antagonists are currently ongoing (Engelhardt H *et al.*, 2009).

Histamine and neurogenic inflammation in the airways

Misery (2008) approached the intriguing parallelism between itch/scratching and cough starting from a common pathophysiology that involves the C fibres, the mast cells, histamine, SP and other tachykinins. **Indeed**, the close proximity between mast cells and the nerve endings in the lung suggests a similar neuro-immuno crosstalk (Misery L, 2008) **as found in the skin**. Sensory nerve endings release neuropeptides such as SP and CGRP, which induce mast cell activation and degranulation. The antidromic release of neuropeptides from nociceptors in the airways causes vasodilatation and oedema associated with nasal obstruction in the upper airways, and broncospasm in the lower airways, as well as plasma protein exudation, mucus secretion and inflammatory cell recruitment. As a proof of neurogenic inflammation in the airways, local anaesthesia with lidocaine improved airway hyperreactivity and reduced capsaicin-induced cough (Muraki M *et al.*, 2008).

A first evidence of the relationship between SP and histamine in the airways came in 1994, when Heaney *et al.* (1994) demonstrated the ability of SP to stimulate human mast cells obtained from bronchoalveolar lavage (Heaney LG *et al.*, 1994). More recently, sputum SP and mast cell tryptase concentrations were remarkably increased in patients with chronic cough, and even increased in

those with cough due to gastro-oesophageal reflux disease (GERD). These latter data suggest that GERD-induced cough may be related to a cough reflex hypersensitivity caused by neurogenic airway inflammation (Qiu Z *et al.*, 2011), and are in keeping with the results obtained by Birring *et al.* (2004), who measured an increased histamine content in the sputum of patients with idiopathic chronic cough and cough variant asthma/eosinophilic bronchitis in comparison with normal subjects (Birring SS *et al.*, 2004). Rat lung mast cells have been found to release histamine in response to high doses of SP *in vitro*; moreover, it has been shown that the NK₁ receptor-mediated mast cell activation partly affords for airway plasma leakage in F344-, but not in BDE-rats, exposed to SP and capsaicin. **The airway responsiveness to tachykinins discriminates these two inbred strains. Actually, only the F344 presents NK₁ receptors mast cells that display a proreleasing effect (Pauwels RA *et al.*, 1995).** More recently, it has been reported that CP-99,994 (5 mg kg⁻¹ i.v.), a NK₁ receptor antagonist, abolished the microvascular leakage elicited in the rat airways by a single inhalation of toluene-2,4-diisocyanate. On the contrary, ketotifen (1 mg kg⁻¹ i.v.), provided with H₁ antagonism and mast cell-stabilizing properties, did not exert any effect in this model (Sakamoto T *et al.*, 2012). In a model of microvascular leakage hypersensitivity induced in the airways of guinea-pigs aerosolised with histamine it has been demonstrated that, while NK_B and the NK₃ receptor agonist senktide, enhanced airway hypersensitivity to histamine. In the same model, both the tachykinin NK₃ receptor antagonists osanetant and the NK₁ receptor antagonist nelpitantium were able to abolish the histamine-induced microvascular leakage. NK₂ ligands were ineffective (Daoui S *et al.*, 2001). Moreover, it has been reported that toluene-2,4-diisocyanate exposure causes an increase in histamine content, HDC activity and gene expression in the nasal mucosa of sensitized rats (Kitamura Y *et al.*, 2004). Olopatadine hydrochloride, an H₁ receptor antagonist, besides inhibiting the capsaicin-induced sneezing response, has been found to inhibit antigen-induced sneeze and nasal rubbing responses in both wild-type and H₁ receptor-deficient mice, although at very high doses (Tamura T *et al.*, 2008), thus suggesting a H₁ receptor independent effect. Indeed, these data, together with the failure of cetirizine to inhibit completely these responses in H₁

receptor-deficient mice (Kayasuga R *et al.*, 2002; Sugimoto Y *et al.*, 2004), could lead to hypothesise the involvement of other receptors in these responses.

Histamine and neurogenic inflammation in the bladder

Interstitial cystitis (IC), or painful bladder syndrome (PBS), numbers among its causes neurogenic inflammation; in fact, an increased density of nerve fibres has been reported (Pecker R *et al.*, 2000).

Evidences from both rodent and humans highlight an important role for mast cells. Activated mast cells have been associated to the rodent neurogenic cystitis induced by the Bartha strain of pseudorabies virus (PRV) (Chen MC *et al.*, 2006; Jasmin L *et al.*, 2000). Moreover, clinical studies have demonstrated elevated mast cell number in the lamina propria of IC bladder biopsies (Leiby BE *et al.*, 2007) and an increased urinary histamine metabolites (el-Mansoury M *et al.*, 1994). Although the central role of mast cells in PBS/IC is still unclear, *in vivo* model of IC pathogenesis suggested a positive feedback loop with SP containing peripheral nerves and mast cells: the activation of the bladder-associated circuits in the CNS initiates SP release by peripheral nerves in the bladder leading SP-mediated mast cell activation. Consequently, mast cell degranulation induces bladder inflammation by acting on urothelium. Histamine contribute to IC/PBS, is not only in evoking an inflammatory response, but seems also to be related to pelvic pain. In fact, histamine and histamine receptors as mediators in pain responses have been described in both animal models and humans (Mobarakeh JI *et al.*, 2006; Thilagarajah R *et al.*, 2001). Although the specific cell type among mast cells, basophils, neutrophils, dendritic cells and histaminergic neurons regulating histamine-mediated pain is unknown, it has been demonstrated, by using a model of PRV-induced pelvic pain in mast cell deficient $\text{Kit}^{\text{W-sh}}/\text{Kit}^{\text{W-sh}}$ mice, that mast cells are required for histamine-mediated pelvic pain (Rudick 2008). However, the same authors demonstrated that $\text{Kit}^{\text{W-sh}}/\text{Kit}^{\text{W-sh}}$ mice reconstituted with $\text{HDC}^{-/-}$ bone marrow exhibited diminished pain, thus suggesting that non-mast cell sources of histamine may also contribute to pain. Moreover, Rudick CN *et al.* (2008) demonstrated that PRV induced pelvic pain is independent from TNF-dependent pathology and

instead is mediated by mast cell histamine, which then induces pain via histamine receptors H₁ and H₂ (Rudick CN *et al.*, 2008). Thus, it has been suggested that antagonists of histamine H₁ and H₂ receptors are candidates for clinical trials in the treatment of chronic pain conditions, such as IC-related pelvic pain. Indeed, pilot clinical studies suggest that antihistamine therapy could be effective on IC-related pelvic pain. In particular, a symptomatic improvement has been reported in 30% of patients treated with the H₁ receptor antagonist hydroxyzine hydrochloride from 25 mg/die at night to 50 mg at night + 25 mg in the morning over a 2-week period (Theoharides TC *et al.*, 1997). Although these positive results suggest a therapeutic role of histamine H₁ receptor antagonist in the treatment of IC/PBS, further studies with newer generation histamine H₁ receptor antagonists, with a lesser sedation component, have to be conducted. Moreover, the histamine H₂ receptor antagonist cimetidine produced significant improvement in pain and nocturia in a limited trial of PBS patients (Thilagarajah R *et al.*, 2001). It has to be stressed that approximately 90% of patients with PBS/IC are women, thus suggesting that the process involving mast cells may be hormonally influenced. In fact, oestrogen receptors (ERs) are expressed on human mast cells and can mediate their degranulation, while the ER antagonist tamoxifen inhibits this phenomenon (Rudick CN *et al.*, 2012).

All together these findings suggest that reproductive hormones may modulate IC symptoms at the mast cell levels. This hypothesis has been recently tested by Rudick CN *et al.* (2012) who assessed the basis of gender specific pelvic pain in the murine model of neurogenic cystitis PRV-induced. The data obtained suggest that pelvic pain in mice with murine neurogenic cystitis is mediated by gender specific responsiveness to mast cells, in contrast pelvic pain severity resulted to be modulated by genetic factors (Rudick CN *et al.*, 2012).

Conclusion

In conclusion, experimental data here included widely show the complex physiopathological mechanism(s) known as neurogenic inflammation and the key role played by histamine (Table 1).

Neurogenic inflammation is involved in several processes of animal and human physiopathology: this review, with no ambition to be completely exhaustive, has anyway outlined the most up-to-date and widest fields of investigation. **A comprehensive evaluation of the histamine role in neurogenic inflammation, as discussed here, highlights how the many experimental evidences have not yet reached a full clinical transferability and not yet support new pharmacotherapeutic approaches.**

Acknowledgements

This work was supported by the University of Turin, Royal College of Anaesthesia/BJA and COST Action BM0806.

Conflict of interest

None.

References

Bell JK, McQueen DS, Rees JL (2004) Involvement of histamine H4 and H1 receptors in scratching induced by histamine receptor agonists in Balb C mice. *Br J Pharmacol* 142: 374-380.

Bileviciute I, Lundeberg T, Ekblom A, Theodorsson E (1994) Substance P-, neurokinin A-, calcitonin gene-related peptide- and neuropeptide Y-like immunoreactivity (-LI) in rat knee joint synovial fluid during acute monoarthritis is not correlated with concentrations of neuropeptide-LI in cerebrospinal fluid and plasma. *Neurosci Lett* 167: 145-148.

Birring SS, Parker D, Brightling CE, Bradding P, Wardlaw AJ, Pavord ID (2004) Induced sputum inflammatory mediator concentrations in chronic cough. *Am J Respir Crit Care Med* 169: 15-19.

Brunelleschi S, Tarli S, Giotti A, Fantozzi R (1991) Priming effects of mammalian tachykinins on human neutrophils. *Life Sci* 48: PL1-5.

Cannon KE, Chazot PL, Hann V, Shenton F, Hough LB, Rice FL (2007) Immunohistochemical localization of histamine H3 receptors in rodent skin, dorsal root ganglia, superior cervical ganglia, and spinal cord: potential antinociceptive targets. *Pain* 129: 76-92.

Chatterjea D, Wetzel A, Mack M, Engblom C, Allen J, Mora-Solano C, et al. (2012) Mast cell degranulation mediates compound 48/80-induced hyperalgesia in mice. *Biochem Biophys Res Commun* 425: 237-243.

Chen MC, Blunt LW, Pins MR, Klumpp DJ (2006) Tumor necrosis factor promotes differential trafficking of bladder mast cells in neurogenic cystitis. *J Urol* 175: 754-759.

Coruzzi G, Adami M, Guaita E, de Esch IJ, Leurs R (2007) Antiinflammatory and antinociceptive effects of the selective histamine H4-receptor antagonists JNJ7777120 and VUF6002 in a rat model of carrageenan-induced acute inflammation. *Eur J Pharmacol* 563: 240-244.

Daoui S, Ahnaou A, Naline E, Emonds-Alt X, Lagente V, Advenier C (2001) Tachykinin NK(3) receptor agonists induced microvascular leakage hypersensitivity in the guinea-pig airways. *Eur J Pharmacol* 433: 199-207.

Dianzani C, Collino M, Lombardi G, Garbarino G, Fantozzi R (2003) Substance P increases neutrophil adhesion to human umbilical vein endothelial cells. *Br J Pharmacol* 139: 1103-1110.

Dunford PJ, Williams KN, Desai PJ, Karlsson L, McQueen D, Thurmond RL (2007) Histamine H4 receptor antagonists are superior to traditional antihistamines in the attenuation of experimental pruritus. *J Allergy Clin Immunol* 119: 176-183.

el-Mansoury M, Boucher W, Sant GR, Theoharides TC (1994) Increased urine histamine and methylhistamine in interstitial cystitis. *J Urol* 152: 350-353.

Engelhardt H, Smits RA, Leurs R, Haaksma E, de Esch IJ (2009) A new generation of anti-histamines: Histamine H4 receptor antagonists on their way to the clinic. *Curr Opin Drug Discov Devel* 12: 628-643.

Fewtrell CM, Foreman JC, Jordan CC, Oehme P, Renner H, Stewart JM (1982) The effects of substance P on histamine and 5-hydroxytryptamine release in the rat. *J Physiol* 330: 393-411.

Foreman JC, Jordan CC (1984) Neurogenic inflammation. *Trends in Pharmacological Sciences* 5: 116-119.

Foreman JC, Jordan CC, Oehme P, Renner H (1983) Structure-activity relationships for some substance P-related peptides that cause wheal and flare reactions in human skin. *J Physiol* 335: 449-465.

Gallicchio M, Benetti E, Rosa AC, Fantozzi R (2008) Substance P-induced cyclooxygenase-2 expression in polymorphonuclear cells. *Inflamm Res* 57 Suppl 1: S17-18.

Gallicchio M, Benetti E, Rosa AC, Fantozzi R (2009) Tachykinin receptor modulation of cyclooxygenase-2 expression in human polymorphonuclear leucocytes. *Br J Pharmacol* 156: 486-496.

Groetzner P, Weidner C (2010) The human vasodilator axon reflex - an exclusively peripheral phenomenon? *Pain* 149: 71-75.

Guillot X, Semerano L, Decker P, Falgarone G, Boissier MC (2012) Pain and immunity. *Joint Bone Spine* 79: 228-236.

Heaney LG, Cross LJ, Stanford CF, Ennis M (1994) Differential reactivity of human bronchoalveolar lavage mast cells to substance P. *Agents Actions* 41 Spec No: C19-21.

Hsieh GC, Chandran P, Salyers AK, Pai M, Zhu CZ, Wensink EJ, et al. (2010a) H4 receptor antagonism exhibits anti-nociceptive effects in inflammatory and neuropathic pain models in rats. *Pharmacol Biochem Behav* 95: 41-50.

Hsieh GC, Honore P, Pai M, Wensink EJ, Chandran P, Salyers AK, et al. (2010b) Antinociceptive effects of histamine H3 receptor antagonist in the preclinical models of pain in rats and the involvement of central noradrenergic systems. *Brain Res* 1354: 74-84.

Janiszewski J, Bienenstock J, Blennerhassett MG (1990) Activation of rat peritoneal mast cells in coculture with sympathetic neurons alters neuronal physiology. *Brain Behav Immun* 4: 139-150.

Jasmin L, Janni G, Ohara PT, Rabkin SD (2000) CNS induced neurogenic cystitis is associated with bladder mast cell degranulation in the rat. *J Urol* 164: 852-855.

Julius D, Basbaum AI (2001) Molecular mechanisms of nociception. *Nature* 413: 203-210.

Kajihara Y, Murakami M, Imagawa T, Otsuguro K, Ito S, Ohta T (2010) Histamine potentiates acid-induced responses mediating transient receptor potential V1 in mouse primary sensory neurons. *Neuroscience* 166: 292-304.

Kashiba H, Fukui H, Morikawa Y, Senba E (1999) Gene expression of histamine H1 receptor in guinea pig primary sensory neurons: a relationship between H1 receptor mRNA-expressing neurons and peptidergic neurons. *Brain Res Mol Brain Res* 66: 24-34.

Kayasuga R, Sugimoto Y, Watanabe T, Kamei C (2002) Participation of chemical mediators other than histamine in nasal allergy signs: a study using mice lacking histamine H(1) receptors. *Eur J Pharmacol* 449: 287-291.

Kitamura Y, Miyoshi A, Murata Y, Kalubi B, Fukui H, Takeda N (2004) Effect of glucocorticoid on upregulation of histamine H1 receptor mRNA in nasal mucosa of rats sensitized by exposure to toluene diisocyanate. *Acta Otolaryngol* 124: 1053-1058.

Leiby BE, Landis JR, Propert KJ, Tomaszewski JE (2007) Discovery of morphological subgroups that correlate with severity of symptoms in interstitial cystitis: a proposed biopsy classification system. *J Urol* 177: 142-148.

Medhurst AD, Atkins AR, Beresford IJ, Brackenborough K, Briggs MA, Calver AR, et al. (2007) GSK189254, a novel H3 receptor antagonist that binds to histamine H3 receptors in Alzheimer's disease brain and improves cognitive performance in preclinical models. *J Pharmacol Exp Ther* 321: 1032-1045.

Medhurst SJ, Collins SD, Billinton A, Bingham S, Dalziel RG, Brass A, et al. (2008) Novel histamine H3 receptor antagonists GSK189254 and GSK334429 are efficacious in surgically-induced and virally-induced rat models of neuropathic pain. *Pain* 138: 61-69.

Misery L (2008) Are pruritus and scratching the cough of the skin? *Dermatology* 216: 3-5.

Mobarakeh JI, Sakurada S, Hayashi T, Orito T, Okuyama K, Sakurada T, et al. (2002) Enhanced antinociception by intrathecally-administered morphine in histamine H1 receptor gene knockout mice. *Neuropharmacology* 42: 1079-1088.

Mobarakeh JI, Sakurada S, Katsuyama S, Kutsuwa M, Kuramasu A, Lin ZY, et al. (2000) Role of histamine H(1) receptor in pain perception: a study of the receptor gene knockout mice. *Eur J Pharmacol* 391: 81-89.

Mobarakeh JI, Takahashi K, Sakurada S, Kuramasu A, Yanai K (2006) Enhanced antinociceptive effects of morphine in histamine H2 receptor gene knockout mice. *Neuropharmacology* 51: 612-622.

Mobarakeh JI, Torkaman-Boutorabi A, Rahimi AA, Ghasri S, Nezhad RM, Hamzely A, et al. (2011) Interaction of histamine and calcitonin gene-related peptide in the formalin induced pain perception in rats. *Biomed Res* 32: 195-201.

Muraki M, Iwanaga T, Haraguchi R, Kubo H, Tohda Y (2008) Continued inhalation of lidocaine suppresses antigen-induced airway hyperreactivity and airway inflammation in ovalbumin-sensitized guinea pigs. *Int Immunopharmacol* 8: 725-731.

Nakanishi M, Furuno T (2008) Molecular basis of neuroimmune interaction in an in vitro coculture approach. *Cell Mol Immunol* 5: 249-259.

Nicolson TA, Bevan S, Richards CD (2002) Characterisation of the calcium responses to histamine in capsaicin-sensitive and capsaicin-insensitive sensory neurones. *Neuroscience* 110: 329-338.

Pauwels RA, Germonpre PR, Kips JC, Joos GF (1995) Genetic control of indirect airway responsiveness in the rat. *Clin Exp Allergy* 25 Suppl 2: 55-60.

Peeker R, Aldenborg F, Dahlstrom A, Johansson SL, Li JY, Fall M (2000) Increased tyrosine hydroxylase immunoreactivity in bladder tissue from patients with classic and nonulcer interstitial cystitis. *J Urol* 163: 1112-1115.

Qiu Z, Yu L, Xu S, Liu B, Zhao T, Lu H (2011) Cough reflex sensitivity and airway inflammation in patients with chronic cough due to non-acid gastro-oesophageal reflux. *Respirology* 16: 645-652.

Raap U, Stander S, Metz M (2011) Pathophysiology of itch and new treatments. *Curr Opin Allergy Clin Immunol* 11: 420-427.

Raffa RB (2001) Antihistamines as analgesics. *J Clin Pharm Ther* 26: 81-85.

Riedel W, Neeck G (2001) Nociception, pain, and antinociception: current concepts. *Z Rheumatol* 60: 404-415.

Rudick CN, Bryce PJ, Guichelaar LA, Berry RE, Klumpp DJ (2008) Mast cell-derived histamine mediates cystitis pain. *PLoS One* 3: e2096.

Rudick CN, Pavlov VI, Chen MC, Klumpp DJ (2012) Gender specific pelvic pain severity in neurogenic cystitis. *J Urol* 187: 715-724.

Sakamoto T, Kamijima M, Miyake M (2012) Neurogenic airway microvascular leakage induced by toluene inhalation in rats. *Eur J Pharmacol* 685: 180-185.

Schmelz M, Petersen LJ (2001) Neurogenic inflammation in human and rodent skin. *News Physiol Sci* 16: 33-37.

Shim WS, Oh U (2008) Histamine-induced itch and its relationship with pain. *Mol Pain* 4: 29.

Shim WS, Tak MH, Lee MH, Kim M, Koo JY, Lee CH, et al. (2007) TRPV1 mediates histamine-induced itching via the activation of phospholipase A2 and 12-lipoxygenase. *J Neurosci* 27: 2331-2337.

Skidgel RA, Kaplan AP, Erdös EG (2011) Histamine, Bradykinin, and Their Antagonists. In: Brunton LL (ed). *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. 12th edn. The McGraw-Hill Companies, Inc. pp 911-935.

Sugimoto Y, Iba Y, Nakamura Y, Kayasuga R, Kamei C (2004) Pruritus-associated response mediated by cutaneous histamine H3 receptors. *Clin Exp Allergy* 34: 456-459.

Summey BT, Yosipovitch G (2005) Pharmacologic advances in the systemic treatment of itch. *Dermatol Ther* 18: 328-332.

Suwa E, Yamaura K, Oda M, Namiki T, Ueno K (2011) Histamine H(4) receptor antagonist reduces dermal inflammation and pruritus in a hapten-induced experimental model. *Eur J Pharmacol* 667: 383-388.

Tamura T, Komai M (2008) Effect of olopatadine hydrochloride, an anti-histamine drug, on rhinitis induced by intranasal instillation of toluene-2,4-diisocyanate in rats. *Int Immunopharmacol* 8: 916-921.

Tani E, Senba E, Kokumai S, Masuyama K, Ishikawa T, Tohyama M (1990) Histamine application to the nasal mucosa induces release of calcitonin gene-related peptide and substance P from peripheral terminals of trigeminal ganglion: a morphological study in the guinea pig. *Neurosci Lett* 112: 1-6.

Theoharides TC, Sant GR (1997) Hydroxyzine therapy for interstitial cystitis. *Urology* 49: 108-110.

Thilagarajah R, Witherow RO, Walker MM (2001) Oral cimetidine gives effective symptom relief in painful bladder disease: a prospective, randomized, double-blind placebo-controlled trial. *BJU Int* 87: 207-212.

Tore F, Tuncel N (2009) Mast cells: target and source of neuropeptides. *Curr Pharm Des* 15: 3433-3445.

Yamaura K, Oda M, Suwa E, Suzuki M, Sato H, Ueno K (2009) Expression of histamine H4 receptor in human epidermal tissues and attenuation of experimental pruritus using H4 receptor antagonist. *J Toxicol Sci* 34: 427-431.

Yoshida A, Mobarakeh JI, Sakurai E, Sakurada S, Orito T, Kuramasu A, et al. (2005) Intrathecally-administered histamine facilitates nociception through tachykinin NK1 and histamine H1 receptors: a study in histidine decarboxylase gene knockout mice. *Eur J Pharmacol* 522: 55-62.

Zuo Y, Perkins NM, Tracey DJ, Geczy CL (2003) Inflammation and hyperalgesia induced by nerve injury in the rat: a key role of mast cells. *Pain* 105: 467-479.

Table 1. Effect of selective histamine receptor antagonism in neurogenic inflammation

Effect	Histamine receptor subtype			
	H₁R	H₂R	H₃R	H₄R
Pain	↓	↓	↓	↓
Hyperalgesia*	↓	↓	↓	↓
Allodynia	?	?	↓	?
Itch	↓	↔	↑	↓
Sneezing response	↓**	?	?	?

*The differential involvement of histamine receptor subtypes has been demonstrated by different experimental models

**only at high doses

Legends

Figure 1. The dual mediator role of mast cells: bidirectional interaction nerves-mast cells.

Neuropeptide released from sensory nerve endings stimulate adjacent mast cells in a receptor dependent manner. Mediators released from mast cells act in both a paracrine and autocrine fashion.

CGRP= calcitonin gene related peptide; SP= substance P; VIP= vasoactive intestinal peptide

Figure 2. The itch system.

Histamine released from mast cells after a local stimulus activates histamine-sensitive pruriceptors, thus generating an actions potential which orthodromically travels, through to the dorsal horn of the spinal cord and the thalamus, to the cortex (blue line). The following antidromic stimulation induces the release of different mediators from sensory endings, substance P (SP) and calcitonin gene related peptide (CGRP). SP and CGRP released causes further mast cells degranulation resulting in vasodilatation (flare) and the recruitment of other pruriceptors.