AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The Grey Goat of Lanzo Valleys (Fiurinà): Breed characteristics, genetic diversity, and quantitative-qualitative milk traits

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/142786> since 2016-01-08T10:24:43Z

Published version:

DOI:10.1016/j.smallrumres.2013.10.006

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in ***Small Ruminant Research*, volume 116(2014), pages 1 – 13, available online 23 October 2013**, <http://dx.doi.org/doi:10.1016/j.smallrumres.2013.10.006>.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ *Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform*]

The Grey Goat of Lanzo Valleys (*Fiurinà*): breed characteristics, genetic diversity, and quantitative-qualitative milk traits

P. Cornale^{a*}, M. Renna^a, C. Lussiana^a, D. Bigi^b, S. Chessa^c, and A. Mimosi^a

^aDipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Grugliasco (TO) 10095, Italy

^bDipartimento di Scienze e Tecnologie Agro-Alimentari – Sezione Allevamenti Zootecnici, Università di Bologna, Reggio Emilia 42123, Italy

^cIstituto di Biologia e Biotecnologia Agraria, CNR Milano, Milano 20133, Italy

E-mail addresses: paolo.cornale@unito.it; manuela.renna@unito.it;

carola.lussiana@unito.it; daniele.bigi@unibo.it; chessa@ibba.cnr.it;

antonio.mimosi@unito.it

*Corresponding author: Paolo Cornale, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, via L. da Vinci, 44 - Grugliasco (Torino) 10095, Italy, +39 011 6708576, +39 011 6708563, paolo.cornale@unito.it

ABSTRACT

The aim of this study was to provide an overview on breed's characteristics of a small dairy goat population recently identified in the Piedmont region (NW Italy): the Grey Goat of Lanzo Valleys. This goat, locally named *Fiurinà*, is composed of about 150 heads. Increasing knowledge on threatened breeds is an effective tool to develop

conservation programs aiming at preserving loss of genetic resources, economically valorizing animal-derived food products, maintaining traditions and cultural values, as well as appropriately managing native habitats. Data and samples were collected in 15 representative farms to obtain information on geographical distribution, breeding systems, morphometric measures, and genetic diversity. Milk traits, including fatty acids profile and caseins polymorphism, were also investigated. The prevalent breeding system is extensive or semi-extensive, mostly with vertical transhumance from lowland to alpine pastures during the grazing season. The breed is characterized by a peculiar color of the fleece (mixture of white, grey and black coarse outer hairs with brown under-down) from which the local name “*Fiurinà*” (=speckled) derives. The breed is of medium size, with a quite high frequency of well-developed and turned backward horns. More than 80% of does have pear-shaped udder, typical of goats, with cylindrical teats directed downward. The microsatellite analysis pointed out significant distances between *Fiurinà* and other goat breeds reared in the same area. Consequently, *Fiurinà* has to be considered as a unique breed. Although highly variable, quite appreciable milk yields were observed. Considering milk gross composition and fatty acids profile, *Fiurinà* goat showed comparable results to other local breeds reared in alpine environment and mainly fed with fresh and conserved forages. A remarkable amount of α -linolenic acid ($0.82 \text{ g } 100 \text{ g}^{-1} \text{ fat}$) was detected. The breed is also characterized by an interesting and wide variability in the casein cluster, with some haplotypes (i.e., *A-C-F-C'*, *E-A-C-B*, *F-C_I-F-C'*) detected only in *Fiurinà*, confirming the genetic uniqueness of the breed. The balanced frequency of medium-strong and weak-null *CSN1S1* alleles could be exploited for different breeding strategies.

Key Words: local goat breed, breeding system, genetic diversity, milk quality.

INTRODUCTION

The positive trend in the number of goats around the world (+58%) occurred from 1980 to 2000 has recently been confirmed in the last 10 years when goat stocks increased by about 23%. These data are even more interesting if compared to the modest increase in cattle and sheep stocks (+9% and +3%, respectively). The growing success in goat farming is noticed worldwide, in developing countries as well as in industrialized areas (FAOSTAT, 2010).

Goats play an important role in rural economy of developing countries, especially in areas with unfavorable environmental and climatic conditions, thanks to their capacity for adaptation to very different environments and because they are important users of marginal and rural lands (Kalantzopoulos et al., 2004). In developed countries, goats are presently considered as 'ecological' animals and seem to adequately respond to increasing consumers' demand concerning product quality, animal welfare, and environment respect (Morand-Fehr et al., 2004). In these contexts, goat dairy products often gain niche market reaching higher prices than other dairy products. Furthermore, goat milk has some distinctive traits, such as high digestibility and low allergenicity, which bring it to be widely considered the alternative to bovine milk, especially for people with allergies and other gastrointestinal disorders (El-Agamy, 2007; Park et al., 2007).

The worldwide increase in goat stocks has also been possible with the diffusion of some specialized breeds (e.g., Saanen). However, the widespread use of a reduced number of high producing breeds led to a dramatic reduction of autochthonous ones, placing most of them in an endangered status. In Europe, for example, although there is only about 4% of the world's goat population, there is the largest share of goat genetic resources, approximately 33% (Galal, 2005). However, the 35% of European

76 local breeds are endangered (Bertaglia et al., 2007).

77 Effective management of farm animal resources requires comprehensive
 78 knowledge of the breeds' characteristics, including data on population size and
 79 structure, geographical distribution, production environment, and within- and
 80 between-breed genetic diversity (Groeneveld et al., 2010). Such an exhaustive
 81 overview on breeds' characteristics, as well as on products' quality, represents the
 82 starting point for the development of a preservation program (Canali, 2006; Lauvie et
 83 al., 2011; Verrier et al., 2005). An effective safeguard of a local breed prevents
 84 extinction and erosion of genetic resources and allows the economic valorization of
 85 derived products, the maintenance of traditions and cultural values as well as the
 86 appropriate management of native habitats (Dubeuf, 2011; Rosa García et al., 2012).

87 In Italy, goats are mainly reared in the Centre-South, where the age-long
 88 dairying tradition counts several cheeses obtained from sheep and goat milk (Pirisi et
 89 al., 2011). Nevertheless, goats are also reared in the Italian alpine regions where dairy
 90 products are recently increasing their relevance on the market. An Italian native goat
 91 dairy population, officially named Grey Goat of Lanzo Valleys (*Capra Grigia delle*
 92 *Valli di Lanzo*) and locally known as "*Fiurinà*" has been recently identified in the
 93 Piedmont region (Cornale et al., 2012, 2010).

94 To prove comprehensive information, the goat population has been the subject
 95 of a multidisciplinary project concerning with population size, geographical
 96 distribution, breeding systems, morphometric measures, and genetic diversity
 97 (microsatellite and mitochondrial DNA analyses). Furthermore, since the main
 98 purpose of the *Fiurinà* breed is milk production, quanti-qualitative traits of milk were
 99 investigated, including fatty acid profile, for its well known human health
 100 implications, and caseins polymorphisms, for their relation with milk composition and

101 technological properties.

102

103

104 MATERIALS AND METHODS

105 The *Fiurinà* goat population is reared in the Piedmont region (NW Italy). All
106 the investigated farms breeding the *Fiurinà* goats are located in uplands and highlands
107 of the northwestern territories of the Torino province.

108 The 3-year research project on the *Fiurinà* breed lasted from the beginning of
109 2009 to the end of 2011, and was developed on two main research lines. The first
110 topic included the collection of data related to population size, geographical
111 distribution, farm management, morphometric characteristics, as well as the analysis
112 of genetic diversity to evaluate if the *Fiurinà* goat could be considered a new breed.
113 These data were collected between 2009 and 2010 by examining 15 farms, breeding
114 approximately the 50% of the *Fiurinà* population. The second topic focused on milk
115 production and quality, including fatty acids and caseins polymorphism analyses.
116 These surveys were carried out between 2010 and 2011, by collecting milk samples in
117 11 representative farms.

118

119 *Morphometric Measurements*

120 Body measurements are of primary importance for breed phenotypical
121 description and they represent a starting point in the definition of breed standards
122 (Zaitoun et al., 2005; Dossa et al., 2007). In the 15 investigated farms, all reproductive
123 females (n = 77), corresponding approximately to 50% of the registered population,
124 were measured.

125 The goats were weighed using a spring balance. Height at withers, rump height,

126 rump width, trunk length, and chest girth were also measured by using a flexible tape
 127 or a Lydtin stick. With the animal standing upright, the height at withers and the rump
 128 height were measured as the distance from the floor to the shoulders and to the
 129 highest point of the rump, respectively. The rump width was measured as the distance
 130 between the pin bones. The trunk length was the distance between the crown and the
 131 sacrococcygeal joint. The chest girth was the circumference of the thoracic cavity
 132 taken just behind the forelimbs.

133 A five-point scale was carried out to assess traits of udder morphology based on
 134 the scoring system proposed by de la Fuente et al. (1996) for ewes and modified for
 135 dairy goats as stated below. The udder depth was evaluated by considering the
 136 position of the udder floor with respect to hock (score 3: at the same hock's level;
 137 score <3: above the hock; score >3: below the hock). The fore udder attachment
 138 evaluates the strength of the attachment of the lateral ligaments to the body wall. The
 139 score can range from extremely loose (score 1) to snug and strong (score 5)
 140 attachment. The rear udder attachment was evaluated through the insertion to the
 141 abdominal wall by assessing the udder height. The cistern shape was assessed through
 142 the degree of separation between the left and right halves (score 1, pronounced
 143 separation; score 5, no separation). Finally, halves' symmetry was recorded.

144 Concerning teats' traits, their shape, inclination, and orientation were assessed.
 145 The teats' shape can range from short and small (score 1) to squat and cone-shaped
 146 (score 5). The physiological inclination of the teats is almost vertical (score 3), but it
 147 can be directed cranially (score 1) or caudally (score 5). The teats' orientation can be
 148 divergent (score 1), parallel (score 3) or convergent (score 5). Furthermore, symmetry
 149 of teats and presence/absence of supernumerary teats were also evaluated. All the
 150 morphometric assessments were performed by the three same operators.

151

152 ***Genetic Diversity***

153 In order to characterize the genetic diversity of the *Fiurina* breed, DNA was
 154 extracted from hair root samples using proteinase k digestion.

155 ***Microsatellite analysis.*** A group of *Fiurina* (n = 26) reared in 15 different farms
 156 was compared with two officially recognized autochthonous Piedmontese breeds,
 157 Sempione (n = 22) and Vallesana (n = 36), reared in other 26 farms located in the
 158 same region. The individuals were selected in order to avoid close relatives (the
 159 number of samples collected per farm varied between 1 and 6). The evaluation of the
 160 genetic structure was investigated using 12 microsatellites (INRA005, MAF65,
 161 INRA063, MCM527, ETH10, SRCRSP5, INRA023, OarFCB20, TGLA53, SRC247,
 162 CSRD247, ILST87), most of them recommended by ISAG/FAO (FAO, 2004).
 163 Genotype determinations were performed on an ABI Prism 3100 DNA Sequencer,
 164 equipped with Genscan and Genotyper software.

165 ***Mitochondrial DNA analysis.*** A subsample of 10 animals belonging to the
 166 *Fiurina* breed was also analyzed for the mitochondrial DNA (mtDNA). The goats
 167 were selected by respecting phenotypic standards, maximizing geographical spread of
 168 the farms, and avoiding relationships among the sampled flocks. The primer pairs
 169 CAP-F (5'-CGTGTATGCAAGTACATAC-3') and CAP-R (5'-
 170 CTGATTAGTCATTAGTCCATC-3') and the amplification conditions were used to
 171 determine the sequence of the mtDNA control region, as described by Luikart et al.
 172 (2001). Amplified products were used for sequencing with the CAP-F or CAP-R
 173 primer. Sequence reactions were performed for both DNA strands by means of an
 174 ABI Prism 3100 DNA Sequencer, according to the manufacturer's instructions. All
 175 sequences were deposited in GenBank (Accession Numbers JQ655153-JQ655162).

176 The sequences of mtDNA d-loop region from 10 sequences of the *Fiurina* goats
 177 belonging to this study were aligned with the mtDNA complete sequence of *Capra*
 178 *hircus* (GenBank NC_005044) using ClustalX, version 2.0.11 (Thompson et al.,
 179 1997). The same software was used to compare the *Fiurina* goats' sequences with the
 180 mtDNA of the 22 reference individuals of the 6 domestic goat haplogroups reported
 181 by Naderi et al. (2007). The alignments were imported in MEGA, version 4.0 (Kumar
 182 et al., 2004) and a neighbour-joining haplotype tree was constructed using Kimura 2-
 183 parameter distance model with 1,000 bootstrap replications.

184

185 ***Milk Yield, Gross Composition and Fatty Acids Analysis***

186 The individual daily milk yield was measured by using recording jars. In the
 187 selected 11 farms, individual milk samples were collected from 52 goats during
 188 lactation. Two aliquots of each individual milk sample were collected during the
 189 morning milking, immediately stored at 4°C in a portable refrigerator, and transported
 190 to the laboratory. One aliquot (50 mL) was then immediately analyzed for fat, protein,
 191 lactose, casein, urea, solids-non-fat (MilkoScan FT 6000, Foss Electric, Hillerød,
 192 Denmark), and somatic cell count (SCC) (Fossomatic 5000, Foss Electric, Hillerød,
 193 Denmark). The other aliquot (150 mL) was frozen at -20°C and successively
 194 analyzed for the FA composition as previously reported by Renna et al. (2012). Milk
 195 fat extraction was obtained by centrifugation at 7,300 rpm for 30 min at -4°C. The
 196 resulting molten butter was filtered through a hydrophobic filter (Whatman 1,
 197 Whatman International Ltd, Maidstone, England). The pure milk fat was then
 198 dissolved in 5 mL of internal standard solution (nonanoic acid in heptane) and fatty
 199 acid methyl esters (FAME) were obtained by trans-esterification of glycerides by
 200 using a solution of potassium hydroxide in methanol (IOfS, 2002). FAME were

analyzed by high-resolution gas chromatography (Shimadzu GC17A, Shimadzu Corporation Analytical Instruments Division, Kyoto, Japan) with flame ionization detector according to Collomb and Bühler (2000). FAME were separated on a CP-Sil 88 capillary column (100 m × 0.25 mm ID, 0.20 µm film thickness; Varian Inc., Lake Forest, CA, USA). The column temperature was held at 45°C for 5 min, then raised 20°C min⁻¹ up to 195°C and maintained for 65 min. The temperatures of the injector and detector were maintained at 250°C and 280°C, respectively. The injection volume was 0.1 µL. Nitrogen constant linear flow rate was set at 40 mL min⁻¹. Peaks were identified by comparing their retention times with pure FAME standards (Matreya Inc., Pleasant Gap, PA, USA and Restek Corporation, Bellefonte, PA, USA). Quantification was assessed by using nonanoic acid as internal standard. The results are expressed as both absolute values (g 100g⁻¹ fat) and percentages of each FAME per total FAME detected.

214

215 ***Genotyping Analysis of Caseins Variability***

Milk was also used as starting material for DNA extraction. Individual milk samples were collected from 52 does randomly chosen in the selected 11 farms. The GFX Genomic Blood DNA Purification kit (Amersham Biosciences, Piscataway, NJ, USA) was used for DNA extraction directly from milk. Extraction was performed starting with 300 µL of milk to recover a 130 µL final volume of genomic DNA solution. All samples were analyzed using a NanoDrop ND-1000 UV-Vis Spectrophotometer (NanoDrop Technologies, Inc., Wilmington, DE, USA) to assess DNA concentration and purity. To analyze most of the 50 alleles described (Küpper et al., 2010) at the casein gene cluster (19, 8, 7 and 16 variants for *CSN1S1*, *CSN2*, *CSN1S2* and *CSN3*, respectively), the samples were typed by 8 different methods, first

screening the major alleles by means of the PCR - Single Strand Conformation Polymorphism (SSCP) methods and then applying the PCR - Allele Specific (AS) and the PCR-RFLP methods available to discriminate alleles grouping together within the PCR-SSCP, as described in Table 1. All the PCR protocols were applied using at least 25 ng of DNA.

231

232 *Statistical Analysis*

The software Arlequin ver. 3.11 (Excoffier et al., 2005) was used to calculate: i) the number of alleles per locus, the observed and expected heterozygosity; ii) to test for Hardy-Weinberg Equilibrium using exact test and sequential Bonferroni correction; iii) to compute Wright's F_{IS} , F_{ST} and F_{IT} fixation indexes (Weir and Cockerham, 1984; Wright, 1965) and to evaluate the significance of genetic differentiation between populations with an Analysis of Molecular Variance (AMOVA, Excoffier et al., 1992). Reynolds's genetic distance (Reynolds et al., 1983) was obtained using GENEDIST, included in the PHYLIP computer package, version 3.65 (Felsenstein, 1989). FSTAT software (Goudet, 2001) was used to calculate the allelic richness (AR) standardized for variation in sample size.

Means and standard deviations (or median and interquartile range, depending on the considered variables) of morphometric data as well as of chemical and hygiene parameters of milk were calculated with the MEANS procedure of SAS (SAS Institute, 2008).

The casein alleles and haplotypes distributions were analyzed by the ALLELE and HAPLOTYPE procedures of SAS (SAS Institute, 2008). The ALLELE procedure uses the notation and concepts described by Weir (1996). Haplotype frequencies were calculated under the null hypothesis of no linkage disequilibrium and under the

alternative hypothesis of associations between casein genes.

Haplotype frequencies were also analyzed by the PRINCOMP procedure of SAS (SAS Institute, 2008) to underline the differences among the *Fiurinà* goats and other Italian goat breeds (Caroli et al., 2006; Martini et al., 2010).

RESULTS AND DISCUSSION

Breed Geographical Distribution and Size

The *Fiurinà* goats are mainly reared for milk production in the territories of Lanzo Valleys (longitude: 7°48'03''E; latitude: 45°27'56''N) from which the official name Grey Goat of Lanzo Valleys (*Capra Grigia delle Valli di Lanzo*) derives. This area, gathering three alpine valleys (*Val Grande di Lanzo*, *Val d'Ala*, and *Valle di Viù*), is located in the northwestern area of the Torino province. Farmers rearing the *Fiurinà* subjects have been also reported in adjacent areas (*Val Susa* and *Canavese*) of the same region. All investigated farms are located between 400 and 1,400 m a.s.l. About 150 heads out of 1,250 goats in the territories of Lanzo Valleys are identified as belonging to the *Fiurinà* breed, distributed in 56 farms, which correspond to the 82% of goat farms in the area. The average herd size is small, ranging from 13 to 42 goats. Only in few farms the flock is mainly composed of *Fiurinà* subjects, otherwise few *Fiurinà* heads are reared with other cosmopolitan (e.g., Camosciata delle Alpi) or local (e.g., Valdostana) goat breeds, or even crossbreds. The presence of few subjects per farm justifies the high percentage of goat farms in the area rearing the *Fiurinà* breed.

Breeding Systems

276 As commonly happens in the alpine territories, all investigated farms are of
 277 small and medium size and the most common types of business are family and single-
 278 worker types. Historically, farms chiefly base their livestock activity on bovine milk
 279 production. Goat milk and meat often represent complementary activities, especially
 280 because of the high ability of small ruminants to exploit marginal mountain areas,
 281 which are otherwise not utilized.

282 Extensive or semi-extensive farming (highly dependent on grazing) is generally
 283 practiced in all farms. In winter months (from November to March-April) the goats
 284 are stabled indoors and mainly fed with local hay. The supplementation, with cereals
 285 (corn and wheat bran) and, occasionally, with other feedstuffs (e.g., chestnuts), is
 286 limited and depends on both availability of hay and nutrient requirements of the goats.
 287 As soon as fresh grass is available, the flocks are moved outdoors and the goats graze
 288 on pastures located near the farms. In just one of the investigated farms, the goats are
 289 always housed and fresh cut grass is offered in through. Vertical transhumance (from
 290 lowland to alpine pastures) is diffused in 87% of the investigated farms. The alpine
 291 grazing season averages from April-May to October-November, in accordance with
 292 weather conditions and fresh forages availability. After alpine grazing, the goats graze
 293 on pastures near the farms until the winter arrival. Consequently, the feeding system
 294 is mostly based on local fresh and conserved forages. Although in free grazing
 295 conditions goats usually utilize trees and shrubs in woodland due to their feeding
 296 behavior (Dwyer, 2009), a regional law generally forbids goats to graze in forests.

297 As stated above, the main purpose of the *Fiurina* goat is milk production. Hand-
 298 milking is practiced in 100% of the investigated farms. The absence of machine-
 299 milking is probably related to the small flock size that does not justify such an
 300 investment. After kids weaning, goat milk, often mixed with bovine milk, is processed

301 at farm level into fresh and matured cheeses and directly sold to consumers.

302 The reproductive career of does is quite long, reaching sometimes 15 years.
 303 Artificial insemination is not used and natural mating occurs with the introduction of
 304 a buck in the flock of all the investigated farms. Average age of does at first kidding is
 305 13 - 14 months. Mating season aims at converging kidding in autumn-winter months
 306 to ensure the availability of kids in Christmas and, particularly, in Easter times. Kids
 307 are maintained with the mother until slaughtering weight (10 - 12 kg). Concerning
 308 reproductive performance, fertility rate (calculated as the percentage of does that kid
 309 per does exposed to buck), fecundity rate (calculated as the percentage of kids born
 310 alive per does exposed to buck), and prolificacy (calculated as the percentage of kids
 311 per does kidding) were 92%, 136%, and 147%, respectively. The *Fiurinà* breed
 312 showed a slightly higher (+2%) fertility rate and lower fecundity rate and prolificacy
 313 (-8% and -13%, respectively) if compared to the Saanen goat reared in the same
 314 region (Bigi and Zanon, 2008). Higher results (95%, 152%, and 160%) were reported
 315 for the Camosciata delle Alpi reared in the same area (Deitos, 2001).

316

317 ***Morphometric Characteristics***

318 The *Fiurinà* breed is of medium size, alertly graceful. The hair is medium to
 319 short. This breed is characterized by a peculiar color of the fleece (mixture of white,
 320 grey and black course outer hairs with brown under-down) from which the local name
 321 “*Fiurinà*” (=speckled) derives (Figure 1). Legs are long and well developed, allowing
 322 the breed to easily move in unfavorable territories such as steep pastures. The head is
 323 straight and light, with quite long upright ears. The neck is long and thin in females,
 324 much short and muscular in males. Beards and wattles are occasionally present in
 325 both bucks and does. All characteristics refer to the typical shape of the dairy goat

breeds (Lucifero, 1981). The breed showed a quite high frequency of well-developed and turned backward horns in males (85.71%), while only 60% of females have horns, usually shorter than in males.

Average morphometric traits measured on adult females of the *Fiurinà* breed are reported in Table 2. By comparing the results with morphometric measurements of other cosmopolitan (Saanen and Camosciata delle Alpi) and local (Sempione and Vallesana) breeds commonly reared in the same area, the *Fiurinà* does showed a slightly reduced size. Such characteristic makes grazing easier in mountain marginal areas and *Fiurinà* appeared to be well adapted to environmental and climatic conditions of alpine valleys.

Udder morphology is normally assessed in dairy ewes (Caja et al. 2000, Casu et al., 2006) and in dairy goats (Capote et al., 2006; Peris et al., 1999; Salama et al., 2004). These authors agreed with the importance of udder morphology assessment in relation to milkability, especially when related to machine milking. Although hand milking is diffused in all investigated farms, the evaluation of udder traits appears to be relevant because of some relationships pointed out between morphological and productive traits (i.e., milk yield) (Casu et al., 2006). Therefore, such information should be taken into account in a future genetic selection program. Concerning udder morphology, more than 80% of the considered subjects had pear-shaped udder, typical of goats, with cylindrical teats directed downward (85.19%). A weak median suspensory ligament, resulting in the udder floor below the hock, rarely occurred in the *Fiurinà* goats (<2%). The udder was usually well supported, with the floor at the same level of the hock (16.67%) or slightly above (>80%). This is important for the udder functionality because, as reported by Altınçekiç and Koyuncu (2011) for dairy ewes, grazing animals with deep udder are more exposed to accidental injuries.

351 Concerning fore udder attachment, none of the goats showed a snug and strong
 352 attachment. Lateral ligaments were intermediate in strength and tightness in 35% of
 353 the goats, while the remaining ones showed a score <3 . The height of the rear udder
 354 attachment is an indication of the goat's potential capacity for milk production.
 355 Almost 90% of the goats obtained a score ≥ 3 . Only 11% of the goats had a low udder
 356 height. Usually the cistern was well shaped and only a small percentage of subjects
 357 showed a pronounced separation of the halves ($<7\%$) or a separation almost absent
 358 ($<9\%$). The symmetry in two halves was detected in almost 80% of the goats.

359 The ease of milking is particularly reflected in teats' evaluation. A good teat
 360 should be cylindrical, with a constant diameter and should have a medium length (Le
 361 Du and Benmederbel, 1984). In the *Fiurinà* population less than 6% of goats had very
 362 narrow teats. The majority (68.52%) showed good cylindrical shape, while tight or
 363 funnel shaped teats occurred in about 1/4 of the goats. Teats' inclination was almost
 364 vertical in more than 85% of goats. The remaining heads showed teats cranially
 365 inclined. Caudal inclination was not detected. Similarly, none of the goats showed
 366 teats pointed inward. Teats were almost parallel (28%) or slightly pointed outward
 367 (72%). More than 85% of goats had symmetric teats and supernumerary ones were
 368 present only in one doe.

369

370 ***Genetic Diversity***

371 ***Microsatellites***. A total of 103 alleles were detected across the 12 microsatellites
 372 loci analyzed, which were polymorphic in the three considered breeds (*Fiurinà*,
 373 Sempione, and Vallesana). The number of alleles ranged between 4 (ET10) to 17
 374 (SRC247). Observed and expected heterozygosities, AR and F_{IS} for each goat breed
 375 are presented in Table 3. The highest value of AR was found in Sempione (6.41) and

the lowest in Vallesana (5.80) while *Fiurinà* showed an intermediate value (6.19). The *Fiurinà* goat showed the lowest level of observed (0.59) and expected (0.63) heterozygosities. F_{IS} value within populations was statistically significant only for Vallesana breed due to a deficiency of heterozygosity. Out of the total of 36 Hardy-Weinberg equilibrium tests, only two (both in Vallesana breed) gave significant deviations at the 1% level.

F statistics per microsatellite locus are shown in Table 4. As expected, because of the close geographical origin of the three goat populations, levels of apparent breed differentiation were quite low. The average F_{ST} , which was significantly different from zero ($P<0.001$), indicated that about 3% of the total genetic variation was explained by differences among breeds, with the remaining 97% corresponding to differences among individuals. Genetic differentiation among breeds was significant ($P<0.001$) only for two loci (SRC247 and ILST87). A significant excess of homozygotes across all breeds ($P<0.001$) was found for MCM527, CSRD247, and ILST87 loci. On average, breeds had a 7.6% ($P<0.001$) deficit of heterozygotes, whereas the total population had a 10.3% ($P<0.001$) deficit of heterozygotes.

Reynolds's genetic distance values ranged from 0.031 between Sempione and Vallesana to 0.062 between *Fiurinà* and Sempione; F_{ST} values expressed similar relationships between the same pairwise breed combinations (Table 5). The distance between *Fiurinà* and the other two breeds is therefore greater than the distance between Sempione and Vallesana. Pairwise F_{ST} values between *Fiurinà* and the other two breeds reached significant levels ($P<0.001$), indicating that *Fiurinà* can be considered as a separate breed. F_{ST} value between Sempione and Vallesana was instead significant only at $P<0.05$. Some other genetic studies about Italian goat breeds have been carried out using microsatellite markers (Iamartino et al., 2005;

Negrini et al., 2012), but no information about Piedmontese goat breeds have been reported till now.

Mitochondrial DNA. The mtDNA fragments of the d-loop region in the 10 *Fiurinà* samples were highly polymorphic, with 51 variable sites over the 481 bp of the alignment (Figure 2). A total of 9 haplotypes were identified.

As shown in Figure 3, using the available goat mtDNA haplogroup classification system, 8 *Fiurinà* goats could be classified into haplogroup A, whereas the remaining 2 animals were assigned to haplogroup C. Six mitochondrial haplogroups A, B, C, D, G, and F have been identified by many authors (Naderi et al., 2007; Royo et al., 2009; Sardina et al. 2006). The most ancient population expansion is probably represented by the haplogroup A, which is observed worldwide with high frequencies (Luikart et al., 2001; Royo et al., 2009) ranging from 89% in Asia to 98% in Europe (Pereira et al., 2005). Haplogroups B and D were found in Asia while haplogroup C was found in Asia and Europe with low frequencies. Haplogroups B and C can be the consequence of a second domestication in Asia, with a more recent expansion (Luikart et al., 2001). A haplogroup G has been found in the Middle East and North Africa (Naderi et al., 2007) while the haplogroup F was limited to Sicily (Sardina et al., 2006).

Haplogroup A is the most observed haplogroup among the 20 Italian goat breeds that have been studied till now (Luikart et al., 2001; Naderi et al., 2007; Vacca et al., 2010). The Girgentana breed represents an exception, because it showed the haplogroup F (Sardina et al., 2006). Apart from *Fiurinà* goat, sequences belonging to the haplogroup C have been recently found in the Sarda breed (Piras et al., 2012), even if with very low frequencies (0.008). This finding gives a further element of

interest on the *Fiurina* breed and its genetic distinctiveness compared to the other breeds reared in the same region.

Milk Yield and Gross Composition

The average length of lactation was about 200 days. A high variability in milk yield was observed: the recorded levels varied from less than 1.0 to approximately 3.5 L head⁻¹ day⁻¹. Although the performance was variable, such result has to be considered quite appreciable, particularly for a local breed usually fed with fresh or conserved forages with only a limited and occasional use of concentrates.

The main milk constituents are reported in Table 6. They were in the range reported by Park et al. (2007) for goat milk, with the exception of the average protein content that showed a slightly lower value (29.9 g kg⁻¹ compared to 34.0 g kg⁻¹). The obtained results agreed also with those reported by Raynal-Ljutovac et al. (2008) for goat milk, including the protein content (reported lower limit: 26.1 g kg⁻¹).

Considering milk gross composition of other goat breeds (Bigi and Zanon, 2008) reared in the same area, *Fiurina* reported comparable protein but higher milk fat levels (*Fiurina*: 35.8 g kg⁻¹; Vallesana: 30.3 g kg⁻¹; Sempione: 31.9 g kg⁻¹). It is worth mentioning that, besides breed, milk main constituents depend on several factors, such as feeding, lactation stage, season, etc.

While in the United States the legal limit established for milk SCC is 1,000 × 10³ cells mL⁻¹, currently there is no legal limit for goat milk in the European Union (Paape et al., 2007). The observed value for the *Fiurina* milk (median: 490 × 10³ cells mL⁻¹) was largely included in the threshold limit of 1,500,000 cells mL⁻¹ proposed by Delgado-Pertinhez et al. (2003) and even in the more restrictive grading scheme

suggested by Leitner et al. (2008): grade A (the best) with $\text{SCC} \leq 840 \times 10^3$ cells mL^{-1} .

451

452 ***Milk Fatty Acids Profile***

453 The individual FA and the groups of FA in *Fiurina* milk are presented in Tables
454 7 and 8, respectively. Results showed that five FA (C10:0, C14:0, C16:0, C18:0, and
455 C18:1 *c9*) accounted for 72% of total FA. Caproic (C6:0), caprylic (C8:0), and capric
456 (C10:0) acids are among the most characteristic FA in goat milk and derived dairy
457 products, being more abundant than in cow milk fat. They are named after the species
458 name (*Capra hircus*) and are generally associated with the characteristic flavor of
459 goat cheeses (Mele et al., 2008). Mainly caprylic and capric acids have become
460 established medical treatments for a wide range of clinical disorders, being
461 consequently considered of particular importance in human nutrition (Haenlein, 2004).
462 In milk fat from the *Fiurina* goat caproic, caprylic, and capric acids accounted on
463 average for 2.23, 2.43, and 7.35 g 100g^{-1} fat, respectively.

464 Among goat milk FA, odd- and branched-chain fatty acids are also responsible
465 for the typical aroma of caprine milk and cheese (Alonso et al., 1999). These FA can
466 be almost exclusively found in dairy products from ruminants and have received
467 increasing attention by researchers in recent years due to their anticancer properties
468 (Oku and Yanagita, 2009; Parodi, 2009). Linear odd pentadecanoic (C15:0) and
469 heptadecanoic (C17:0) acids accounted for the majority of total detected odd- and
470 branched-chain fatty acids; their average concentrations were equal to 0.80 and 0.71 g
471 100g^{-1} milk fat. Among branched-chain FA, the most abundant ones in *Fiurina* milk
472 fat were the *iso* and *aiso* forms of pentadecanoic and heptadecanoic acids, the *iso*
473 form of hexadecanoic (C16:0) acid and the *aiso* form of octadecanoic (C18:0) acid;

474 such results confirm previous findings for goat milk fat (Massart-Leën et al., 1981;
475 Alonso et al., 1999; Žan et al., 2006).

476 Considering unsaturated FA, either FA of nutritional interest such as conjugated
477 linoleic acids (CLA) and omega-3 FA or presumably negative FA such as *trans* fatty
478 acids (TFA) can be found in milk and dairy products from ruminants. The acronym
479 CLA refers to a mixture of positional and geometric isomers of octadecadienoic acid,
480 with double bonds located in adjacent carbon atoms. Many beneficial biological
481 effects have been attributed to CLA in animal models of human diseases, including
482 anticarcinogenic, antidiabetic, antiaterogenic, and antiinflammatory properties (Park,
483 2009). Appreciable amounts of CLA are usually reported in goat milk as well as in
484 other ruminant-derived food products. Results of FA analysis showed that total CLA
485 in *Fiurina* milk was 0.64 g 100g⁻¹ fat. In the applied chromatographic conditions the
486 most represented among CLA isomers, rumenic acid (C18:2 *c9t11*), coeluted with
487 other two isomers (CLA *t7c9* and *t8c10*). As usually occurs in milk fat from
488 ruminants (Parodi, 2009), the sum of these three isomers accounted for more than
489 90% of total CLA.

490 Omega-3 FA have been shown to possess positive health effects in chronic
491 diseases including cancer, insulin resistance and cardiovascular diseases (Anderson
492 and Ma, 2009). Increasing levels of these FA in dairy food products are thus pursued.
493 The sum of omega-3 FA in *Fiurina* milk was equal to 1.25 g 100g⁻¹ fat. Remarkable
494 amounts of α -linolenic acid (C18:3 *c9c12c15*, ALA), the most abundant among
495 omega-3 FA in ruminant-derived food products, were found. Detected ALA levels, in
496 fact, averaged 1.02% of total detected FA; such value was more than doubled
497 compared to the mean value (0.42%) reviewed by Park et al. (2007) for goat milk fat.

498 In ruminant milk fat, the predominant TFA have a chain length of 18 carbon atoms.
 499 Both mono- and diunsaturated TFA, especially the latter ones, have been reported to
 500 increase risk factors for coronary heart diseases (Baylin et al., 2003). In milk from
 501 *Fiurina* they accounted for 2.70 and 1.47 g 100 g⁻¹ fat. The most abundant among
 502 TFA in ruminant derived food products is vaccenic acid (Precht et al., 2001).
 503 Differently from other TFA, vaccenic acid has been reported to exert protective
 504 effects against cardiovascular diseases (Wang et al., 2012). In the applied
 505 chromatographic conditions, vaccenic acid's peak coeluted with those of other *trans*-
 506 octadecenoic isomers (C18:1 *t*6 - 10), showing a concentration equal to 1.98 g 100 g⁻¹
 507 fat. Detected monounsaturated TFA with less or more than 18 carbon atoms were
 508 C14:1 *t*, C16:1 *t*, C17:1 *t*, and C20:1 *t*. These FA were found only in low or very low
 509 concentrations in *Fiurina* milk fat, similarly to previous observations in cow and
 510 sheep milk (Abilleira et al., 2009; Collomb et al., 2008).

511 The obtained results regarding the FA composition of *Fiurina* milk are in
 512 accordance with range values previously reported for goat milk in the literature (Park
 513 et al., 2007). Of particular note is the remarkable ALA level found in *Fiurina* milk,
 514 which deserves positive considerations. Similar ALA amounts were previously
 515 observed in milk from dairy goat breeds reared in the alpine environment and
 516 managed according to extensive pasture-based systems (Žan et al., 2006).

517

518 ***Milk Caseins Polymorphism***

519 The extraction of DNA from milk samples gave DNA of varying quality and
 520 quantity. The estimated concentration of the samples ranged from 4.39 to 71.57 ng/μL.
 521 The 260/280 ratios ranged from 1.22 to 1.93 with 13 samples having a 260/280 ratio

inferior to 1.5 and the remaining a mean 260/280 ratio of 1.67. Despite the limited purity and concentration of DNA, all the samples but one gave good PCR products.

A total of 19 alleles were found in the breed and 14 had a frequency higher than 0.05 (Table 9). In particular, 7 variants were found in the *CSN1S1*, and 4 in each of the other 3 casein genes. Genetic equilibrium was generally found at each gene, except for a significant deviation at *CSN2*, in which an excess of homozygotes CC occurred, due to the linkage of this variant with the θ_I allele at the *CSN1S1*.

It is well known that goat caseins are characterized by different expression levels, distinguishing alleles responsible for strong, medium, weak or null casein content in milk, depending on the casein fraction (Chiatti et al., 2007; Küpper et al., 2010). The 4 strong alleles found at the *CSN1S1* had a total frequency of 0.135, even lower than the frequency of the null allele 0 (0.198). The medium allele E and the weak allele F occurred with the same frequency (0.333), giving a population with a quite balanced proportion of medium-strong and weak-null alleles at the *CSN1S1*. A null allele was found also at the *CSN2*, even if only at the heterozygous status in one individual. The same distribution was found for the rare alleles *CSN1S1**A' (0.010) and *CSN1S1**B' (0.010), thus far described only in German (Küpper et al., 2010) and African breeds (Caroli et al., 2007), respectively.

Of the 240 haplotypes expected from the possible combinations of casein genes, only 18 and 5 showed association frequencies higher than 0.008 and 0.05, respectively. The predominant haplotype was the *CSN1S1**E-*CSN2**A-*CSN1S2**A-*CSN3**B, whereas the ancestral one B-A-A-B (Caroli et al., 2006) was found only with a frequency 0.029, due to the low frequency of the *CSN1S1**B allele (0.063) in the analyzed breed. Other two haplotypes were quite frequent: θ_I -C-B-A (0.194) and F- C_I -F-A (0.171). Other two haplotypes carrying the weak *CSN1S1**F allele associated

547 with *CSN2*CI* complete the list of the haplotypes occurred with a frequency higher
 548 than 0.05: *F-C_I-F-B* (0.061) and *F-C_I-F-C* (0.051). The 5 haplotypes represent more
 549 than 76% of the casein cluster variability in the *Fiurinà* breed. Interestingly the
 550 *CSN1S1*F* allele was found always in association with the *CSN2*C_I* allele and the
 551 association of the *CSN1S2*B* allele with the *CSN1S1*0_I* allele, previously described
 552 in the Frisa Valtellinese breed (Caroli et al., 2006), was confirmed also in *Fiurinà*.
 553 The individual heterozygous for the *CSN2*0'* allele carries the haplotypes *A-0'-A-*
 554 *A/F-C_I-F-B* with a probability of 0.963, thus including the *CSN2*0'* allele in a
 555 haplotype combination never described before.

556 The *Fiurinà* breed is characterized by an interesting and wide variability in the
 557 casein cluster, represented by alleles and haplotypes spread in different breeds (i.e., *F-*
 558 *C-F-A*, *F-C-F-B*, *E-A-A-B* haplotypes), rarely found in other breeds (i.e., *F-C-F-C* in
 559 the White Shorthaired, as described by Sztankóová et al. in 2009 or *F-C-C-B* in the
 560 Bunte Deutsche Edelziege, as described by Küpper et al. in 2010), and detected only
 561 in the *Fiurinà* breed (i.e., *A-C-F-C'*, *E-A-C-B*, *F-C_I-F-C'* haplotypes). Haplotypes
 562 frequencies found in *Fiurinà* were also compared with haplotypes frequencies of three
 563 breeds reared in the Lombardy region (Frisa Valtellinese, Orobica, and Verzaschese),
 564 one in the Tuscan region (Garfagnina), and the Italian cosmopolitan Camosciata delle
 565 Alpi breed analyzed in previous studies (Caroli et al., 2006; Martini et al., 2010).
 566 Only alleles with a frequency higher than 0.01 were considered for the haplotype
 567 reconstruction and *CSN2*C* and *CSN2*CI* were considered together because in all
 568 but the *Fiurinà* and Garfagnina breeds no test for the *CSN2*CI* was carried out. As it
 569 can be seen in Figure 4, the Principal Component Analysis clearly separates *Fiurinà*
 570 from all the other breeds. The first three principal components (Prin1, Prin2 and
 571 Prin3) accounted for the 0.33, the 0.31 and the 0.16 of the variability, respectively.

Both the presence of rare and unique haplotypes in the *Fiurinà* breed and the results of the Principal Component Analysis confirm the genetic uniqueness of this breed found in the genetic diversity analysis carried out in the present study.

The balanced frequency of medium-strong and weak-null *CSN1S1* alleles in *Fiurinà* could be exploited for different breeding strategies. Because almost all *Fiurinà* milk is currently processed into cheese, the cheesemaking aptitude could be improved by selecting haplotypes carrying medium-strong alleles. Otherwise, the selection of haplotypes carrying weak-null alleles could be used for fresh milk consumption. Indeed, even if further studies will be needed to clearly assess the relationship between goat casein genotypes and milk protein tolerability, it has already been proven that particular *CSN1S1* genotypes carrying null or weak alleles can reduce the intolerance of allergic subjects in specific cases (Ballabio et al., 2011).

CONCLUSIONS

Despite the limited number of *Fiurinà* goats, the results of the present paper support the increasing interest of local farmers towards this native goat population. The genetic uniqueness of the *Fiurinà* breed was found with genetic diversity analysis and also supported by the presence of rare and unique haplotypes discovered in caseins polymorphism analysis. The adaptability to the native territories as well as its ability of exploiting local feed resources, are promising aspects that can be improved through an appropriate breeding selection. Moreover, the appreciable milk yields, the average gross composition, and the good fatty acids profile, characterized by a remarkable amount of α -linolenic acid, are interesting results for dairy products made with *Fiurinà* milk.

The results of the present study constitute a preliminary but fundamental step in the rescuing process of this autochthonous goat breed. An important result in the success of this process was the recognition by Regional authority. In a recent modification of the Rural Development Programme (Regione Piemonte, 2012) the *Fiurinà* goat has been inserted in the list of the threatened breeds and breeder can now receive financial support for their *Fiurinà* heads.

The *in situ* preservation program of the *Fiurinà* goats can help the safeguard of this population and give real economic opportunities to local farmers, consequently promoting the native territories with their traditions.

ACKNOWLEDGMENTS

The authors wish to gratefully thank the farmers for their willingness to take part in the project. The research was funded by Regione Piemonte (*Direzione Regionale Agricoltura - Settore Sviluppo delle Produzioni Zootecniche*).

REFERENCES

- Abilleira, E., Collomb, M., Schlichtherle-Cerny, H., Virto, M., de Renobales, M., Barron, L.J.R., 2009. Winter/spring changes in fatty acid composition of farmhouse Idiazabal cheese due to different flock management systems. *J. Agric. Food Chem.* 57, 4746–4753.
- Alonso, L., Fontecha, J., Lozada, L., Fraga, M.J., Juarez, M., 1999. Fatty acid composition of caprine milk: major, branched-chain, and trans fatty acids. *J. Dairy Sci.* 82, 878–884.

- Altınçekiç, Ş.Ö., Koyuncu M., 2011. Relationship between udder measurements and the linear scores for udder morphology traits in Kıvrıkcık, Tahirova and Karacabey Merino ewes. *Kafkas Univ. Vet. Fak. Derg.* 17(1), 71–76.
- Anderson, B. M., Ma, D.W.L., 2009. Are all n-3 polyunsaturated fatty acids created equal? *Lipids Health Dis.* doi:10.1186/1476-511X-8-33.
- Ballabio, C., Chessa, S., Rignanese, D., Gigliotti, C., Pagnacco, G., Terracciano, L., Fiocchi, A., Restani, P., Caroli, A.M., 2011. Goat milk allergenicity as a function of α S1-casein genetic polymorphism. *J. Dairy Sci.* 94, 998–1004.
- Baylin, A., Kabagambe, E.K., Ascherio, A., Spiegelman, D., Campos, H., 2003. High 18:2 trans-fatty acids in adipose tissue are associated with increased risk of nonfatal acute myocardial infarction in Costa Rican adults. *J. Nutr.* 133, 1186–1191.
- Bertaglia, M., Joost, S., Roosen, J., the Econgene Consortium, 2007. Identifying European marginal areas in the context of local sheep and goat breeds conservation: A geographic information system approach. *Agric. Syst.* 94, 657–670.
- Bigi, D., Zanon, A., 2008. *Atlante delle razze autoctone. Bovini, equini, ovicaprini, suini allevati in Italia*, first ed. Edagricole, Bologna, Italy.
- Caja, G., Such, X., Rovai, M., 2000. Udder morphology and machine milking ability in dairy sheep. In: *Proceedings of the 6th Great Lakes Dairy Sheep Symposium*, 2–4 November 2000, Guelph, Canada, pp. 17–40.
- Canali, G., 2006. Common agricultural policy reform and its effects on sheep and goat market and rare breed conservation. *Small Rumin. Res.* 62, 207–213.
- Capote, J., Argüello, A., Castro, N., Lopez, J.L., Caja, G., 2006. Short communication: correlations between udder morphology, milk yield, and milk

- 647 ability with different milking frequencies in dairy goats. *J. Dairy Sci.* 89, 2076–
648 2079.
- 649 Caroli, A., Chiatti, F., Chessa, S., Rignanese, D., Bolla, P., Pagnacco, G., 2006.
650 Focusing on the goat casein gene complex. *J. Dairy Sci.* 89, 3178–3187.
- 651 Caroli, A., Chiatti, F., Chessa, S., Rignanese, D., Ibeagha-Awemu, E., Erhardt, G.,
652 2007. Characterization of the casein gene complex in West Africa goats and
653 description of a new α S1-casein polymorphism. *J. Dairy Sci.* 90, 2989–2996.
- 654 Casu, S., Pernazza, I., Carta, A., 2006. Feasibility of a linear scoring methods of udder
655 morphology for the selection scheme of Sardinian sheep. *J. Dairy Sci.* 89, 2200–
656 2209.
- 657 Chessa, S., Rignanese, D., Chiatti, F., Radeghieri, A., Gigliotti, C., Caroli, A., 2008a.
658 Technical note: simultaneous identification of CSN1S2 A, B, C, and E alleles in
659 goats by polymerase chain reaction-single strand conformation polymorphism. *J.*
660 *Dairy Sci.* 91, 1214–1217.
- 661 Chessa, S., Rignanese D., Küpper J., Pagnacco G., Erhardt G., Caroli A., 2008b.
662 □ Short communication: the β -casein (CSN2) silent allele C1 is highly spread □ in
663 goat breeds. *J. Dairy Sci.* 91, 4433–4436.
- 664 Chiatti, F., Chessa, S., Bolla, P., Cigalino, G., Caroli, A., Pagnacco, G., 2007. Effect
665 of κ -casein polymorphism on milk composition in Orobica goat. *J. Dairy Sci.* 90,
666 1962–1966.
- 667 Collomb, M., Bühler, T., 2000. Analyse de la composition en acides gras de la graisse
668 de lait Optimisation et validation d’une méthode générale à haute résolution.
669 *Travaux de chimie alimentaire et d’hygiène* 91, 306–332.
- 670 Collomb, M., Bisig, W., Butikofer, U., Sieber, R., Bregy, M., Etter, L., 2008. Fatty
671 acid composition of mountain milk from Switzerland: comparison of organic and

- intergrated farming systems. *Int. Dairy J.* 18, 976–982.
- Cornale, P., Renna, M., Battaglino, J.J., Lussiana, C., Battaglini, L.M., Mimosi, A., Bianchi, M. 2010. La capra Grigia delle Valli di Lanzo: una nuova razza del territorio alpino piemontese, in: Bovolenta, S. (Ed.), *Zootecnia e Montagna: quali strategie per il futuro?* 6° Quaderno SoZooAlp. Nuove Arti Grafiche, Trento, pp. 207–215.
- Cornale, P., Renna, M., Lussiana, C., Bigi, D., Chessa, S., Mimosi, A. 2012. Characterization of the Grey Goat of Lanzo Valleys (*Fiurinà*), a new Italian dairy population. Books of Abstracts of the XI International Conferences on Goats, Gran Canaria, Spain, September 23–27, p. 184.
- Cosenza, G., Illario, R., Rando, A., Di Gregorio, P., Masina, P., Ramunno, L., 2003. Molecular characterization of the goat CSN1S101 allele. *J. Dairy Res.* 70, 237–240.
- Deitos, E., 2001. L'allevamento caprino nelle Valli di Lanzo: dalle antiche tradizioni alle nuove tecniche. MS Thesis. University of Torino, Italy.
- de la Fuente, L.F., Fernandez, G., San Primitivo, F., 1996. A linear evaluation system for udder traits of dairy ewes. *Livest. Prod. Sci.* 45, 171–178.
- Delgado-Pertiñez, M., Alcalde, M.J., Guzmán-Guerrero, J.L., Castel, J.M., Mena, Y., Caravaca, F., 2003. Effect of hygiene-sanitary management on goat milk quality in semi-extensive systems in Spain. *Small Rumin. Res.* 47, 51–61.
- Dossa, L.H., Wollny, C., Gauly, M., 2007. Spatial variation in goat populations from Benin as revealed by multivariate analysis of morphological traits. *Small Rumin. Res.* 73, 150–159.
- Dubeuf, J.-P., 2011. The social and environmental challenges faced by small livestock local activities: present contribution of research–development and stakes for the

- 697 future. *Small Rumin. Res.* 98, 3–8.
- 698 Dwyer, C., 2009. The Behaviour of Sheep and Goats, in: Jensen, P. (Ed.), The
 699 Ethology of Domestic Animals. An Introductory Text. CAB International,
 700 Wallingford, UK, pp. 161–176.
- 701 El-Agamy, E.I., 2007. The challenge of cow milk protein allergy. *Small Rumin. Res.*
 702 68, 64–72.
- 703 Excoffier, L., Smouse, P., Quattro, J., 1992. Analysis of molecular variance inferred
 704 from metric distances among DNA haplotypes: Application to human
 705 mitochondrial DNA restriction data. *Genetics* 131, 479–491.
- 706 Excoffier, L., Laval, G., Schneider, S., 2005. Arlequin ver. 3.0: An integrated
 707 software package for population genetics data analysis. *Evolutionary*
 708 *Bioinformatics* 1, 47–50.
- 709 FAO, 2004. Secondary Guidelines: Measurement of Domestic Animal Diversity
 710 (MoDAD): New Recommended Microsatellite Markers. Food and Agriculture
 711 Organization, Rome, Italy. Accessed Feb. 15, 2012. [http://dad.fao.org/cgi-](http://dad.fao.org/cgi-bin/getblob.cgi?sid=-1,50006220)
 712 [bin/getblob.cgi?sid=-1,50006220](http://dad.fao.org/cgi-bin/getblob.cgi?sid=-1,50006220).
- 713 FAOSTAT, 2010. Live animals. Stocks (Heads). Accessed Dec. 20, 2010.
 714 <http://faostat.fao.org/site/573/DesktopDefault.aspx?PageID=573#ancor>.
- 715 Felsenstein, J., 1989. PHYLIP - Phylogeny Inference Package (Version 3.2).
 716 *Cladistics* 5, 164–166.
- 717 Galal, S., 2005. Biodiversity in goats. *Small Rumin. Res.* 60, 75–81.
- 718 Goudet, J., 2001. FSTAT, a program to estimate and test gene diversities and fixation
 719 indices (version 2.9.3). Université de Lausanne, Lausanne, Suisse. Accessed Feb.
 720 15, 2012. [http:// www2.unil.ch/popgen/softwares/fstat.htm](http://www2.unil.ch/popgen/softwares/fstat.htm).
- 721 Groeneveld, L.F., Lenstra, J.A., Eding, H., Toro, M.A., Scherf, B., Pilling, D.,

- 722 Negrini, R., Finlay, E.K., Jianlin, H., Groeneveld, E., Weigend, S., 2010.
 723 GLOBALDIV Consortium: Genetic diversity in farm animals - a review. *Anim.*
 724 *Genet.* 41, 6–31.
- 725 Haenlein, G.F.W., 2004. Goat milk in human nutrition. *Small Rumin. Res.* 51:155–
 726 163.
- 727 Iamartino, D., Bruzzone, A., Lanza, A., Blasi, M., Pilla, F., 2005. Genetic diversity of
 728 Southern Italian goat populations assessed by microsatellite markers. *Small*
 729 *Rumin. Res.* 57: 249–255.
- 730 IOFS, 2002. Milk fat, preparation of fatty acid methyl ester. ISO Standard 15884.
- 731 Jansà-Pérez, M., Leroux, C., Sánchez Bonastre, A., Martin, P., 1994. Occurrence of a
 732 LINE sequence in the 3' UTR of the goat *cs1*-casein E-encoding allele associated
 733 with reduced protein synthesis level. *Gene* 147:179–187.
- 734 Kalantzopoulos, G., Dubeuf, J.-P., Vallerand, F., Pirisi, A., Casalta, E., Lauret, A.,
 735 Trujillo, T., 2004. Characteristics of sheep and goat milks: quality and hygienic
 736 factors for the sheep and goat dairy sectors. *Int. Dairy Fed. Bul.* 390, 17–28.
- 737 Kumar, S., Tamura, K., Nei, M., 2004. MEGA3: integrated software for molecular
 738 evolutionary genetics analysis and sequence alignment. *Brief. Bioinform.* 5, 150–
 739 163.
- 740 Küpper, J., Chessa, S., Rignanese, D., Caroli, A., Erhardt, G., 2010. Divergence at the
 741 casein haplotypes in dairy and meat goat breeds. *J. Dairy Res.* 77, 56–62.
- 742 Lauvie, A., Audiot, A., Couix, N., Casabianca, F., Brives, H., Verrier, E., 2011.
 743 Diversity of rare breed management programs: Between conservation and
 744 development. *Livestock Sci.* 140, 161–171.
- 745 Le Du, J., Benmederbel, B., 1984. Aptitude des chèvres de race Saanen à la traite
 746 mécanique. Relations avec les caractéristiques physiques du trayon. *Ann.*

- 747 Zootech. 33, 375–384.
- 748 Leitner, G., Silanikove, N., Merin, U., 2008. Estimate of milk and curd yield loss of
 749 sheep and goats with intramammary infection and its relation to somatic cell
 750 count. *Small Rumin. Res.* 74, 221–225.
- 751 Lucifero, M., 1981. *Allevamento moderno della capra*. first ed. Edagricole, Bologna,
 752 Italy.
- 753 Luikart, G., Gielly, L., Excoffier, L., Vigne, J.D., Bouvet, J., Taberlet, P., 2001.
 754 Multiple maternal origins and weak phylogeographic structure in domestic goats.
 755 *Proc. Natl. Acad. Sci. USA*, 98, 5927–5932.
- 756 Martini, M., Salari, F., Altomonte, I., Rignanese, D., Chessa, S., Gigliotti, C., Caroli,
 757 A., 2010. The Garfagnina goat: a zootechnical overview of a local dairy
 758 population. *J. Dairy Sci.* 93, 4659–4667.
- 759 Massart-Leën, A.M., De Pooter, H., Decloedt, M., Schamp, N., 1981. Composition
 760 and variability of the branched chain fatty acid fraction in the milk of goats and
 761 cows. *Lipids* 16, 286–292.
- 762 Mele, M., Buccioni, A., Serra, A., Antongiovanni, M., Secchiari, P., 2008. Lipids of
 763 goat's milk: origin, composition and main sources of variation, in: Cannas, A.,
 764 Pulina, G., (Eds), *Dairy goats feeding and nutrition*. CAB International,
 765 Wallingford, pp. 47–70.
- 766 Morand-Fehr, P., Boutonnet, J.P., Devendra, C., Dubeuf, J.P., Haenlein, G.F.W.,
 767 Holst, P., Mowlem, L., Capote, J., 2004. Strategy for goat farming in the 21st
 768 century. *Small Rumin. Res.* 51, 175–183.
- 769 Naderi, S., Rezaei, H.R., Taberlet, P., Zundel, S., Rafat, S.A., Naghash, H.R.,
 770 Elbarody, M.A.A., Ertugrul, O., Pompanon, F., 2007. Large-scale mitochondrial
 771 DNA analysis of the domestic goat reveals six haplogroups with high diversity.

- 772 PLoS ONE, 2, e1012. doi:10.1371/journal.pone.0001012.
- 773 Negrini, R., D'Andrea, M., Crepaldi, P., Colli, L., Nicoloso, L., Guastella, A.M.,
 774 Sechi, T., Bordonaro, S., Ajmone-Marsan, P., Pilla, F., the Econogene
 775 Consortium, 2012. Effect of microsatellite outliers on the genetic structure of
 776 eight Italian goat breeds. *Small Rumin. Res.* 103, 99–107.
- 777 Noè, L., Gaviraghi, A., D'Angelo, A., Bonanno, A., Di Trana, A., Sepe, L., Claps, S.,
 778 Annichiarico, G., Bacciu, N., 2005. Le razze caprine in Italia, in: Pulina, G. (Ed.),
 779 L'alimentazione della capra da latte. Avenue Media, Bologna, Italy, pp. 381–435.
- 780 Oku, H., Yanagita, T., 2009. Branched-chain fatty acids as a functional lipid, in: Hou,
 781 C.T., Shaw, J.-F., (Eds), *Biocatalysis and agricultural biotechnology*. CRC Press,
 782 Inc., Boca Raton, FL., pp. 259–267.
- 783 Paape, M.J., Wiggans, G.R., Bannerman, D.D., Thomas, D.L., Sandersb, A.H.,
 784 Contreras, A., Moroni, P., Miller, R.H., 2007. Monitoring goat and sheep milk
 785 somatic cell counts. *Small Rumin. Res.* 68, 114–125.
- 786 Park, Y.W., 2009. Conjugated linoleic acid (CLA): good or bad *trans* fat? *J. Food*
 787 *Compos. Anal.* 22S, S4–S12.
- 788 Park, Y.W., Juárez, M., Ramos, M., Haenlein, G.F.W., 2007. Physico-chemical
 789 characteristics of goat and sheep milk. *Small Rumin. Res.* 68, 88–113.
- 790 Parodi, P.W., 2009. Milk fat nutrition, in: Tamime, A.Y., (Ed), *Dairy fats and related*
 791 *products*. Blackwell Publishing Ltd., Oxford, pp. 28–51.
- 792 Pereira, F., Pereira, L., Van Asch, B., Bradley, D.G., Amorim, A., 2005. The mtDNA
 793 catalogue of all Portuguese autochthonous goat (*Capra hircus*) breeds: high
 794 diversity of female lineages at the western fringe of European distribution. *Mol.*
 795 *Ecol.* 14, 2313–2318.
- 796 Peris, S., Caja, S., Such, X., 1999. Relationships between udder and milking traits in

- 797 Murciano-Granadina dairy goats. *Small Rumin. Res.* 33, 171–179.
- 798 Piras, D., Doro, M.G., Casu, G., Melis, P.M., Vaccargiu, S., Piras, I., Parracciani, D.,
 799 Stradoni, R., Frongia, B., Lai, G., Sale, S., Cattari, W., Piras, R., Querci, O.,
 800 Demuro, P., Cui, S., Atzori, F., Mancosu, M., Marchiori, F., Cammelli, R., Spiga,
 801 A., Loddo, P.P., Pili, G., Boi, R., Argiolas, G., Mereu, P., Leoni, G.G., Naitana,
 802 S., Pirastu, M., Novelletto, A., 2012. Haplotype Affinities Resolve a Major
 803 Component of Goat (*Capra hircus*) MtDNA D-Loop Diversity and Reveal
 804 Specific Features of the Sardinian Stock. *PLoS One*. 2012;7(2):e30785.
- 805 Pirisi, A., Comunian, R., Urgeghe, P.P., Scintu, M.F., 2011. Sheep's and goat's dairy
 806 products in Italy: Technological, chemical, microbiological, and sensory aspects.
 807 *Small Rumin. Res.* 101, 102–112.
- 808 Precht, D., Molkentin, J., Destailats, F., Wolf, R.L., 2001. Comparative studies on
 809 individual isomer 18:1 acids in cow, goat and ewe milk fat by low-temperature
 810 high-resolution capillary gas-liquid chromatography. *Lipids* 36, 827–832.
- 811 Prinzenberg, E.M., Gutscher, K., Chessa, S., Caroli, A., Erhardt, G., 2005. Caprine
 812 κ -Casein (CSN3) Polymorphism: New Developments in molecular knowledge.
 813 *J. Dairy Sci.* 88, 1490–1498.
- 814 Ramunno, L., Cosenza, G., Pappalardo, M., Longobardi, E., Gallo, D., Pastore, N.,
 815 2001a. Characterization of two new alleles at the goat CSN1S2 locus. *Anim.*
 816 *Genetics* 32, 264–268.
- 817 Ramunno, L., Longobardi, E., Pappalardo, M., Rando, A., Di Gregorio, P., Cosenza,
 818 G., 2001b. An allele associated with a non-detectable amount of α_s2 casein in
 819 goat milk. *Anim. Genetics* 32, 19–26.
- 820 Raynal-Ljutovac, K., Lagriffoul, G., Paccard, P., Guillet, I., Chilliard, Y., 2008.
 821 Composition of goat and sheep milk products: an update. *Small Rumin. Res.* 79,

- 822 57–72.
- 823 Regione Piemonte, 2012. Modifications of the Rural Development Programme 2007-
 824 2013. Roundup report. Accessed Jul. 18, 2012.
 825 http://www.regione.piemonte.it/agri/psr2007_13/dwd/misure/2011/scheda_compl
 826 [essiva.pdf](http://www.regione.piemonte.it/agri/psr2007_13/dwd/misure/2011/scheda_compl).
- 827 Renna, M., Cornale, P., Lussiana, C., Malfatto, V., Fortina, R., Mimosi, A., Battaglini
 828 L.M., 2012. Use of *Pisum sativum* (L.) as alternative protein resource in diets for
 829 dairy sheep: effects on milk yield, gross composition and fatty acid profile. Small
 830 Rumin. Res 102, 142–150.
- 831 Reynolds, J.B., Weir, B.S., Cockerham, C.C., 1983. Estimation of the coancestry
 832 coefficient: basis for a short-term genetic distance. Genetics 105, 767–779.
- 833 Rosa García, R., Celaya, R., García, U., Osoro, K. 2012. Goat grazing, its interactions
 834 with other herbivores and biodiversity conservation issues. Small Rumin. Res.
 835 107, 49–64.
- 836 Royo, L.J., Traorè, A., Tambourà, H.H., Alvarez, I., Kaborè, A., Fernández, I.,
 837 Ouédraogo-Sanou, G., Toguyeni, A., Sawadogo, L., Goyache, F., 2009. Analysis
 838 of mitochondrial DNA diversity in Burkina Faso populations confirm the
 839 maternal genetic homogeneity of the West African goat. Anim. Genet. 40, 344–
 840 347.
- 841 Salama, A.A.K., Caja, G., Suck, X., Peris, S., Sorensen, A., Knight, C.H., 2004.
 842 Changes in cisternal udder compartment induced by milking interval in dairy
 843 goats milked once or twice daily. J. Dairy Sci. 87, 1181–1187.
- 844 Sardina, M.T., Ballester, M., Marmi, J., Finocchiaro, R., van Kaam, J.B.C.H.M.,
 845 Portolano, B., Folch, J.M., 2006. Phylogenetic analysis of sicilian goats reveals a
 846 new mtDNA lineage. Anim. Genet. 37, 376–378.

- 847 SAS Institute, 2008. SAS/STAT User's Guide. Release 9.2. SAS Institute Inc., Cary,
848 NC.
- 849 Sztankóová, Z., Mátlová, V., Kyselová, J., Jandurová, O.M., Riha, J., Senese, C.,
850 2009. Polymorphism of casein cluster genes in Czech local goat breeds—Short
851 communication. J. Dairy Sci. 92, 6197–6201.
- 852 Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997.
853 The ClustalX windows interface: flexible strategies for multiple sequence
854 alignment aided by quality analysis tools. Nucleic Acids Res. 24, 4876–4882.
- 855 Vacca, G.M., Daga, C., Pazzola, M., Carcangiu, V., Dettori, M.L., Cozzi, M.C., 2010.
856 D-loop sequence mitochondrial DNA variability of Sarda goat and other goat
857 breeds and populations reared in the Mediterranean area. J. Anim. Breed. Genet.
858 27, 352–360.
- 859 Verrier, E., Tixier-Boichard, M., Bernigaud, B., Naves M., 2005. Conservation and
860 value of local livestock breeds: usefulness of niche products and/or adaptation to
861 specific environments. Anim. Genet. Resour. Inform. 36, 21–31.
- 862 Wang, Y., Jacome-Sosa, M.M., Proctor, S.D., 2012. The role of ruminant *trans* fat as
863 a potential nutraceutical in the prevention of cardiovascular disease. Food Res.
864 Int. 46, 460–468.
- 865 Weir, B.S., 1996. Genetic Data Analysis II, Sunderland, MA: Sinauer Associates, Inc.
- 866 Weir, B.S., Cockerham, C., 1984. Estimating F-statistics for the analysis of
867 population structure. Evolution 38, 1358–1369.
- 868 Wright, S., 1965. The interpretation of population structure by F-statistics with special
869 regard to systems of mating. Evolution 19, 395–420.
- 870 Zaitoun, I.S., Tabbaa, M.J., Bdour, S., 2005. Differentiation of native goat breeds of
871 Jordan on the basis of morphostructural characteristics. Small Rumin. Res. 56,

872 173–182.

873 Žan, M., Stibilj, V., Rogelj, I., 2006. Milk fatty acid composition of goats grazing on

874 alpine pasture. Small Rumin. Res. 64, 45–52.

875

Accepted Manuscript

875 **Figure captions**

876 Figure 1. A small flock of *Fiurinà* goats.

877

878 Figure 2. Nucleotide substitutions detected in 10 *Fiurinà* goat mtDNA sequences
879 (481-bp fragment) compared to the reference sequence (GenBank NC_005044
880 abbreviated as REF). The positions are given with respect to the reference sequence.
881 Dots (.) denote identical sites.

882

883 Figure 3. Neighbour-joining tree of domestic goat based on 10 mtDNA sequences of
884 the *Fiurinà* breed and on 22 reference mtDNA haplotypes (Ref. Seq.). Distances were
885 calculated using Kimura 2-parameter with 1,000 bootstrap replications.

886

887 Figure 4. Principal Component Analysis of the haplotypes frequencies in *Fiurinà* and
888 other five Italian goat breeds.

Figure 2

	15728	
	15744	
	15747	
	15763	
	15771	
	15797	
	15807	
	15811	
	15822	
	15831	
	15843	
	15862	
	15864	
	15866	
	15870	
	15873	
	15883	
	15887	
	15893	
	15913	
	15920	
	15930	
	15947	
	15950	
	15965	
	15969	
	15973	
	15974	
	15975	
	15976	
	15977	
	15978	
	15981	
	15982	
	15983	
	15984	
	15992	
	15994	
	15999	
	16000	
	16003	
	16006	
	16010	
	16011	
	16019	
	16022	
	16026	
	16027	
	16037	
	16038	
	16040	
	16043	
	16045	
	16054	
	16062	
	16066	
	16103	
	16104	
	16151	
REF	AAGGGGAATTAAAAATATATAAAAAAAGTAATTTTAGAGTTTATAAAAAATAATTGTAAGA	
Seq1	G.A...TA..G....G..AG..TTTT...TA..G..A..A..ATTG..GA.T.A...T..	
Seq2	G.A...TA..G....AG.TTTTTT.ATTA.AGA.A....ATTG..GA.T.A...T..	
Seq3	G.A...TA..G....TAG.TT.TT.A.TA..G..A....ATTG..GA.T.A...TT..	
Seq4	G.A...TA..G....GA.AG.TTTTTT...TA..GA.A....ATTG..GA.T.A...T..	
Seq5	G.A...TA..G....GA.AG.TTTTTT...TA..GA.A....ATTG..GA.T.A...T..	
Seq6	.TAAAG..A..GTGA...AG.TT.T.A...A.AGAT.A.AT....GG.AGTA.AA.TAT	
Seq7	G.A...TA..G....G..AGG.TTTTT.A.TAA.G..A....ATTG..GA.T.A...T..	
Seq8	.TAAAG..A..GTGA...AG.TT.T.A...A.AGAT.A.AT....GG.AG.A.AA.TAT	
Seq9	G.A...TA..G....AG.TTTTTT...TA..G..A....ATTG.....T.....T..	
Seq10	G.A...TA.GG.....AGGTTTTT..ATTA.AGA.A....ATTG.GGA.TAA...T..

Figure 4

uscrip

Table 1. Analyses applied to screen most of the genetic variability of goat caseins at the DNA level. A slash in the allele list indicates that they are not discernable with the method applied. The analyses available to separate part of the alleles grouping together within the first screening method are in the fourth column.

Gene ^a	DNA method ^b	Discernable alleles	Further analysis and discernable alleles
<i>CSN1S1</i>	PCR-SSCP ^b (Küpper et al., 2010)	A/0 ₁ , A', B/E, B', F, N	AS ^b -PCR - B, E (Jansà-Pérez et al., 1994) AS-PCR - A, 0 ₁ (Cosenza et al., 2003)
<i>CSN2</i>	PCR-SSCP (Chessa et al., 2008b)	A, C, A ₁ , C ₁ , E, 0, 0'	
<i>CSN1S2</i>	PCR-SSCP (Chessa et al., 2008a)	A/D/0/F, B, C, E	PCR-RFLP - A, D, 0 (Ramunno et al. 2001b) PCR-RFLP - A, F (Ramunno et al. 2001a)
<i>CSN3</i>	PCR-SSCP (Prinzenberg et al., 2005)	A, B/B', B'', C, C', E, D/I/K/L, G, H, J, M	

^a *CSN1S1* = α_{S1} -CN; *CSN2* = β -CN; *CSN1S2* = α_{S2} -CN; *CSN3* = κ -CN.

^bMethods: SSCP = Single Strand Conformation Polymorphism; AS = Allele Specific.

- 1 Table 2. Average morphometric data^a of adult females in the *Fiurinà* breed and in
 2 other goat breeds reared in the same Italian region.

	<i>Fiurinà</i>	Camosciata delle Alpi ^b	Saanen ^b	Sempione ^b	Vallesana ^b
Height at withers (cm)	73±6.1	74	74	72	75
Rump height (cm)	75±4.8	77	77	74	78
Rump width (cm)	17±3.2	19	20	-	17
Trunk length (cm)	77±1.9	85	89	82	87
Chest girth (cm)	87±6.4	91	96	88	95
Chest height (cm)	34±7.8	34	35	-	-
Weight (kg)	50±12.9	70	60	60	70

- 3 ^aStandard deviations are reported only for *Fiurinà* breed since for the other considered breeds only mean values of morphometric
 4 data are available in the literature.

- 5 ^bNoè et al. (2005).

- 1 Table 3. Basic information, values, and significance for parameters^a of microsatellites
 2 polymorphism observed in the *Fiurinà*, Sempione, and Vallesana breeds.

Breed	Approximate breed size	Sample size	H _O	H _E	AR	F _{IS}
<i>Fiurinà</i>	150	26	0.59	0.63	6.19	0.05
Sempione	155	22	0.66	0.69	6.41	-0.01
Vallesana	100	36	0.69	0.70	5.80	0.12 ***

- 3 ^aH_O = observed heterozygosity; H_E = expected heterozygosity; AR = allelic richness; F_{IS} = heterozygote deficiency coefficient.

- 4 Probability: *** $P < 0.001$.

- 1 Table 4. F-statistics F_{IS} , F_{ST} , F_{IT} and their P -values calculated for each microsatellite
- 2 in the *Fiurinà*, Sempione, and Vallesana breeds.

Locus	F_{IS}	F_{ST}	F_{IT}
INRA005	-0.008	-0.003	-0.011
MAF65	0.002	0.016	0.018
INRA063	-0.006	0.002	-0.004
MCM527	0.187 ***	0.029	0.211 ***
ET10	-0.086	0.026	-0.057
SRCRSP5	-0.003	0.009	0.006
INRA023	-0.017	-0.001	-0.018
FCB20	0.140	0.025	0.162 ***
TGLA53	-0.021	0.021	-0.001
SRC247	-0.048	0.042 ***	0.088 *
CSRD247	0.320 ***	0.026	0.338 ***
ILST87	0.258 ***	0.119 ***	0.347 ***
Average	0.076 ***	0.029 ***	0.103 ***

- 3 Probability: * $P < 0.05$; *** $P < 0.001$.

- 1 Table 5. Reynolds's distance (above the diagonal) and pairwise comparisons using
 2 F_{ST} estimates (below the diagonal) among the *Fiurinà*, Sempione, and Vallesana
 3 breeds based on the microsatellites loci.

	<i>Fiurinà</i>	Sempione	Vallesana
<i>Fiurinà</i>	-	0.062	0.055
Sempione	0.041***	-	0.031
Vallesana	0.038***	0.013*	-

- 4 Probability: * $P < 0.05$; *** $P < 0.001$.

- 1 Table 6. Main constituents and somatic cell count in milk from *Fiurina* goats.

	Mean	SD
Fat (g kg ⁻¹)	35.8	7.4
Solids-non-fat (g kg ⁻¹)	80.5	3.7
Lactose (g kg ⁻¹)	44.1	2.4
Protein (g kg ⁻¹)	29.9	3.0
Casein (g kg ⁻¹)	24.5	2.3
Urea (mg dL ⁻¹)	34.66	7.65
Somatic cell count ^a ($\times 10^3$ cells mL ⁻¹)	490.00	(217.00 – 917.25)

- 2 ^a Results are expressed as median and interquartile range.

1 Table 7. Individual fatty acids (g 100g⁻¹ fat and % of total FAME^a) in milk from
 2 *Fiurina* goats.

	g 100g ⁻¹ fat		% total FAME	
	mean	SD	mean	SD
C4	2.29	0.32	2.91	0.41
C5	<0.01	<0.01	<0.01	<0.01
C6	2.23	0.27	2.85	0.46
C7	0.02	0.01	0.02	0.01
C8	2.43	0.40	3.12	0.62
C10	7.35	1.34	9.39	1.84
C10:1	0.20	0.06	0.26	0.08
C12	2.88	0.66	3.68	0.87
C13 <i>iso</i>	0.02	0.01	0.02	0.01
C13 <i>aiso</i>	0.02	0.01	0.03	0.02
C12:1 <i>c</i> + C13	0.10	0.03	0.13	0.04
C14 <i>iso</i>	0.07	0.04	0.09	0.05
C14	7.31	1.24	9.30	1.49
C15 <i>iso</i>	0.17	0.06	0.21	0.08
C14:1 <i>t</i>	<0.01	<0.01	<0.01	<0.01
C15 <i>aiso</i>	0.30	0.10	0.38	0.11
C14:1 <i>c</i>	0.07	0.03	0.09	0.04
C15	0.80	0.17	1.02	0.20
C16 <i>iso</i>	0.18	0.07	0.22	0.08
C16	19.02	3.10	24.10	3.14
C17 <i>iso</i>	0.37	0.14	0.47	0.16
C16:1 <i>t</i>	0.11	0.05	0.13	0.05
C17 <i>aiso</i>	0.61	0.13	0.77	0.13
C16:1 <i>c</i>	0.35	0.09	0.44	0.10
C17	0.71	0.15	0.89	0.15
C18 <i>iso</i>	<0.01	<0.01	<0.01	<0.01
C17:1 <i>t</i>	0.05	0.03	0.06	0.03
C18 <i>aiso</i>	0.27	0.11	0.34	0.14
C18	9.12	2.33	11.54	2.63
C18:1 <i>t5</i>	<0.01	<0.01	<0.01	<0.01
C18:1 <i>t6</i> - 11	1.98	0.71	2.51	0.88
C18:1 <i>t12</i> - 14 + <i>c6</i> - 8	0.71	0.28	0.89	0.34
C18:1 <i>c9</i>	13.94	2.79	17.65	3.03
C18:1 <i>c11</i>	0.37	0.09	0.47	0.10
C18:1 <i>c12</i>	0.13	0.04	0.16	0.05
C18:1 <i>c14</i> + <i>t16</i>	0.38	0.12	0.49	0.14
C19	0.11	0.03	0.13	0.04
C18:2 <i>t,t</i> - NMID + <i>t9t12</i>	0.11	0.04	0.14	0.05
C18:2 <i>c9t13</i> + <i>t8c12</i>	0.05	0.03	0.06	0.03
C18:2 <i>c9t12</i>	0.22	0.09	0.28	0.11
C18:2 <i>c,c</i> - MID + <i>t8c13</i>	0.22	0.08	0.27	0.09
C18:2 <i>t11c15</i> + <i>t9c12</i>	0.24	0.09	0.30	0.11
C18:2 <i>c9c12</i> (LA)	1.17	0.32	1.47	0.35
C18:2 <i>c9c15</i>	0.02	0.02	0.02	0.02
C20	0.26	0.13	0.33	0.15
C20:1 <i>t</i>	0.01	0.01	0.01	0.01
C18:3 <i>c6c9c12</i> (GLA)	0.01	0.01	0.01	0.01
C20:1 <i>c5</i>	0.01	0.01	0.02	0.01

C20:1 <i>c</i> 9	0.03	0.02	0.04	0.03
C20:1 <i>c</i> 11	0.03	0.02	0.03	0.02
C18:3 <i>c</i> 9 <i>c</i> 12 <i>c</i> 15 (ALA)	0.82	0.26	1.02	0.28
CLA <i>c</i> 9 <i>t</i> 11 + <i>t</i> 7 <i>c</i> 9 + <i>t</i> 8 <i>c</i> 10	0.61	0.25	0.77	0.31
CLA <i>t</i> 11 <i>c</i> 13 + <i>c</i> 9 <i>c</i> 11	0.02	0.02	0.02	0.02
CLA <i>t</i> 9 <i>t</i> 11	0.01	0.01	0.01	0.01
C20:2 <i>c,c</i> n6	0.01	0.01	0.01	0.01
C22	0.04	0.03	0.05	0.03
C20:3 n6	0.01	0.01	0.01	0.01
C20:3 n3	0.01	0.01	0.01	0.01
C20:4 n6 (AA)	0.08	0.04	0.09	0.05
C20:5 n3 (EPA)	0.06	0.03	0.07	0.03
C22:5 n3 (DPA)	0.09	0.05	0.11	0.06
C22:6 n3 (DHA)	0.03	0.03	0.04	0.04

^aFAME = fatty acid methyl esters; SD = standard deviation; *c* = *cis*; *t* = *trans*; NMID = non methylene interrupted diene; MID = methylene interrupted diene; LA = linoleic acid; GLA = γ -linoleic acid; ALA = α -linoleic acid; CLA = conjugated linoleic acid; AA = arachidonic acid; EPA = eicosapentaenoic acid; DPA = docosapentaenoic acid; DHA = docosahexaenoic acid.

Table 8. Groups of fatty acids (g 100g⁻¹ fat and % of total FAME^a) in milk from *Fiurina* goat.

	g 100g ⁻¹ fat		% total FAME	
	mean	SD	mean	SD
Σ short chain ^b	14.52	1.94	18.55	2.93
Σ medium chain ^c	32.39	4.44	41.10	4.53
Σ long chain ^d	31.85	5.82	40.30	5.85
Σ saturated ^e	56.59	4.88	71.88	3.88
Σ branched chain ^f	2.01	0.50	2.54	0.56
Σ monounsaturated ^g	18.47	3.22	23.39	3.34
Σ C18:1 ^h	17.51	3.16	22.18	3.31
Σ C18:1 <i>trans</i> ⁱ	2.70	0.88	3.42	1.06
Σ polyunsaturated ^j	3.75	0.95	4.72	0.98
Σ C18:2 ^k	2.65	0.66	3.35	0.70
Σ C18:2 <i>trans</i> ^l	1.47	0.47	1.86	0.55
Σ <i>trans</i> without CLA ^m	6.77	2.06	8.58	2.45
Σ n3 FA ⁿ	1.25	0.39	1.57	0.42
Σ n6 FA ^o	2.47	0.56	3.08	0.59
Σ n6 / Σ n3	2.08	0.61	2.68	0.91
Σ CLA ^p	0.64	0.27	0.81	0.33
Σ unsaturated ^q	22.21	3.90	28.12	3.88

^a FAME = fatty acid methyl esters; SD = standard deviation; CLA = conjugated linoleic acid; FA = fatty acids.

^b C4, C5, C6, C7, C8, C10, C10:1.

^c C12, C13 *iso*, C13 *aiso*, C12:1 *c* + C13, C14 *iso*, C14, C15 *iso*, C14:1 *t*, C15 *aiso*, C14:1 *c*, C15, C16 *iso*, C16, C17 *iso*, C16:1 *t*, C17 *aiso*, C16:1 *c*.

^d C17, C18 *iso*, C17:1 *t*, C18 *aiso*, C18, Σ C18:1, C19, Σ C18:2, C20, C20:1 *t*, C18:3 *c6c9c12*, C20:1 *c5*, C20:1 *c9*, C20:1 *c11*, C18:3 *c9c12c15*, C18:2 *c9t11* + *t7c9* + *t8c10*, C18:2 *t11c13* + *c9c11*, C18:2 *t9t11*, C20:2 *c,c* n6, C22, C20:3 n6, C20:3 n3, C20:4 n6, C20:5 n3, C22:5 n3, C22:6 n3.

^e C4, C5, C6, C7, C8, C10, C12, Σ branched chain, C14, C15, C16, C17, C18, C19, C20, C22.

^f C13 *iso* + *aiso*, C14 *iso*, C15 *iso* + *aiso*, C16 *iso*, C17 *iso* + *aiso*, C18 *iso* + *aiso*.

^g C10:1, C12:1 *c* + C13, C14:1 *ct*, C16:1 *ct*, C17:1 *t*, Σ C18:1, C20:1 *t*, C20:1 *c5*, C20:1 *c9*, C20:1 *c11*.

^h C18:1 *t5*, *t6* - 11, *t12* - 14 + *c6* - 8, *c9*, *c11*, *c12*, *c14* + *t16*.

ⁱ C18:1 *t5*, *t6* - 11, *t12* - 14 + *c6* - 8.

^j Σ C18:2, C18:3 *c6c9c12*, C18:3 *c9c12c15*, C20:2 *c,c* n6, C20:3 n3, C20:3 n6, C20:4 n6, C20:5 n3, C22:5 n3, C22:6 n3.

^k C18:2 *t,t* - NMID + *t9t12*, *c9t13* + *t8c12*, *c9t12*, *c,c* - MID + *t8c13*, *t11c15* + *t9c12*, *c9c12*, *c9c15*, *c9t11* + *t7c9* + *t8c10*, *t11c13* + *c9c11*, *t9t11*.

^l C18:2 *t,t* - NMID + *t9t12*, *c9t13* + *t8c12*, *c9t12*, *c,c* - MID + *t8c13*, *t11c15* + *t9c12*, C18:2 *c9t11* + *t7c9* + *t8c10*, C18:2 *t11c13* + *c9c11*, C18:2 *t9t11*.

^m C14:1 *t*, C16:1 *t*, C17:1 *t*, Σ C18:1 *t*, Σ C18:2 *t* (without CLA *trans*), C20:1 *t*.

ⁿ C18:2 *t11c15* + *t9c12*, C18:2 *c9c15*, C18:3 *c9c12c15*, C20:3 n3, C20:5 n3, C22:5 n3, C22:6 n3.

^o C18:1 *t12*, C18:1 *c12*, C18:2 *t,t* - NMID + *t9t12*, C18:2 *c9t12*, C18:2 *t11c15* + *t9c12*, C18:2 *c9c12*, C18:3 *c6c9c12*, C20:2 *c,c* n6, C20:3 n6, C20:4 n6.

^p C18:2 *c9t11* + *t7c9* + *t8c10*, *t11c13* + *c9c11*, *t9t11*.

^q C10:1, C12:1 *c* + C13, C14:1 *ct*, C16:1 *ct*, C17:1 *t*, Σ C18:1, Σ C18:2, C20:1 *t*, C18:3 *c6c9c12*, C20:1 *c5*, C20:1 *c9*, C20:1 *c11*, C18:3 *c9c12c15*, C18:2 *c9t11* + *t7c9* + *t8c10*, C18:2 *t11c13* + *c9c11*, C18:2 *t9t11*, C20:2 *c,c* n6, C20:3 n6, C20:3 n3, C20:4 n6, C20:5 n3, C22:5 n3, C22:6 n3.

Table 9. Casein allele and haplotype frequencies in the *Fiurina* breed. Haplotype frequencies were calculated both under hypothesis of loci independence (H0) and taking association into account (H1) by the HAPLOTYPE procedure of SAS (SAS Institute Inc., 2008). Only haplotypes with H1 higher than 0.008 are shown.

Casein gene ^a	Allele	Frequency	<i>CSN1S1</i>	<i>CSN2</i>	<i>CSN1S2</i>	<i>CSN3</i>	H0	H1
<i>CSN1S1</i>	<i>A</i>	0.052	<i>E</i>	<i>A</i>	<i>A</i>	<i>B</i>	0.033	0.310
	<i>A'</i>	0.010	<i>O_I</i>	<i>C</i>	<i>B</i>	<i>A</i>	0.003	0.172
	<i>B</i>	0.063	<i>F</i>	<i>C_I</i>	<i>F</i>	<i>A</i>	0.017	0.171
	<i>B'</i>	0.010	<i>F</i>	<i>C_I</i>	<i>F</i>	<i>B</i>	0.022	0.061
	<i>E</i>	0.333	<i>F</i>	<i>C_I</i>	<i>F</i>	<i>C</i>	0.002	0.051
	<i>F</i>	0.333	<i>E</i>	<i>A</i>	<i>F</i>	<i>B</i>	0.028	0.034
	<i>O_I</i>	0.198	<i>B</i>	<i>A</i>	<i>A</i>	<i>B</i>	0.006	0.029
<i>CSN2</i>	<i>A</i>	0.406	<i>F</i>	<i>C_I</i>	<i>A</i>	<i>B</i>	0.027	0.023
	<i>C</i>	0.219	<i>A</i>	<i>C</i>	<i>A</i>	<i>B</i>	0.003	0.023
	<i>C_I</i>	0.365	<i>O_I</i>	<i>C_I</i>	<i>B</i>	<i>A</i>	0.005	0.022
	<i>O'</i>	0.010	<i>A</i>	<i>C</i>	<i>A</i>	<i>A</i>	0.002	0.020
<i>CSN1S2</i>	<i>A</i>	0.388	<i>B</i>	<i>A</i>	<i>C</i>	<i>B</i>	0.001	0.020
	<i>B</i>	0.204	<i>B</i>	<i>A</i>	<i>A</i>	<i>A</i>	0.004	0.013
	<i>C</i>	0.041	<i>F</i>	<i>C_I</i>	<i>F</i>	<i>C'</i>	0.001	0.012
	<i>F</i>	0.367	<i>E</i>	<i>A</i>	<i>C</i>	<i>B</i>	0.003	0.012
<i>CSN3</i>	<i>A</i>	0.388	<i>A</i>	<i>C</i>	<i>F</i>	<i>B</i>	0.002	0.010
	<i>B</i>	0.541	<i>F</i>	<i>C_I</i>	<i>C</i>	<i>B</i>	0.003	0.009
	<i>C</i>	0.051	<i>A</i>	<i>C</i>	<i>F</i>	<i>C'</i>	0.000	0.008
	<i>C'</i>	0.020						

^a *CSN1S1* = α_{S1} -CN; *CSN2* = β -CN; *CSN1S2* = α_{S2} -CN; *CSN3* = κ -CN.