

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Functional properties of ion channels and transporters in tumour vascularization

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/143024> since 2017-05-12T18:30:45Z

Published version:

DOI:10.1098/rstb.2013.0103

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Philos Trans R Soc Lond B Biol Sci. 2014 Feb 3;369(1638):20130103. doi:
10.1098/rstb.2013.0103. Print 2014 Mar 19. Review.

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://rstb.royalsocietypublishing.org/content/369/1638/20130103.long>



FUNCTIONAL PROPERTIES OF ION CHANNELS AND TRANSPORTERS IN TUMOR VASCULARIZATION

Journal:	<i>Philosophical Transactions B</i>
Manuscript ID:	RSTB-2013-0103.R1
Article Type:	Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Fiorio Pla, Alessandra; University of Torino, Life Sciences and Systems Biology Munaron, Luca; University of Torino, Life Sciences and Systems Biology
Issue Code: Click here to find the code for your issue.:	ION
Subject:	Physiology < BIOLOGY, Biophysics < BIOLOGY, Cellular Biology < BIOLOGY
Keywords:	endothelial cells, channels, transporters

SCHOLARONE™
Manuscripts



FUNCTIONAL PROPERTIES OF ION CHANNELS AND TRANSPORTERS IN TUMOR VASCULARIZATION

Alessandra Fiorio Pla^{1*} & Luca Munaron¹

¹Department of Life Sciences & Systems Biology, Center for Complex Systems in Molecular Biology and Medicine (SysBioM), Nanostructured Interfaces and Surfaces Centre of Excellence (NIS), University of Torino

*Correspondence to:

Alessandra Fiorio Pla, Ph.D.
Dept. Life Sciences & Systems Biology
University of Torino
Via Accademia Albertina 13
10123 Torino
ITALY
alessandra.fiorio@unito.it

Running title: channels/transporters in tumor angiogenesis

Keywords: tumor vascularization, endothelial cells, VOCS, TRP channels, aquaporins, transporters, nicotinic receptors.

ABSTRACT

Vascularization is crucial for solid tumor growth and invasion, providing metabolic support and sustaining metastatic dissemination.

It is now accepted that ion channels and transporters play a significant role in driving the cancer growth at all stages. They may represent novel therapeutic, diagnostic, and prognostic targets for anti-cancer therapies. On the other hand, although the expression and role of ion channels and transporters in the vascular endothelium is well recognized and subject of recent reviews, only recently their involvement in tumor vascularization have been recognized.

Here we review the current literature on ion channels and transporters directly involved in angiogenic process. Particular interest will be focused on tumor angiogenesis *in vivo*, as well as in the different steps that drive this process *in vitro*, such as endothelial cell proliferation, migration, adhesion and tubulogenesis.

Moreover, we compare the 'transportome' system of tumor vascular network with the physiological one.

INTRODUCTION

Endothelium is a multifaceted and dynamic interface between blood components and tissues. Endothelial cells (EC) mediate a great number of physiological functions, including control of metabolism, water supply, inflammation and immune response. Consequently, several diseases are causally due to the deregulation of normal EC functions. The importance of vascularization in the tumor progression sparked hopes that manipulating this process could offer therapeutic opportunities [1,2]. Consequently, so far hundreds of thousands of patients benefit of antiangiogenic therapies that use VEGF as major drug target and approved by the US Food and Drug Administration. The anti-VEGF antibody (bevacizumab [Avastin]) is used in combination with chemotherapy, cytokine therapy or radiotherapy for several advanced metastatic cancers. Additionally, four multitargeted pan-VEGF receptor tyrosine kinase inhibitors have been approved: Sunitinib (Sutent), Pazopanib (Votrient) for metastatic Renal Cancer Carcinoma (RCC), Sorafenib (Nexavar) for metastatic RCC, unresectable hepatocellular carcinoma and advanced pancreatic neuroendocrine tumors, and Vandetanib (Zactima) for medullary thyroid cancer [3]. On the other

1
2
3 hand, despite promising results, emerging data indicate that responses to vascular targeting therapy
4 (VTT) are short-lived and resistance develops in the majority of patients. The discovery of new
5 therapeutic targets is therefore necessary to provide a new input to the antiangiogenic therapy.

6 Being involved in nearly all of the 'hallmarks of cancer' as defined by Hanahan and Weinberg [4],
7 there is an increasing consensus on the idea that ion channels and transporters could play a
8 significant role in driving cancer progression at all stages. Therefore they may be seen as potential
9 novel therapeutic, diagnostic, and prognostic targets for anti-cancer therapies.

10 Nonetheless, although the expression and role of ionic channels and transporters (collectively
11 indicated as "transportome") in the vascular endothelium is well recognized and subject of a number
12 of recent reviews [5–8], 'transportome' entered only recently as major players in tumor
13 vascularization [9,10].

14 Here we collect and discuss current literature focused on ion channels and transporters directly
15 involved in angiogenic process. Moreover, starting from a critical review of the experimental data
16 obtained so far *in vitro* and *in vivo*, we will try to define the most promising checkpoints at which
17 tumor vascular 'transportome' differs from the physiological one.
18
19

20 **VOLTAGE-GATED CHANNELS**

21 Although EC are generally described as non excitable cells, a number of experimental evidences
22 suggest a role for voltage dependent channels (VOCs) in both cultured and freshly isolated EC [10].
23 On the other hand, the role of VOCs in tumor progression has been largely described and different
24 data point to Na⁺, K⁺ and Ca²⁺ channels as key players, suitable to be specifically potential target
25 in clinical treatments [11].

26 K⁺ channels (K_v) attracted most of the work in oncology since the early discovery unveiling their
27 role in the control of cell proliferation [12,13]. Ether-á-go-go-1 (EAG1, KCNH1, K_v10.1) is a
28 CNS-localized voltage-gated K⁺ channel that is found ectopically expressed in many solid tumors.
29 Most of the interest in K_v10.1 arises from its expression in up to 70% of tumor cell lines and human
30 cancers [13]. Monoclonal antibodies against human EAG1, developed by Stuhmer's and Pardo's
31 groups, might represent a suitable tool in cancer therapy [14]. K_v10.1 expression might offer an
32 advantage to tumors through increased vascularization and resistance to hypoxia: indeed, EAG1
33 regulates cellular oxygen homeostasis, increasing HIF-1 activity, and thereby VEGF secretion and
34 tumor vascularization [15]; accordingly EAG1 silencing inhibits tumor growth and angiogenesis in
35 osteosarcoma *in vivo* [16] (Table 1 and Fig.1).

36 A promising issue is related to other K⁺ channels, such as human ether-a-gogo related gene-1
37 (hERG1)-K_v11 [13,17,18]; Pillozzi and coworkers showed that hERG1 channels regulate vegf-a
38 expression and VEGF-A secretion in cancer cells potentially promoting angiogenesis [19].
39 Moreover, it has been discovered a correlation between the levels of VEGF-A, hERG1 and
40 microvessel density and proliferation-related parameters in two cases of bilateral retinoblastoma
41 patients [20]. Beside the role of K_v10 and K_v11, K_v 1.3 channels are involved in VEGF-mediated
42 HUVEC proliferation: VEGF-mediated hyperpolarization via Margatoxin (MTX)-sensitive K_v
43 channels causes a Ca²⁺ entry, leading to an increase in NO synthesis, finally resulting in EC growth
44 enhancement [21] (Table 1 and Fig.1). All together, the data point out an important role for K⁺
45 channels in the cross talk between cancer cells and tumor endothelium by induction of VEGF
46 release that in turn promotes neovascularization. This particular function of K⁺ channels makes
47 them clinically interesting as potential targets to promote vascular "normalization" by interfering
48 with VEGF signaling during a critical window of the antiangiogenic treatments (see also the
49 conclusion section).
50

51 Voltage-gated Na⁺ channels (VGSCs) are also expressed in non excitable cells and functionally up-
52 regulated in metastatic tumor cells [22–24]. Recently, a clear relationship between functional
53 expression and biological role of VGSCs in EC has been described [25]. Molecular expression
54 analyses and electrophysiology revealed consistently that the main functional VGSC isoforms in
55 HUVEC are Nav1.5 and Nav1.7. VGSC activity potentiates VEGF-induced ERK1/2 activation by
56
57
58
59
60

1
2
3 attenuating membrane depolarization, altering $[Ca^{2+}]_i$ kinetics and PKC activity with a consequent
4 increase in cellular proliferation, chemotaxis, and tubulogenesis [25] (Table 1 and Fig.1). Although
5 the question on the specificity of VGSC on VEGF signaling pathway remains to be elucidated, the
6 data unveil an intriguing mechanism for the control of V_m in non-excitabile cells by VGSCs in
7 response to physiological stimuli *in vitro*.

8 As regarding voltage-gated Ca^{2+} channels (VGCCs), most of the studies have been conducted on
9 human breast carcinoma cell lines, which actually express VGCCs, mainly of the T-type [26–28].
10 Nevertheless, the role and expression of VGCCs in endothelium is still debated [3,7,24].
11 Conflicting data could arise from the use of different cultured EC lines and their well known
12 variable behaviour (see also the conclusion section for a more detailed discussion). In human
13 umbilical vein EC (HUVEC) Angiotensin II stimulates Ca^{2+} influx via Ca_v and promotes cell
14 migration [30]. On the other hand, Ca_v expressed by VSMCs could play an anti-angiogenic role
15 through indirect effects on EC: nifedipine, a widely used inhibitor of L-type calcium channels,
16 stimulates VEGF production from human coronary smooth muscle cells, an effect abolished by
17 PKC inhibitors and a bradykinin B2 receptor antagonist [31] (Table 1 and Fig.1).

20 **TRANSIENT RECEPTOR POTENTIAL (TRP) PROTEINS AND STIM1-ORAI1** 21 **COMPLEX.**

22 Transient Receptor Potential (TRP) channels trigger Ca^{2+} signals that control intracellular events
23 involved in the initiation and progression of cancer. It is not therefore surprising that the expression
24 and function of some TRP proteins are altered during tumor growth and metastasis [9,32].

25 TRPs are widely expressed in endothelium and their activity has been related to normal and tumor
26 vascularization [5,6]. TRP -mediated Ca^{2+} influx can be triggered by the release from intracellular
27 Ca^{2+} stores giving rise to store-operated Ca^{2+} entry (SOCE). An alternative route is the store-
28 independent Ca^{2+} entry (NSOCE) [33].

29 VEGF mediates NSOCE through TRPC6 channels in human microvascular EC [34,35]. Dominant
30 negative TRPC6 significantly reduces EC number, migration and sprouting [36]. Moreover, TRPC6
31 promotes both proliferation and tubulogenesis induced by VEGF, but not bFGF, in HUVEC [37].
32 Phosphatase and tensin homolog (PTEN) regulates cell surface expression of TRPC6, and
33 consequently Ca^{2+} entry, endothelial permeability, and angiogenesis in human pulmonary EC [38].
34 Nonetheless, TRPC6 can also exert its proangiogenic role indirectly through its activity on cancer
35 cells being a key mediator hypoxia-mediated notch-driven growth and invasiveness of glioblastoma
36 multiforme (GBM): inhibition of the hypoxia upregulated TRPC6 expression and NFAT activation
37 in glioma cells, markedly reduced the number of branch points in EC grown in conditioned medium
38 harvested from glioma cells, indicating that TRPC6 is essential for the angiogenic potential of
39 glioma cells [39] (Table 1 and Fig.1).

40 Other groups reported a role of VEGF-mediated SOCE due to TRPC1 in the enhancement of
41 HMVEC and HUVEC permeability [40–42]. Remarkably, TRPC1 is proangiogenic *in vivo*.
42 Knockdown of zebrafish TRPC1 by morpholinos caused severe angiogenic defects in intersegmental
43 vessel sprouting, presumably due to impaired filopodia extension during EC migration [43] (Table
44 1 and Fig. 1).

45 This apparently surprising ability of VEGF to couple to different channels responsible for SOCE or
46 NSOCE could simply depend on tissue variability, especially between small capillaries and large
47 vessels. Accordingly, the pattern of TRPC channels expressed in HMVEC and HUVEC is different,
48 TRPC4 being undetectable in HMVEC [36].

49 Besides TRPC1 and TRPC6, also Orai1 and STIM1, components of the so-called calcium release
50 activated currents (CRAC) channels, concur to the VEGF-mediated SOCE in HUVEC [44,45].
51 VEGF stimulation promotes STIM1 clustering which in turn activates Orai1 [45]. Moreover,
52 knock-down of Orai1 inhibits VEGF-mediated HUVEC migration, proliferation and tubulogenesis
53 [44–46]. On the other hand, Trebak and coworkers reported recently that the thrombin-induced
54 decrease in EC permeability requires STIM1, but is unrelated to Orai1 and Ca^{2+} entry across the
55
56
57
58
59
60

1
2
3 plasma membrane [47] (Table 1).

4 Interestingly, STIM1, as well as TRPC1 and TRPC4 knockdown, inhibits tube formation in both
5 HUVEC and EA.hy926 cells, an EC line derived from HUVEC fused with human lung
6 adenocarcinoma cell line A549 [48]. Since Orai1 and TRPC1 can functionally interact at least in
7 some models, the TRP- and Stim/Orai- pathways may give rise to a complex signaling network
8 underlying proangiogenic calcium signals [49].

9 Since VEGF regulates several activities in EC, the discovery of a specific role for the different
10 channels in selected cell functions, such as migration and proliferation on one side or permeability
11 on the other, could be a more useful molecular target than the broad VEGFR inhibitors (see also
12 Conclusion section).

13 TRPV4 is another emerging player in angiogenesis. The availability of high selective antagonists
14 for this channel makes it a promising molecular target for antiangiogenic treatments [50]. TRPV4 is
15 widely expressed in the vascular endothelium where it acts as a mechanosensor during changes in
16 cell morphology, cell swelling and shear stress [50–53]. A study conducted both *in vivo* and in
17 cultured EC reports that both shear stress and agonist-activation of TRPV4 enhance EC
18 proliferation as well as collateral growth after arterial occlusion [54]. Recently, we showed a key
19 role for TRPV4 in tumor-derived EC (TEC) migration (better discussed below) [55] (Table 1 and
20 Fig.1). It is worth noting that the dynamics of a single TRP should be considered in a more
21 integrated framework: for instance, the trafficking to the plasma membrane of TRPV4-TRPC1
22 heteromeric complex is enhanced by Ca²⁺ store depletion in HUVEC, resulting in an enhanced
23 Ca²⁺ influx upon exposure to shear flow [56].

24 A number of cellular stress factors, including hypoxia, nutrient deprivation, and reactive oxygen
25 species, are important stimuli for angiogenic signaling [57]. TRPM2 promotes macrovascular
26 pulmonary EC permeability in a H₂O₂-dependent manner. TRPM2 knockdown or overexpression
27 of the TRPM2 short isoform (that acts as dominant negative for TRPM2 long isoform)
28 significantly reduced the H₂O₂/Ca²⁺-mediated increase of paracellular permeability and cell
29 death in H5V EC [58,59] (Table 1 and Fig.1). These data open the exciting possibility of targeting
30 TRPM2 for endothelial protection against ROS-induced cell damage. Additionally, the same
31 strategy could be employed for treatment of malignant tumors, because TRPM2 isoforms are
32 expressed in different tumors, and at least one of them may function as a tumor enhancer [60].

33 Finally, TRPM7, a Ca²⁺ and Mg²⁺ permeable channel that regulates Mg²⁺ homeostasis, is
34 involved in a number of vascular disorders such as hypertension and dysfunction of endothelial
35 and smooth muscle cells [61]. A notable structural feature of TRPM7 is the presence of a kinase
36 domain at its C-terminus, making TRPM7 unique amongst ion channels, and allowing its
37 involvement both in cellular Mg²⁺ homeostasis and broad signaling [62]. TRPM7 acts negatively
38 on HUVEC proliferation and migration, whereas its functions on HMEC seem to be different [63–
39 65] (Table 1 and Fig.1). Once again, more studies are required to better understand the variability
40 of the effects induced by TRPM7 silencing in vascular endothelium.

41 In addition to the canonical angiogenesis, tumor vascularization may be supported by bone
42 marrow (BM)-derived endothelial progenitor cells (EPCs) incorporating within sprouting
43 neovessels. This feature hinted at EPC inhibition as a novel therapeutic target to pursue along
44 with anti-angiogenic treatments [1,57]. Suppression of Orai1 in EPC prevents SOCE and tubule
45 formation [45,66]. Moreover, EPCs isolated from RCC patients (RCC-EPCs) display an increased
46 SOCE, which correlates with Orai1, Stim1, and TRPC1 overexpression as compared to EPCs
47 from healthy patients: genetic suppression of Stim1, Orai1, and TRPC1 affects SOCE in RCC-
48 EPCs [67]. TRPC1 regulates proliferation and migration of EPCs isolated from rats bone marrow
49 [68] (see also Table 1 and Fig.1).

56 NICOTINIC RECEPTORS

57 nAChR are homo- or hetero-pentameric ion channels activated by endogenous acetylcholine or
58 exogenous agonists like nicotine [69]. EC express most of the known mammalian nAChR subunits
59
60

1
2
3 [70–72]. In particular $\alpha 7$ nAChR mediates the main effects of nicotine on EC, such as proliferation,
4 survival, migration, tube formation, and intracellular signaling (calcium and NO signals,
5 phosphorylation events and gene transcription). Interestingly, $\alpha 9$ and $\alpha 7$ nAChRs exert opposing
6 effects on nicotine-induced cell proliferation and survival [72–74].

7 Exposure to nicotine up-regulates $\alpha 7$ -nAChR and pharmacological inhibition of $\alpha 7$ -nAChR by
8 Mecamylamine or α -Bungarotoxin significantly and reversibly reduces EC tubulogenesis *in vitro*.
9 Even more importantly, pharmacological inhibitors or genetic disruption of $\alpha 7$ -nAChR significantly
10 suppress neo-angiogenesis in inflammation, ischemia, and neoplasia in several models. The
11 angiogenic effect of nAChR is exerted through MAPK, PI3K/Akt, and NF- κ B pathway; however,
12 since nAChR-mediated angiogenesis is only partially inhibited in $\alpha 7$ -nAChR-deficient mouse, other
13 nAChR isoforms are presumably involved [72]. Nicotine triggers neo-angiogenesis in breast, colon
14 and lung tumor cells implanted in chick chorioallantoic membranes and promotes b-FGF release
15 through the recruitment of nicotinic receptor, $\alpha v \beta 3$ integrin, and MAPK pathway [75–77]. The
16 ability of nicotine to promote late EPCs proliferation, migration, adhesion, and tubulogenesis
17 strongly suggests that its role is not restricted to mature EC [78] (Table 1 and Fig. 1)
18
19

20 **VOLUME-REGULATED ANION CHANNELS**

21 Resting normal EC expresses volume-regulated anion channels (VRACs), mainly permeable to
22 chloride ions and activated by osmotic cell swelling and shear stress. Endothelial VRACs are open
23 in resting conditions and contribute to the maintenance of the resting potential in non-stimulated
24 cells, in addition to K⁺ channels [10].

25 VRAC blockers (Mibefradil, NPPB, Tamoxifen, and Clomiphene) inhibit tube formation of rat and
26 human microvascular EC and are strongly antiangiogenic *in vivo* [79] (Table 1). Although the
27 mechanism of VRACs involvement in angiogenesis has not been clarified yet, one possible
28 explanation is that VRAC activation could lead to an increase of the driving force for Ca²⁺ entry
29 into the cell, thus affecting the intracellular Ca²⁺ concentration.
30
31

32 **WATER CHANNELS**

33 Aquaporins (AQPs) allow passive water flow in response to local osmotic gradients. They
34 contribute to epithelial secretion and absorption, and cell volume regulation. Ectopic AQP
35 expression is associated with several human cancers [12,80]. A number of reports point to AQP,
36 mainly AQP1, involvement in cell motility and tumor vascularization [81–83]: its expression in
37 tumor cells and their vasculature is variable being dependent not only on the origin of the tumor,
38 but also on its location in the host animal. This observation strengthens the strong inductive role of
39 the microenvironment on tumor features.
40

41 Interestingly, AQP1 is upregulated in human brain tumors: little or no AQP1 expression is found in
42 normal human brain microvessel endothelium, consistently with its general low permeability. On
43 the other hand, AQP1 expression in the vasculature increases with the progression from normal
44 brain to low-grade to high-grade astrocytoma [84].

45 Verkman and coworkers provided direct evidence for AQP1 role in angiogenesis *in vivo* by
46 implanting melanoma cells in AQP1 null mice and syngenic mice lacking AQP1 [85]. In both cases
47 the authors observed a markedly lower density of microvessels and the presence of islands of viable
48 tumor cells surrounded by necrotic tissue compared to control mice. Functional analyses on mouse
49 aortic EC isolated from AQP1 null mice and wild type mice revealed an impaired migration,
50 invasiveness and capability to form capillary-like structures in matrigel [85]. On the other hand,
51 RNA interference experiments performed by intratumoral injections of AQP1 siRNAs in a mouse
52 model of melanoma suggest that AQP1 inhibition can hamper tumor growth significantly lowering
53 microvessel density [86]. AQP1 is also overexpressed in both human and rodent chronic liver
54 disease. Its overexpression during cirrhosis is localized to the altered neovasculature. AQP1
55 promotes angiogenesis, fibrosis, and portal hypertension through mechanisms dependent on
56 osmotically sensitive microRNAs, as revealed on human and mouse hepatic EC [87]. Finally,
57
58
59
60

1
2
3 microvessel overexpression of AQP1 is associated with bone marrow angiogenesis in patients with
4 active multiple myeloma [88] (tab 1 and Fig.1).
5

6 **CARRIERS**

7 Beside the role of ion channels, extensive evidence points out the involvement of carriers and
8 transporters in tumor progression [89,90]. We will focus on sodium-proton exchanger and sodium-
9 calcium exchanger, the best studied so far for their involvement in tumor progression and
10 vascularization (Table 1 and Fig.1).

11 *Sodium-proton exchanger (NHE)*. It is well recognised that pathological elevations of pHi can
12 concur to some functional features of malignant cells [91]. All tumors share an altered regulation of
13 hydrogen ion dynamics and tumor progression correlates with the peculiar acid-base balance in
14 cancer cells: an extracellular acid microenvironment (pHe) linked to an alkaline intracellular pH
15 (pHi). Indeed, tumor cells have alkaline pHi values in the range of 7.12-7.7 vs 6.99-7.05 of normal
16 cells, while producing acidic pHe values of 6.2-6.9 vs 7.3-7.4 of normal cells. This reversed pH
17 gradient across the cell membrane increases with tumor progression. Since NHE is a universal and
18 conserved regulator of cellular proton balance, it received great attention. Through its action the
19 inwardly directed Na⁺ gradient can drive the uphill extrusion of protons that drives pHi
20 alkalinization and pHe acidification [91].
21

22 The highly hypoxic tumor microenvironment hyperactivates NHE1 and, since specific NHE1
23 inhibitors (Cariporide) are available, some authors propose them for innovative combination trials
24 with antiangiogenic drugs. Low concentrations of Cariporide can lead to a decrease in pHi and
25 down-regulation of VEGF. Moreover, exposure to cariporide inhibits HUVEC proliferation and
26 migration promoted by conditional medium from K562 leukemia cells. *In vivo* experiments directly
27 confirmed that inhibition of NHE1 by Cariporide could affect tumor growth and angiogenesis.
28 Tumor regression is thus presumably a result of the decreased microvessel density, which causes
29 insufficient oxygen and nutrients supply [92]. Blocking NHE1 reduces VEGF release from the
30 tumor cells suggesting that, in addition to being stimulated by hypoxia, VEGF production and
31 angiogenesis are linked to acidic pHe and to the NHE1-dependent changes in pH [93]. Systemic
32 Amiloride perfusion also reduced neovascularization experimentally induced in an animal model,
33 probably through inhibition of NHE1 [94].
34

35 *Sodium-calcium exchanger (NCX)*. Sodium influx mediated by non-selective cation channels can
36 drive to its accumulation beneath the plasma membrane. This event may increase [Ca²⁺]_i by locally
37 inverting (3Na⁺ out : 1 Ca²⁺ in) the operation mode of NCX [95].
38

39 An intriguing example has been described in HUVEC, in which a coupling between NCX and
40 voltage-dependent sodium channels (VGSCs) occurs. As previously stated, VGSC activity
41 promotes VEGF-induced proliferation, chemotaxis, and tubular differentiation and decreases
42 adhesion to substrate [25]. Moreover, Ca²⁺ inflow through reverse mode NCX is required for PKC
43 activation and targeting to the plasma membrane, as well as for VEGF-induced ERK1/2
44 phosphorylation and downstream EC functions in angiogenesis [96].
45
46

47 **CA²⁺ SIGNALS, ION CURRENTS AND CHANNELS IN TUMOR-DERIVED** 48 **ENDOTHELIAL CELLS.**

49 As previously stated, a possible reason for the failure of the antiangiogenic therapies may be the
50 high instability of EC within the tumor. It is now well established that normal and altered EC are
51 highly heterogeneous in structure and function, due to genetic modifications and the variability of
52 the local microenvironment [97–100]. The basic properties of EC obtained from different human
53 tumors (tumor-derived EC, TEC) have been investigated only recently by a limited number of
54 groups [101–103]. Breast tumor vessels display differential expression of over 1000 genes when
55 compared with normal vessels, as revealed by gene array analysis [104]. Affymetrix microarray
56 analysis of laser-captured CD31-positive blood vessels identified 63 genes that are upregulated
57
58
59
60

1
2
3 significantly (5–72 fold) in angiogenic blood vessels associated with human invasive ductal
4 carcinoma of the breast as compared with blood vessels in normal human breast [105].

5 On the other hand, TEC have been isolated and cultured from human kidney and breast carcinomas
6 on the basis of membrane markers and exhibit altered genotype, gene expression, phenotype, and
7 function. They are often aneuploid and display chromosomal instability. In addition, TEC avoid
8 senescence, a process typical of normal EC, and display enhanced proliferation, motility, and ability
9 to organize into capillary-like tubules [101,106,107]. Moreover, EC from human breast cancer are
10 significantly more radiosensitive than their normal counterparts from the same patients [108]. A
11 recent report compared the characteristics of two types of human TEC from high-metastatic (HM)
12 and low-metastatic (LM) tumors: HM-TEC showed higher proliferative and invasive activity than
13 LM-TEC [109].

14 Tumor-derived blood vessels are capillary structures and therefore TEC can be truly considered as
15 altered microvascular EC. They can be compared to ‘physiological’ microvascular EC and thus the
16 best choice would be the use to human microvascular EC obtained from the same ‘healthy’ tissue of
17 TEC. Unfortunately, it is often very difficult to isolate and maintain in culture microvascular EC
18 from all human healthy tissues: therefore dermal human microvascular EC (HMEC) are often used
19 as a physiological counterpart. Conversely, macrovascular EC, such as HUVEC, are a less suitable
20 choice, due to their features highly divergent from microvascular endothelium.

21 In the last years, our group provided substantial evidence that TEC-mediated (mainly Breast cancer-
22 derived TEC, BTEC, and more recently renal-TEC, RTEC) intracellular signaling pathways linked to
23 Ca²⁺ signals are quite different from that observed in normal human microvascular EC (Fig. 2). We
24 investigated in detail the differential effects of intracellular Ca²⁺ signaling regulated by the
25 complex and networking pathways involving arachidonic acid (AA), Nitric Oxide (NO) and
26 Hydrogen Sulfide (H₂S), key-intracellular messengers triggered by proangiogenic factors (VEGF,
27 bFGF) in vascular EC [6]. Low micromolar AA concentrations trigger NO release and protein
28 kinase A (PKA)-dependent Ca²⁺ entry which in turn stimulate BTEC migration and tubulogenesis
29 *in vitro* [110,111]. AA-dependent Ca²⁺ signals are intriguingly related to the tubule maturation
30 stage, being downregulated in the late phases of the process [55,112]. On the other hand, AA failed
31 to induce any pro-migratory effect in HMEC, with consistent significantly smaller Ca²⁺ signals
32 compared with BTEC [113] (Fig. 2).

33 Notably, both the tubulogenic and promigratory effects induced by AA are highly sensitive to
34 carboxyamidotriazole (0.1 μM CAI), a well known inhibitor of agonist-activated Ca²⁺ entry
35 [112,114]. CAI affects proliferation, invasion, metastasis, and neovascularization both *in vitro* and
36 *in vivo*. Combined to other compounds, it reduces the growth of cholangiocarcinoma, melanoma,
37 colorectal, lung, pancreatic, ovarian and breast cancer [6]. Since CAI is effective from 1 μM on
38 normal EC, the higher sensitivity of BTEC to this compound could be suitable to increase the
39 efficacy of antiangiogenic agents and to reduce their secondary effects in combination therapies.
40 Higher doses of CAI exert antiangiogenic activity in different systems such as mouse presenting
41 ischemic retinopathy, rat aortic ring culture, or chorioallantoic membrane [112].

42 H₂S is a recently discovered gasotransmitter [115,116] involved in angiogenesis regulation,
43 particularly via VEGF signaling [117]. H₂S activates Ca²⁺-permeable non-selective channels in a
44 subpopulation of BTEC and the following Ca²⁺ is enhanced in BTEC compared to HMEC.
45 Remarkably H₂S mediates tumor proangiogenic signaling triggered by VEGF: B-TEC pretreated
46 with DL-propargylglycine, an inhibitor of the H₂S-producing enzyme cystathionine γ-lyase,
47 showed drastically reduced migration and Ca²⁺ signals induced by VEGF [117] (Fig. 2). H₂S
48 donors also activate ATP-dependent K⁺ (KATP) channels both in normal EC and in BTEC [117–
49 120]. This evidence is of particular interest since during ischemic/hypoxic conditions, typical of the
50 initial phases of cancer progression, KATP channels act as ATP sensors.

51 We recently provided strong evidences about the role of TRPV4 channels in promoting AA-
52 mediated TEC migration: TRPV4 channels, are upregulated in BTEC and RTEC as compared with
53 dermal HMEC and normal kidney glomerular EC [55]. AA-activated TRPV4 is essential for BTEC
54
55
56
57
58
59
60

1
2
3 migration: loss of TRPV4 expression results in decreased Ca²⁺ responses to the TRPV4-specific
4 agonist 4 α -phorbol 12,13-didecanoate and in complete inhibition of AA-induced BTEC migration.
5 The mechanism by which AA regulates TRPV4 was also revealed in BTEC. AA induces actin
6 remodeling, which triggers TRPV4 recruitment in the plasma membrane: the consequent Ca²⁺
7 entry finally leads to BTEC migration [55].

8 However, as previously stated, TRPV4 is ubiquitous in healthy vascular endothelium and plays a
9 physiological role both in large arteries and microvessels: these relevant activities require careful
10 consideration of its therapeutic potential. On the other hand, an overexpression on TEC could be
11 exploited for a tumor targeted therapy based on lower inhibiting doses of TRPV4 antagonists which
12 could selectively affect TEC and not normal EC.
13
14
15

16 CONCLUSIONS

17 Since the seminal hypothesis proposed by Judas Folkman in '70, interference with tumor
18 vascularization is considered a key therapeutic opportunity in cancer treatment [2].

19 Unfortunately, despite promising results, vascular targeted therapy (VTT) appear short lived and
20 resistance develops in the majority of patients [121]. The relative inefficacy of VTT maybe due to
21 several reasons.
22

23 More suitable preclinical cancer models are needed in oncological practice. As previously stated,
24 vessels in cancer significantly differ from normal vasculature and the instability of EC within the
25 tumor is a relevant feature. To this purpose, the use of TEC seems a more appropriate model
26 compared to the normal EC. We expect that more detailed studies on the "transportome" in tumor
27 vascularization using the aforementioned models (beside the EC models already in use) will give
28 new input in unveiling the differences in signaling, transcriptome profiles, and vascular "ZIP
29 codes" and will likely prove to be important for understanding the conversion of normal
30 endothelial cells into tumor-associated endothelial cells. As a preliminary example, overexpression
31 of TRPV4 in TEC [55] could be useful for selectively targeted therapy using lower doses of channel
32 antagonists which affect TEC reducing secondary undesired effects on normal EC.
33

34 Another high priority challenge is the research of novel molecular anti-vascular targets (related or
35 unrelated to VEGF signaling). The evaluation of their clinical potential, in particular as combination
36 therapy with current VEGF (receptor) inhibitors, is likely to expand the antiangiogenic
37 armamentarium. In particular it could be useful to narrow the field of action for VEGF-mediated
38 targeted therapy. In this context, the recent interest on human 'transportome' involvement in tumor
39 vascularization is a promising field, since several members are activated downstream the
40 recruitment of VEGF receptors. For example, whereas the interference with the bulk VEGF
41 signaling alters the activity of a multitude of different cells and functions, targeting TRPC6 or Orail
42 may only affect EC migration and proliferation [36,37,39,45,66], while TRPC1 and STIM1 may
43 selectively influence vascular permeability [40–42,47].
44

45 It is worth noting that channels and transporters are widely distributed and ubiquitous. This feature
46 has to be carefully taken in account when considering them as clinical targets. This problem could
47 be overcome by directed targeted therapies taking advantage from nano-biomedicine: for example,
48 nanoparticle functionalization with peptide cyclic RGD for angiogenesis-specific targeting [122]
49 together with a specific channel modulator could be successfully employed.
50

51 On the other hand, the ubiquitous expression of the channels could be used as a positive feature, due
52 to the redundancy of the signaling pathways which regulates the different hallmarks of cancer: in
53 other words, the use of specific channels to selective co-target different key steps of carcinogenesis
54 beside tumor vascularization, could result in more effective and long lasting therapies. For example,
55 TRPC6 channels targeting could affect VEGF release from tumor cells as well as EC migration and
56 tumor vascularization [36,37,39].
57

58 Another important issue is the therapeutic potential of sustained vessel normalization to suppress
59 metastasis and enhance chemotherapy. Indeed, several preclinical studies have revealed that the
60

high levels of VEGF in tumors induce vessel abnormalities. It is reasonable to postulate that these vessel abnormalities could be decreased by lowering VEGF signaling. VEGF-targeted therapy induces characteristic features of vessel normalization, including reduced number and size of immature tumor vessels and increased pericyte coverage, together with decreased permeability, oedema and interstitial fluid pressure [123]. Interfering with K⁺ channels, such as EAG1 and hERG1, TRPC6 channels or NHE exchanger on tumor cells could be useful to promote vascular “normalization” by interfering with VEGF signaling during a critical window of the antiangiogenic treatments .

Finally, even if big efforts have been produced in the last years in order to characterize and study the involvement of transportome in cancer cell biology, and in particular in tumor vascularization, the field is relatively novel. The scientific interest on this topic is largely increasing as pointed out by PubMed search. The research on transportome and cancer is expected to expand even more in the next decade, and we believe that the oncogenic roles of channels, as well as the molecular mechanisms responsible for their regulation, will be largely unveiled.

FIGURE LEGEND

Table 1

Ion Channels and carriers involved in the different phases of angiogenesis. HMEC, human microvascular EC; HPAEC, human pulmonary artery EC; HUVEC, human umbilical vein EC; EA.hy926, EC line derived from HUVEC fused with human lung adenocarcinoma cell line A549; PAEC, porcine aortic endothelial cells; BTEC, tumor derived EC from breast carcinoma; H5VEC, heart endothelioma (H5V) EC; MAEC, mouse aortic EC; EPC, endothelial progenitor cells; RCC-EPC, EPC isolated from renal carcinoma patients; Numbers in parenthesis indicate the respective reference number.

Figure 1

Schematic representation of channels/transporters role in the different key steps of tumor vascularization. The mechanisms are presented in representative EC, SMC, EPC and tumors without any tissue specification. EC, endothelial cells; EPC, endothelial progenitor cells; VSM, vascular smooth muscle cells; MAPK, mitogen-activated protein kinase; PI3K, Phosphatidylinositide 3-kinases; AKT, protein kinase B; NF- κ B, nuclear factor kappa-light-chain-enhancer of activated B cells; bFGF, basic Fibroblast Growth Factor; VEGF, Vascular Endothelium Growth Factor; VEGFR, VEGF Receptor; NFAT, Nuclear factor of activated T-cells; PAR, protease-activated receptors; PTEN, Phosphatase and tensin homolog; PKC, protein kinase C.

Figure 2

A. Schematic representation of the differences between normal endothelial cells (EC) and tumor derived endothelial cells (TEC) in terms of Ca²⁺-related intracellular signaling pathways. Arachidonic Acid (AA), Nitric Oxide (NO) and Hydrogen sulfide (H₂S)-promoted Ca²⁺ signals are significantly upregulated in TEC compared with EC. These differences are at least in part due to TRPV4 overexpression and consequent TEC migration. **B.** Schematic representation of the signal transduction pathway involved in proangiogenic Ca²⁺ signals in TEC: (1) AA-mediated actin-remodeling promotes TRPV4 vesicles to traffic and insert in the plasma membrane; as a consequence, more functional channels allow Ca²⁺ entry required for TEC migration. (2) Activation of endothelial NO synthase (eNOS) mediated by AA-mediated protein kinase A (PKA) promotes NO release and consequent Ca²⁺ entry via unknown channels. (3) VEGF promotes promigratory Ca²⁺ signals mediated by H₂S via cystathionine γ -lyase (CSE).

Acknowledgements

We thank Daniele Avanzato (PhD student in Complex Systems in Life Sciences, University of Torino) for art graphics.

REFERENCES

- 1 Carmeliet, P. 2005 Angiogenesis in life, disease and medicine. *Nature* **438**, 932–6. (doi:10.1038/nature04478)
- 2 Folkman, J. 1971 Tumor angiogenesis: therapeutic implications. *The New England journal of medicine* **285**, 1182–6. (doi:10.1056/NEJM197111182852108)
- 3 Potente, M., Gerhardt, H. & Carmeliet, P. 2011 Basic and therapeutic aspects of angiogenesis. *Cell* **146**, 873–87. (doi:10.1016/j.cell.2011.08.039)
- 4 Hanahan, D. & Weinberg, R. A. 2011 Hallmarks of cancer: the next generation. *Cell* **144**, 646–674.
- 5 Fiorio Pla, A., Avanzato, D., Munaron, L. & Ambudkar, I. S. 2012 Ion channels and transporters in cancer. 6. Vascularizing the tumor: TRP channels as molecular targets. *American journal of physiology. Cell physiology* **302**, C9–15. (doi:10.1152/ajpcell.00280.2011)
- 6 Munaron, L., Genova, T., Avanzato, D., Antoniotti, S. & Fiorio Pla, A. 2013 Targeting calcium channels to block tumor vascularization. *Recent patents on anti-cancer drug discovery* **8**, 27–37.
- 7 Yao, X. & Garland, C. J. 2005 Recent developments in vascular endothelial cell transient receptor potential channels. *Circulation research* **97**, 853–63. (doi:10.1161/01.RES.0000187473.85419.3e)
- 8 Watanabe, H., Murakami, M., Ohba, T., Takahashi, Y. & Ito, H. 2008 TRP channel and cardiovascular disease. *Pharmacology & therapeutics* **118**, 337–51. (doi:10.1016/j.pharmthera.2008.03.008)
- 9 Nilius, B., Owsianik, G., Voets, T. & Peters, J. A. 2007 Transient receptor potential cation channels in disease. *Physiological reviews* **87**, 165–217. (doi:10.1152/physrev.00021.2006)
- 10 Nilius, B. & Droogmans, G. 2001 Ion channels and their functional role in vascular endothelium. *Physiological reviews* **81**, 1415–59.
- 11 Becchetti, A. 2011 Ion channels and transporters in cancer. 1. Ion channels and cell proliferation in cancer. *American journal of physiology. Cell physiology* **301**, C255–65. (doi:10.1152/ajpcell.00047.2011)
- 12 Arcangeli, A., Crociani, O., Lastraioli, E., Masi, A., Pillozzi, S. & Becchetti, A. 2009 Targeting ion channels in cancer: a novel frontier in antineoplastic therapy. *Current medicinal chemistry* **16**, 66–93.

- 1
2
3 13 Wulff, H., Castle, N. A. & Pardo, L. A. 2009 Voltage-gated potassium channels as
4 therapeutic targets. *Nature reviews. Drug discovery* **8**, 982–1001. (doi:10.1038/nrd2983)
5
6 14 Gómez-Varela, D. et al. 2007 Monoclonal antibody blockade of the human Eag1 potassium
7 channel function exerts antitumor activity. *Cancer research* **67**, 7343–9. (doi:10.1158/0008-
8 5472.CAN-07-0107)
9
10 15 Downie, B. R., Sánchez, A., Knötgen, H., Contreras-Jurado, C., Gymnopoulos, M., Weber,
11 C., Stühmer, W. & Pardo, L. A. 2008 Eag1 expression interferes with hypoxia homeostasis
12 and induces angiogenesis in tumors. *The Journal of biological chemistry* **283**, 36234–40.
13 (doi:10.1074/jbc.M801830200)
14
15 16 Wu, J., Wu, X., Zhong, D., Zhai, W., Ding, Z. & Zhou, Y. 2012 Short Hairpin RNA
17 (shRNA) Ether à go-go 1 (Eag1) Inhibition of Human Osteosarcoma Angiogenesis via
18 VEGF/PI3K/AKT Signaling. *International journal of molecular sciences* **13**, 12573–83.
19 (doi:10.3390/ijms131012573)
20
21 17 Munaron, L. & Arcangeli, A. 2013 Editorial: ion fluxes and cancer. *Recent patents on anti-
22 cancer drug discovery* **8**, 1–3.
23
24 18 D'Amico, M., Gasparoli, L. & Arcangeli, A. 2013 Potassium channels: novel emerging
25 biomarkers and targets for therapy in cancer. *Recent patents on anti-cancer drug discovery* **8**,
26 53–65.
27
28 19 Pillozzi, S. et al. 2007 VEGFR-1 (FLT-1), beta1 integrin, and hERG K⁺ channel for a
29 macromolecular signaling complex in acute myeloid leukemia: role in cell migration and
30 clinical outcome. *Blood* **110**, 1238–50. (doi:10.1182/blood-2006-02-003772)
31
32 20 Fortunato, P., Pillozzi, S., Tamburini, A., Pollazzi, L., Franchi, A., La Torre, A. & Arcangeli,
33 A. 2010 Irresponsiveness of two retinoblastoma cases to conservative therapy correlates with
34 up- regulation of hERG1 channels and of the VEGF-A pathway. *BMC cancer* **10**, 504.
35 (doi:10.1186/1471-2407-10-504)
36
37 21 Erdogan, A. et al. 2005 Margatoxin inhibits VEGF-induced hyperpolarization, proliferation
38 and nitric oxide production of human endothelial cells. *Journal of vascular research* **42**,
39 368–76. (doi:10.1159/000087159)
40
41 22 Yildirim, S., Altun, S., Gumushan, H., Patel, A. & Djamgoz, M. B. A. 2012 Voltage-gated
42 sodium channel activity promotes prostate cancer metastasis in vivo. *Cancer letters* **323**, 58–
43 61. (doi:10.1016/j.canlet.2012.03.036)
44
45 23 Djamgoz, M. B. A. & Onkal, R. 2013 Persistent current blockers of voltage-gated sodium
46 channels: a clinical opportunity for controlling metastatic disease. *Recent patents on anti-
47 cancer drug discovery* **8**, 66–84.
48
49 24 House, C. D. et al. 2010 Voltage-gated Na⁺ channel SCN5A is a key regulator of a gene
50 transcriptional network that controls colon cancer invasion. *Cancer research* **70**, 6957–67.
51 (doi:10.1158/0008-5472.CAN-10-1169)
52
53
54
55
56
57
58
59
60

- 1
2
3 25 Andrikopoulos, P. et al. 2011 Angiogenic functions of voltage-gated Na⁺ Channels in human
4 endothelial cells: modulation of vascular endothelial growth factor (VEGF) signaling. *The*
5 *Journal of biological chemistry* **286**, 16846–60. (doi:10.1074/jbc.M110.187559)
6
7 26 Bertolesi, G. E., Shi, C., Elbaum, L., Jollimore, C., Rozenberg, G., Barnes, S. & Kelly, M. E.
8 M. 2002 The Ca(2+) channel antagonists mibefradil and pimoziide inhibit cell growth via
9 different cytotoxic mechanisms. *Molecular pharmacology* **62**, 210–9.
10
11 27 Asaga, S., Ueda, M., Jinno, H., Kikuchi, K., Itano, O., Ikeda, T. & Kitajima, M. In press.
12 Identification of a new breast cancer-related gene by restriction landmark genomic scanning.
13 *Anticancer research* **26**, 35–42.
14
15 28 Panner, A. & Wurster, R. D. 2006 T-type calcium channels and tumor proliferation. *Cell*
16 *calcium* **40**, 253–9. (doi:10.1016/j.ceca.2006.04.029)
17
18 29 Kuo, I. Y.-T., Wölfle, S. E. & Hill, C. E. 2011 T-type calcium channels and vascular
19 function: the new kid on the block? *The Journal of physiology* **589**, 783–95.
20 (doi:10.1113/jphysiol.2010.199497)
21
22 30 Martini, A., Bruno, R., Mazzulla, S., Nocita, A. & Martino, G. 2010 Angiotensin II regulates
23 endothelial cell migration through calcium influx via T-type calcium channel in human
24 umbilical vein endothelial cells. *Acta physiologica (Oxford, England)* **198**, 449–55.
25 (doi:10.1111/j.1748-1716.2009.02070.x)
26
27 31 Miura, S.-I., Fujino, M., Matsuo, Y., Tanigawa, H. & Saku, K. 2005 Nifedipine-induced
28 vascular endothelial growth factor secretion from coronary smooth muscle cells promotes
29 endothelial tube formation via the kinase insert domain-containing receptor/fetal liver kinase-
30 1/NO pathway. *Hypertension research* □: *official journal of the Japanese Society of*
31 *Hypertension* **28**, 147–53. (doi:10.1291/hypres.28.147)
32
33 32 Gkika, D. & Prevarskaya, N. 2011 TRP channels in prostate cancer: the good, the bad and
34 the ugly? *Asian journal of andrology* **13**, 673–6. (doi:10.1038/aja.2011.18)
35
36 33 Ambudkar, I. S. & Ong, H. L. 2007 Organization and function of TRPC channelosomes.
37 *Pflügers Archiv* □: *European journal of physiology* **455**, 187–200. (doi:10.1007/s00424-007-
38 0252-0)
39
40 34 Pocock, T. M., Foster, R. R. & Bates, D. O. 2004 Evidence of a role for TRPC channels in
41 VEGF-mediated increased vascular permeability in vivo. *American journal of physiology.*
42 *Heart and circulatory physiology* **286**, H1015–26. (doi:10.1152/ajpheart.00826.2003)
43
44 35 Cheng, H.-W., James, A. F., Foster, R. R., Hancox, J. C. & Bates, D. O. 2006 VEGF
45 activates receptor-operated cation channels in human microvascular endothelial cells.
46 *Arteriosclerosis, thrombosis, and vascular biology* **26**, 1768–76.
47 (doi:10.1161/01.ATV.0000231518.86795.0f)
48
49 36 Hamdollah Zadeh, M. A., Glass, C. A., Magnussen, A., Hancox, J. C. & Bates, D. O. 2008
50 VEGF-mediated elevated intracellular calcium and angiogenesis in human microvascular
51 endothelial cells in vitro are inhibited by dominant negative TRPC6. *Microcirculation (New*
52 *York, N.Y.* □: *1994)* **15**, 605–14. (doi:10.1080/10739680802220323)
53
54
55
56
57
58
59
60

- 1
2
3 37 Ge, R., Tai, Y., Sun, Y., Zhou, K., Yang, S., Cheng, T., Zou, Q., Shen, F. & Wang, Y. 2009
4 Critical role of TRPC6 channels in VEGF-mediated angiogenesis. *Cancer letters* **283**, 43–51.
5 (doi:10.1016/j.canlet.2009.03.023)
6
7 38 Kini, V., Chavez, A. & Mehta, D. 2010 A new role for PTEN in regulating transient receptor
8 potential canonical channel 6-mediated Ca²⁺ entry, endothelial permeability, and
9 angiogenesis. *The Journal of biological chemistry* **285**, 33082–91.
10 (doi:10.1074/jbc.M110.142034)
11
12 39 Chigurupati, S. et al. 2010 Receptor channel TRPC6 is a key mediator of Notch-driven
13 glioblastoma growth and invasiveness. *Cancer research* **70**, 418–27. (doi:10.1158/0008-
14 5472.CAN-09-2654)
15
16 40 Mehta, D., Ahmmed, G. U., Paria, B. C., Holinstat, M., Voyno-Yasenetskaya, T., Tiruppathi,
17 C., Minshall, R. D. & Malik, A. B. 2003 RhoA interaction with inositol 1,4,5-trisphosphate
18 receptor and transient receptor potential channel-1 regulates Ca²⁺ entry. Role in signaling
19 increased endothelial permeability. *The Journal of biological chemistry* **278**, 33492–500.
20 (doi:10.1074/jbc.M302401200)
21
22 41 Paria, B. C., Vogel, S. M., Ahmmed, G. U., Alamgir, S., Shroff, J., Malik, A. B. &
23 Tiruppathi, C. 2004 Tumor necrosis factor-alpha-induced TRPC1 expression amplifies store-
24 operated Ca²⁺ influx and endothelial permeability. *American journal of physiology. Lung*
25 *cellular and molecular physiology* **287**, L1303–13. (doi:10.1152/ajplung.00240.2004)
26
27 42 Jho, D., Mehta, D., Ahmmed, G., Gao, X.-P., Tiruppathi, C., Broman, M. & Malik, A. B.
28 2005 Angiopoietin-1 opposes VEGF-induced increase in endothelial permeability by
29 inhibiting TRPC1-dependent Ca²⁺ influx. *Circulation research* **96**, 1282–90.
30 (doi:10.1161/01.RES.0000171894.03801.03)
31
32 43 Yu, P., Gu, S., Bu, J. & Du, J. 2010 TRPC1 is essential for in vivo angiogenesis in zebrafish.
33 *Circulation research* **106**, 1221–32. (doi:10.1161/CIRCRESAHA.109.207670)
34
35 44 Abdullaev, I. F., Bisailon, J. M., Potier, M., Gonzalez, J. C., Motiani, R. K. & Trebak, M.
36 2008 Stim1 and Orai1 mediate CRAC currents and store-operated calcium entry important
37 for endothelial cell proliferation. *Circulation research* **103**, 1289–99.
38 (doi:10.1161/01.RES.0000338496.95579.56)
39
40 45 Li, J. et al. 2011 Orai1 and CRAC channel dependence of VEGF-activated Ca²⁺ entry and
41 endothelial tube formation. *Circulation research* **108**, 1190–8.
42 (doi:10.1161/CIRCRESAHA.111.243352)
43
44 46 Beech, D. J. 2012 Orai1 calcium channels in the vasculature. *Pflügers Archiv*: *European*
45 *journal of physiology* **463**, 635–47. (doi:10.1007/s00424-012-1090-2)
46
47 47 Shinde, A. V et al. 2013 STIM1 controls endothelial barrier function independently of Orai1
48 and Ca²⁺ entry. *Science signaling* **6**, ra18. (doi:10.1126/scisignal.2003425)
49
50 48 Antigny, F., Girardin, N. & Frieden, M. 2012 Transient receptor potential canonical channels
51 are required for in vitro endothelial tube formation. *The Journal of biological chemistry* **287**,
52 5917–27. (doi:10.1074/jbc.M111.295733)
53
54
55
56
57
58
59
60

- 1
2
3 49 Cheng, K. T., Liu, X., Ong, H. L., Swaim, W. & Ambudkar, I. S. 2011 Local Ca²⁺ entry via
4 Orail regulates plasma membrane recruitment of TRPC1 and controls cytosolic Ca²⁺ signals
5 required for specific cell functions. *PLoS biology* **9**, e1001025.
6 (doi:10.1371/journal.pbio.1001025)
7
- 8
9 50 Everaerts, W., Nilius, B. & Owsianik, G. 2010 The vanilloid transient receptor potential
10 channel TRPV4: from structure to disease. *Progress in biophysics and molecular biology*
11 **103**, 2–17. (doi:10.1016/j.pbiomolbio.2009.10.002)
12
- 13 51 Vriens, J., Watanabe, H., Janssens, A., Droogmans, G., Voets, T. & Nilius, B. 2004 Cell
14 swelling, heat, and chemical agonists use distinct pathways for the activation of the cation
15 channel TRPV4. *Proceedings of the National Academy of Sciences of the United States of*
16 *America* **101**, 396–401. (doi:10.1073/pnas.0303329101)
17
- 18 52 Hartmannsgruber, V., Heyken, W.-T., Kacik, M., Kaistha, A., Grgic, I., Harteneck, C.,
19 Liedtke, W., Hoyer, J. & Köhler, R. 2007 Arterial response to shear stress critically depends
20 on endothelial TRPV4 expression. *PloS one* **2**, e827. (doi:10.1371/journal.pone.0000827)
21
- 22 53 Thodeti, C. K., Matthews, B., Ravi, A., Mammoto, A., Ghosh, K., Bracha, A. L. & Ingber,
23 D. E. 2009 TRPV4 channels mediate cyclic strain-induced endothelial cell reorientation
24 through integrin-to-integrin signaling. *Circulation research* **104**, 1123–30.
25 (doi:10.1161/CIRCRESAHA.108.192930)
26
- 27 54 Troidl, C. et al. 2009 Trpv4 induces collateral vessel growth during regeneration of the
28 arterial circulation. *Journal of cellular and molecular medicine* **13**, 2613–21.
29 (doi:10.1111/j.1582-4934.2008.00579.x)
30
- 31 55 Fiorio Pla, A., Ong, H. L., Cheng, K. T., Brossa, A., Bussolati, B., Lockwich, T., Paria, B.,
32 Munaron, L. & Ambudkar, I. S. 2012 TRPV4 mediates tumor-derived endothelial cell
33 migration via arachidonic acid-activated actin remodeling. *Oncogene* **31**, 200–12.
34 (doi:10.1038/onc.2011.231)
35
- 36 56 Ma, X., Cao, J., Luo, J., Nilius, B., Huang, Y., Ambudkar, I. S. & Yao, X. 2010 Depletion of
37 intracellular Ca²⁺ stores stimulates the translocation of vanilloid transient receptor potential
38 4-c1 heteromeric channels to the plasma membrane. *Arteriosclerosis, thrombosis, and*
39 *vascular biology* **30**, 2249–55. (doi:10.1161/ATVBAHA.110.212084)
40
- 41 57 North, S., Moenner, M. & Bikfalvi, A. 2005 Recent developments in the regulation of the
42 angiogenic switch by cellular stress factors in tumors. *Cancer letters* **218**, 1–14.
43 (doi:10.1016/j.canlet.2004.08.007)
44
- 45 58 Hecquet, C. M., Ahmmed, G. U., Vogel, S. M. & Malik, A. B. 2008 Role of TRPM2 channel
46 in mediating H₂O₂-induced Ca²⁺ entry and endothelial hyperpermeability. *Circulation*
47 *research* **102**, 347–55. (doi:10.1161/CIRCRESAHA.107.160176)
48
- 49 59 Sun, L., Yau, H.-Y., Wong, W.-Y., Li, R. A., Huang, Y. & Yao, X. 2012 Role of TRPM2 in
50 H₂O₂-induced cell apoptosis in endothelial cells. *PloS one* **7**, e43186.
51 (doi:10.1371/journal.pone.0043186)
52
53
54
55
56
57
58
59
60

- 1
2
3 60 Orfanelli, U., Wenke, A.-K., Doglioni, C., Russo, V., Bosserhoff, A. K. & Lavorgna, G. 2008 Identification of novel sense and antisense transcription at the TRPM2 locus in cancer. *Cell research* **18**, 1128–40. (doi:10.1038/cr.2008.296)
- 4
5
6
7 61 Yogi, A., Callera, G. E., Antunes, T. T., Tostes, R. C. & Touyz, R. M. 2011 Transient
8 receptor potential melastatin 7 (TRPM7) cation channels, magnesium and the vascular
9 system in hypertension. *Circulation journal* □: *official journal of the Japanese Circulation*
10 *Society* **75**, 237–45.
- 11
12 62 Paravicini, T. M., Chubanov, V. & Gudermann, T. 2012 TRPM7: a unique channel involved
13 in magnesium homeostasis. *The international journal of biochemistry & cell biology* **44**,
14 1381–4. (doi:10.1016/j.biocel.2012.05.010)
- 15
16 63 Inoue, K. & Xiong, Z.-G. 2009 Silencing TRPM7 promotes growth/proliferation and nitric
17 oxide production of vascular endothelial cells via the ERK pathway. *Cardiovascular*
18 *research* **83**, 547–57. (doi:10.1093/cvr/cvp153)
- 19
20 64 Baldoli, E. & Maier, J. A. M. 2012 Silencing TRPM7 mimics the effects of magnesium
21 deficiency in human microvascular endothelial cells. *Angiogenesis* **15**, 47–57.
22 (doi:10.1007/s10456-011-9242-0)
- 23
24 65 Baldoli, E., Castiglioni, S. & Maier, J. A. M. 2013 Regulation and function of TRPM7 in
25 human endothelial cells: TRPM7 as a potential novel regulator of endothelial function. *PloS*
26 *one* **8**, e59891. (doi:10.1371/journal.pone.0059891)
- 27
28 66 Dragoni, S. et al. 2011 Vascular endothelial growth factor stimulates endothelial colony
29 forming cells proliferation and tubulogenesis by inducing oscillations in intracellular Ca²⁺
30 concentration. *Stem cells (Dayton, Ohio)* **29**, 1898–907. (doi:10.1002/stem.734)
- 31
32 67 Lodola, F. et al. 2012 Store-operated Ca²⁺ entry is remodelled and controls in vitro
33 angiogenesis in endothelial progenitor cells isolated from tumoral patients. *PloS one* **7**,
34 e42541. (doi:10.1371/journal.pone.0042541)
- 35
36 68 Kuang, C., Yu, Y., Wang, K., Qian, D., Den, M. & Huang, L. 2012 Knockdown of transient
37 receptor potential canonical-1 reduces the proliferation and migration of endothelial
38 progenitor cells. *Stem cells and development* **21**, 487–96. (doi:10.1089/scd.2011.0027)
- 39
40 69 Taly, A., Corringer, P.-J., Guedin, D., Lestage, P. & Changeux, J.-P. 2009 Nicotinic
41 receptors: allosteric transitions and therapeutic targets in the nervous system. , 1–18.
- 42
43 70 Cardinale, A., Nastrucci, C., Cesario, A. & Russo, P. 2012 Nicotine: specific role in
44 angiogenesis, proliferation and apoptosis. *Critical Reviews in Toxicology* **42**, 68–89.
- 45
46 71 Egleton, R. D., Brown, K. C. & Dasgupta, P. 2009 Angiogenic activity of nicotinic
47 acetylcholine receptors: Implications in tobacco-related vascular diseases. *Pharmacology*
48 *& therapeutics* **121**, 205–223.
- 49
50 72 Heeschen, C., Weis, M., Aicher, A., Dimmeler, S. & Cooke, J. P. 2002 A novel angiogenic
51 pathway mediated by non-neuronal nicotinic acetylcholine receptors. *Journal of Clinical*
52 *Investigation* **110**, 527–536.
- 53
54
55
56
57
58
59
60

- 1
2
3 73 Wu, J. C. F., Chruscinski, A., De Jesus Perez, V. A., Singh, H., Pitsiouni, M., Rabinovitch,
4 M., Utz, P. J. & Cooke, J. P. 2009 Cholinergic modulation of angiogenesis: Role of the 7
5 nicotinic acetylcholine receptor. *Journal of cellular biochemistry* **108**, 433–446.
6
- 7 74 Ng, M. K. C., Wu, J., Chang, E., Wang, B., Katzenberg-Clark, R., Ishii-Watabe, A. & Cooke,
8 J. P. 2007 A central role for nicotinic cholinergic regulation of growth factor-induced
9 endothelial cell migration. *Arteriosclerosis, thrombosis, and vascular biology* **27**, 106–12.
10 (doi:10.1161/01.ATV.0000251517.98396.4a)
11
- 12 75 Mousa, S. & Mousa, S. A. 2006 Cellular and molecular mechanisms of nicotine's pro-
13 angiogenesis activity and its potential impact on cancer. *Journal of cellular biochemistry* **97**,
14 1370–1378.
15
- 16 76 Arias, H. R., Richards, V. E., Ng, D., Ghafoori, M. E., Le, V. & Mousa, S. A. 2009 Role of
17 non-neuronal nicotinic acetylcholine receptors in angiogenesis. *The international journal of*
18 *biochemistry & cell biology* **41**, 1441–51. (doi:10.1016/j.biocel.2009.01.013)
19
- 20 77 Cardinale, A., Nastrucci, C., Cesario, A. & Russo, P. 2012 Nicotine: specific role in
21 angiogenesis, proliferation and apoptosis. *Critical Reviews in Toxicology* **42**, 68–89.
22
- 23 78 Yu, M., Liu, Q., Sun, J., Yi, K., Wu, L. & Tan, X. 2011 Nicotine improves the functional
24 activity of late endothelial progenitor cells via nicotinic acetylcholine receptors. *Biochemistry*
25 *and cell biology = Biochimie et biologie cellulaire* **89**, 405–10. (doi:10.1139/o11-032)
26
- 27 79 Manolopoulos, V. G., Liekens, S., Koolwijk, P., Voets, T., Peters, E., Droogmans, G.,
28 Lelkes, P. I., De Clercq, E. & Nilius, B. 2000 Inhibition of angiogenesis by blockers of
29 volume-regulated anion channels. *General pharmacology* **34**, 107–16.
30
- 31 80 Monzani, E., Shtil, A. A. & La Porta, C. A. M. 2007 The water channels, new druggable
32 targets to combat cancer cell survival, invasiveness and metastasis. *Current drug targets* **8**,
33 1132–7.
34
- 35 81 Verkman, A. S. 2012 Aquaporins in clinical medicine. *Annual review of medicine* **63**, 303–
36 16. (doi:10.1146/annurev-med-043010-193843)
37
- 38 82 Alleva, K., Chara, O. & Amodeo, G. 2012 Aquaporins: another piece in the osmotic puzzle.
39 *FEBS letters* **586**, 2991–9. (doi:10.1016/j.febslet.2012.06.013)
40
- 41 83 Endo, M., Jain, R. K., Witwer, B. & Brown, D. 1999 Water channel (aquaporin 1) expression
42 and distribution in mammary carcinomas and glioblastomas. *Microvascular research* **58**, 89–
43 98. (doi:10.1006/mvre.1999.2158)
44
- 45 84 Saadoun, S., Papadopoulos, M. C., Davies, D. C., Bell, B. A. & Krishna, S. 2002 Increased
46 aquaporin 1 water channel expression in human brain tumours. *British journal of cancer* **87**,
47 621–3. (doi:10.1038/sj.bjc.6600512)
48
- 49 85 Saadoun, S., Papadopoulos, M. C., Hara-Chikuma, M. & Verkman, A. S. 2005 Impairment
50 of angiogenesis and cell migration by targeted aquaporin-1 gene disruption. *Nature* **434**,
51 786–92. (doi:10.1038/nature03460)
52
53
54
55
56
57
58
59
60


- 1
2
3 86 Nicchia, G. P., Stigliano, C., Sparaneo, A., Rossi, A., Frigeri, A. & Svelto, M. 2013
4 Inhibition of aquaporin-1 dependent angiogenesis impairs tumour growth in a mouse model
5 of melanoma. *Journal of molecular medicine (Berlin, Germany)* **91**, 613–23.
6 (doi:10.1007/s00109-012-0977-x)
7
8
9 87 Huebert, R. C., Jagavelu, K., Hendrickson, H. I., Vasdev, M. M., Arab, J. P., Splinter, P. L.,
10 Trussoni, C. E., Larusso, N. F. & Shah, V. H. 2011 Aquaporin-1 promotes angiogenesis,
11 fibrosis, and portal hypertension through mechanisms dependent on osmotically sensitive
12 microRNAs. *The American journal of pathology* **179**, 1851–60.
13 (doi:10.1016/j.ajpath.2011.06.045)
14
15 88 Vacca, A., Frigeri, A., Ribatti, D., Nicchia, G. P., Nico, B., Ria, R., Svelto, M. & Dammacco,
16 F. 2001 Microvessel overexpression of aquaporin 1 parallels bone marrow angiogenesis in
17 patients with active multiple myeloma. *British journal of haematology* **113**, 415–21.
18
19 89 Cardone, R. A., Casavola, V. & Reshkin, S. J. 2005 The role of disturbed pH dynamics and
20 the Na⁺/H⁺ exchanger in metastasis. *Nature reviews. Cancer* **5**, 786–95.
21 (doi:10.1038/nrc1713)
22
23 90 Monteith, G. R., Davis, F. M. & Roberts-Thomson, S. J. 2012 Calcium channels and pumps
24 in cancer: changes and consequences. *The Journal of biological chemistry* **287**, 31666–73.
25 (doi:10.1074/jbc.R112.343061)
26
27 91 Reshkin, S. J., Cardone, R. A. & Harguindey, S. 2013 Na⁺-H⁺ exchanger, pH regulation and
28 cancer. *Recent patents on anti-cancer drug discovery* **8**, 85–99.
29
30 92 Gao, W. et al. 2011 Inhibition of K562 leukemia angiogenesis and growth by selective
31 Na⁺/H⁺ exchanger inhibitor cariporide through down-regulation of pro-angiogenesis factor
32 VEGF. *Leukemia research* **35**, 1506–11. (doi:10.1016/j.leukres.2011.07.001)
33
34 93 Xu, L., Fukumura, D. & Jain, R. K. 2002 Acidic extracellular pH induces vascular
35 endothelial growth factor (VEGF) in human glioblastoma cells via ERK1/2 MAPK signaling
36 pathway: mechanism of low pH-induced VEGF. *The Journal of biological chemistry* **277**,
37 11368–74. (doi:10.1074/jbc.M108347200)
38
39 94 Avery, R. L., Connor, T. B. & Farazdaghi, M. 1990 Systemic amiloride inhibits
40 experimentally induced neovascularization. *Archives of ophthalmology* **108**, 1474–6.
41
42 95 Blaustein, M. P. & Lederer, W. J. 1999 Sodium/calcium exchange: its physiological
43 implications. *Physiological reviews* **79**, 763–854.
44
45 96 Andrikopoulos, P., Baba, A., Matsuda, T., Djamgoz, M. B. A., Yaqoob, M. M. & Eccles, S.
46 A. 2011 Ca²⁺ influx through reverse mode Na⁺/Ca²⁺ exchange is critical for vascular
47 endothelial growth factor-mediated extracellular signal-regulated kinase (ERK) 1/2
48 activation and angiogenic functions of human endothelial cells. *The Journal of biological*
49 *chemistry* **286**, 37919–31. (doi:10.1074/jbc.M111.251777)
50
51 97 Aird, W. C. 2012 Endothelial cell heterogeneity. *Cold Spring Harbor perspectives in*
52 *medicine* **2**, a006429.
53
54
55
56
57
58
59
60

- 1
2
3 98 Regan, E. R. & Aird, W. C. 2012 Dynamical systems approach to endothelial heterogeneity. *Circulation Research* **111**, 110–130.
4
5
6 99 Yano, K. et al. 2007 Phenotypic heterogeneity is an evolutionarily conserved feature of the
7 endothelium. *Blood* **109**, 613–5.
8
9 100 Chi, J.-T. et al. 2003 Endothelial cell diversity revealed by global expression profiling. *Proc*
10 *Natl Acad Sci USA* **100**, 10623–10628.
11
12 101 Bussolati, B., Grange, C. & Camussi, G. 2011 Tumor exploits alternative strategies to
13 achieve vascularization. *FASEB journal*: official publication of the Federation of American
14 *Societies for Experimental Biology* **25**, 2874–82. (doi:10.1096/fj.10-180323)
15
16 102 Ghilardi, C., Chiorino, G., Dossi, R., Nagy, Z., Giavazzi, R. & Bani, M. 2008 Identification
17 of novel vascular markers through gene expression profiling of tumor-derived endothelium.
18 *BMC genomics* **9**, 201.
19
20 103 Allport, J. R. & Weissleder, R. 2003 Murine Lewis lung carcinoma-derived endothelium
21 expresses markers of endothelial activation and requires tumor-specific extracellular matrix
22 in vitro. *Neoplasia (New York, NY)* **5**, 205–217.
23
24 104 Bhati, R. et al. 2008 Molecular characterization of human breast tumor vascular cells. *The*
25 *American Journal of Pathology* **172**, 1381–1390.
26
27 105 Ohga, N. et al. 2012 Heterogeneity of tumor endothelial cells: comparison between tumor
28 endothelial cells isolated from high- and low-metastatic tumors. *The American Journal of*
29 *Pathology* **180**, 1294–1307.
30
31 106 Bussolati, B., Deambrosio, I., Russo, S., Deregibus, M. C. & Camussi, G. 2003 Altered
32 angiogenesis and survival in human tumor-derived endothelial cells. *FASEB journal*:
33 *official publication of the Federation of American Societies for Experimental Biology* **17**,
34 1159–61. (doi:10.1096/fj.02-0557fje)
35
36 107 Grange, C., Bussolati, B., Bruno, S., Fonsato, V., Sapino, A. & Camussi, G. 2006 Isolation
37 and characterization of human breast tumor-derived endothelial cells. *Oncol Rep* **15**, 381–
38 386.
39
40 108 Park, M.-T. et al. 2012 The radiosensitivity of endothelial cells isolated from human breast
41 cancer and normal tissue in vitro. *Microvascular research* **84**, 140–8.
42 (doi:10.1016/j.mvr.2012.06.002)
43
44 109 Ohga, N. et al. 2012 Heterogeneity of tumor endothelial cells: comparison between tumor
45 endothelial cells isolated from high- and low-metastatic tumors. *The American Journal of*
46 *Pathology* **180**, 1294–1307.
47
48 110 Fiorio Pla, A., Grange, C., Antoniotti, S., Tomatis, C., Merlino, A., Bussolati, B. & Munaron,
49 L. 2008 Arachidonic acid-induced Ca²⁺ entry is involved in early steps of tumor
50 angiogenesis. *Mol Cancer Res* **6**, 535–545.
51
52
53
54
55
56
57
58
59
60


- 1
2
3 111 Fiorio Pla, A., Genova, T., Pupo, E., Tomatis, C., Genazzani, A., Zaninetti, R. & Munaron,
4 L. 2010 Multiple roles of protein kinase a in arachidonic acid-mediated Ca²⁺ entry and
5 tumor-derived human endothelial cell migration. *Molecular Cancer Research* **8**, 1466–1476.
6
7 112 Fiorio Pla, A., Grange, C., Antoniotti, S., Tomatis, C., Merlino, A., Bussolati, B. & Munaron,
8 L. 2008 Arachidonic acid-induced Ca²⁺ entry is involved in early steps of tumor
9 angiogenesis. *Mol Cancer Res* **6**, 535–545.
10
11 113 Fiorio Pla, A., Genova, T., Pupo, E., Tomatis, C., Genazzani, A., Zaninetti, R. & Munaron,
12 L. 2010 Multiple roles of protein kinase a in arachidonic acid-mediated Ca²⁺ entry and
13 tumor-derived human endothelial cell migration. *Molecular Cancer Research* **8**, 1466–1476.
14
15 114 Kohn, E. C., Felder, C. C., Jacobs, W., Holmes, K. A., Day, A., Freer, R. & Liotta, L. A.
16 1994 Structure–function analysis of signal and growth inhibition by carboxyamido-triazole,
17 CAI. *Cancer research* **54**, 935–42.
18
19 115 Mancardi, D., Pla, A. F., Moccia, F., Tanzi, F. & Munaron, L. 2011 Old and new
20 gasotransmitters in the cardiovascular system: focus on the role of nitric oxide and hydrogen
21 sulfide in endothelial cells and cardiomyocytes. *Curr Pharm Biotechnol* **12**, 1406–1415.
22
23 116 Wang, R. 2012 Physiological implications of hydrogen sulfide: a whiff exploration that
24 blossomed. *Physiol Rev* **92**, 791–896.
25
26 117 Pupo, E., Pla, A. F., Avanzato, D., Moccia, F., Cruz, J. E., Tanzi, F., Merlino, A., Mancardi,
27 D. & Munaron, L. 2011 Hydrogen sulfide promotes calcium signals and migration in tumor-
28 derived endothelial cells. *Free Radic Biol Med* **51**, 1765–1773.
29
30 118 Cai, W.-J., Wang, M.-J., Moore, P. K., Jin, H.-M., Yao, T. & Zhu, Y.-C. 2007 The novel
31 proangiogenic effect of hydrogen sulfide is dependent on Akt phosphorylation.
32 *Cardiovascular research* **76**, 29–40. (doi:10.1016/j.cardiores.2007.05.026)
33
34 119 Tang, G., Wu, L. & Wang, R. 2010 Interaction of hydrogen sulfide with ion channels.
35 *Clinical and experimental pharmacology & physiology* **37**, 753–63.
36
37 120 Munaron, L. & Fiorio Pla, A. 2009 Endothelial calcium machinery and angiogenesis:
38 understanding physiology to interfere with pathology. *Curr Med Chem* **16**, 4691–4703.
39
40 121 Ebos, J. M. L. & Kerbel, R. S. 2011 Antiangiogenic therapy: impact on invasion, disease
41 progression, and metastasis. *Nature Reviews Clinical Oncology* **8**, 210–221.
42 (doi:10.1038/nrclinonc.2011.21)
43
44 122 Cheng, J., Gu, Y.-J., Wang, Y., Cheng, S. H. & Wong, W.-T. 2011 Nanotherapeutics in
45 angiogenesis: synthesis and in vivo assessment of drug efficacy and biocompatibility in
46 zebrafish embryos. *International journal of nanomedicine* **6**, 2007–21.
47 (doi:10.2147/IJN.S20145)
48
49 123 Carmeliet, P. & Jain, R. K. 2011 Principles and mechanisms of vessel normalization for
50 cancer and other angiogenic diseases. *Nature reviews. Drug discovery* **10**, 417–27.
51 (doi:10.1038/nrd3455)
52
53
54
55
56
57
58
59
60

Channels/transporters	TRPC1	TRPC6	TRPC3,4,5	TRPV4	TRPM2	TRPM7	Orai1/Stim1	Nav	Cav	Kv	VRAC	NHE1	NCX	AQP1	nAChR	
Migration	RCC-EPC [67] EPC[65]	HMEC [36]		BTEC [55]			RCC-EPC [67] HUVEC [44-46]	HUVEC [25]	HUVEC [30]			HUVEC [92]	HUVEC [25, 97]	MAEC from KO mice [85]	HMEC [73, 74] EPC [78]	
Survival and Proliferation	RCC-EPC [67] EPC [45, 68]	HMEC [36] HUVEC [37]		PAECs [54]	HSVEC [58]	HUVEC, HMEC [63-65]	RCC-EPC [67] HUVEC [42]	HUVEC [25]		Kv.1.3-HUVEC[21]		HUVEC [92]	HUVEC [25, 97]	human hepatic sinusoidal ECs [87]	Human pulmonary artery ECs, human retinal microvascular ECs; HUVEC, HMEC [73] EPC [78]	
In vitro Tube Formation	RCC-EPC [67] HUVEC, EA.hy926 [48]	HMEC [36, 39]	EA.hy926 [48]				RCC-EPC [67] HUVEC, EPC [45, 48]	HUVEC [25]			Microvascular ECs from the rat adrenal medulla (RAMECs), HMEC [79]	HUVEC [92]	HUVEC [25, 97]	MAEC from KO mice [85]	HUVEC, HMEC [72, 73] EPC [78]	
Permeability	HMEC, HUVEC [40-42]	HPAEC [38] Frog mesenteric microvessels [34]			HSVEC [58]		HUVEC [47]									
In vivo Angiogenesis	Zebrafish [43]	CAM [37]		Collateral growth [54]			CAM [43]			HERG-1- Retinoblastoma [20] EAG1-Xenograft in SCID mice and human osteosarcoma [15, 16]		Xenograft in nude mice [92] Rabbit cornea [94]		Human mammary carcinoma, glioblastoma [83, 84] AQP1 KO mice and C57BL/6 mice [85, 86, 87] Bone marrow angiogenesis in patients with active multiple myeloma [88]	Disc angiogenesis system, hind limb ischemia [72] Breast, colon and lung tumor cells implanted in CAM [75]	



AQUAPORINS & CARRIERS




NICOTINIC RECEPTORS




VOLTAGE - GATED CHANNELS




TRP channels




channels in EPC



TRP channels



TRP channels



Blood Vessel

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

