



## AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Novel approaches for the design and discovery of quorum-sensing inhibitors.

This is the author's manuscript	
Original Citation:	
Availability:	
This version is available http://hdl.handle.net/2318/143270	since 2020-11-03T10:22:45Z
Published version:	
DOI:10.1517/17460441.2014.894974	
Terms of use:	
Open Access	
Anyone can freely access the full text of works made available a under a Creative Commons license can be used according to the of all other works requires consent of the right holder (author or	as "Open Access". Works made available e terms and conditions of said license. Use

(Article begins on next page)

protection by the applicable law.



# UNIVERSITÀ DEGLI STUDI DI TORINO

$\frac{2}{3}$	
5 4	
5	
6	This is an author version of the contribution published on:
7	Questa è la versione dell'autore dell'opera:
, 8	[Fynert Onin Drug Discov 9(4) 2014 DOI 10 1517/17460441 2014 8949741
0	owero [Scutera S Zucca M Savoja D 9 Informa Healthcare 2014 page 353-66]
10	The definitive version is available at:
10	I a versione definitiva è disponibile alla UPI :
11	$La \ versione \ aejiniliva \ e \ aisponibile \ alla \ OKL.$
12	[nttp://injormanealthcare.com/aol/abs/10.151//1/400441.2014.8949/4]
13	
14 15	
16	
17	
18	
19	
20	
21	
22	
23	
24 25	
25	
27	
28	
29	
30	
31	
32	
33	
54 25	
36	
37	
38	

1 2

4

#### Novel approaches for the design and discovery of quorum-sensing inhibitors

3 Sara Scutera, Mario Zucca & Dianella Savoia†

<sup>5</sup> †University of Torino, S. Luigi Gonzaga Hospital, Department of Clinical and Biological Sciences,
<sup>6</sup> Orbassano (To), Italy

7 Abstract

8 Introduction: The spread of antibiotic resistance, together with the lack of antibiotics based on 9 novel molecular scaffolds, mark the so-called "post-antibiotic era." Among the potential strategies 10 to develop new anti-infective drugs, interference with bacterial virulence is an attractive approach. 11 The discovery that virulence gene expression is mostly regulated by quorum sensing (QS) raised a 12 lot of interest and prompted a lot of research aimed at finding inhibitors of this mechanism. 13 *Areas covered:* This paper deals with the most recent strategies aimed at discovering new inhibitors 14 able to disrupt the different steps of the OS system, targeting signal production, signal molecules 15 and signal receptors. We provide an overview of the literature including research papers, mostly 16 dealing with inhibitors of the S. aureus and P. aeruginosa QS systems, and a number of 17 comprehensive reviews dealing with the application of the newest technologies to the field. 18 Prospects and emerging concerns regarding the possible clinical applications are also discussed. 19 *Expert opinion:* QS inhibition is a promising strategy against infections. However, despite the 20 discovery of a huge number of QS inhibitors, with about 40 patents, the potential of QS inhibition is 21 still to be fully assessed. The current validation methods of QS inhibitors must be optimized, and 22 the discovery that QS disruption may favour or select more virulent strains must be investigated in 23 depth. Given the current market-dependent situation, the possibility to develop hits into licensed 24 drugs is likely to be made possible by joint ventures between private companies, academia and 25 public institutions.

26

Keywords: Antibodies; autoinducers; *P. aeruginosa*; quorum sensing; receptors; *S. aureus*,
virulence factors.

# 1 1. Introduction

2 Intercellular communication by means of secreted hormone-like molecules (pheromones) is an 3 important natural feature, which in different bacterial species is involved in bioluminescence, 4 virulence, biofilm formation, antibiotic production, competence, conjugation, swarming, motility, 5 and sporulation [1]. For most pathogenic bacteria, the ability to implement this kind of multicellular 6 coordination is critical for disease pathogenesis and host colonization. Quorum sensing (QS) is the 7 term used to describe a density-dependent intercellular communication that allows a coordinated 8 multicellular behaviour in prokaryotes, based on the constitutive production and secretion of small 9 molecules defined quorum sensors or autoinducers (AI) [2]. When the AI concentration reaches a 10 critical detection threshold that depends upon bacterial population density (quorum), the AI 11 interaction with species-specific receptors triggers a signal transduction cascade leading to an 12 alteration in gene expression. The production of a number of virulence factors involved in bacterial 13 pathogenesis and in the formation of bacterial sessile communities called biofilms is based on this 14 mechanism. Biofilms account for more than 80% of human infections [3] and their treatment is 15 highly problematic, because sessile bacteria are up to 1000 times more resistant to antibiotics in 16 comparison to their planktonic counterparts, and are less exposed to the host immune system [4]. 17 Since the pioneering work on AI by Tomasz and Hotchkiss [5] and Nealson and co-workers, 18 [6], several AI molecules have been identified, which may be classified as follows: i) AI-1, or N-19 acyl homoserine lactones (AHLs), which are produced by over 70 species of Gram-negative 20 bacteria; ii) oligopeptides, consisting of 5-17 amino acids, which are generally produced by Gram-21 positive bacteria; iii) AI-2, a class of furanone-based AI derived from 4,5-dihydroxy-2,3-22 pentanedione (DPD), which can be produced by both Gram-positive and Gram-negative bacteria 23 and may be considered universal interspecies signalling molecules; and iv) other small molecules 24 that include the Pseudomonas quinolone signal (PQS), the Vibrio cholerae autoinducer (S)-3-

25 hydroxytridecan-4-one, 3-hydroxypalmitic acid methyl ester, the AI-3 (a small AI with unknown

structure, used by enterohemorrhagic *E. coli*), bradyoxetin, and other molecules that do not fall
 within one of the major groups [7].

3 QS inhibition, also called quorum quenching (QQ), is considered an attractive modality of 4 therapeutic intervention for a series of reasons: i) it would overcome the conventional mechanisms 5 of resistance to antibiotics; ii) a single inhibitor could have multiple effects, being able to interfere 6 with the production of a number of virulence factors and with biofilm formation; iii) in principle, 7 anti-virulence factors should not exert an evolutionary pressure on the targeted bacterial population, 8 avoiding the emergence of resistant strains; iv) avoiding or limiting biofilm formation, QS 9 inhibitors could enhance the efficacy of existing antibiotics; v) except AI-2 members, most AI are 10 species-specific. Consequently, the narrow spectrum of activity of inhibitors should avoid the 11 disruption of beneficial microbiota [8]. It is well known that both prokaryotic and eukaryotic 12 organisms have evolved molecular mechanisms able to carry on QQ activity. The study of the 13 signal degradation systems enacted by different bacteria, plants, and mammals may lead to the 14 design of novel approaches for quenching bacterial QS [9]. Ideally, a QS inhibitor should meet the 15 following criteria: it should be a small molecule, highly specific, non toxic for eukaryotic cells, and 16 chemically stable [10].

17 QQ compounds could be used in many fields, such as human and veterinary medicine, 18 agriculture, and aquaculture. The therapy of chronic infections, which are often due to multiresistant 19 bacterial strains, could benefit particularly from the use of antivirulence drugs [11]. By means of *in* 20 *vitro* and *in vivo* experiments, it has been shown that *P. aeruginosa* QS blockage restores the 21 immune response efficiency favouring the clearance of bacterial biofilms [12].

QS systems generally offer three points of attack: the signal generation process, the signal molecule, and the signal receptor [13]. During the last three decades, many natural and synthetic agents belonging to the categories of small non-peptide molecules, peptides, and enzymes, have been identified as QS inhibitors [14]. A very comprehensive review on QS inhibitors has recently been published by Kalia [10]. The field is rapidly evolving, and many inhibitors are being

1 discovered. However, clinical applications are still lacking. To the best of our knowledge, only two 2 clinical trials on QS inhibitors have been performed. In the first one (ClinicalTrials.gov Identifier: 3 NCT00610623), the effect of azithromycin, used as a QS blocker at subinhibitory doses for the 4 prevention of *P. aeruginosa* ventilator-associated pneumonia was evaluated. The authors of the 5 study concluded that the major effect of QS inhibition was the selection of more virulent strains. 6 The second trial (ClinicalTrials.gov Identifier: NCT01201577), completed but with no study results 7 posted, was aimed at studying the effect of orally administered combinations of prebiotics, 8 probiotics and antibiotics on QS signalling molecules and on innate and adaptive immunity in 9 humans.

In this article, we focus on modern approaches for the development and validation of QS
inhibitors, providing the reader with a comprehensive overview of the recent literature on the
subject.

13

## 14 **2. Strategies targeting QS signal production**

15 The studies aimed to hit at targeting this step are relatively few, and the field is open for further 16 investigation. Compounds affecting the AHL biosynthesis or efflux pumps are likely to behave as 17 QQ. Substrate analogues of AI, like butyryl-S-adenosylmethionine, holo-acyl carrier protein, 18 sinefungin and L/D-S-adenosylhomocysteine can block AHL production in vitro. However, in vivo 19 experiments have not been performed, because these homologues are likely to affect the central 20 pathways of amino acid and fatty acid metabolism [15]. Small-molecule inhibitors of the AHL 21 synthase have been identified by Chung et al. [16]. The involvement of multidrug efflux pumps in 22 the QS of *Burkholderia pseudomallei* has been established, suggesting that this pathway could also 23 be exploited [17].

Methylthioadenosine/S-adenosyl homocysteine nucleosidase (MTAN) is the product of the *pfs* gene, which is highly conserved across bacterial species. This enzyme is present in bacteria but not in mammals, in which its activity is performed by two different enzymes: methylthioadenosine

1 phosphorylase and purine nucleotide phosphorylase [18]. Beyond playing a crucial role in 2 maintaining homeostasis, MTAN is directly involved in the biosynthesis of AI-1 and AI-2 [19]. For 3 this reason, QS inhibitors belonging to the class of MTAN transition state analogues could be used 4 to block both AI-1 and AI-2 production in bacteria without interfering with host cell metabolism. 5 This approach has been explored by Gutierrez et al. [19], and Longshaw et al. [20], who found that 6 both sulphur-containing and sulphur-free MTAN transition state analogues are potent inhibitors of 7 *E. coli* QS. Recently, the Schramm laboratory patented two different types of transition state 8 analogues and a pharmaceutically acceptable carrier for treating bacterial infections [21]. These 9 developments indicate the interest that MTAN has aroused as a possible target for bacterial anti-10 infective drug design [1]. 11 It has been demonstrated that anthranilate (Figure 1, 1) is a precursor of the PQS (Figure 1, 12 3), which is a secondary metabolite maximally produced during the late stationary phase of growth. 13 An anthranilate analogue (methyl anthranilate) (Figure 1, 2) inhibits the production of PQS, with a 14 negative effect on the production of elastase by *P. aeruginosa* [22]. These data suggest that 15 pseudomonas PQS targeting may have a therapeutic value, but it must be taken into account that P. 16 aeruginosa has two other QS systems, based on N-(3-oxododecanoyl)-L-homoserine lactone (3-17 oxo-C12-HSL) (Figure 2, 4) and N-butyryl-L-homoserine lactone (C4-HSL) (Figure 2, 5), 18 respectively. In this regard, it has recently been shown that eugenol (Figure 3, 6), the major 19 constituent of clove extract, affects P. aeruginosa OS decreasing the transcriptional activation of 20 both las and pqs systems, which together regulate the expression of numerous virulence-related 21 genes [23]. Further studies are needed to assess the relative importance of each system on the 22 virulence of *P. aeruginosa*, and to examine the possibility of simultaneously turning off all three 23 systems.

24

## 25 **3. Strategies targeting QS signal**

Neutralization of QS signals has been extensively investigated [24]. Biological inactivation of AHLs
 can be achieved by enzymatic degradation or antibody-mediated inactivation of the signal molecule.
 3

4 3.1 Enzymes

QQ enzymes can be grouped in two classes: class I includes the enzymes that break the AHL
molecule, i.e. AHL-lactonase, AHL-acylase and paraoxonase, whereas class II includes enzymes
that reduce carbonyl to hydroxyl, i.e. oxidoreductases [25].

8 About 20 AHL lactonases have been discovered to date, most of which are supposedly 9 valuable for the biocontrol of plant diseases. Recently, Migiyama et al. reported the effect of the 10 AiiM lactonase on a mouse model of acute pneumonia by *P. aeruginosa* [26]. The study supports 11 the therapeutic potential of AHL lactonases in therapy, demonstrating that AiiM is a potent inhibitor 12 of *P. aeruginosa* QS and reduces bacterial virulence. The use of this molecule could represent a 13 new strategy to cure chronic pulmonary infections by multiresistant strains. However, the 14 problematic issue is the way of administration: in their experiments, the authors used lactonase-15 overexpressing plasmid vector-transformed P. aeruginosa to infect mice. This technique can be a 16 good experimental tool, but, of course, it is not of any use for therapeutic purposes. The authors 17 suggest two possible solutions to this impasse: the genetic modification of probiotics, or the local 18 administration of the purified AiiM protein. In both cases, the authors agree that further studies are 19 needed to evaluate the possibility of developing AiiM lactonase into a drug molecule. 20 One of the best studied acylases is the PvdQ protein, a member of the N-terminal 21 nucleophile hydrolase superfamily. This enzyme is produced by the P. aeruginosa PAO1 strain, 22 suggesting that it may participate to the regulation of its own QS-dependent pathogenic potential. 23 PvdQ behaved as a QQ *in vitro* in a number of phenotypic assays in which the *pvdQ* gene was 24 overexpressed in *P. aeruginosa*, and its activity was demonstrated *in vivo* in a *Caenorhabditis* 

25 *elegans* model [27]. The development of the molecule has reached the stage of the production of a

stable and inhalable powder formulation for the treatment of *P. aeruginosa* pulmonary infection
 [28].

Paraoxonases (PON) are a group of  $Ca^{2+}$ -dependent esterases with AHL lactonase activity 3 4 widely conserved in mammals, but not present in chicken and fish [29]. In humans, the family 5 comprises three members, whose genes are located on chromosome 7: PON1 and PON3, mainly 6 expressed in liver and kidney, and PON2, expressed in various tissues but not present in plasma 7 [30]. These enzymes are supposed to play an important role in the defence against the formation of 8 bacterial biofilms. The protective effect of PON1 against P. aeruginosa infection has been 9 demonstrated in an experimental model of transgenic Drosophila melanogaster expressing human 10 PON1 [31]. Recently, by in vitro and in vivo experiments Devarajian et al. demonstrated that PON2 11 plays an important role in the mouse innate immune response [32].

12 Oxidoreductases catalyze the functional inactivation of AHLs by oxidating or reducing their 13 acyl side chain. Two oxidoreductases have been isolated from cultivated bacteria, one from Bacillus 14 megaterium CYP102A1 and the other from *Rhodococcus erythropolis* W2 [33, 34]. Recently, a novel 15 oxidoreductase named BpiB09, able to inactivate the 3-oxo-C12-homoserine lactone, was obtained 16 by screening a soil metagenomic library. The expression of this molecule in *P. aeruginosa* PAO1 17 significantly reduced pyocyanin production, bacterial motility, and biofilm formation. In addition, 18 transformed bacteria resulted non pathogenic in a *C. elegans* model of infection [35]. The possibility 19 to screen metagenomic libraries opens wide horizons for the discovery of new active QQ enzymes. 20 However, their value for clinical purposes needs to be assessed. It can be expected that they will 21 find application in the treatment of skin and airway infections, and in the inhibition of bacterial 22 biofilm formation on catheters and prosthetic devices.

23

#### 24 **3.2 Antibodies**

Most AHLs exert potent biochemical effects, such as induction of apoptosis and modulation of NFKB activity, behaving as small-molecule toxins on mammalian cells. Consequently, specific

1 antibodies can achieve the dual purpose of neutralizing both the QS activity and the toxic effect. 2 The immunological approach includes the induction of antibodies by vaccination, and the administration of preformed mono- or poly-clonal antibodies.

3

4 Following the first demonstration of the efficacy of vaccination with a homoserinelactone-5 carrier protein conjugate in a mouse model of *P. aeruginosa* lung infection by Miyairi et al. [36], the 6 possibility to produce protective antibodies has been extensively investigated, especially by the 7 Janda laboratory at The Scripps Research Institute [37-39]. The attractivity of this approach is based 8 on the high evolutionary conservation of AI, their extracellular distribution, and the established 9 knowledge of antibody pharmacodynamic and kinetic properties. The chemical structure and the 10 low molecular weight make the AI poor antigens, which could be more properly defined haptens [7]. 11 The efficacy of the immunopharmacotherapeutic approach to the inhibition of QS has been demonstrated in vitro and in vivo against S. aureus and P. aeruginosa. Park et al. reported the 12 13 production of a monoclonal antibody, AP4-24H11, elicited against the autoinducing peptide AIP-4 14 produced by S. aureus RN4850. The antibody was able to inhibit QS in vitro, suppressed S. aureus 15 pathogenicity in an abscess formation mouse model, provided complete protection against a lethal 16 S. aureus challenge [38], and its structure was subsequently further investigated in detail [40]. 17 Very recently, Palliyil et al. reported the development of monoclonal antibodies against

18 homoserine lactones produced by P. aeruginosa. These antibodies were produced by using a 19 combination of sheep immunisation and phage antibody display library construction/selection, and 20 were characterized by high sensitivity (100-1000 times higher than that of any published antibodies 21 raised to the same target). Their protective effect was assessed in a slow-killing model of C. 22 elegans, and in a mouse model of *P. aeruginosa* infection [41]. These findings provide a strong 23 foundation for further investigations of the potential of AHL monoclonal antibodies in the 24 immunopharmacotherapy of antibiotic resistant strains of *P. aeruginosa*.

#### **4.** Strategies targeting autoinducer-receptor interaction and/or receptor-

#### 2 mediated signal.

The possibility to inhibit the expression of QS signal receptors by means of antisense RNAs has been demonstrated by Hirakawa et al. [42]. For a comprehensive review on the inhibition of signal detection, the reader is referred to [43]. In the following paragraphs, we will discuss some recent insights on the QS of *S. aureus* and *P. aeruginosa* that may suggest the possibility of interfering with the receptor side of the system.

8 The clarification of the QS system in Gram-positive bacteria allowed the identification of 9 molecules inhibiting different steps of the mechanism. The QS system of S. aureus, which is the 10 most studied, consists of the accessory gene regulator (agr) locus, which is composed of two 11 transcripts called RNAII and RNAIII, driven by the P2 and P3 promoters, respectively. The RNAIII 12 transcript is the effector of the agr response, which up-regulates a number of toxins and multiple 13 exoenzymes (proteases, lipases, and nucleases), and down-regulates the expression of numerous 14 surface protein genes. The RNAII transcript is an operon of four genes, agrBDCA. AgrD is the 46 15 aminoacid precursor of the secreted AI, a cyclic thiolactone peptide (AIP) of 7-9 aminoacids, which 16 is processed, cyclized and exported through the transmembrane protein AgrB. The extracellular AI 17 binds to AgrC, the transmembrane receptor, which is a histidine kinase that following the binding 18 phosphorylates the AgrA cytoplasmic response regulator, activating the P2 and P3 promoters. More 19 advanced strategies targeting S. aureus virulence gene regulation make use of inhibitors of the 20 sensor kinase AgrC, of the transcriptional activator AgrA, or of the RNAIII [44].

Due to allelic variations within the agr gene system, *S. aureus* can be subdivided into four agr specific groups, agr I–IV, each secreting a distinct AIP with different primary amino acid sequence. Most cross-group AIP-AgrC interactions are inhibitory, with AIPs activating their cognate receptors and competitively inhibiting non-cognate receptors. The competitive AIP inhibition constitutes a promising therapeutic approach for attenuating *S. aureus* infections.

1 Recently, it has been demonstrated by *in vitro* experiments that oxidized low-density lipoproteins 2 bind all four *S. aureus* agr AIPs and antagonize agr signalling by each agr allele [45]. 3 Molecules interfering with the *S. aureus* agr system have been obtained by a marine 4 *Photobacterium*. Due to the structural similarity of these molecules, called solonamides, to the agr 5 AI, the proposed mechanism of action is competitive inhibition [46]. The solonamides are the first 6 reported natural antagonists with a structure resembling that of native S. aureus AI. Another natural 7 product, hamamelitannin, obtained from Hamamelis virginiana (witch hazel), was identified as an 8 inhibitor of RNAIII and  $\delta$ -hemolysin production in *S. aureus* by Kiran et al [47]. 9 Two of the three *P. aeruginosa* QS systems consist of the *lasRI* and *rhlRI* genes, where 10 LasR and RhlR belong to the LuxR family of transcriptional regulators that specifically bind to N-11 (3-oxododecanoylhomoserine lactone (3-oxo-C12-HSL) and N-butanoylhomoserine lactone (C4-12 HSL), respectively [48]. The las and rhl systems regulate over 10% of the P. aeruginosa genome 13 and are hierarchically organized. The 3-oxo-C12-HSL signal is the key factor that exerts 14 transcriptional control over *lasI*, *rhlR*, *rhlI* and the genes of the third system, the PQS system. 15 Despite the positive auto-regulation of *lasI*, levels of 3-oxo-C12-HSL reach a steady level long 16 before cultures reach the stationary phase, due to the intervention of the transcriptional repressor 17 RsaL, whose activity was originally reported in 1999 [49; 50]. It is now known that the QS systems 18 are controlled by a complex regulatory network at both the transcriptional and post-transcriptional 19 level, and RsaL is just one of over 15 QS regulators that have been identified so far [51]. RsaL is a 20 DNA binding protein that competes against LasR, directly repressing both lasI and rsaL 21 transcription by binding to the bi-directional *rsaL-lasI* promoter. In addition, Rampioni et al. 22 observed that, independently of its effects on the production of QS signal molecules, RsaL directly 23 represses 120 genes, many of which code for virulence and antibiotic resistance, and activates 10 24 genes (undefined) [50].

A recently identified novel LasR-specific antiactivator is QslA, which binds to LasR and
prevents it from binding to its target promoter [52]. The investigation of bacterial autoregulation of

QS may open new horizons for the therapeutic exploitation of QS control. Studying the interaction
between QslA and the LasR ligand-binding domain, Fan et al. elucidated the crystal structure of
QslA and demonstrated that its mechanism of action consists in the disruption of LasR dimers by
occupying the LasR dimerization interface, which is a previously unknown mechanism of QS
inactivation [53].

6 Other QS negative regulators produced by *P. aeruginosa* are QscR, which is an "orphan" 7 receptor activated by the 3-oxo-C12-HSL signal [54], and QteE, which reduces LasR protein 8 stability without affecting LasR transcription or translation [55]. It has been shown that negative 9 regulation of QS by QteE and QscR has a major impact on the absolute and relative fitness of P. 10 aeruginosa [56]. The possibility to interact with these regulator systems for therapeutic purposes is 11 currently being investigated. In a recent paper, Weng et al. report on a novel QS inhibitor, called 12 C2, which causes a 375.4% upregulation of the *qscR* gene, resulting in the attenuation of elastase 13 and protease activity, swarming motility and biofilm formation in the *P. aeruginosa* strain 14 PAO1[57].

15 The *P. aeruginosa* PQS system may be inhibited by anthranilic acid halogenated analogues, 16 whose development, however, has not been pursued [58; 59]. However, novel insights on the 17 structure of PqsE, a product of the pqs operon, could lay the foundations for the computer-aided 18 design of PqsE inhibitors [60]. Antagonists of the PqsR were reported for the first time in 2012 [61], 19 and a peculiar mechanism of functional inversion of a PqsR antagonist has been recently described 20 by Lu et al. [59]. These authors observed that compound 1 (Figure 4, 7), which behaves as a pure 21 PqsR antagonist in an *E. coli* reporter gene assay, displayed a dose-dependent agonistic activity 22 when tested in a *P. aeruginosa* functional assay. This unusual behaviour can be explained by the 23 transformation of compound 1 into compound 2 (Figure 4, 8). A product of the pqs operon, the 24 enzyme PqsH, which is present in *P. aeruginosa* only, carries out the transformation. The analysis 25 of the molecular structure of the antagonist and of the chemical modification that turns it into an 26 agonist allowed the synthesis of a small library of substituted compounds, among which compound

3 (Figure 4, 9) showed high potency in the *E. coli* reporter gene assay, but retained its antagonistic
activity in *P. aeruginosa* without displaying any agonistic activity [59]. In a recent paper, Ilangovan
et al. report the determination of the PqsR crystal structure, which allowed the visualization of the
shape of the PQS-binding site. This work in conjunction with the chemical synthesis of PQS
analogues resulted in the discovery of potent quinazolinone inhibitors of PqsR. These novel insights
into the structure of PqsR provide further opportunities for targeting *P. aeruginosa* QS [62].

7 Farnesol (Figure 5, 10) is a sesquiterpene alcohol produced by *Candida albicans* and 8 present in the essential oils of citrus fruits. Beyond its activity as fungal QS molecule [63], it affects 9 PQS production by inhibiting the transcription of the pqs operon. According to the proposed model, 10 farnesol stimulates PqsR interaction with a *pqsA* promoter fragment in the region of the PqsR 11 binding site, causing a non-productive interaction with the pqsA promoter that does not lead to 12 transcriptional activation of the pqsA-E operon [64]. The result is the impairment of production of 13 pyocyanin, a pseudomonas QS-controlled virulence factor. Farnesol is an interesting molecule, 14 because it is devoid of toxic effects, non mutagenic, and affects biofilm formation by C. albicans, 15 Streptococcus mutans, and Staphylococcus aureus [65]. For an interesting discussion on the 16 interactions between C. albicans and P. aeruginosa, the reader is referred to [66]. However, farnesol 17 use is hampered by its low solubility in water. In a very recent paper, Bhattacharyya et al. describe a 18 new efficient strategy for the local simultaneous delivery of farnesol and vancomycin in the 19 treatment of multi-resistant S. aureus (MRSA) infections [67]. Following the observation that 20 sufficient quantities of vancomycin and farnesol can be incorporated into sol-gel silica applied as 21 thin films on an implant surface, they demonstrated a potent adjuvant effect of farnesol on 22 vancomycin in the inhibition of in vitro MRSA experimental infection. Although in this case the 23 mechanism of action proposed for farnesol does not involve the inhibition of QS, the technique is 24 attractive and could be used to prevent or limit the formation of bacterial biofilms on the surfaces of 25 catheter and joint implants.

#### **5. Modern approaches based on innovative technologies.**

Innovative technologies are being tested in the field of QS control, opening up possibilities that
were unthinkable a few years ago.

4

## 5 **5.1 Chemical methods.**

6 Following the original discovery of the anti-QS activity of natural halogenated furanones 7 (fimbrolides) isolated from *Delisea pulchra*, many analogues have been and are produced by 8 chemical synthesis [68]. Traditional synthetic approaches usually yield small panels of synthetic AI 9 analogues, some of which have proved potent QS modulators. A significant improvement of the QS 10 disrupting activity has been recently obtained by the combination of fimbrolides with NO-donor 11 molecules in hybrid compounds (Figure 6, 11, 12) [69]. The combination, which represents the first 12 example of dual-acting molecules with high activity against QS and biofilm formation, is worth of 13 further development. The chemical process typically requires different steps such as individual 14 isolation, purification, and reaction optimization, which for large sets of analogues can rapidly 15 become cumbersome. On the other hand, it usually allows the establishment of correlations between 16 structure and activity. In this regard, Tsuchikama et al. recently reported the synthesis of a panel of analogues of (4S)-4,5-dihydroxy-2,3-pentanedione (DPD), called C4-alkoxy-HPDs, which are more 17 18 potent than natural AI-2 molecules. These findings highlight how manipulation of the DPD scaffold 19 can provide valuable tools for in-depth studies of the ligand-receptor interactions involved in AI-2-20 mediated QS, and lay the foundation for future chemical structure-based studies aimed at 21 identifying and developing antagonistic analogues [70]. For a comprehensive review focused on 22 chemical methods including combinatorial synthesis, affinity chromatography, and electrochemical 23 sensing of QS signals, the reader is referred to [71].

24

#### 25 **5.2** Computer-aided investigations.

1 Among modern approaches, high throughput screening maintains its validity yielding hit molecules 2 often based on new molecular scaffolds [72, 73], but in silico investigations are considered more convenient in terms of cost, speed of execution, and potency, being able to screen up to millions of 3 4 compounds. Examples of results obtained by the use of structure-based virtual screening are the 5 recent identification of inhibitors of *P. aeruginosa* QS from a natural-derivative database [74], and 6 from the ChemBridge library [75]. The *in silico* approach has been used not only for screening 7 purposes, but also for studying QS dynamics [76], and to identify bacterial species that contained 8 both QS and aromatics degrading systems, in order to acquire information for developing novel bio-9 processing techniques [77].

10 An innovative approach is the production by Schaadt et al. of a multi-level logical model 11 based on computational analysis of a combined regulatory and metabolic network for the three P. 12 *aeruginosa* QS systems. The model may be used to analyse how enzyme inhibitors and receptor 13 antagonists affect the formation of AI and virulence factors. In addition, it allows almost 14 quantitative predictions about the effect of inhibitors of AI biosynthesis and antagonists of their 15 corresponding receptor, and the investigation of the effect of reviewed network topologies. In the 16 words of the authors, this approach can serve as a basis for further integrating the effect of random 17 mutations in various parts of the network [78].

18 In order to get a better insight into the regulation of S. aureus QS system, Audretsch et al 19 used an innovative approach to model the whole system [79]. Instead of creating a mathematical 20 model, these authors used a Boolean network of nodes centred on the agr locus of S. aureus. The in 21 silico node activation patterns were compared with gene activation patterns obtained from 22 microarray, northern blot, and transcriptome data. The network can easily be manipulated and 23 studied, and has two different steady states: one representing an invasive, toxic phenotype, and the 24 other representing a biofilm producing phenotype. By manipulating the nodes of the simulated 25 network, the model may be used to test theories about mutant strains and to predict the effects of 26 QS inhibitors.

1 Sahner et al. have recently devised a novel approach based on the combination of *in silico* and 2 biophysical methods (surface plasmon resonance, isothermal titration calorimetry, saturation 3 transfer difference, and nuclear magnetic resonance) for the development of P. aeruginosa QS 4 inhibitors. They demonstrated that the use of the two methods in combination represents a powerful 5 complement to co-crystallography, and allows the rapid and efficient development of inhibitors. By 6 using this combination technique, these authors identified an irreversible inhibitor that covalently 7 binds to the active site of PqsD, which mediates the formation of the precursor of the Pseudomonas 8 quinolone signal (PQS) [80].

9

#### 10 **5.3 Nano-and micro-technologies.**

11 Nanotechnologies can be used to direct bacterial communication, by means of "nanofactory"-loaded 12 biopolymer capsules placed at the midst of bacterial populations [81, 82]. It can be expected that this 13 type of instruments, once developed, could be used to assist or replace the activity of the classic 14 antibiotics, and to extend our knowledge of the QS systems and modulate them, so as to direct the 15 microbiome to operate in a host-friendly manner. The possibility to modulate QS has also been 16 demonstrated by means of titanium dioxide bead coated with AHL [83]. The implications of these 17 experiments include the production of medical devices (catheters, joint and cardiac prosthetic 18 devices) with coated surfaces able to interrupt QS signalling and to avoid biofilm formation.

19 The application of novel technologies derived from quite a different field like 3D printing to 20 study bacterial interactions opens new horizons. By means of a microscopic three-dimensional (3D) 21 printing strategy that enables multiple populations of bacteria to be organized within essentially any 22 3D geometry, including adjacent, nested, and free-floating colonies, it has been possible to obtain 23 the rapid growth of fully enclosed cellular populations that release a number of biologically active 24 molecules, including polypeptides, antibiotics, and QS signals. Using this approach, Connell et al. 25 showed that picoliter-volume aggregates of S. aureus can display substantial resistance to  $\beta$ -lactam 26 antibiotics by enclosure within a shell composed of P. aeruginosa [84]. The technique takes

advantage of thermally set gelatine mixtures as a reagent for micro-3D printing, which provides the
ability to print enclosures around any bacterial cell of interest suspended within the hydrogel
matrix. In the words of the authors, this manufacturing strategy provides a versatile base for
exploring the mechanisms involved in cell communication, which allow bacterial adaptation to the
natural environment by means of social behaviour. This technology could perhaps be exploited at
its best if coupled with the use of bacteria programmed to sense and destroy highly pathogenic
species.

8

## 9 5.4 Bacterial bio-engeneering

10 In a recent paper, Gupta et al. describe the engineering of sentinel E. coli cells, capable of 11 specifically detecting the QS AI molecule 3OC<sub>12</sub>HSL, produced by *P. aeruginosa*, and then to 12 synthesize and secrete the chimeric bacteriocin CoPy, specific for PAO1 P. aeruginosa [85]. The 13 engineering of *E. coli* consists of three interconnected modules: the detection module, which is 14 activated by the PAO1 QS signal; the destruction module, which produces the toxin; and the 15 secretion module that allows the secretion of CoPy. The effectiveness of the system was assessed by 16 experiments in which the two bacterial species were co-cultured on semi-solid agar plates. The 17 system has two advantages: first, it is possible to change the type of toxin produced; second, in the 18 absence of the pathogen, the system remains in stand-by, but the production starts immediately after 19 the detection of the pathogen.

Bacterial QS is also involved in extracellular cell death and in the action of antibiotics, through its interference in a system also called "addiction module", which can be found in many different Gram-positive and -negative bacteria. The system, which has mainly been studied in *E. coli*, consists of a pair of genes whose products are a stable toxin and an unstable antitoxin, which prevents the lethal action of the toxin [86]. In this situation, bacterial survival depends upon the continuous production of the short-lived antitoxin, to which the cell is "addicted". It has been shown that in *E. coli* the effect of some antibiotics such as rifampicin, chloramphenicol,

1 spectinomycin, and nalidixic acid is due to the inhibition of the antitoxin production, and that this 2 action is dependent upon the presence in the culture of high concentrations of a QS mediator called 3 "extracellular death factor" (EDF) [87, 88]. The E.coli EDF is a linear pentapeptide with the amino 4 acid sequence Asn-Asn-Trp-Asn-Asn, which is able to induce programmed cell death also in 5 conditions of cell stress, like amino acid starvation or DNA damage. Recently, it has been shown 6 that one hexapeptide produced by *Bacillus subtilis* and three peptides, one nonapeptide and two 7 hexadecapeptides produced by *P. aeruginosa*, have the same effect of the *E. coli* EDF on *E. coli* 8 cultures [89]. These findings establish the existence of a growing family of EDFs that behave as AI 9 and exert both intra- and inter-specific activity. These molecules may provide a basis for the 10 development of a new class of antibacterials, which could synergize with antibiotics.

11

## 12 **6. Expert opinion**

13 Despite the huge amount of molecules so far discovered that inhibit the QS mechanisms, the 14 potential of QS inhibition as a future treatment strategy has yet to be fully assessed. In addition, 15 there are obstacles, some obvious, others in the process of being recognized, which hinder the 16 transformation of QS inhibitors into real drugs. The most obvious roadblock, which QS inhibitors 17 share with antibiotics, is the lack of interest of pharmaceutical companies to bear the costs related to 18 the development of hits into drugs that are inevitably bound to have a very limited market. This 19 issue is critical to the future of the treatment of infectious diseases in this so-called post-antibiotic 20 era. In accordance with what has already been proposed for the development of new antibiotics and 21 drugs for neglected diseases, probably the best way to overcome the impasse is the union of skills 22 and funds from private companies, academia and public institutions. In the short term, the 23 knowledge gained so far, especially that concerning plant-derived inhibitors, could perhaps be used 24 in the field of dietary advice and food supplements. Since the pioneering work of Givskov et al. [90] 25 reporting the inhibitory effect of brominated furanones from the Australian macroalga D. pulchra 26 on Serratia liquefaciens QS-mediated swarming, many plant-derived inhibitors have been

1 identified. For a comprehensive review, the reader is referred to [91]. To improve the characteristics 2 of natural derivatives, a number of synthetic molecules have also been developed, and more than 40 3 have been patented [92, 93, 15, 1]. However, we are still far from having a licensed drug available on 4 the market, if we exclude some antibiotics, such as azithromycin, ceftazidime and ciprofloxacin, 5 which at subinhibitory concentrations block the expression of QS-regulated virulence factors in P. 6 *aeruginosa* [94]. In this regard, we must stress the importance of distinguishing between the direct 7 toxic effect of a substance and its inhibiting activity on QS. To this aim, it is essential to verify the 8 effect of QS inhibitors on bacterial viability by using sensitive methods and appropriate controls. In 9 the case of pyrogallol, it has recently been shown that its previously highlighted QS-disrupting 10 activity is a side effect of the peroxide production induced by this compound rather than true QS 11 inhibition [95]. According to Defoirdt et al., many of the results obtained in experiments that make 12 use of QS signal molecule reporter strains may be questioned, due to the possibility that the tested 13 compounds may be toxic to the reporter strains [96]. These authors suggest that toxicity tests more 14 stringent than those implemented so far should be used on molecules proposed as QS inhibitors. 15 Another emerging problem is the possibility that QS inhibitors can select more virulent 16 strains. Resistance to QS inhibitors has been studied by means of mathematical models [97-101], and 17 of in vitro experiments [102-104]. Bacterial resistance to brominated furanones is mediated by 18 increased efflux activities. As this kind of resistance overlaps with resistance to antibiotics, the 19 treatments with antibiotics endowed with a strong selective pressure may result in improved 20 resistance to antivirulence compounds (and perhaps vice versa). The finding that some strains 21 isolated from cystic fibrosis patients are resistant to QS inhibitors, corroborates this hypothesis. 22 Additional clinical evidence of the ability of bacteria to evolve resistance to QQ compounds was 23 provided by studying the resistance of *P. aeruginosa* isolated from urine, blood, and catheter tip 24 specimens obtained from Mexican children [104].

Other *in vivo* and *in vitro* results reported by Kohler et al. demonstrate that QS inhibition
interferes with natural selection towards reduced virulence, and therefore may increase the

prevalence of more virulent genotypes [105]. Some concerns about the widespread use of QQ
compounds as part of an antibacterial therapy have been raised by the possibility that targeting a
specific bacterial group may predispose the patient to infection by other groups. For example, in *S. aureus*, but not in *S. epidermidis*, biofilm production is under the agr system control. In this
situation, agr inhibition would favour *S. epidermidis*, suggesting that agr antagonists would not be
indicated for the control of staphylococcal biofilm infections [106].

Interestingly, resistance to QQ compounds may be enhanced by the complex interplay of
bacteria with bacteriophages, and it seems that QS constitutes a significant, but so far overlooked,
determinant of bacterial susceptibility to phage attack [107]

10 The AHLs often exert toxin-like effects on mammalian cells. Hence, neutralizing antibodies 11 would achieve the double goal of disrupting bacterial QS and protecting the host from toxicity. As 12 discussed above, the disrupting efficacy of antibodies was demonstrated both in *P. aeruginosa* and 13 S. aureus QS systems. Based on these results, and taking into account the evolutionary high 14 conservation and extracellular distribution of AI, QS can be considered a good target for active and 15 passive immunotherapy approaches [39]. However, here again some caution is necessary. In a recent 16 paper, Michael-Gayego et al. report that SilCR, a QS peptidic AI produced by Group A (GAS) and 17 G streptococci, effectively decreases bacterial virulence in vivo. The effect was also observed in 18 SilCR-vaccinated mice, which developed a more severe disease than non-vaccinated ones [108]. The 19 paradox is that SilCR is the product of a gene belonging to the streptococcal invasion locus, sil, 20 which is a virulence factor involved in GAS spreading into deeper tissues. This is perhaps the first 21 demonstration that antibodies directed against a bacterial component increase the severity of the 22 disease. On the other hand, QS activation can be used to improve the antigenicity and the efficacy 23 of an *E. coli* bacterin preparation to be used for vaccination purposes, as reported by Sturbelle et al. 24 [109]. These examples underline both the potentials of QS exploitation for therapeutic purposes and 25 the need to clarify all aspects of the involved mechanisms.

1 In conclusion, QS inhibition can be considered a promising strategy against infections, especially 2 those associated with biofilm formation, and the discovery of new inhibitory molecules is rapidly 3 progressing. However, to identify a new molecule as a pure QS inhibitor, it is necessary to develop 4 more standardized methods of evaluation, with particular attention to rule out direct toxicity that 5 may mimic a QQ effect. Moreover, concerns about the possible emergence of resistance and 6 adverse effects resulting from the alteration of the microbiome must be properly addressed in-depth. 7 In our view, the current challenge in the field of QS research is not so much to find new inhibitory 8 molecules, but to refine and deepen our knowledge on the QS-based interactions between different 9 bacterial species and between bacteria and the host. Important issues that need further investigation 10 are the impact of QS on the social evolution of bacteria, as well as the impact of bacterial social-11 driven evolution on the infection of the host. Studies performed in mice [110] and humans [111] 12 demonstrated that cultures containing mixtures consisting of QS cooperators, i.e. autoinducer-13 producing cells, and cheats, i.e. cells that do not produce autoinducers but benefit of cooperator-14 produced autoinducers, are less virulent than cultures containing pure cooperators, suggesting that 15 asocial cheats could be exploited to exert a potential therapeutic role [112]. An interesting discussion 16 on bacterial cooperation and its possible relevance in the clinical field can be found at [113]. 17 The last, but not least important aspect that must be taken into account is that so far the 18 transformation of laboratory results into viable drugs is almost non-existent [114]. This situation is 19 not expected to improve in the near future, because, as is the case for antibiotics, market forces are 20 insufficient to drive the development of antibacterial drugs based on new scaffolds. The need for 21 long-term, huge investments, and the prospect to see the novel drug indications limited to small 22 number of cases, made many large pharmaceutical companies to quit antibiotic discovery for more 23 profitable therapeutics [115]. The political, medical and public concern about the rising innovation 24 gap in 2009 prompted the U.S. and European Community presidencies to establish a Transatlantic 25 Task Force to address antimicrobial resistance, and the Infectious Diseases Society of America 26 called for a global commitment to develop 10 novel antimicrobials by 2020 [116]. We can envisage

1 that in the near future the synergy between new technical developments and public-private 2 industrial partnerships will bring into being a new harvest of badly needed novel antimicrobials, including antibiotics and antivirulence agents. 3 4 5 Article highlights: 6 7 The increased frequency of infections caused by multiresistant strains indicates the need for new 8 antibacterial drugs with a low selection pressure. The quenching of bacterial virulence by disrupting 9 QS-dependent intercellular communication is one of the best options to achieve this goal. 10 11 Many natural and synthetic QS inhibitors have been identified by means of modern methods that 12 include high throughput screenings and *in silico* structure-function analysis. About 40 molecules 13 with QS inhibitory activity have been patented. 14 15 The possibility to interrupt QS by means of active or passive immunologic interventions is being 16 evaluated and appears promising. 17 18 The most recent approaches in the field of QS research span from mathematical modelling of the 19 QS systems to the combination of *in silico* and biophysical methods, to the creation of hybrid 20 compounds by fusing fimbrolides with NO-donor molecules, and to the exploitation of the 3D 21 printing technology to study short-distance interspecific bacterial interactions. 22 23 The impact of the social behaviour of bacteria on the microbiome and on the establishment of 24 infections needs further investigation. Some concerns are emerging following the someway 25 unexpected discovery that QS inhibitors can exert a selective pressure favouring the establishment

of more virulent strains. This aspect should be thoroughly investigated before any QS inhibitor is
 licensed and put on the market.

3

	4	None of the inhibitors	identified to date	have reached the	market yet. T	This can be attributed in par	t
--	---	------------------------	--------------------	------------------	---------------	-------------------------------	---

- 5 to the necessity of further basic research and development, and in part to the lack of interest of
- 6 pharmaceutical companies for the development of drugs with limited market prospects.

7

2 The authors declare no conflict of interest.

3

# 4 **Bibliography**

- 5 Papers of special note have been highlighted as either of interest  $(\bullet)$  or of considerable interest  $(\bullet\bullet)$
- 6 to readers.
- Jiang T, Minyong L. Quorum sensing inhibitors: a patent review. Exp Opin Ther Pat 2013;23:867 94
- 9 2. Fuqua WC, Winans SC, Greenberg EP. Quorum sensing in bacteria: the LuxR-LuxI family of cell
- 10 density-responsive transcriptional regulators. J Bacteriol 1994;176:269-75
- Davies D. Understanding biofilm resistance to antibacterial agents. Nat Rev Drug Discov 2003;
   2:114-22
- 13 4. Blackledge MS, Worthington RJ, Melander C. Biologically inspired strategies for combating
- 14 bacterial biofilms. Curr Opin Pharmacol 2013;13:699-706
- 15 5. Tomasz A, Hotchkiss RD. Regulation of the transformability of pneumococcal cultures by
- 16 macromolecular cell products. Proc Natl Acad Sci USA 1964;51:480-7
- Nealson KH, Platt T, Hastings JVV. Cellular control of the synthesis and activity of the bacterial
   luminescent system. J Bacteriol 1970;104:313-22
- 19 7. Amara N, Krom BP, Kaufmann GF, et al. Macromolecular inhibition of quorum sensing:
- 20 enzymes, antibodies, and beyond. Chem Rev 2011;111:195-208
- 8. Williams P. Quorum sensing: an emerging target for antibacterial chemotherapy. Exp Opin Ther
   Targets 2002;6:257-74
- 23 9. Zhang LH, Dong YH. Quorum sensing and signal interference: diverse implications. Mol
- 24 Microbiol 2004;53:1563-71
- 25 10. Kalia VC. Quorum sensing inhibitors: an overview. Biotechnol Adv 2013;31:224-45

1	• An updated and very comprehensive overview on natural and synthetic QS inhibitors and their
2	biotechnological applications.

- 3 11. Hentzer M, Givskov M. Pharmacological inhibition of quorum sensing for the treatment of 4 chronic bacterial infections. J Clin Invest 2003;112:1300-7 5 12. Jacobsen TH, van Gennip M, Phipps RK, et al. Ajoene, a sulfur-rich molecule from garlic, inhibits 6 genes controlled by Quorum Sensing. Antimicrob Agents Chemother 2012;56:2314-25 7 13. Rasmussen TB, Givskov M. Quorum-sensing inhibitors as anti-pathogenic drugs. Intern J Med 8 Microbiol 2006;296:149-61 9 14. Kaufmann GF, Park J, Janda KD. Bacterial quorum sensing: a new target for anti-infective 10 immunotherapy. Exp Opin Biol Ther 2008;8:719-24 11 15. Bhardwaj AK, Vinothkumar K, Rajpara N. Bacterial quorum sensing inhibitors: attractive 12 alternatives for control of infectious pathogens showing multiple drug resistance. Recent Pat 13 Antiinfect Drug Discov 2013;8:68-83 14 •• A clear exposition of the different QS systems and of their inhibitors, with a detailed analysis of 15 the most relevant patents. 16 16. Chung J, Goo E, Yu S, et al. Small-molecule inhibitor binding to an N-acyl-homoserine lactone 17 synthase. Proc Natl Acad Sci USA 2011;108:12089-94 18 17. Chan YY, Chua KL. The Burkholderia pseudomallei BpeAB-OprB efflux pump: expression and 19 impact on quorum sensing and virulence. J Bacteriol 2005;187:4707-19 20 18. Li X, Chu S, Feher VA, et al. Structure-based design, synthesis, and antimicrobial activity of 21 indazole-derived SAH/MTA nucleosidase inhibitors. J Med Chem. 2003;46:5663-73 22 19. Gutierrez JA, Crowder T, Rinaldo-Matthis A, et al. Transition state analogs of 5'-23 methylthioadenosine nucleosidase disrupt quorum sensing. Nat Chem Biol 2009;5:251-7 24 20. Longshaw AI, Adanitsch F, Gutierrez JA, et al. Design and synthesis of potent "sulfur-free" 25 transition state analogue inhibitors of 5'-methylthioadenosine nucleosidase and 5'-
- 26 methylthioadenosine phosphorylase. J Med Chem 2010;53:6730-46

- Schramm VL. Methods and compositions for treating bacterial infections by inhibiting quorum
   sensing. US20110190265; 2011
- 22. Calfee MW, Coleman JP, Pesci EC. Interference with Pseudomonas quinolone signal synthesis
  inhibits virulence factor expression by *Pseudomonas aeruginosa*. Proc Natl Acad Sci USA
  2001; 98:11633-7
- 6 23. Zhou L, Zheng H, Tang Y, et al. Eugenol inhibits quorum sensing at sub-inhibitory
  7 concentrations. Biotechnol Lett 2013;35:631-7
- 8 24. Kalia VC, Purohit HJ. Quenching the quorum sensing system: potential antibacterial drug
  9 targets. Crit Rev Microbiol 2011;37:121-40
- 10 25. Chen F, Gao Y, Chen X, Yu Z, Li X. Quorum quenching enzymes and their application in
- 11 degrading signal molecules to block quorum sensing-dependent infection. Int J Mol Sci 2013;14:
  12 17477-500
- 13 26. Migiyama Y, Kaneko Y, Yanagihara K, et al. Efficacy of AiiM, an N-acylhomoserine lactonase,
- 14 against *Pseudomonas aeruginosa* in a mouse model of acute pneumonia. Antimicrob Agents
- 15 Chemother 2013;57:3653-8
- 16 27. Papaioannou E, Wahjudi M, Nadal-Jimenez P, et al. Quorum-Quenching acylase reduces the
- 17 virulence of *Pseudomonas aeruginosa* in a *Caenorhabditis elegans* infection model. Antimicrob
- 18 Agents Chemother 2009;53:4891-7
- 19 28. Wahjudi M, Murugappan S, van Merkerk R, et al. Development of a dry, stable and inhalable
- 20 acyl-homoserine-lactone-acylase powder formulation for the treatment of pulmonary
- 21 *Pseudomonas aeruginosa* infections. Europ J Pharmac Sciences 2013;48:637–43
- 22 29. Yang F, Wang LH, Wang J, et al. Quorum quenching enzyme activity is widely conserved in the
   23 sera of mammalian species. FEBS Lett 2005;579:3713-7
- 24 30. Camps J, Pujol I, Ballester F, et al. Paraoxonases as potential antibiofilm agents: their relationship
- 25 with Quorum-Sensing signals in Gram-Negative bacteria. Antimicrob Agents Chemother 2011;
- 26 55:1325-31

- Stoltz DA, Zabner J. Paraoxonase 1, Quorum Sensing, and *P. aeruginosa* infection: a
   novel model. Adv Exp Med Biol 2010;660:183-93
- 3 32. Devarajan A, Bourquard N, Grijalva VR, et al. Role of PON2 in innate immune response in an
  acute infection. Mol Genet Metab 2013;110:362-70
- 5 33. Chowdhary PK, Keshavan N, Nguyen HQ, et al. *Bacillus megaterium* CYP102A1 oxidation of
  6 acyl homoserine lactones and acyl homoserines. Biochemistry 2007;46:14429-37
- 34. Uroz S, Chhabra SR, Camara M, et al. N-Acylhomoserine lactone quorum-sensing molecules are
  modified and degraded by *Rhodococcus erythropolis* W2 by both amidolytic and novel
- 9 oxidoreductase activities. Microbiology 2005;151:3313-22
- 10 35. Bijtenhoorn P, Mayerhofer H, Muller-Dieckmann J, et al. A novel metagenomic short-chain
- 11 dehydrogenase/reductase attenuates *Pseudomonas aeruginosa* biofilm formation and virulence on

12 *Caenorhabditis elegans*. PLoS ONE 2011;6:e26278

- 13 36. Miyairi S, Tateda K, Fuse ET, et al. Immunization with 3-oxododecanoyl-L-homoserine lactone-
- protein conjugate protects mice from lethal *Pseudomonas aeruginosa* lung infection. J Med
   Microbiol 2006;55:1381-7
- 37. Debler EW, Kaufmann GF, Kirchdoerfer RN, et al. Crystal structures of a quorum-quenching
  antibody. J Mol Biol 2007;368:1392-402
- 38. Park J, Jagasia R, Kaufmann GF, et al. Infection control by antibody disruption of bacterial
  quorum sensing signaling. Chem Biol 2007;14:1119-27
- 20 39. Kaufmann GF, Park J, Mee JM, et al. The quorum quenching antibody RS2-1G9 protects
- 21 macrophages from the cytotoxic effects of the *Pseudomonas aeruginosa* quorum sensing
- signalling molecule *N*-3-oxo-dodecanoyl-homoserine lactone. Mol Immunol 2008;45:2710-4
- 40. Kirchdoerfer RN, Garner AL, Flack CE, et al. Structural basis for ligand recognition and
- discrimination of a quorum-quenching antibody. J Biol Chem 2011;286:17351-8

1	41. Palliyil S, Downhama C, Broadbent I, et al. High sensitivity monoclonal antibodies specific for
2	homoserine lactones protect mice from lethal Pseudomonas aeruginosa infections. Appl Environ
3	Microbiol 2014;80:462-9
4	42. Hirakawa H, Harwood CS, Pechter KB, et al. Antisense RNA that affects Rhodopseudomonas
5	palustris quorum-sensing signal receptor expression. Proc Natl Acad Sci USA 2012;109:12141-6
6	43. LaSarre B, Federle MJ. Exploiting quorum sensing to confuse bacterial pathogens. Microbiol Mol
7	Biol Rev 2013;77:73-111
8	44. Gordon CP, Williams P, Chan WC. Attenuating Staphylococcus aureus virulence gene regulation:
9	a medicinal chemistry perspective. J Med Chem 2013;56:1389-404
10	45. Hall PR, Elmore BO, Spang CH, et al. Nox2 modification of LDL is essential for optimal
11	apolipoprotein B-mediated control of agr Type III Staphylococcus aureus Quorum-sensing. PLOS
12	Pathogens 2013;9:e1003166
13	46. Mansson M, Nielsen A, Kjærulff L, et al. Inhibition of virulence gene expression in
14	Staphylococcus aureus by novel depsipeptides from a marine Photobacterium. Mar Drugs
15	2011;9:2537-52
16	47. Kiran MD, Adikesavan NV, Cirioni O, et al. Discovery of a quorum-sensing inhibitor of drug-
17	resistant staphylococcal infections by structure-based virtual screening. Mol Pharmacol
18	2008;73:1578-86
19	48. Williams P, Cámara M. Quorum sensing and environmental adaptation in Pseudomonas
20	aeruginosa: a tale of regulatory networks and multifunctional signal molecules. Curr Opin
21	Microbiol 2009;12:182-91
22	49. de Kievit T, Seed PC, Nezezon J, et al. RsaL, a novel repressor of virulence gene expression in
23	Pseudomonas aeruginosa. J Bacteriol 1999;181:2175-84
24	50. Rampioni G, Schuster M, Greenberg EP, et al. RsaL provides quorum sensing homeostasis and
25	functions as a global regulator of gene expression in Pseudomonas aeruginosa. Mol Microbiol
26	2007;66:1557-65

1	51. Liang H, Duan J, Sibley CD, et al. Identification of mutants with altered phenazine production
2	in Pseudomonas aeruginosa. J Med Microbiol 2011;60:22-34
3	52. Seet Q, Zhang LH. Anti-activator QslA defines the quorum sensing threshold and response in
4	Pseudomonas aeruginosa. Mol Microbiol 2011;80:951-65
5	53. Fan H, Dong Y, Wu D, et al. QsIA disrupts LasR dimerization in antiactivation of bacterial
6	quorum sensing. Proc Natl Acad Sci U S A 2013;110:20765-70
7	54. Chugani SA, Whiteley M, Lee KM, et al. QscR, a modulator of quorum-sensing signal
8	synthesis and virulence in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A 2001;98:2752-7
9	55. Siehnel R, Traxler B, An DD, et al. A unique regulator controls the activation threshold of
10	quorum-regulated genes in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A
11	2010;107:7916-21
12	56. Gupta R, Schuster M. Negative regulation of bacterial quorum sensing tunes public goods
13	cooperation. ISME J 2013;7:2159-68
14	57. Weng LX, Yang YX, Zhang YQ, et al. A new synthetic ligand that activates QscR and blocks
15	antibiotic-tolerant biofilm formation in Pseudomonas aeruginosa. Appl Microbiol Biotechnol
16	2013;Dec 11. PMID:24327212 [Epub ahead of print]
17	58. Lesic B, Lépine F, Déziel E, et al. Inhibitors of pathogen intercellular signals as selective anti-
18	infective compounds. PLoS Pathog 2007;3:1229-39
19	59. Lu C, Maurer CK, Kirsch B, et al. Overcoming the Unexpected Functional Inversion of a PqsR
20	Antagonist in Pseudomonas aeruginosa: An In Vivo Potent Antivirulence Agent Targeting pqs
21	Quorum Sensing. Angew Chem Int Ed Engl 2014;53:1109-12
22	60. Folch B, Déziel E, Doucet N. Systematic mutational analysis of the putative hydrolase PqsE:
23	toward a deeper molecular understanding of virulence acquisition in Pseudomonas aeruginosa.
24	PLoS One 2013;8:e73727.

1	61. Lu C, Kirsch B, Zimmer C, de Jong JC, et al. Discovery of antagonists of PqsR, a key player in
2	2-alkyl-4-quinolone-dependent quorum sensing in Pseudomonas aeruginosa. Chem Biol
3	2012;19:381-90
4	62. Ilangovan A, Fletcher M, Rampioni G, et al. Structural basis for native agonist and synthetic
5	inhibitor recognition by the Pseudomonas aeruginosa quorum sensing regulator PqsR (MvfR).
6	PLoS Pathog 2013;9:e1003508.
7	63. Décanis N, Tazi N, Correia A, et al. Farnesol, a fungal quorum-sensing molecule triggers
8	Candida albicans morphological changes by downregulating the expression of different
9	secreted aspartyl proteinase genes. Open Microbiol J 2011;5:119-26.
10	64. Cugini C, Calfee MW, Farrow JM 3rd, et al. Farnesol, a common sesquiterpene, inhibits PQS
11	production in Pseudomonas aeruginosa. Mol Microbiol 2007;65:896-906
12	65. Jabra-Rizk MA, Meiller TF, James CE, et al. Effect of farnesol on Staphylococcus aureus biofilm
13	formation and antimicrobial susceptibility. Antimicrob Agents Chemother 2006;50:1463-9
14	66. Méar JB, Kipnis E, Faure E, et al. Candida albicans and Pseudomonas aeruginosa interactions:
15	more than an opportunistic criminal association? Med Mal Infect 2013;43:146-51
16	67. Bhattacharyya S, Agrawal A, Knabe C, et al. Sol-gel silica controlled release thin films for the
17	inhibition of methicillin-resistant Staphylococcus aureus. Biomaterials 2014;35:509-17
18	• An interesting report on an innovative biomaterial.
19	68. Sabbah M, Bernollin M, Doutheau A, et al. A new route towards fimbrolide analogues:
20	importance of the exomethylene motif in LuxR dependent quorum sensing inhibition. Med Chem
21	Commun 2013;4:363-6
22	69. Kutty SK, Barraud N, Pham A, et al. Design, synthesis and evaluation of fimbrolide-nitric oxide
23	donor hybrids as antimicrobial agents. J Med Chem 2013;56:9517-29
24	• The paper describes an innovative fusion of two molecules with antibacterial activity.
25	70. Tsuchikama K, Zhu J, Lowery CA, et al. C4-alkoxy-HPD: a potent class of synthetic modulators
26	surpassing nature in AI-2 quorum sensing. J Am Chem Soc 2012;134:13562-4

1	71. Praneenararat T, Palmer AG, Blackwell HE. Chemical methods to interrogate bacterial quorum
2	sensing pathways. Org Biomol Chem 2012;10:8189-99

- 72. Desouki SE, Nishiguchi K, Zendo T, et al. High-throughput screening of inhibitors targeting
   *Agr/Fsr* quorum sensing in *Staphylococcus aureus* and *Enterococcus faecalis*. Biosci Biotechnol
   Biochem 2013;77:923-7
- 6 73. Christensen QH, Groveb TL, Booker SJ, et al. A high-throughput screen for quorum-sensing
  7 inhibitors that target acyl-homoserine lactone synthases. Proc Natl Acad Sci 2013;110: 13815-20
- 8 74. Tan SY-Y, Chua S-L, Chen Y, et al. Identification of five structurally unrelated quorum-sensing
- 9 inhibitors of *Pseudomonas aeruginosa* from a natural-derivative database. Antimicrob Agents
- 10 Chemother 2013;57:5629-41
- 75. Skovstrup S, Le Quement ST, Hansen T, et al. Identification of LasR ligands through a virtual
   screening approach. Chem Med Chem 2013;8:157-63
- 76. Weber M, Buceta J. Dynamics of the quorum sensing switch: stochastic and non-stationary
  effects. BMC Syst Biol 2013;7:6.
- 15 77. Huang Y, Zeng Y, Yu Z, et al. In silico and experimental methods revealed highly diverse
- 16 bacteria with quorum sensing and aromatics biodegradation systems--a potential broad
- 17 application on bioremediation. Bioresour Technol 2013;148:311-6
- 78. Schaadt NS, Steinbach A, Hartmann RW, et al. Rule-based regulatory and metabolic model for
  Quorum sensing in *P. aeruginosa*. BMC Systems Biology 2013;7:81
- 20 79. Audretsch C, Lopez D, Srivastava M, et al. A semi-quantitative model of Quorum-Sensing in
- 21 *Staphylococcus aureus*, approved by microarray meta-analyses and tested by mutation studies.
- 22 Mol BioSyst 2013;9:2665-80
- 23 80. Sahner JH, Brengel C, Storz MP, et al. Combining in silico and biophysical methods for the
- 24 development of *Pseudomonas aeruginosa* quorum sensing inhibitors: an alternative approach for
- structure-based drug design. J Med Chem 2013;56:8656-64

1	81. Hebert CG, Gupta A, Fernandes R, et al. Biological nanofactories target and activate epithelial
2	cell surfaces for modulating bacterial quorum sensing and interspecies signaling. ACS Nano
3	2010;4:6923-31
4	82. Gupta A, Terrell JL, Fernandes R, et al. Encapsulated fusion protein confers "Sense and
5	Respond'' activity to chitosan-alginate capsules to manipulate bacterial quorum sensing.
6	Biotechnol Bioeng 2013;110:552-62
7	83. Gomes J, Grunau A, Lawrence AK, et al. Bioinspired, releasable quorum sensing modulators.
8	Chem Commun 2013;49:155-7
9	84. Connell JL, Ritschdorff ET, Whiteley M, et al. 3D printing of microscopic bacterial communities.
10	Proc Natl Acad Sci 2013;110:18380-5
11	85. Gupta S, Bram EE, Weiss R. Genetically programmable pathogen sense and destroy. ACS Synth
12	Biol 2013;PMID 23763381
13	• This article demonstrates the feasibility of a futuristic technology that uses bacteria programmed to
14	protect the host.
15	86. Guglielmini J, Van Melderen L. Bacterial toxin-antitoxin systems. Mobile Genetic Elements
16	2011;1:283-90
17	87. Kolodkin-Gal I, Hazan R, Gaathon A, et al. A linear pentapeptide is a quorum-sensing factor
18	required for mazEF-mediated cell death in Escherichia coli. Science 2007;318:652-5
19	88. Kolodkin-Gal I, Sat B, Keshet A, et al. The communication factor EDF and the toxin-antitoxin
20	module <i>mazEF</i> determine the mode of action of antibiotics. PLoS Biology 2008;6:e319
21	89. Kumar S, Kolodkin-Gal I, Engelberg-Kulka H. Novel quorum-sensing peptides mediating
22	interspecies bacterial cell death. mBio 2013;4:e00314-13
23	90. Givskov M, de Nys R, Manefield M, et al. Eukaryotic interference with homoserine lactone-
24	mediated prokaryotic signalling. J Bacteriol 1996;178:6618-22
25	91. Koh CL, Sam CK, Yin WF, et al. Plant-Derived Natural Products as Sources of Anti-Quorum
26	Sensing Compounds. Sensors 2013;13:6217-28

1	92. Pan J, Ren D. Quorum sensing inhibitors: a patent overview. Expert Opin Ther Pa
2	2009;19:1581-601

3	93. Romero M, Acuña L, Otero A. Patents on quorum quenching: interfering with bacterial
4	communication as a strategy to fight infections. Recent Pat Biotechnol 2012;6:2-12
5	94. Skindersoe ME, Alhede M, Phipps R, et al. Effects of antibiotics on quorum sensing of
6	Pseudomonas aeruginosa. Antimicrob Agents Chemother 2008;52:3648-63
7	95. Defoirdt T, Pande GS, Baruah K, et al. The apparent quorum-sensing inhibitory activity of
8	pyrogallol is a side effect of peroxide production. Antimicrob Agents Chemother
9	2013;57:2870-3
10	96. Defoirdt T, Brackman G, Coenye T. Quorum sensing inhibitors: how strong is the evidence?
11	Trends Microbiol 2013;21:619-24
12	• This is perhaps the first paper highlighting the need for a more stringent evaluation of QS
13	inhibitors.
14	97. Dockery JD, Keener JP. A mathematical model for quorum sensing in Pseudomonas
15	aeruginosa. Bull Math Biol 2001;63:95-116
16	98. Anguige K, King JR, Ward JP. A multi-phase mathematical model of quorum sensing in a
17	maturing Pseudomonas aeruginosa biofilm. Math Biosci 2006;203:240-76
18	99. Janakiraman V, Englert D, Jayaraman A, et al. Modeling growth and quorum sensing in
19	biofilms grown in microfluidic chambers. Ann Biomed Eng 2009;37:1206-16
20	100. Fozard JA, Lees M, King JR, et al. Inhibition of quorum sensing in a computational biofilm
21	simulation. Biosystems 2012;109:105-14
22	101. Beckmann BE, Knoester DB, Connelly BD, et al. Evolution of resistance to quorum
23	quenching in digital organisms. Artif Life 2012;18:291-310
24	102. Maeda T, García-Contreras R, Pu M, et al. Quorum quenching quandary: resistance to
25	antivirulence compounds. ISME J 2012;6:493-501

1	103.	Kalia VC, Wood TK, Kumar P. Evolution of Resistance to Quorum-Sensing Inhibitors.
2	М	ficrob Ecol 2013; PMID:24194099

- 3 104. García-Contreras R, Martínez-Vázquez M, Velázquez Guadarrama N, et al. Resistance to
- 4 the quorum-quenching compounds brominated furanone C-30 and 5-fluorouracil in
- 5 *Pseudomonas aeruginosa* clinical isolates. Pathog Dis 2013;68:8-11
- Köhler T, Perron GG, Buckling A, et al. Quorum sensing inhibition selects for virulence and
  cooperation in *Pseudomonas aeruginosa*. PLoS Pathog. 2010;6:e1000883.
- 8 106. Harraghy N, Kerdudou S, Herrmann M. Quorum-sensing systems in staphylococci as
- 9 therapeutic tergets. Anal Bioanal Chem 2007;387:437-44
- 10 107. Høyland-Kroghsbo NM, Maerkedahl RB, Svenningsen SL. A quorum-sensing-induced
  11 bacteriophage defense mechanism. mBio 2013;4:e00362-12
- 12 108. Michael-Gayego A, Dan-Goor M, Jaffe J, et al. Characterization of sil in invasive group A
- 13 and G streptococci: antibodies against bacterial pheromone peptide SilCR result in severe
- 14 infection. Infect Immun 2013;81:4121-7
- 15 109. Sturbelle RT, Conceição RC, Da Rosa MC, et al. The use of quorum sensing to improve
  vaccine mmune response. Vaccine 2014;32:90-5
- 17 110. Rumbaugh KP, Diggle SP, Watters CM, et al. Quorum sensing and the social evolution of
  18 bacterial virulence. Curr Biol 2009;19:341-45
- 19 111. Köhler T, Buckling A, van Delden C. Cooperation and virulence of clinical *Pseudomonas* 20 *aeruginosa* populations. Proc Natl Acad Sci U S A 2009;106:6339-44
- 21 112. Diggle SP. Microbial communication and virulence: lessons from evolutionary theory.
   22 Microbiology 2010;156:3503-12
- 23 113. Harrison F. Bacterial cooperation in the wild and in the clinic: are pathogen social
- behaviours relevant outside the laboratory? Bioessays 2013;35:108-12
- 25 114. Zhu J, Kaufmann GF. Quo vadis quorum quenching? Curr Opin Pharmacol 2013;13:688-98

1	115. Davies J. How to discover new antibiotics: harvesting the parvome. Curr Opin Chem Biol		
2	2011;15:5-10		
3	116. Gwynn MN, Portnoy A, Rittenhouse SF, et al. Challenges of antibacterial discovery		
4	revisited. Ann NY Acad Sci 2010;1213:5-19		
5			
6	Figure legends		
7			
8	Figure 1		
9	Chemical structures of anthranilate (1), methyl anthranilate (2), and PQS (3)		
10	Figure 2		
11	Chemical structures of N-(3-oxododecanoyl)-L-homoserine lactone (3-oxo-C12-HSL) (4), and N-		
12	butyryl-L-homoserine lactone (C4-HSL) (5)		
13	Figure 3		
14	Chemical structure of eugenol (6)		
15	Figure 4		
16	Chemical structures of two PqsR antagonists (7) and (9), and of a PqsR agonist (8) (from Lu et al		
17	2014)		
18	Figure 5		
19	Chemical structure of farnesol (10)		
20	Figure 6		
21	Chemical structures of examples of fimbrolides-NO-donor hybrid compounds (11), (12) (from		
22	Kutty et al., 2013)		
23			
24			
25			
26			





