

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Biocontrol activity of an alkaline serine protease from *Aureobasidium pullulans* expressed in *Pichia pastoris* against four postharvest pathogens on apple

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/144278> since 2016-10-20T17:29:56Z

Published version:

DOI:10.1016/j.ijfoodmicro.2014.05.001

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in [Banani H., Spadaro D., Zhang D., Matic S., Garibaldi A., Gullino M.L. (2014) *Biocontrol activity of an alkaline serine protease from Aureobasidium pullulans expressed in Pichia pastoris against four postharvest pathogens on apple. International Journal of Food Microbiology, 182-183, 1-8. DOI: 10.1016/j.ijfoodmicro.2014.05.001*].

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [10.1016/j.ijfoodmicro.2014.05.001]

23 **Biocontrol activity of an alkaline serine protease from *Aureobasidium pullulans* expressed in**
24 ***Pichia pastoris* against four postharvest pathogens on apple**

25

26 Houda Banani¹, Davide Spadaro^{1,2*}, Dianpeng Zhang¹, Slavica Matic¹, Angelo Garibaldi¹, Maria
27 Lodovica Gullino^{1,2}

28

29 ¹ Centre of Competence for the Innovation in the Agro-environmental Sector - AGROINNOVA,
30 University of Torino, via L. da Vinci 44, I-10095 Grugliasco (TO), Italy.

31

32 ² DiSAFA – Dept. Agricultural, Forestry and Food Sciences, University of Torino, via L. da Vinci
33 44, I-10095 Grugliasco (TO), Italy.

34

35 * Corresponding Author.

36 Tel.: +39 011 6708942; fax: +39 011 6709307.

37 *E-mail address:* davide.spadaro@unito.it (D. Spadaro).

38 **Abstract**

39 The yeast-like fungus *Aureobasidium pullulans* PL5 is a microbial antagonist against postharvest
40 pathogens of fruits. The strain is able to produce hydrolases, including glucanases, chitinases and
41 proteases. The alkaline serine protease gene *ALP5* from *A. pullulans* was cloned, inserted into the
42 vector pPIC9 to construct pPIC9/PL5, and then expressed in *Pichia pastoris* strain KM71. *ALP5*
43 had a molecular mass of 42.9 kDa after 5 days growth with 1% methanol induction at 28 °C. The
44 recombinant protease expressed in *P. pastoris* showed its highest activity under alkaline conditions
45 (at pH 10) and temperature of 50 °C. The antifungal activity of the recombinant protease was
46 investigated against *Penicillium expansum*, *Botrytis cinerea*, *Monilinia fructicola* and *Alternaria*
47 *alternata* *in vitro* and on apple. The recombinant protease reduced significantly the spore
48 germination and the germ tube length of the tested pathogens in PDB medium. The highest level of
49 protease efficacy was observed against *M. fructicola* and *B. cinerea*, whereas a lower efficacy was
50 observed against *P. expansum* and *A. alternata* indicating a possible effect of the pathogen cell wall
51 composition on the proteolytic activity of the recombinant protease. The presence of protease was
52 able to cause swelling of the hyphae of *B. cinerea*, under optical microscope. The recombinant
53 protease expressed in the *P. pastoris* was more active against the pathogens *in vitro* than the same
54 enzyme expressed in *E. coli* in previous studies. The efficacy of *ALP5* was also evaluated against
55 the pathogens *in vivo* on apples cv Golden delicious. The protease was more efficient in controlling
56 *M. fructicola*, *B. cinerea* and *P. expansum* than *A. alternata*. However, the extent of the activity was
57 dependent on the enzyme concentration and the length of fruit storage. This study demonstrated the
58 capacity of the alkaline serine protease to keep its enzymatic activity for some days in the
59 unfavourable environment of the fruit wounds. The alkaline serine protease could be developed as a
60 postharvest treatment with antimicrobial activity for fruit undergoing a short shelf life.

61
62 **Keywords:** *Alternaria alternata*, *Botrytis cinerea*, *Monilinia fructicola*,
63 *Penicillium expansum*, postharvest, recombinant expression.

64

65 **1. Introduction**

66

67 *Penicillium expansum*, *Botrytis cinerea*, *Monilinia fructicola* and *Alternaria alternata* are among
68 the most severe postharvest pathogens on apples in production areas where the most advanced
69 storage technologies are available (Khamis et al., 2012; Martini et al., 2013; Snowdon, 1990). To
70 control postharvest diseases of fruits, few synthetic fungicides are admitted (Singh and Sharma,
71 2007; Zhu, 2006). However, pathogen resistance to fungicides (Holmes and Eckert, 1999), and the
72 willingness to use safer and eco-friendly treatments, have generated interest in the development of
73 alternative non-chemical methods to reduce postharvest losses (Lopez-Reyes et al., 2010; Nunes et
74 al., 2012). Biological control using microbial antagonists has emerged as one of the most promising
75 alternatives to fungicides, either alone or as part of an integrated pest management (Janisiewicz and
76 Korsten, 2002). A clear understanding about the mode of action of biocontrol agents is important
77 for a successful implementation of postharvest biocontrol technology (Droby et al., 2009; Zhang et
78 al. 2011).

79 Among the different biocontrol agents, yeasts are promising and gaining popularity (Jamalizadeh et
80 al., 2011; Janisiewicz et al., 2010; Spadaro et al., 2008). In particular, the yeast-like fungus
81 *Aureobasidium pullulans* De Bary (Arnaud), showed to be effective against *B. cinerea*, *P.*
82 *expansum* and *Rhizopus stolonifer* on various fruit, including apple, grapes, sweet cherry,
83 strawberry and peach (Bencheqroun et al., 2007; Ippolito et al., 2000; Lima et al., 1997; Schena et
84 al., 2003). Moreover, the strain PL5 of *A. pullulans* showed high efficacy in the control of *B.*
85 *cinerea* and *P. expansum* on apples, in addition to *Monilinia laxa* on plums and peaches (Zhang et
86 al., 2010a).

87 Several mechanisms have been reported to play a significant role in the biocontrol activity of *A.*
88 *pullulans* strains, including induction of defence responses (Ippolito et al., 2000) and competition
89 for nutrients (Bencheqroun et al., 2007). Recently, it was demonstrated that the strain PL5 secretes

90 β -1,3-glucanase, exo-chitinase and endo-chitinase, in addition to the secretion of alkaline serine
91 protease (Zhang et al., 2010a; 2012).

92 In mycoparasitism, fungal proteases may be significantly involved in antagonistic activity, because
93 they may play a significant role in fungal cell wall lysis, which is composed of chitin and glucan
94 polymers embedded in, and covalently linked to a protein matrix (Wessels, 1986). The inner layer
95 of fungal cell walls is primarily composed of glucans and chitin arranged as interwoven
96 microfibrils, while the outer electron dense layer is mainly composed of covalently bound
97 mannosylated proteins (Klis et al. 2002). Proteases catalyse the cleavage of peptide bonds in
98 proteins. In recent years, there has been an increasing interest in the study of proteolytic enzymes,
99 because they constitute one of the most important group of industrial enzymes due to their
100 commercial value and potential application in several fields, including food science and technology,
101 pharmaceutical industries and detergent manufactories (Feijoo-Siota and Villa, 2011).

102 The protease gene *ALP5* of *A. pullulans* strain PL5 was previously cloned and expressed in
103 *Escherichia coli* BL21 (Zhang et al., 2012), showing a low enzymatic activity. Prokaryotic
104 expression systems could present some drawbacks, including incorrect protein processing, folding
105 and posttranslational modification, lower heterologous protein expression levels, and lower activity.
106 *Pichia pastoris* has recently emerged as an important yeast host for heterologous protein expression
107 (Cregg et al., 1993; Macauley et al., 2005). As an eukaryote, *P. pastoris* has many of the advantages
108 of higher eukaryotic expression systems, such as protein processing and folding, and
109 posttranslational modifications, while being as easy to manipulate as *Escherichia coli*
110 (Balamurugan et al., 2007). In the yeast expression system, the secreted heterologous protein is the
111 vast majority in the medium, and, if there are glycosylation sites, glycosylation may occur at these
112 sites. For this reason, yeast genes could be better expressed in eukaryotic expression systems, such
113 as *P. pastoris*.

114 Therefore, the objectives of this research were to clone the protease gene *ALP5* from *A. pullulans*
115 strain PL5 and to express it in *P. pastoris* to evaluate its activity. A second objective was to

116 demonstrate the antifungal activity of the recombinant protease in controlling different postharvest
117 pathogens *in vitro* and *in vivo* on fruits, and to prove its involvement in the biocontrol activity of the
118 yeast-like fungus PL5.

119

120 **2. Materials and Methods**

121

122 *2.1. Microorganisms, plasmids and molecular kits*

123 *Aureobasidium pullulans* strain PL5 was isolated from the carposphere of plum cv Angeleno and
124 selected for its efficacy (Zhang et al., 2010b). It was identified through microscopic observation of
125 cell and colony morphology, and by sequencing of the ribosomal region ITS (Genbank accession
126 number: FJ919775).

127 Strains of *Botrytis cinerea*, *Penicillium expansum*, *Monilinia fructicola* and *Alternaria alternata*
128 were isolated from rotten apples or peaches (Pellegrino et al., 2009; Saravanakumar et al., 2008),
129 then selected throughout this work for their virulence by inoculation in artificially wounded apples
130 cv Golden delicious. Each strain was maintained at 4 °C on PDA (potato dextrose agar, Merck,
131 Germany) slants.

132 The *Escherichia coli* strain DH5 α used in this study as host for plasmids, was obtained from
133 Invitrogen (Life Technologies, Carlsbad, USA). The oligonucleotides, pGEM-T vector and the *E.*
134 *coli* strain JM109 were purchased from Promega (Madison, USA). *Pichia pastoris* KM71 strain
135 (Invitrogen) was used as host for transformations with the plasmid pPIC9 (Invitrogen).

136

137 *2.2. Total RNA isolation and first-strand cDNA synthesis*

138 The strain PL5 was grown in liquid medium YPD (20 g D-glucose, 20 g peptone casein, and 10 g
139 yeast extract per litre) at 25 °C. After 48h shaking at 200 rpm on a rotatory shaker (ASAL, Italy),
140 the culture was centrifuged for 10 min at 5,000 g. RNA was extracted from the pellet with
141 RNeasy® extraction kit (Qiagen, Hilden, Germany), then the first-strand cDNA was synthesized

142 using Reverse Transcript kit according to the manufacturer's instructions (Qiagen). RNA
143 concentration and purity were checked by spectrophotometer (Nanodrop 2000, Thermo Scientific,
144 Wilmington, USA).

145

146 2.3. Cloning of ALP5-Encoding cDNA gene

147 In order to amplify the cDNA of the *ALP5* gene encoding the alkaline serine protease by PCR, we
148 designed the forward primer (Pf-*ALP5*) as 5'-ACTGAATTCATGTGGAAGAAGAGTGTTGC-3'
149 and reverse primer (Pr-*ALP5*) as 5'-AATGAATTCTAACGACCGCTGTTGTTGTAAAC-3'; (bases
150 underlined encode *EcoRI* restriction site) according to the sequence of the protease gene obtained
151 from the genomic DNA (GenBank accession number HQ113460.1). PCR conditions were as
152 follows: an initial step at 94 °C for 3 min, and 35 cycles at 94 °C for 30 s, 58 °C for 45 s, and 72 °C
153 for 90 s. The purified PCR amplicons were ligated into pGEM-T-Easy cloning vector, then
154 transformed into chemically competent cells of *E. coli* strain DH5 α .

155

156 2.4. Construction of the expression vector pPIC9-ALP5

157 The product was subsequently digested with *EcoRI* and ligated into the pPIC9 vector, which was
158 previously digested with the same restriction enzyme. The consequent plasmid pPIC9-*ALP5* was
159 transformed into *E. coli* JM109 (Promega). The recombinant plasmid was isolated from the positive
160 transformants using Qiaprep Spin Miniprep Kit (Qiagen). The presence and correct orientation of
161 the insert sequence was confirmed by DNA sequencing at BMR Genomics (Padova, Italy).

162

163 2.5. Transformation of *P. pastoris*

164

165 The pPIC9-*ALP5* construct was linearized by *Stu I* enzyme for efficient integration into the *P.*
166 *pastoris* genome, then it was transformed into *P. pastoris* strain KM71. The empty vector (pPIC9)
167 was also transformed in *P. pastoris* for negative control tests following the manufacturer's

168 instructions. After transformation with plasmid pPIC9-ALP5, His⁺ transformants of *P. pastoris*
169 KM71 were purified on minimal medium plates without histidine to ensure pure clonal isolates,
170 then the genomic DNA of the transformants were isolated and PCR amplification and sequencing
171 were done to confirm whether the protease cDNA was integrated into the genomic DNA of *P.*
172 *pastoris*. Primers used for PCR, 5'AOX1 (5'-GACTGGTTCCAATTGACAGC-3') and 3'AOX1
173 (5'-GCAAATGGCATTCTGACATCC-3') were provided by the manufacturer. The PCR screening
174 of the positive recombinants produced a 1.7 kb fragment, while the control yeast transformed with
175 pPIC9 produced a 492 bp product (data not shown), which confirmed the integration of the insert
176 into *P. pastoris* genome.

177

178 2.6. Expression and purification of *A. pullulans* strain PL5 protease

179 Transformed *P. pastoris* isolates were cultured in 100 mL of BMGY medium (1% yeast extract, 2%
180 peptone, 100 mM potassium phosphate, pH 6.0, 1.34% YNB, 4×10^{-5} % biotin and 1% glycerol)
181 for approximately 24 h at 28 °C with constant shaking till OD 600 nm reached about 2-6. Cells
182 were centrifuged and the cell mass was resuspended in 20 ml of BMMY medium (1% yeast extract,
183 2% peptone, 100 mM potassium phosphate, pH 6.0, 1.34% YNB, 4×10^{-5} % biotin and 0.5%
184 methanol) to induce expression of the recombinant proteins. The culture was supplemented daily
185 with 100% methanol to a final concentration of 1% to maintain induction. Supernatants were then
186 harvested by centrifugation at 3,000 g at 4 °C. The production of the protease in the supernatant of
187 the culture medium was analysed by sodium dodecyl sulphate polyacrylamide gel electrophoresis
188 SDS-PAGE (Laemmli, 1970) (Amersham ECL Gel 10%, GE Healthcare Life science, Uppsala,
189 Sweden) of aliquots taken at different times (0, 24, 48,72, 96, 120 and 144h). Large-scale
190 production was performed under similar conditions using the isolate that rendered the best yield in
191 the small-scale experiments. To purify the recombinant protease, the supernatant was collected and
192 protein content was precipitated by adding ammonium sulphate (Sigma Aldrich, Milan, Italy,
193 approximately 80% saturation). After centrifugation at 8,000 g for 15 min, the protein precipitate

194 was dissolved in 50 mM Tris-HCl (pH 8.0) and dialyzed overnight against three changes of the
195 same buffer at 4 °C. The dialyzed sample was clarified by centrifugation and then applied to an
196 anion-exchange column of DEAE Sepharose 1×20 cm (GE Healthcare, United Kingdom),
197 previously equilibrated with 50 mM Tris-HCl (pH 8.0). To elute the bound proteins, 200 mL
198 equilibration buffer were used with a linear gradient from 0 to 0.3 M NaCl at a flow rate of 40
199 mL/h. Purity of the recombinant protease was determined by SDS-PAGE.

200

201 *2.7. Protease assay*

202 The recombinant protease activity was assayed against azocasein according to Inamura et al. (1985)
203 with minor modifications. Concisely, a linear rate of increase in activity was obtained up to an
204 absorbance of 0.2 at 440 nm. One unit of the enzyme activity was defined as the quantity of
205 protease which gave an absorbance of 0.001 at 440 nm. The protein concentration was determined
206 by using the Bradford assay (Bradford, 1976). Bovine serum albumin (Sigma-Aldrich, Milan, Italy)
207 was used as a standard. All the experiments and measurements were performed at least in triplicate.

208

209 *2.8. The effect of temperature and pH on the activity and stability on the recombinant protease*

210 To check the effect of temperature on the activity of the recombinant alkaline serine protease, the
211 enzyme was incubated between 20 °C and 60 °C using the standard assay conditions for the
212 determination of the protease activity. Similarly, the effect of pH on the enzyme activity was
213 studied by incubating the protease at different pH values (ranging from 4.0 to 12.0). The
214 recombinant *ALP5* activity obtained at temperature of 50 °C and pH of 10 was used as a reference
215 (relative activity was 100%), in order to calculate the relative activity at different pH values and
216 temperatures. The experiment was performed three times.

217

218 *2.9. Effect on pathogen mycelium growth inhibition*

219 The activity of the recombinant protease against postharvest pathogens was assayed in Petri dishes
220 containing PDA for *B. cinerea*, *P. expansum*, *M. fructicola* and *A. alternata* according to Zhang et
221 al. (2012) with some modifications. In brief, the mycelial plugs (5 mm in diameter) of the
222 pathogens were corked from a PDA culture and fixed in Petri dish. Ten μL of protease was streaked
223 into PDA at 30 mm from the pathogen plug, and after 4, 8, and 16 days of pathogens growing at 25
224 $^{\circ}\text{C}$, direct interaction *in vitro* was observed. The inhibitory effect of the protease on the pathogens
225 mycelial growth was calculated by following the formula:

226 Percentage of mycelium growth inhibition (%) = $\left[\frac{DC - DP}{DC} \right]$

227 DC and DP refers to the average diameters of fungal mycelia of control and protease respectively.
228 The experiment was performed three times.

229
230 *2.10. Effect on pathogen spore germination and germ tube elongation*

231 The effects of the protease on spores germination and germs tube elongation of *B. cinerea*, *P.*
232 *expansum*, *M. fructicola* and *A. alternata* were assayed in potato dextrose broth (PDB, Merck), by
233 using the method of Zhang et al. (2012). Briefly, 300 μL of 1×10^6 conidia/mL Ringer solution
234 (Merck) of each pathogen were transferred to a glass tube containing 2.4 mL PDB. Then 300 μL of
235 purified recombinant protease (62,5 ng/ μl) was added to 45 $^{\circ}$ sloping tubes and they were incubated
236 on a rotary shaker (200 rpm) at 25 $^{\circ}\text{C}$ for 9 h and 18 h. In addition, the control was incubated in
237 similar conditions, and it contained a PDB medium inoculated with the pathogens and treated with
238 300 μL inactivated enzymes boiled at 100 $^{\circ}\text{C}$ for 10 min. After 9h and 18 h of incubation, the
239 germination rate and germ tube length were measured on 100 conidia per replication by optical
240 microscopy (Eclipse 55i, Nikon, Tokyo, Japan). For each treatment and pathogen, three replications
241 of three tubes were prepared, and the experiment was performed twice.

242
243 *2.11. Antifungal activity of recombinant protease against postharvest pathogens on apple*

244 The antifungal activity of recombinant protease against *B. cinerea*, *P. expansum*, *M. fructicola* and
245 *A. alternata* was assessed following the method of Yan et al. (2008) with modifications. The
246 conidial suspension of each pathogen was prepared by flooding 14 day-old Petri dish cultures
247 incubated at 25 °C, and adjusted to 10⁵ conidia/mL with sterile Ringer solution using a
248 haemocytometer (Zhang et al., 2011). Freshly harvested apples (cv Golden delicious) were surface-
249 sterilized with 1% sodium hypochlorite for 1 min, rinsed with tap water, air-dried, and punctured
250 with a sterile needle at the equatorial region (3 mm depth; three wounds per fruit).

251 In order to evaluate the effect of the protease concentration on the control of postharvest pathogens,
252 two concentrations of the protease PL5 were used: 62.5 ng/μL and 6.25 ng/μL, where 20 μL aliquot
253 of crude protease was pipetted into each wound. Heat inactivated crude protease and water served
254 as a control. Two hours later, 20 μL of conidial suspension (10⁵ conidia/mL) of each pathogen was
255 applied into each wound. In order to compare the biocontrol activity of *A. pullulans* PL5 with the
256 efficacy of the protease against the pathogens, the antagonist PL5 was grown in YPD medium for
257 48 h at 25 °C on a rotary shaker at 250 rpm, then the cells were harvested by centrifugation at
258 5000×g for 10 min and adjusted to final concentration of 10⁸ cells/mL, from which 20 μl was
259 pipetted into each wound. Two hours later, 20 μL of the pathogen suspension was inoculated into
260 each wound. The treated fruits were incubated at 23 °C, and the rot diameter was measured 4 and 7
261 days after inoculation (DAI) for *B. cinerea* and *P. expansum*, at 8 and 12 DAI for *M. fructicola*, and
262 at 12 and 21 DAI for *A. alternata*. Each treatment contained three replicates with ten fruits per
263 replicate and the experiment was performed three times.

264

265 2.12. Data analysis

266 All statistical analyses were performed with SPSS software (SPSS Inc., version 20.0, Chicago, IL,
267 USA). Data from all the experiments were analysed using analysis of variance (ANOVA), and the
268 treatment means were separated at 5% significance level by using Duncan's multiple range tests.

269

270 **3. Results**

271

272 *3.1. Expression and purification of recombinant protease*

273 The expression of purified recombinant protease from a transformed isolate of *P. pastoris* was
274 analysed through SDS-PAGE (Figure 1). After 120 h induction, the protease band was observed in
275 some isolates at 42.9 kDa, which corresponds to the same molecular weight of the *ALP5* protease
276 (Zhang et al., 2012), while no band was present in the negative control (non-insert control: lane
277 number 0). After small-scale production, the best producer isolates (colonies 4 and 5, Fig. 1) were
278 selected for large-scale expression. The recombinant protein was easily purified with DEAE-
279 Sepharose column protein purification system.

280

281 *3.2. Effect of temperature and pH on protease activity*

282 The effect of temperature and pH on the enzyme activity is shown in table 1. The recombinant
283 protease had its maximal activity at 50 °C, however the activity started decreasing when the enzyme
284 reaction temperature exceeded 50 °C.

285 Results on the effect of pH on the enzyme activity showed that the maximum activity of the
286 protease was observed at pH 10, then it was reduced at higher pH values.

287

288 *3.3 Effect on pathogen mycelium growth inhibition*

289 After several days of pathogen growth in PDA plates streaked with the protease, the effect of the
290 recombinant enzyme on pathogen mycelium growth was assessed. *M. fructicola* (A) and *B. cinerea*
291 (B) mycelial growth were significantly inhibited by the presence of the recombinant protease (fig.
292 2a). After 4 days of incubation, the mean inhibition of mycelia growth of *M. fructicola* and *B.*
293 *cinerea* were 43.3 % and 33.7 % respectively. The mycelial growth inhibition was progressively
294 reduced with the incubation time and it decreased up to 37.7 % (A) and 12.2 % (B), respectively,
295 after 16 days of incubation (Fig. 2b). On the opposite, the recombinant protease did not show any

296 inhibition of the mycelial growth of *P. expansum* (Fig. 2a (C)) and *A. alternata* (Fig. 2a (D)),
297 starting from the first assessment.

298

299 3.3. Effect on pathogen spore germination and germ tube elongation

300 The effect of protease on spore germination and germ tube elongation of *B. cinerea*, *P. expansum*,
301 *M. fructicola*, and *A. alternata* is shown in Figures 3 and 4. The data shows that after 9 hours of co-
302 culturing (pathogen+enzyme) in liquid medium (PDB), the protease completely inhibited spore
303 germination of *B. cinerea* (Fig 3a) and *M. fructicola*, (Fig 3b). After 18h of incubation, the
304 germination percentages of *B. cinerea* (Fig 3a) and *M. fructicola* (Fig 3b) were significantly
305 reduced in presence of the recombinant protease from 98% to 54%, and from 98% to 37%,
306 respectively. No significant differences in germination percentage comparing to the control (heat
307 inactivated crude protease) were observed on *P. expansum* (Fig 3c) and *A. alternata* (Fig 3d) treated
308 with the recombinant protease either at 9h or 18h of incubation.

309 In addition, the average germ tube lengths of *B. cinerea* (Fig 4a) and *M. fructicola* (Fig 4b) conidia
310 were 88 and 57 μm after 18 h of incubation, respectively, while the average germ tube lengths of
311 their untreated controls were 372 and 206 μm , respectively. At 9 h and 18 h of incubation, no
312 significant differences in germ tube length were observed on *P. expansum* and *A. alternata* treated
313 with the recombinant protease or with heat inactivated crude protease.

314

315 3.4. Efficacy of recombinant protease against postharvest pathogens on apple

316 The antifungal activity of the recombinant protease was investigated on apple (Table 2). Two
317 protease concentrations, 62.5 ng/ μL and 6.25 ng/ μL were applied in order to study the effect of the
318 enzyme concentration on the biocontrol of the postharvest pathogens.

319 After 4 days of inoculation of *B. cinerea* conidia (Fig. 5), the protease was highly efficient in
320 controlling the pathogen compared to the controls (heat inactivated enzyme and water: 24.0 mm and
321 23.4 mm). The highest protease concentration significantly controlled the lesion diameter on apple

322 (2.8 mm), similarly to the whole antagonistic cells of *A. pullulans* PL5 (0.0 mm). However, the
323 lowest protease concentration was statistically less effective (10.8 mm) than the highest one or the
324 whole antagonist *A. pullulans*. At 7 DAI, the lesion diameters of grey mould rot increased to 28.3
325 mm and 37.3 mm, respectively, on apples treated with the highest and the lowest protease
326 concentration, compared to 15.0 mm on *A. pullulans* PL5 treated fruits, but the protease was still
327 significantly more efficient than the controls, which showed 59.3 mm (heat inactivated protease)
328 and 61.2 mm (water) rot diameter.

329 The two protease concentrations similarly reduced *P. expansum* growth on apple at 4 DAI, with a
330 rot diameter of 8.3 mm and 9.0 mm respectively, compared to 15.0 mm and 17.1 mm in the control
331 fruits (heat inactivated protease and water treated) and no rot in PL5 treated fruit. At 7 DAI, the
332 blue mould lesion diameter on apple fruit treated with both protease concentrations increased to
333 become similar to the controls. The biocontrol agent PL5 showed more efficacy in controlling *P.*
334 *expansum* compared to the other treatments.

335 At 8 DAI with *M. fructicola*, the rot diameter of fruits treated with the highest protease
336 concentration (33.3 mm) was lower than the rots in apples treated with the lowest concentration
337 (43.7 mm), with water or with the inactivated enzyme (50.6 mm and 49.8 mm). The best control
338 against brown rot caused by *M. fructicola* was observed in *A. pullulans* PL5 treated fruits (20.8
339 mm). At 12 DAI, the rot diameters enlarged, and *A. pullulans* PL5 remained the best treatment with
340 41.0 mm rot, although the protease at the highest concentration controlled better *M. fructicola* (54.5
341 mm) than its lowest concentration (72.5 mm) or the control fruits (treated with heat inactivated
342 protease or water: 73.2 mm and 75.8 mm).

343 Only *A. pullulans* PL5 significantly reduced the rot diameter caused by *A. alternata* on apple, and
344 neither protease concentrations were efficient in controlling the pathogen at 12 or 21 DAI.

345

346 4. Discussion

347

348 *Aureobasidium pullulans* is a yeast-like fungus that resides in different environments, such as
349 woody tissues and leaves (Gonzalez and Tello, 2011), the surface of fruits from the early stages of
350 their development to maturity (Janisiewicz et al., 2010), and also human skin (Hawkes et al., 2005).
351 Different strains of *A. pullulans* have shown significant control of postharvest pathogens of fruits
352 (Bencheqroun et al., 2007, Ippolito et al., 2000; Zhang et al., 2010a). Recently, *Aureobasidium*
353 *pullulans* DSM 14940 and DSM 14941 have been registered as antimicrobial agents to control fire
354 blight *Erwinia amylovora* on pomefruit (EFSA, 2011). According to toxicological studies, *A.*
355 *pullulans* DSM 14940 and DSM 14941 are not acutely toxic, pathogenic or infective, and not able
356 to replicate within the human body. Anyway, some strains of this species have been recognized as
357 etiologic agents of unusual mycoses in immunosuppressed patients (Bolognino and Criseo, 2003; de
358 Oliveira et al., 2013; Huang et al., 2008).

359 In the present study, we cloned and expressed the protease gene *ALP5* of the antagonistic yeast-like
360 fungus *A. pullulans* in the methylotrophic yeast *P. pastoris* in order to verify its involvement in the
361 biocontrol activity of *A. pullulans* against four pathogens both *in vitro* and *in vivo*. Microbial
362 alkaline proteases are generally considered as safe (GRAS) substances and they dominate the world
363 enzyme market, because of their extensive use in the detergent and food industry, especially the
364 ones produced by some *Bacillus* species which (Schallmey et al., 2004).

365 The alkaline serine protease gene *ALP5* was successfully expressed in *P. pastoris* and, as expected,
366 the proteolytic activity of the recombinant protease showed its highest activity under alkaline
367 conditions (at pH 10) and temperature of 50 °C, which was very similar to the activity of the
368 recombinant protease expressed in *E. coli* and to other alkaline serine proteases of bacterial (Gupta
369 et al., 2002) and fungal (Macchione et al., 2008) origin. These results are consistent with a previous
370 report indicating that the optimal pH and temperature of a related protein (alkaline α -amylase gene)
371 from alkaliphilic *Alkalimonas amylolytica* expressed both in *E. coli* and *P. pastoris* are similar
372 (Yang et al., 2012).

373 This study demonstrated that the recombinant protease gene *ALP5* expressed in *P. pastoris* is
374 effective in reducing spore germination and germ tube length of some postharvest pathogens. The
375 dual culture procedure was adopted to study the enzymatic activity of the alkaline serine protease,
376 showing that this *in vitro* method could be applied not only to hypothesize an antimicrobial activity
377 due to release of antibiotics, but also to evaluate the presence of active hydrolases. The presence of
378 protease was able to cause swelling of the hyphae of *B. cinerea*, under optical microscope.

379 Different levels of *ALP5* efficacy were observed, indicating a possible effect of the pathogen cell
380 wall composition on the proteolytic activity of the recombinant protease. The highest levels of
381 protease efficacy were observed in *M. fructicola* and *B. cinerea*. Conversely, *ALP5* protease was
382 less effective against *P. expansum* and *A. alternata*. These data suggest that the postharvest
383 pathogens display differential sensitivity to protease, as shown in previous studies which revealed
384 that despite having similar cell wall constituents (Bartnicki-Garcia, 1968), the filamentous fungi are
385 differentially sensitive to the cell wall degrading enzymes (chitinases and β -1,3-glucanases;
386 Schlumbaum et al., 1986; Saravanakumar et al., 2009). Proteins represent an important component
387 of the fungal cell wall. Many fungal cell wall proteins identified by tandem mass-spectroscopy have
388 putative carbohydrate-modifying functions involved in cell wall synthesis and remodeling, others
389 are classified as adhesins or heme-binding proteins involved in iron uptake (De Groot et al. 2005).

390 Proteins are the second major component of the cell wall prepared from *B. cinerea* cultures (Cantu
391 et al., 2009). In *Penicillium* spp. (Pessoni et al., 2005) the percentage of proteins is lower (24%),
392 and carbohydrate represent the major component (43%). In species of *Alternaria*, the protein
393 composition has not been characterized, but it is lower than in other fungal species, and melanin, a
394 dark pigment, plays a major role in strengthening the cell wall resistance to proteases (Kishore et
395 al., 2005). The different composition of the cell wall of different pathogens could justify the results
396 of the protease activity both *in vivo* and *in vitro*.

397 In addition, the current study confirms that the recombinant protease *ALP5* expressed in *P. pastoris*
398 exhibited higher proteolytic activity than the protease expressed in *E. coli*, since it is more efficient

399 in reducing spore germination of *M. fructicola* and *B. cinerea* than the protease expressed in the
400 prokaryotic expression system (Zhang et al., 2012). Similar results were obtained by Morton and
401 Potter (2000) which performed a comparative analysis between different model systems including
402 *E. coli* and *P. pastoris* for the expression of a carboxylesterase enzyme. Although the recombinant
403 carboxylesterase protein was observed in *E. coli*, little or no enzymatic activity was detected. In
404 contrast, active protein was produced in *P. pastoris*. These results demonstrate that *P. pastoris* is
405 more convenient than *E. coli* for an efficient expression of the protease from the antagonist *A.*
406 *pullulans* PL5.

407 As a prosecution of these studies, the biocontrol activity of the recombinant protease was evaluated
408 *in vivo* on apple wounds. The results demonstrated the capacity of the alkaline serine protease to
409 keep its enzymatic activity for some days in the unfavourable environment of the fruit wounds. The
410 recombinant protease reduced the activity of *M. fructicola*, *B. cinerea* and *P. expansum* for the first
411 days of the treatment, after which the pathogen lesion diameters increased. The protease reduced the
412 lesion diameter compared with the controls (the heat inactivated enzyme or water), but the extent
413 was dependent on the enzyme concentration and the temporal distance from the protease treatment.
414 This may be due to the loss of the proteolytic activity with increasing the number of storage days.
415 Our results are in accordance with the results obtained by Yan et al.(2008), which demonstrated that
416 the efficacy of recombinant rice chitinase expressed in *P. pastoris* against *B. cinerea* of loquat
417 fruits, is dependent on the concentration of the enzyme and the time of chitinase treatment and
418 pathogen inoculation.

419 Concerning *A. alternata*, the protease was totally ineffective, and only the biocontrol agent could
420 control the pathogen growth. The protease did not reduce the rot diameter caused by *A. alternata*,
421 which has a slower growth rate compared to the other pathogens (Kader, 2002); after 12 and 21
422 DAI the activity of the protease in degrading the cell wall of the pathogen was lost. Another reason
423 could be also the chemical composition of the cell wall (Kawamura et al., 1999; Wang et al., 1996).
424 The microorganism PL5 better reduced the rot diameter of the pathogens studied, compared to the

425 protease, probably because other mechanisms of action, and possibly hydrolases, are synergistically
426 involved in pathogen control. Previous work reported the broad efficacy of *A. pullulans*, as the
427 result of a multicomponent action, involving competition for nutrients (Bencheqroun et al., 2007),
428 induction of defense responses (Ippolito et al., 2000), and production of lytic enzymes (Zhang et al.,
429 2010; 2012). To our knowledge, this work provides the clarification of the protease role in the
430 antagonistic activity of the biocontrol agent *A. pullulans* PL5.

431 In conclusion, either the use of microbes or enzymes as biocontrol approaches presents
432 advantages and drawbacks. The biocontrol agent guarantees higher and longer activity, but it could
433 imply human safety issues, while the enzyme is generally regarded as safe compound, also for the
434 food industry, it has been deeply studied and characterized, but it can be used for limited storage
435 periods. The alkaline serine protease could be developed as a postharvest treatment with
436 antimicrobial activity for fruit undergoing a short shelf life.

437

438 **Acknowledgements**

439

440 Research funded by the project “SafeFoodControl – Development of innovative systems and
441 technologies for the production, storage, processing and valorisation of Piedmontese fruit and
442 vegetables” granted by the Piedmont Region. The current address of Dianpeng Zhang is: Institute of
443 Plant and Environment Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing
444 100097, China.

445

446 **References**

447

448 Bartnicki-Garcia, S., 1968. Cell wall chemistry, morphogenesis, and taxonomy of fungi. Annual
449 Reviews of Microbiology 22, 87–108.

450 Balamurugan, V., Reddy, G.R., Suryanarayana V.V.S., 2007. *Pichia pastoris*: A notable
451 heterologous expression system for the production of foreign proteins—Vaccines. Indian Journal of
452 Biotechnology 6, 175–186.

453 Bencheqroun, S.K., Bajji, M., Massart, S., Labhilili, M., El-Jafari, S., Jijakli, M.H., 2007. *In*
454 *vitro* and *in situ* study of postharvest apple blue mold biocontrol by *Aureobasidium pullulans*:
455 evidence for the involvement of competition for nutrients. Postharvest Biology and Technology 46,
456 128–135.

457 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities
458 of protein utilizing the principle of protein-dye binding. Analytical Biochemistry 72, 1105–1112.

459 Bolignano, G., Criseo, G., 2003. Disseminated Nosocomial Fungal Infection by *Aureobasidium*
460 *pullulans* var. *melanigenum*: a Case Report. Journal of Clinical Microbiology 41, 4483-4485.

461 Cantu, D., Greve, C., Labavitch, J.M., Powell, A.L.T., 2009. Characterization of the cell wall of
462 the ubiquitous plant pathogen *Botrytis cinerea*. Mycological research 113, 1396-1403.

463 Cregg, J.M., Vedvick, T.S., Raschke, W.C., 1993. Recent advances in the expression of foreign
464 genes in *Pichia pastoris*, Biotechnology 11, 905–910.

465 Cregg, JM., Tolstorukov, I., Kusari, A., Sunga, J., Madden, K., Chappell, T., 2009. Expression in
466 the yeast *Pichia pastoris*. Methods in Enzymology 463, 169–189.

467 Damasceno, LM., Huang, CJ., Batt, CA., 2012. Protein secretion in *Pichia pastoris* and advances
468 in protein production. Applied Microbiology and Biotechnology 93, 31–39.

469 De Groot, P.W.J., Ram, A.F., Klis, F.M., 2005. Features and functions of covalently linked
470 proteins in fungal cell walls. Fungal Genetics and Biology 42, 657–675.

471 De Oliveira, L.R., Moraes-Souza, H., Maltos, A.L., Santos, K.C., Molina, R.J., Barata, C.H.,
472 2013. *Aureobasidium pullulans* infection in a patient with chronic lymphocytic leukemia. Revista
473 da Sociedade Brasileira de Medicina Tropical 46, 660-662.

474 De Pourcq, K., De Schutter, K., Callewaert, N., 2010. Engineering of glycosylation in yeast and
475 other fungi: current state and perspectives. *Applied Microbiology and Biotechnology* 87, 1617–
476 1631.

477 Droby, S., Wisniewski, M., Macarisin, D., Wilson, C., 2009. Twenty years of postharvest
478 biocontrol research: is it time for a new paradigm? *Postharvest Biology and Technology* 52, 137–
479 145.

480 EFSA, 2011. Potential inclusion of the microorganism *Aureobasidium pullulans* strains DSM
481 14940 and DSM 14941 in Annex IV of Regulation (EC) No 396/2005. *EFSA Journal* 9 (11), 2435.

482 Feijoo-Siota, L., Villa, T., 2011. Native and Biotechnologically Engineered Plant Proteases with
483 Industrial Applications. *Food and Bioprocess Technology* 4, 1066–1088.

484 Gonzalez, V., Tello, M.L., 2011. The endophytic mycota associated with *Vitis vinifera*
485 in central Spain. *Fungal Diversity* 47, 29–42.

486 Gupta, R., Beg, O.K., Lorenz, P., 2002. Bacterial alkaline proteases: molecular approaches and
487 industrial applications. *Applied Microbiology and Biotechnology* 59, 15–32.

488 Hawkes, M., Rennie, R., Sand, C., Vaudry, W., 2005. *Aureobasidium pullulans* infection:
489 Fungemia in an infant and a review of human cases. *Diagnostic Microbiology and Infectious*
490 *Disease* 51, 209-213.

491 Holmes, G.J., Eckert, J.W., 1999. Sensitivity of *Penicillium digitatum* and *P. italicum* to
492 postharvest citrus fungicides in California. *Phytopathology* 89, 716–721.

493 Huang, Y.T., Liaw, S.J., Liao, C.H., Yang, J.L., Lai, D.M., Lee, Y.C., Hsueh, P.R., 2008. Catheter-
494 related septicemia due to *Aureobasidium pullulans*. *International Journal of Infectious Diseases* 12:
495 137-139.

496 Inamura, H., Nakai, T., Muroga, K., 1985. An extracellular protease produced by *Vibrio*
497 *anguillarum*. *Bulletin of the Japanese Society of Scientific Fisheries* 51, 1915–1920.

498 Ippolito, A., Ghaouth, A.E., Wilson, C.L., Wisniewski, M., 2000. Control of postharvest decay
499 of apple fruit by *Aureobasidium pullulans* and induction of defense responses. *Postharvest Biology*
500 *and Technology* 19, 265–272.

501 Jamalizadeh, M., Etebarian, H.R., Aminian, H., Alizadeh, A., 2011. A review of mechanisms of
502 action of biological control organisms against post-harvest fruit spoilage. *EPPO Bulletin* 41, 65-71.

503 Janisiewicz, W.J., Korsten, L., 2002. Biological control of postharvest diseases of fruits. *Annual*
504 *Review of Phytopathology* 40, 411–441.

505 Janisiewicz, W.J., Kurtzman, C.P., Buyer, J.S., 2010. Yeasts associated with nectarines and their
506 potential for biological control of brown rot. *Yeast* 27, 389-398.

507 Kader, A.A., 2002. *Postharvest Technology of Horticultural Crops*. UCANR Publications.
508 California.

509 Kawamura, C., Tsujimoto, T., Tsuge, T., 1999. Targeted disruption of a melanin biosynthesis
510 gene affects conidial development and UV tolerance in the Japanese pear pathotype of *Alternaria*
511 *alternata*. *Molecular Plant Microbe Interactions* 12, 59–63.

512 Khamis, Y., Ligorio, A., Sanzani, S.M., Nigro, F., Ippolito, A., 2012. Control of storage diseases
513 of citrus by pre- and postharvest application of salts. *Postharvest Biology and Technology* 72, 57-
514 63.

515 Kishore, K.H., Kanjilal, S., Misra, S., Reddy, C.R., Murty, U.S., 2005. Comparative chemical
516 characterization of pigmented and less pigmented cell walls of *Alternaria tenuissima*. *Current*
517 *Microbiology* 51, 399-401.

518 Klis, F.M., Mol, P., Hellingwerf, K., Brul, S., 2002. Dynamics of cell wall structure in
519 *Saccharomyces cerevisiae*. *FEMS Microbiological Reviews* 26, 239–256.

520 Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head of
521 bacteriophage T4. *Nature* 227, 680-685.

522 Lima, G., Ippolito, A., Nigro, F., Salerno, M., 1997. Effectiveness of *Aureobasidium pullulans*
523 and *Candida oleophila* against postharvest strawberry rots. *Postharvest Biology and Technology* 10,
524 169–178.

525 Lopez-Reyes, J.G., Spadaro, D., Garibaldi, A., Gullino, M.L., 2010. Efficacy of plant essential
526 oils on postharvest control of rot caused by fungi on four cultivars of apples *in vivo*. *Flavour and*
527 *Fragrance Journal* 25, 171-177.

528 Macauley-Patrick, S., Fazenda, M.L., McNeil, B., Harvey, L.M., 2005. Heterologous protein
529 production using the *Pichia pastoris* expression system. *Yeast* 22, 249–270.

530 Macchione, M.M., Merheb, C.W., Eleni Gomes, E., Silva, R., 2008. Protease production by
531 different thermophilic fungi. *Applied Biochemistry and Biotechnology* 146, 223–230.

532 Martini C., Spadoni A., Mari M., 2013. First Report of Brown Rot Caused by *Monilinia*
533 *fruticola* on Apple in Italy. *Plant Disease* 97, 689.

534 Morton, CL., Potter, PM., 2000. Comparison of *Escherichia coli*, *Saccharomyces cerevisiae*,
535 *Pichia pastoris*, *Spodoptera frugiperda*, and *COS7* cells for recombinant gene expression:
536 Application to a rabbit liver carboxylesterase. *Molecular Biotechnology* 16, 193-202.

537 Nunes, C., 2012. Biological control of postharvest diseases of fruit. *European Journal of Plant*
538 *Pathology* 133, pp. 181-196.

539 Pellegrino, C., Gullino, M.L., Garibaldi, A., Spadaro, D., 2009. First report of brown rot of stone
540 fruit caused by *Monilinia fruticola* in Italy. *Plant Disease*, 93, 668.

541 Pessoni, R.A.B., Freshour, G., Figueiredo-Ribeiro, R.C.L., Hahn, M.G., Braga, M.R., 2005. Cell-
542 wall structure and composition of *Penicillium janczewskii* as affected by inulin. *Mycologia* 97, 304–
543 311.

544 Saravanakumar, D., Ciavarella, A., Spadaro, D., Garibaldi, A., Gullino, M.L., 2008.
545 *Metschnikowia pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and
546 *Penicillium expansum* in apples through iron depletion. *Postharvest Biology and Technology* 49,
547 121–128.

548 Saravanakumar, D., Spadaro, D., Garibaldi, A., Gullino, M.L., 2009. Detection of enzymatic
549 activity and partial sequence of a chitinase gene in *Metschnikowia pulcherrima* strain MACH1 used
550 as post-harvest biocontrol agent used as post-harvest biocontrol agent. *European Journal of Plant*
551 *Pathology* 123, 183-193.

552 Schallmey, M., Singh, A., Ward, O.P., 2004. Developments in the use of *Bacillus* species in
553 industrial production. *Canadian Journal of Microbiology* 50, 1-17.

554 Schena, L., Nigro, F., Pentimone, I., Ligorio, A., Ippolito, A., 2003. Control of postharvest rots
555 of sweet cherries and table grapes with endophytic isolates of *Aureobasidium pullulans*. *Postharvest*
556 *Biology and Technology* 30, 209–220.

557 Schlumbaum, A., Mauch, F., Vogeli, U., Boller, T., 1986. Plant chitinases are potent inhibitors
558 of fungal growth. *Nature* 324, 365–367.

559 Singh, D., Sharma, R.R., 2007. Postharvest diseases of fruit and vegetables and their
560 management. In: Prasad, D. (Ed.), *Sustainable Pest Management*. Daya Publishing House, New
561 Delhi, pp. 321-334.

562 Snowdon, A.L., 1990. A color atlas of post-harvest diseases and disorders of fruits and
563 vegetables: General Introduction and Fruits, Vol. 1. CRC Press, Boca Raton.

564 Spadaro, D., Sabetta, W., Acquadro, A., Portis, E., Garibaldi, A., Gullino, M.L., 2008. Use of
565 AFLP for differentiation of *Metschnikowia pulcherrima* strains for postharvest disease biological
566 control. *Microbiological Research* 163, 523-530.

567 Wang, H., Li, J., Bostock, R.M., Glichrist, D.G., 1996. Apoptosis: A Functional paradigm for
568 programmed plant cell death induced by a host-selective phytotoxin and invoked during
569 development. *Plant Cell* 8, 375–391.

570 Wessels, JGH., 1986. Cell wall synthesis in apical hyphal growth. *International Review of*
571 *Cytology* 104, 37–79.

572 Yan, R., Ding, D., Guan, W., Hou, J., Li, M., 2008. Control of grey mould rot of loquat with
573 chitinase expressed in *Pichia pastoris*. *Crop Protection* 27, 1312– 1317.

574 Yang, H., Liu, L., Shin, HD., Chen, RR., Li, J., Du, G., Chen, J., 2012. Comparative analysis of
575 heterologous expression, biochemical characterization optimal production of an alkaline α -amylase
576 from alkaliphilic *Alkalimonas amylolytica* in *Escherichia coli* and *Pichia pastoris*. *Biotechnology*
577 *Progress* 29, 39-47.

578 Zhang, H.Y., Zheng, X.D., Fu, C.X., Xi, Y.F., 2005. Postharvest biological control of gray mold
579 rot of pear with *Cryptococcus laurentii*. *Postharvest Biology and Technology* 35, 79–86.

580 Zhang, D., Spadaro, D., Garibaldi, A., Gullino, M.L., 2010a. Selection and evaluation of new
581 antagonists for their efficacy against postharvest brown rot of peaches. *Postharvest Biology and*
582 *Technology* 55, 174-181.

583 Zhang, D., Spadaro, D., Garibaldi, A., Gullino, M.L., 2010b. Efficacy of the antagonist
584 *Aureobasidium pullulans* PL5 against postharvest pathogens of peach, apple and plum and its
585 modes of action. *Biological Control* 54, 172–180.

586 Zhang, D., Spadaro, D., Garibaldi, A., Gullino, M.L., 2011. Potential biocontrol activity of a
587 strain of *Pichia guilliermondii* against grey mould of apples and its possible modes of action.
588 *Biological Control* 57, 193-201.

589 Zhang, D., Spadaro, D., Valente, S., Garibaldi, A., Gullino, M.L., 2012. Cloning,
590 characterization, expression and antifungal activity of an alkaline serine protease of *Aureobasidium*
591 *pullulans* PL5 involved in the biological control of postharvest pathogens. *International Journal of*
592 *Food Microbiology*. 153, 453-464.

593 Zhu, S.J., 2006. Non-chemical approaches to decay control in postharvest fruit. In: Nouredine,
594 B., Norio, S. (Eds.), *Advances in Postharvest Technologies for Horticultural Crops*. Research
595 Signpost, Trivandrum, pp. 297–313.

596 **Tables**

597

598 **Table 1** - Determination of the effect of temperature (a) and pH (b) on the enzymatic reaction of the
599 recombinant alkaline serine protease. The recombinant *ALP5* activity obtained at 50 °C and pH 10.0
600 was used as a reference in order to calculate the relative activity at different temperatures and pH
601 values. Experiments were performed three times and the data are presented as mean \pm standard
602 error.

603

604 **a)**

Temperature (°C)	Relative activity (%)
20	42 \pm 2.4
30	51 \pm 1.8
40	77 \pm 2.2
50	100 \pm 3.1
60	62 \pm 3.9

605 **b)**

pH value	Relative activity (%)
5	33 \pm 2.7
6	42 \pm 2.3
7	67 \pm 4.6
8	79 \pm 3.9
9	90 \pm 4.5
10	100 \pm 4.8
11	83 \pm 5.1

606 **Table 2** Antifungal activity of the recombinant protease ALP5 in controlling the decay development of *B. cinerea*, *P. expansum*, *M. fructicola*, and
607 *A. alternata* in wound-inoculated apples cv Golden Delicious (20 µL of a suspension at 10⁵ conidia/mL). The protease was applied at 62.5 ng/µL
608 and 6.25 ng/µL . The results are the mean of three independent experiments.
609

Treatment	Rot lesion diameters (mm) _a							
	<i>Botrytis cinerea</i>		<i>Penicillium expansum</i>		<i>Monilinia fructicola</i>		<i>Alternaria alternata</i>	
	4 days	7 days	4 days	7 days	8 days	12 days	12 days	21 days
Protease (62.5 ng/µL)	2.8±1.0a	28.3±3.2b	8.3±1.5b	31.0±4.6b	33.3±3.5b	54.5±4.1b	8.2±1.8b	48.5±4.9b
Protease (6.25 ng/µL)	10.8± 2.3b	37.3± 3.8b	9.0±1.3b	31.7±3.8b	43.7±4.7c	72.5±9.5c	9.5c1.3b	52.0±7.2b
<i>A. pullulans</i> PL5 10 ⁸ cells/ml	0.0±0.0a	15.0±2.3a	0.0±0.0a	8.2 1±1.4a	20.8±2.6a	41.0±4.6a	0.0±0.0a	35.2±3.8a
Control (inactivated protease)	24.0±2.6c	59.3±7.2c	15.0±3.6c	35.0±6.6b	49.8±6.1c	73.2±8.6c	11.8±2.6b	53.0±4.4b
Control (water)	23.4±3.0c	61.2±5.3c	17.1±2.9c	34.2±5.8b	50.6±6.4c	75.8±10.1c	12.1±3.2b	55.6±3.9b

610
611 “ ± ” stands for standard error of the means. Values of each column followed by different letters show significant difference (P < 0.05) according to
612 Duncan’s multiple range Test (SPSS 20.0).

613 **Figure captions**

614 **Figure 1** - Figure 1: SDS-PAGE analysis of the recombinant protease expressed in *P. pastoris*.
615 Supernatants of the yeast culture were taken from different isolates after 120 h of induction. Lanes:
616 M: molecular weight marker (Precision Plus Protein Dual Color Standards, BIO RAD); 0: *P.*
617 *pastoris* KM71 isolate transformed with PPIC9 (Control); 1, 2, 3, 4 and 5: some transformed *P.*
618 *pastoris* isolates with PPIC9-ALP5.

619
620 **Figure 2** - Antifungal activity of the recombinant protease from *A. pullulans* PL5 expressed in *P.*
621 *pastoris* against postharvest pathogens. (a): inhibition of pathogen growth *in vitro*: A) *M.*
622 *fructicola*; B): *B. cinerea*; C) *P. expansum*, D) *A. alternata*, (b): Mycelial growth inhibition (%) of
623 *B. cinerea*, *P. expansum*, *M. fructicola* and *A. alternata* after, respectively, 4, 4, 8, and 16 days of
624 growth at 25 °C.

625
626 **Figure 3** - Effect of the recombinant protease ALP5 on spore germination of *B. cinerea*, *P.*
627 *expansum*, *M. fructicola* and *A. alternata* in potato dextrose broth medium. Treatments followed by
628 different letters are statistically different following the Duncan's multiple range test ($p < 0.05$).
629 Spore germination were measured microscopically after 9 h and 18 h of incubation at 25 °C.

630
631 **Figure 4** - Effect of the recombinant protease ALP5 on germ tube length of *B. cinerea*, *P.*
632 *expansum*, *M. fructicola* and *A. alternata* in potato dextrose broth medium. Treatments followed by
633 different letters are statistically different following the Duncan's multiple range test ($p < 0.05$).
634 Germ tube length were measured microscopically after 9h and 18 h of incubation at 25 °C.

635
636 **Figure 5** – Grey mould on apple cv Golden Delicious after 4 days of *B. cinerea* (20 μL of a
637 suspension at 10^5 conidia/mL) inoculation and treatment with (a): heat-inactivated protease as a
638 control, (b): Protease at 6.25 ng/ μL , (c): Protease at 62.5 ng/ μL .