AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cropping system intensification grading using an agro-environmental indicator set in northern Italy

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/144457> since 2016-10-05T16:43:40Z

Published version:

DOI:10.1016/j.ecolind.2014.01.004

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *Ecological Indicators*, Volume 40, May 2014, Pages 76-89, <http://dx.doi.org/10.1016/j.ecolind.2014.01.004>.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), <http://www.sciencedirect.com/science/article/pii/S1470160X14000065>

1 CROPPING SYSTEM INTENSIFICATION GRADING USING AN AGRO-
2 ENVIRONMENTAL INDICATOR SET IN NORTHERN ITALY

3 Author names and affiliations: Stefano Gaudino, Irene Goia, Giorgio Borreani, Ernesto
4 Tabacco, Dario Sacco

5 University of Torino - Department of Agricultural, Forest and Food Sciences (DISAFA) – via
6 L. Da Vinci 44, 10095 Grugliasco (TO) - ITALY

7

8 Corresponding author:

9 Stefano Gaudino, University of Torino - Department of Agricultural, Forest and Food
10 Sciences (DISAFA) – via L. Da Vinci 44, 10095 Grugliasco (TO) – ITALY. Tel.:
11 +390116708905, Fax: +390116708798, email: stefano.gaudino@unito.it.

Title: CROPPING SYSTEM INTENSIFICATION GRADING USING AN AGRO-
ENVIRONMENTAL INDICATOR SET IN NORTHERN ITALY

Abstract

The term agro-environmental sustainability in agriculture usually refers to farming intensity. Lower intensity farming can be managed by reducing chemical and energy inputs. Beyond ethical issues and having in mind only agronomic aspects, cropping systems are defined by regulations that classify them according to their different input levels as conventional (most intensive), integrated (intermediate intensity), and organic (least intensive).

Among organic cropping systems, it is expected that the most intense cropping level would be arable farms where there is a greater need to import input factors, and the least intense level would be livestock farms. This research aims to systematically grade conventional, integrated, and organic cropping systems using a set of 22 indicators of input and environmental pressure. The grading results will then be compared to regulation-defined intensities.

Eight cropping systems belonging to four intensification levels were analysed by an indicator set classified as driving force or pressure indicators per the DPSIR schema. Driving forces represented farmer management decisions; pressures represented stressors to the environment resulting from agricultural activities not directly modifiable by the farmer. The 22 indicators analyse five aspects of cropping system: land use, fertiliser use, pesticide use, energy use and gaseous emissions.

Study results showed that most indicators were able to accurately grade the cropping system intensities. Specific driving forces and pressures indicators that failed to grade the cropping systems as expected related to several explainable factors. For driving force indicators, conventional systems demonstrated the highest impact on the environment and arable organic cropping systems the lowest. For pressure indicators, conventional cropping system presented

the highest impact, followed by integrated cropping systems. In this case the arable organic cropping system presented a higher impact than did the livestock organic system. This level of discrimination showed that pressure indicators performed better at grading system intensification than did driving force indicators.

As a consequence, the analysis showed that higher input levels do not always result in higher pressures on the environment. Therefore, the environment would be better served by regulations that set thresholds for pressures rather than system inputs. The results also underlined that practices such as manure use and meadow presence improve the environmental performances of cropping systems.

Key words: Agro-environmental sustainability assessment, environmental impact, organic farming, integrated farming, conventional farming.

1 Introduction

Over the past 60 years, European agriculture has undergone a period of rapid intensification achieved through an increased application of chemical fertilisers and pesticides, combined with implementation of best management practices, mechanisation, irrigation, and with the use of improved seed varieties (Tilman et al., 2002). Today, the term “agro-environmental sustainability” has come to imply high dry matter (DM) yields and society’s expectation for ecological service while complying with European environmental programs (Cross-compliance 73/2009/EC (EC, 2009a), Water Framework Directive 60/2000/EC (EC, 2000), Sustainable use of pesticides Directive 128/2009/EC (EC, 2009b), Birds Directive 147/2009/EC (EC, 2009c), and Habitats Directive 43/1992/EEC (Council of the European Communities, 1992)). These changes have led public and scientific communities to turn their attention to alternative farming systems including, among others, integrated farming, precision farming, conservation agriculture, and organic farming.

All of the above distinguish themselves from intensive conventional systems in their improved resource use efficiencies, rather than on external inputs to maintain productivity and profitability (Liebman et al., 2008). Low external-input and organic cropping systems could provide a good compromise between intensity (level of input used per unit of surface) and efficiency (quantity of product obtained per level of input used) (Alluvione et al., 2011; Michos et al., 2012; Pointereau et al., 2012).

Cropping system intensity is defined by European, national, and regional level regulations. This paper considers only the agronomic aspects, contained in the different regulations and do not consider the different ethical aspects that have led to them. Conventional cropping systems must satisfy statutory management requirements defined in the cross compliance system (73/2009/EC (EC, 2009a)), which represent the minimum legal limits. In Italy, the regional Rural Development Program (RDP) determines regulations for integrated farming

systems, whereas organic agriculture is governed by European regulations 834/2007/EC (EC, 2007) and 889/2008/EC (EC, 2008). Among low-input cropping systems, integrated agriculture has been promoted for its reduced environmental impact and increased sustainable resource use (Alluvione et al., 2011; Morris and Winter, 1999). Organic farming has also been advocated as more sustainable than conventional systems over the long-term (Pimentel et al., 2005), as it uses the fewest inputs and therefore, is the least intense. Banned chemical products, improved nutrient recycling, and “minimisation of the use of non-renewable resources and off-farm inputs” are keys to its sustainability (Regulation 834/2007/EC (EC, 2007)).

When livestock production systems are paired with organic systems, further efficiency and sustainability is achieved. Regulation 834/2007/EC has defined livestock production as “fundamental to organization of agricultural production...” because it can provide organic nutrients to the cropping system through within-the-farm recycling, and allows for partitioning between low sustainability/externally- and high sustainability/internally-produced inputs (Nemecek et al., 2011). From this follows that in organic farms the highest intensification level should be on arable ones because they require more imported inputs; conversely, the lowest intensification level should be on livestock organic farms as they utilise nutrient recycling to meet many of their input needs.

Several authors have confirmed the relationship between lower intensification level and lower environmental pressures (i.e. Flessa et al., 2002; Kramer et al., 2006; Liu et al., 2007). Environmental pressures, however, have not always corroborated the expectations associated with the intensification levels described above, with organic cropping systems being less sustainable than conventional systems (Kirchmann and Bergström, 2001; Eltun et al., 2002; Basset-Mens and van der Werf, 2005). Finally, van der Werf et al. (2007), comparing many assessment methods applied to farms producing crops and pigs, found that the rank between

organic and conventional farms depends on the assessment method applied and on the aspect analysed.

Field experiments and farm measures are two ways to evaluate directly the agro-environmental sustainability of different cropping systems, however, these methodologies are time-consuming when many aspects are analysed. “Indicators are an alternative when it is not possible to carry out direct measurements” (Bookstaller et al., 1997). They allow not only an understanding of complex systems (Mitchell and al., 1995), but also compare different situations, two characteristics that make them highly useful in the analysis of agricultural managements and their environmental pressures.

Different authorities — at both the European and worldwide scales — have created lists of indicators. Among them there are: EU Agro-Environmental indicators AEI (COM (2006) 508 (EC, 2006)), OECD agro-environmental indicators (OECD 1999), and FAO agro-environmental indicators (FAO, 2012). At the European level indicators are also used to evaluate environmental policy effects. Some indicators are suitable to analyse different levels of complexities, such as Input Output Account (IOA) (Halberg et al., 2005), the Life Cycle Assessment (LCA) (ISO 2006), and the Ecological Footprint (EF) (Rees, 2000). The IOA has been applied to different sustainability aspects, but in particular, to nutrient balances (Bassanino et al., 2007; Oenema et al., 2003; Schröder and Neetson, 2008) and energy balances (Alluvione et al., 2011; Meul et al., 2007). In the case of the LCA and EF, they analyse the sustainability of the entire production system via pressure category assessment. Analysis of specific pressures related to different agricultural managements is most useful when performed by single indicators or indicator sets.

This work analyses different cropping systems at various intensification levels (conventional, integrated, and organic) using an agro-environmental indicator set built of different indicators derived from literature. The investigation aims to grade these cropping systems on both input

level and environmental pressures; thereafter, the results will be compared to the expected grade derived from the intensification levels as defined by regulation.

2 Materials and methods

2.1 Description of the area

The study was carried out in the western Po Valley (Piemonte Region, NW Italy). The climate is temperate sub-continental, characterised by two main rain periods in spring and autumn, with an annual mean precipitation of 850 mm and an annual mean temperature of 11.8°C. The soil types are Inceptisols, Entisols and Alfisols (Bassanino et al., 2007), mainly characterized by silt-loam and silt texture.

According to the regional administrative database (Regione Piemonte, 2010), arable and livestock farms cover most of the Utilized Agricultural Area (UAA). Conventional arable farms are in the majority (94.5%) while integrated and organic farms represent just 4.9% and 0.6%, respectively. The main arable farm crops were maize (*Zea mays* L.), winter cereals (*Triticum aestivum* L., *Hordeum vulgare* L.), soybean (*Glycine max* (L.) Merr.), and meadows (Sacco et al., 2003). Livestock farms bred principally bovine and swine. Bovine livestock farms fell into one of three breeding types: beef, dairy cows, or suckling cows (Bassanino et al., 2007), with suckling cows comprising the largest share at 47%, of which 1.2% were organic farms. Bovine livestock farm main crops included maize (for grain and silage production), winter cereals, lucerne (*Medicago sativa* L.), Italian ryegrass (*Lolium multiflorum* Lam.), and hay-producing meadows (mixed grasses and legumes).

2.2 Farm types

Conventional, integrated, and organic cropping systems of farms were considered in this study. Organic farms were further divided into arable organic farms and livestock organic

farms according to their external input levels, which created four different farm intensification groups:

- conventional arable farms (CONV)
- integrated arable farms (INT)
- organic arable farms (ORG)
- organic livestock farms (LIV)

Two farms were selected at each intensification level, to represent the variability of farm managements and input use levels. Organic livestock farms were selected from the suckling cow breeding type. We further focused our work on cropping systems alone. From conventional and integrated farms, only those that applied mineral fertiliser were chosen to represent typical farmer behaviour in the area.

2.2.1 Farm survey and data collection

Farm management and cropping system data included farm characteristics, crop production and management, farm inputs and outputs, and animal production. They were collected using a structured questionnaire, progressively completed during an average of two face-to-face interviews of about two hours each. Subsequently, the information was organized and stored in a Microsoft Excel© file for later calculation of the agro-environmental indicators.

Soil samples were taken from four representative fields at each farm at a depth of 0.3 m. Official Italian soil analysis methods (MIPAAF, 2000) were used to analyse sample texture, pH, organic carbon content, total N, Olsen P, and exchangeable K.

2.2.2 Farm descriptions

Table 1 reports a general description of the farms. The average UAA was 48 ha. The two organic arable farms were the smallest at 19 and 24 ha, while the other farms were more

variable. Soil textures were loam or silt-loam; other soil characteristics varied more. Organic matter content was higher in livestock organic farms, followed by arable organic farms. The other arable farms were the lowest, except for one conventional farm that had a previous presence of permanent grassland. Total N content nearly tracked the organic matter trend as C/N ratio did not show a large variability. Olsen P levels were high in all the farms, and homogeneous among the groups. Exchangeable K was usually low.

Table 2 presents the crops and their yields of each farm. As expected for the area, the major crops were maize and winter cereals, followed by soybean. In addition to these crops, organic farms also included various legumes (mostly lucerne) in their crop rotation. Meadows and other forages were present on organic farms only.

The organic livestock farms bred 120 and 89 Livestock Units (LSU) with stocking rates of 3.4 and 1.7 LSU ha⁻¹, respectively. Manure was managed by a permanent litter made of barley straw and maize stalk residue. Manure was spread mainly inside the farms, but farmyard manure quotas of 22% and 13%, respectively, were still exported to neighbouring farms. The spread manure limit of 170 kg N ha⁻¹ was accomplished on both farms.

2.3 Application of agro-environmental indicators

The selected farms were analysed using the set of 22 indicators derived from literature and reported in Figure 1. Those selected, according to the DPSIR schema (Kristensen, 2004; EEA, 2005), can be classified as driving force or pressure indicators. Driving force indicators represent system inputs related to land use planning, agricultural managements, chemical, and energy; pressure indicators represent the result of these practices and are usually not directly modifiable by farmers. Oenema et al. (2011) considers the AEI “soil cover” indicator a driving force indicator, however, we considered it a pressure indicator to recognize that

farmers actively select the number and type of crops to grow based on economic strategies rather than on simply covering the soil for longer.

Figure 1 makes clear how driving forces and pressures relate. Some pressure indicators (soil cover, fertiliser, and pesticide indicators) relate to just one or few driving force indicators, while others (gaseous emissions and energy indicators) relate to most. Separating driving force indicators and pressure indicators allows analysis of the critical points of cropping systems and makes evident the agricultural managements that cause the pressures.

Indicators were selected to evaluate the agro-environmental sustainability of cropping system managements from five aspects (land use, fertiliser use, pesticide use, energy use and gaseous emissions). To each aspect corresponded a group of indicators. The different indicators, with the exception of *Number of practices*, derived from the literature and international methodologies. Most came from Agro-Environmental indicators (AEI) that have been defined in Communication COM (2006) 508 of the European Commission (EC, 2006). *Number of Crops*, *Tillage Practices*, and *Irrigation* were directly calculated at the cropping system scale, while all others were calculated at the crop level and related to the cropping system scale using a weighted average based on the surface of each crop.

Pressure indicators were calculated using standardized international methodologies and were not directly measured. Pressure indicators have the advantage that they are based on information easily collectable from farm interviews and official databases. The relationship between indicator results and effective impact on the system is described in the cited literature.

The majority of indicators represented system inputs (driving force indicators) or system impacts (pressure indicators), and therefore, results were generally considered to have lower sustainability when their values were high. However, *Number of Crops*, *Number of Practices*,

Soil Cover, Gross nutrient balances, Net Energy, and Energy Use Efficiency have different interpretations, which have been detailed in the specific section.

Due to the large pedological and climatic variability that affects crop production, indicators results were presented only per unit of surface and not per unit of production.

2.3.1 Driving force indicators

2.3.1.1 Land use

Three indicators comprise the Land use driving forces group: *Number of Crops*, *Tillage Practices*, and *Irrigation*, all of which were derived from AEI indicators (Oenema et al., 2011). *Number of Crops* defines the number of different species cultivated without regard to final use (grain, silage, green forage, or hay). It indicates the structural biodiversity of a cropping system. *Number of Crops* indicator show higher sustainability when values are high. *Tillage Practices* highlights the different practices applied on a farm, and is calculated as the percentage of the UAA cultivated with conventional practices. *Irrigation* does not consider the potential irrigable land; rather, it indirectly measures water consumption as the percentage of the UAA that is effectively irrigated.

2.3.1.2 Fertiliser use

Five indicators belong to the Fertiliser use driving forces group: *Mineral fertilisers*, *Organic fertilisers*, *N fertilisers*, *P fertilisers*, and *K fertilisers*. All five were derived from the AEI fertiliser consumption indicator (Oenema et al., 2011), and each was calculated as the total amount of fertiliser or nutrient applied to a hectare (kg ha^{-1}). The nutrient quantities applied through farmyard manure were calculated using a mass balance (Amon et al., 2011) that considered feed and litter nutrient content as inputs and nutrients exiting the system via pathways other than excreta as outputs.

248

249 2.3.1.3 Pesticides use

250 The two indicators in the Pesticides use driving forces group are *Consumption of Pesticides*
251 and *Equivalent Treatment*. The former, an AEI indicator (Oenema et al., 2011) is the total
252 active ingredient quantity applied to a hectare (kg ha^{-1}), while the latter is the number of
253 average treatments used and is quantified as the ratio between actual applied pesticide
254 quantity and average quantity suggested by the manufacturer (Dennis et al., 2010).

255

256 2.3.1.4 Energy use

257 The two indicators that belong to the Energy use driving forces group include *Number of*
258 *Practices* (not reported in the literature) and *Energy Input* (Alluvione et al., 2011), which
259 corresponds to the AEI indicator Energy Use as defined by Oenema et al. (2011). *Number of*
260 *Practices* equals the number of tillage, sowing, fertilisation, weeding, ridging, irrigation,
261 harvesting, silaging, and drying events performed per crop. Each operation counts as a unit
262 regardless of the time or energy consumed. *Energy Input* (EI) is the sum of direct and indirect
263 energy inputs. Fertilisers, pesticides, seeds, diesel, and lubricant constitute direct energy
264 inputs, while indirect energy inputs are those used to produce, package, and transport the
265 direct inputs and energy embedded in farm machinery. Notably absent from the EI are
266 environmental and labour inputs (Alluvione et al., 2011).

267 All energy inputs, both direct and indirect, were calculated through mass flow and determined
268 by multiplying inputs by the equivalent energy shown in Table 3, that represents the energy
269 embedded in each product. The value for fertiliser energy input was computed by multiplying
270 various N forms, P, and K quantities by their specific energy equivalent, and then the product-
271 specific Formulation Packaging Transport coefficient (FPT) was added. Manure has no
272 energy equivalent because it is a livestock farming by-product. Pesticide energy input was

determined by multiplying the quantity of each active ingredient by its specific energy equivalent (Green, 1987), and then adding the pesticide FPT coefficient. Average herbicide, fungicide, and insecticide energy values were employed when necessary. Seed energy equivalents included the energy required for selecting, packaging, and transporting the seeds. Fuel energy input values were based on farmer reported diesel consumption; total lubricant energy (direct + indirect) and machine-embedded energy were considered to be proportional to diesel consumption. Table 4 lists the maximum and minimum values for each practice.

2.3.2 Pressure indicators

2.3.2.1 Land use

Soil Cover was the land use pressure indicator used in the present study. It is from the AEI indicator set (Oenema et al., 2011), and when combined with *Tillage Practices* (AEI), can be used to evaluate soil erosion risk (Bockstaller et al., 1997; Vereijken, 1995; Castoldi and Bechini, 2006). *Soil Cover* (SC) is defined as the number of days (expressed as year percentage) during which the crop is present. High values (long soil coverage period) equate to more system sustainability.

2.3.2.2 Fertiliser use

Three indicators belong to the fertiliser use pressure group: *Gross N Balance* (GNB), *Gross P Balance* (GPB), and *Gross K Balance* (GKB). GNB and GPB were calculated according to AEI indicators (Oenema et al., 2011); GKB was calculated following Bassanino et al. (2011).

The gross nutrient balances were calculated as:

$$\text{GNB, GPB and GKB} = F_c + F_o + A_d + B_{fx} + S_e - \text{Off}$$

where F_c was the mineral fertiliser nutrient supply, F_o was the organic fertiliser nutrient supply, A_d was the N and P atmospheric depositions, B_{fx} was the biological nitrogen fixation

by legumes, Se was the seeds nutrient content, and Off was the crop nutrient off-take. The values utilised for nutrient content both in crops and seeds are shown in Table 5. The values used for atmospheric deposition were $26 \text{ kg N ha}^{-1} \text{ y}^{-1}$ (Bassanino et al., 2011) and $1.8 \text{ kg P ha}^{-1} \text{ y}^{-1}$ (study area value, Experimental Centre, University of Turin). The legume fixation value was calculated as:

$$Bfx = Off - (Fc + Fo + Ad + Se)$$

on pure legume crops (soybean, lucerne, beans) (Bassanino et al., 2007; Grignani et al., 2003); in meadows and permanent grassland (composed of grasses and legumes), the N fixed value considered was 40 kg N ha^{-1} (Regione Piemonte, 2009). This assumption derives from the simplified ideas that these crops tend to use N from fertilisers, before fixing atmospheric N (Meisinger and Randall, 1991) and that their balance is equal to zero (Bassanino et al., 2007).

Gross nutrient balances were difficult to evaluate for agro-environmental sustainability as they could result in either positive or negative values. Although the surplus of gross nutrient balances includes potential soil immobilisation, they also indicate nutrient loss potential due to gaseous emissions, leaching, and run-off. Therefore, a higher surplus suggests higher losses and higher environmental impact. On the contrary, negative values or deficits, imply nutrient use from immobilised soil pools, potentially leading to a depletion of soil nutrients. In summary, gross nutrient balances were considered “better” when closer to zero and “worse” when high (absolute value), as it would imply greater losses or soil depletion.

2.3.2.3 Pesticide use

The two indicators in the pesticides use pressure group are *Load Index* and *Environmental Impact Quotient*. The *Load Index* (LI) (Bechini and Castoldi, 2009; OECD, 2004) indicates potential effect on a non-target organism class. It is calculated by dividing the application rate

by the LD50 or the LC50 of each active ingredient. The *Environmental Impact Quotient* (EIQ) value (Kovach et al., 1992) is more complex to calculate as it takes into account active ingredient properties and analyses the potential impact on three different components: farmers, consumers, and environment. The present work used the active ingredient properties defined by the Pesticide Property Database (University of Hertfordshire, 2012) and the Italian Ministry of Agriculture database (MIPAAF, 2012).

2.3.2.4 Energy use

Two indicators in the energy use pressure group are *Net Energy* and *Energy Use Efficiency*. *Net Energy* (NE) and *Energy Use Efficiency* (EUE) indicators (Alluvione et al., 2011) allow evaluation of energy output as well as the relationship between yield and plant production energy used. The data needed to calculate these indicators are energy input and energy output. *Net Energy* is the difference between energy output and energy input, while *Energy Use Efficiency* is the ratio between energy output and energy input. Inputs were determined per the *Energy Input* indicator described earlier. Energy outputs were defined as the gross energy contained in crops and residues removed from the field (Table 3). The *Net Energy* represents the amount of energy gained per unit of area, while the *Energy Use Efficiency* represents the energy gained per unit of energy input. Therefore, larger values correspond to lower impact.

2.3.2.5 Gaseous emissions

Ammonia Emission and *GHG Emission* are the two indicators in the Gaseous emissions pressure group. According to AEI (Oenema et al., 2011), the methodologies used for gaseous emissions are those internationally recognized by law. These methodologies are EMEP/EEA for ammonia (EEA, 2009) and IPCC for greenhouse gases (GHG) (IPCC, 2006).

A Tier 2 approach (EMEP/EEA methodology) was used for mineral and organic fertiliser calculations of the *Ammonia Emissions* (AE) indicator. The mineral fertiliser calculation relies on the average spring temperature, which was 17.2°C computed according to the methodology. It was obtained from 10 years of data measured at the Experimental Center of the University of Turin in Carmagnola (TO).

The Tier 2 methodology for organic fertiliser addresses three different NH₃ loss phases: housing, storage, and spreading. As this paper focuses on only cropping systems, ammonia emissions during housing and storage were not considered. The amount of nitrogen available for spreading was calculated as N excreted minus N lost during housing and storage. N losses during housing and storage were calculated by the EMEP methodology, while N₂O losses during manure storage were calculated using the IPCC methodology (IPCC 2006), adjusted with a localized EF value of 0.02 for cattle solid manure (ISPRA, 2011).

In the case of imported manure, only the spreading phase was considered. The cattle solid manure total ammonia nitrogen (TAN) used for calculation was 20% (CRPA, 1993).

GHG Emissions were calculated per the IPCC methodology (IPCC 2006) and expressed as CO₂ equivalents. According to the methodology and without a change in land use, the emissions considered were those from diesel consumption and from direct and indirect N₂O losses from agricultural soils. Diesel fuel combustion accounts for CO₂, CH₄, and N₂O emissions. To calculate those emissions, a diesel density of 0.855 kg l⁻¹ was used (Bosch, 1996).

Direct N₂O losses consider all the nitrogen added to the system as fertiliser and as crop residues. A Tier 1 approach was applied because of a lack of specific emission factors. Indirect N₂O losses were calculated with Tier 2, applying EMEP/EEA methodologies for NH₃ and NO losses.

2.4 Data analysis

The expected grade of the different cropping systems was defined through ranking them from 1 to 4 to represent a progressive environmental sustainability from conventional (1) to livestock organic (4) cropping systems. Only for pesticides use indicators, ORG and LIV were set to 3 as in both these two cropping systems chemicals are not permitted in the same way. The association between the different agro-environmental indicators and the grade assigned to each cropping system represents the ability of the indicator to correctly grade the cropping systems and was assessed through Kendall Tau-b rank correlation (Kendall, 1938). The test was carried out using SPSS ver. 20.

To better summarise results and to underline the grading of different cropping system groups, the indicators were presented as radar graphs, one for driving force indicators and one for pressure indicators. Radar graphs were elaborated using R software ver. 2.15.1. Each axis represented an indicator. To evaluate the cropping systems in radar graph, values were presented as the average of each farm group. Each indicator was rescaled between the minimum and maximum values.

Most indicators indicated higher environmental sustainability with low values. However, some indicators had opposite meaning. Consequently, to standardise results, the *Number of Crops* and *Net Energy* indicators were multiplied by -1, and *Soil Cover* and *Energy Use Efficiency* were represented as their reciprocals. Finally, gross nutrient balances were considered as absolute values. Therefore, on the graphs, the cropping systems showing higher sustainability and lower impact occupy a smaller area.

Among driving force indicators *Mineral fertilisers* and *Organic fertilisers* were not presented in radar graph, since their results were redundant when compared to *N*, *P*, and *K fertiliser* indicators.

3 Results

3.1 Driving force indicators

Driving force indicators (Table 6) describe the cropping system characteristics through four agricultural management aspects: land use, fertiliser use, pesticides use, and energy use.

Table 7 represents the ability of each indicator to correctly grade the different cropping systems through the Kendall $\tau(b)$ correlation test.

Tillage Practices allowed to grade the different cropping systems and to differentiate organic cropping systems from the other two systems (Kendall $\tau(b)$ -0.87, $P(\tau)$ 0.006). Although *Number of Crops* presented higher values in organic cropping systems, the grading was not significant (Kendall $\tau(b)$ 0.60, $P(\tau)$ n.s.). *Irrigation* demonstrated more homogeneity between the different farm types and also in this case Kendall correlation was not significant.

Fertiliser use clearly separated organic cropping systems from the other two systems as the former used only organic fertiliser and the latter only mineral. Moreover, LIV showed higher values than ORG due to farmyard manure application, while INT showed a lower value than CONV due to RDP restrictions.

N fertilisers decreased from CONV through INT to ORG systems. Values for LIV were higher than in INT due to the greater nutrient availability from recycling internal manure. If LIV is removed from the correlation analysis, the grading of the other systems is significant (Kendall $\tau(b)$ is -0.89, $P(\tau)$ 0.017, not shown in table 7). Even though LIV2 stayed within the 170 kg ha⁻¹ organic regulation limitation, the methodology used to calculate N excreta showed nitrogen fertiliser input surpassed this limit. *P* and *K fertilisers* were higher in LIV due to tied N/P and N/K ratios and to the large amount of supplied manure. *P* and *K fertilisers* showed no trends in the other cropping systems (*P fertilisers* Kendall $\tau(b)$ 0.15, $P(\tau)$ n.s. and *K fertilisers* Kendall $\tau(b)$ 0.31, $P(\tau)$ n.s.).

Pesticides were only applied in non-organic cropping systems. *Consumption of Pesticides* highlighted the low pesticide use in INT *versus* CONV (Kendall $\tau(b)$ -0.95, $P(\tau)$ 0.004). *Equivalent Treatments* indicator was also able to grade correctly CONV and INT (Kendall $\tau(b)$ -0.86, $P(\tau)$ 0.009).

The *Number of Practices* was higher on organic farms, both for ORG and LIV, which arose primarily from the high frequency of operations required for hay production (Kendall $\tau(b)$ 0.69, $P(\tau)$ 0.022). Secondly, the presence of another crop on a portion of the UUA increased the average practice number.

Energy Input was higher in CONV and INT than in organic cropping systems (Kendall $\tau(b)$ -0.69, $P(\tau)$ 0.022). Figure 2 shows the energy inputs considered and their related values. The greatest energy inputs were those related to mechanisation and fertiliser use, followed by seed energy inputs. Pesticides showed very low values.

The rank of mechanisation energy input use were, on average, high for CONV and INT, followed by LIV, and lowest for ORG. Notably, INT1 presented a lower value than LIV. Fertiliser energy inputs were very high in CONV and INT, very low in ORG, and zero in LIV. While only a small amount of commercial organic fertiliser was used in ORG, the energy input necessary for its production was included. The absence of fertiliser energy inputs in LIV stems from its manure use considered as by-product, and consequently, requiring no energy input. Seed energy inputs were higher in INT due to an elevated wheat seed use, and highest in ORG2, in which transplanted tomato seedlings were used.

3.2 Pressure indicators

3.2.1 Land use

Soil Cover (Figure 3) was higher in organic cropping systems due to the presence of meadows, other forages, and double crops (Kendall $\tau(b)$ 0.69, $P(\tau)$ 0.022).

3.2.2 Fertiliser use

Figure 4 lists the nutrient inputs and their nutrient gross balances for N, P, and K. The main N inputs were mineral fertilisers for CONV and INT, biological fixation for ORG, and organic fertilisers for LIV. The *Gross N Balances* showed CONV had a higher surplus than the other systems due to its high input use and low off-take. The second highest surplus was found in INT (approximately 50 kg N ha⁻¹) as opposed to the low LIV values (near zero). LIV underwent higher fertiliser inputs and legume fixation, but it had a lower surplus due to more crop off-take from meadow and double crop presence. Low levels of inputs in ORG led to a negative *Gross N Balance*. In general this indicator is able to correctly grade the different cropping systems (Kendall $\tau(b)$ -0.77, $P(\tau)$ 0.011).

Gross P Balances were about zero or negative. CONV and INT presented higher variability within their groups, which made differentiation between them impossible. ORG had the most negative values due to its lower fertiliser input level. For LIV, the balances were slightly negative due to a high input of manure fertilisation. The lack of a correct grading was confirmed by Kendall correlation that was not significant.

Gross K Balances were positive for INT, negative for CONV1, and lower for CONV *versus* INT. GKB were negative for all organic cropping systems. The very low fertiliser input levels in ORG, was reflected in a very low GKB also. LIV too had a negative balance; its higher level of potassium input partially compensated the off-take. Kendall correlation was not significant, thus confirming the high variability of GKB (Kendall $\tau(b)$ 0.39, $P(\tau)$ n.s.).

3.2.3 Pesticide use

Load Index graded correctly CONV and INT (Figure 5) for each class of non-target organism (algae Kendall $\tau(b)$ -0.95, $P(\tau)$ 0.004, fishes Kendall $\tau(b)$ -0.76, $P(\tau)$ 0.021, bees Kendall $\tau(b)$ -0.86, $P(\tau)$ 0.009, earthworms Kendall $\tau(b)$ -0.76, $P(\tau)$ 0.021, mammals Kendall $\tau(b)$ -0.86,

P(τ) 0.009, birds Kendall $\tau(b)$ -0.95, P(τ) 0.004). *Load Index* trended in a like pattern on all farms for each class of non-target organism. The values were lower for birds and mammals, while the highest values were for fishes and algae. The *Environmental Impact Quotient* (EIQ) differentiated the cropping systems better, and it made evident a lower potential impact of pesticide use in INT than in CONV (Figure 6) (Kendall $\tau(b)$ -0.95, P(τ) 0.004). Analysis of the three *EIQ* components (farmers, consumers and environment) trended like total *EIQ*. INT had the lowest impact values in each. The environmental component was the most impacted; consumers were impacted the least.

3.2.4 Energy use

Figure 7 presents *Net Energy* and *Energy Use Efficiency* indicator results. *Net Energy* was higher for LIV, with values nearly double those of the other cropping systems. The incorrect grading was confirmed by a not significant Kendall correlation. *Energy Use Efficiency* resulted in similar values for CONV and INT. All organic cropping systems had higher values of *Energy Use Efficiency*, and LIV systems had the highest (Kendall $\tau(b)$ 0.72, P(τ) 0.011).

3.2.5 Gaseous emissions

In arable cropping systems, the *Ammonia Emissions* (Figure 8) indicator trended similarly to nitrogen fertiliser inputs; that is, values decreased from CONV through INT to ORG. LIV showed the highest values. Kendall correlation was not significant. Figure 9 displays *GHG Emissions* as the sum of two sources, expressed in CO₂ equivalent. The total *GHG Emissions* presented values that distinguished between cropping system groups. The highest values were in CONV; INT and LIV had similar intermediate values, and the lowest values were those calculated for ORG (Kendall $\tau(b)$ -0.62, P(τ) 0.041). N₂O emissions trended like the total *GHG emissions*.

497

498 4 Discussion

499 The grading of the cropping systems has been analysed according to the indicator groups to
500 describe the existing relationships between input levels (driving force indicators) and
501 environmental pressures (pressure indicators). Results were compared with the expected
502 grading derived from the different intensification levels as defined by regulations.

503

504 4.1 Land Use

505 Although biodiversity is an important issue that should be analysed, the majority of the crops
506 here explored were renewed each year with industrial selected seeds and therefore within
507 species diversity is not expected. Meadows are also usually renewed each 3-5 years, and only
508 in one case a small surface is permanent grassland. The analysis of within-species diversity
509 could give interesting information that completes the analysis of crop biodiversity, but the
510 level of detail required to obtain this information is beyond the aims of this works, that is to
511 analyse data collected through interviews and database.

512 The analysed farms mainly cultivated arable crops typical of the study area: maize, winter
513 cereals, and soybean (Sacco et al., 2003; Bassanino et al., 2007). Organic cropping systems,
514 however, varied their crop rotations more to include meadows, double crops, and legumes
515 (soybean, bean, lucerne). The fact that the organic systems had a larger number of crops in
516 rotation met several needs: to control pests, to increase N addition through N fixation, and to
517 grow fodder crops in the case of livestock production systems.

518 Lampkin (2002) reported that crop rotation helped control pests in organic systems, and a
519 recent review by Gomiero et al. (2011) showed that crop rotation is an effective farming
520 practice to reduce the negative impact of weeds. European regulation 834/2007/EC (EC,
521 2007) also suggests crop rotation as one preventive measure to maintain plant health.

522 Introducing legumes into the crop rotation is aimed at increasing N supply into the system as
523 crop uptake of N fixing crops is, at the least, balanced by N biological fixation. Practices such
524 as these allow systems to overcome the imposed 170 kg N ha⁻¹ limit on fertiliser use.

525 All organic systems introduce meadows into the farm area. On livestock organic farms, they
526 are necessary to feed animals; in arable organic farms, they produce hay, which can be sold to
527 livestock organic farms or exchanged for manure fertiliser. The presence of meadows (3-5
528 year duration) permits a no-till area to be present without adopting no-tillage practices.
529 Furthermore, meadows and double crops lead to longer periods of soil cover during the year.
530 These two aspects have a minor environmental impact, and result in higher sustainability
531 characteristic of organic cropping systems *versus* the less sustainable conventional and
532 integrated cropping systems.

533 The DM yields declared by organic system farmers are generally in the range of conventional
534 and integrated cropping systems. However, according to the literature (Kirchmann and Ryan,
535 2005; Eltun et al., 2002; Basset-Mens and van der Werf, 2005), organic cropping systems
536 usually produce less than conventional cropping systems, although manure fertilisation could
537 reduce the yield gap (Kirchmann and Ryan, 2005). The analysed cropping systems were
538 selected for their regional representativeness. As such, they came from a wide area
539 characterised by different pedological and climatic conditions with high production variability
540 that makes crop DM yield comparisons not feasible. Consequently, indicators were calculated
541 only per unit of surface and not per unit of production. An assessment per unit of production
542 could give additional information about the sustainability of the different systems, but
543 requires more homogeneous pedological and climatic conditions.

544 When the land use pressure indicators were employed to grade the different farms organic
545 cropping systems were shown to impact the environment less than conventional and
546 integrated cropping systems.

547

548 4.2 Fertilisers use

549 Organic cropping systems that paired manure with meadows in the crop rotation showed
550 higher soil organic matter content. Between the two organic cropping systems considered,
551 livestock systems had the highest soil organic matter values consequent to their higher
552 manure input. Similar results were observed by Bertora et al. (2009) in manure-based
553 conventional and integrated cropping systems and by Fließbach et al. (2007) in livestock-
554 based bio-organic and bio-dynamic cropping systems.

555 In livestock organic cropping systems, the manure amount applied depends on the stocking
556 rate. Per European and regional regulations, N input is calculated from stocking rate using
557 tabular data, while respecting the 170 kg N ha⁻¹ limit. However, in the present study, the real
558 amount of N supplied in the livestock organic cropping systems has been calculated using
559 nutrient mass balance, which resulted in a higher N input, even in farms that complied with
560 European regulations.

561 As livestock organic farms manure fertilisation is calibrated on N loads, P and K inputs are
562 defined by N/P and N/K ratios in manure and not on actual crop need (Bassanino et al., 2011).
563 For this reason, P and K amounts were the highest in the livestock organic cropping systems
564 (Spear et al., 2003; Bassanino et al., 2011).

565 Arable organic cropping systems used the lowest levels of fertiliser inputs not only because it
566 is difficult to retrieve manure, but also because of the high cost of organic fertiliser. On both
567 of the farms of this group, legume fixation was the main source of N, which made it essential
568 to compensate for the very low N from fertilisers.

569 Therefore, in terms of N fertiliser inputs, the farms decreased in intensity from conventional
570 to integrated to arable organic systems. Livestock organic cropping systems demonstrated an
571 input level similar to integrated cropping systems. Conventional and integrated system

differences related to fertiliser use limits defined by the RDP for integrated cropping systems. If the analysis had considered all N additions, including N from legume fixation, the trend would be altered to show the highest values for conventional and livestock organic systems, and the lowest values for integrated and arable organic systems.

Gross nutrient balances did not always trend like nutrient inputs as crop off-take introduced large differences among farms types. N balance of conventional, integrated, and arable organic systems reflected the trend of N fertiliser input. Although livestock organic systems showed higher fertiliser input with the highest input derived from legume fixation, they produced lower surpluses than did integrated and conventional systems due to large crop off-takes from meadows and double crops. Arable organic systems were the only that resulted in negative N balances.

P balances were negative for all systems, which helped to offset the large soil P content. Arable organic systems showed the most negative balances due to their low nutrient supply. Livestock systems had the highest input from their high manure fertilisation, but it failed to compensate for the high off-take from the presence of meadows.

Finally, the K balances clearly diverged between organic and non-organic cropping systems; in fact, they showed positive values only in the latter group. For the studied area, Bassanino et al. (2011) demonstrated that manure fertilisation usually balances K off-take, however, they found wide crop variances (positive balances for maize and negative balances for meadows). This variability also explains the negative K balances of livestock organic systems given their high meadow portion. Torstensson et al. (2006), who studied the nutrient use efficiencies of organic and conventional cropping systems in Sweden, found negative K balances in all cropping systems, especially in animal manure organic systems (-36 kg K ha^{-1} per year). This result was attributed to the large amounts of K taken up by forage crops.

4.3 Pesticide use

The organic farms analysed in this work did not use pesticides. *Consumption of Pesticides* allowed the farms to be graded according to their relative intensification levels, with conventional cropping systems using higher input quantities. Integrated cropping systems fell below these levels due to limits set by the RDP.

Load Index underlined the impact on non-target organisms shared across the cropping systems. In all cases, the lowest impacts were on birds and mammals, while the highest impacts were on the aquatic environment (fishes and algae). Bechini and Castoldi (2009) had also indicated that algae have the highest *Load Index* values.

The *Environmental Impact Quotient* clearly distinguished between conventional (higher values) and integrated cropping systems (lower values). Integrated system pesticide limits, introduced and monitored regionally by the RDP, have been confirmed by the IPLA (2012) to reduce the potential impact of pesticide applications. Farmer and environmental components of the indicator made evident the differences between conventional and integrated systems. The main impact was to the environment in all systems, but integrated management did severely lessen its environmental impact achieving the goal of the regulation.

4.4 Energy use

Fertilisation and mechanisation are the two main components that characterise *Energy Input* on the eight farms, in agreement with other studies (Alluvione et al., 2011; Meul et al., 2007; Fumagalli et al., 2011). According to Castoldi and Bechini (2010), cropping system energy input depends mainly on the crops in the system and their relative shares of the farm surface. The highest energy inputs correspond to maize, followed by meadows, and finally to winter cereals. Notwithstanding, energy input values are also closely linked to the fertilisation management used for each crop (Bechini and Castoldi, 2009).

In this study, *Energy Input* enabled system grading by expected intensification level. The lowest level, recorded in organic cropping systems, depended mostly on two factors — the presence of meadows and organic fertilisation that has a zero energy cost (as a by-product of breeding activity). The energy input derived from mechanisation was also low on organic farms due to the very low fuel amount required for tedding, raking, and baling forage crops, even if they used a great number of passes. In conventional and integrated cropping systems, fertiliser and mechanisation inputs differentiated the two and proved conventional cropping systems to have the highest values. Alluvione et al. (2011) demonstrated this same rank in a field experiment conducted in the same agricultural area in two cropping systems fertilised with only mineral fertilisers. Cruse et al. (2010) conducted a six-year study that compared energy use in a conventional two-year rotation system (maize and soybean) to two low input cropping systems that used more diverse crops (maize, soybean, small grains, and red clover or lucerne), manure, less fertiliser and herbicides. They found that the two low-input systems used 23% to 56% less fossil energy than did the conventional system.

The driving force indicator *Number of Practices* showed unexpected and contradictory results relative to *Energy Input*. The high number of operations associated with forage field drying in hay production yielded high *Number of Practices* for organic cropping, yet the relatively small amount of fuel consumed for each pass kept *Energy Input* low.

The two pressure indicators, *Net Energy* and *Energy Use Efficiency*, identified three different situations:

- high *Net Energy* and high *Energy Use Efficiency* in livestock organic systems;
- low *Net Energy*, but high *Energy Use Efficiency* in arable organic systems;
- low *Net Energy* and low *Energy Use Efficiency* in conventional and integrated systems.

The higher values of *Energy Use Efficiency* recorded in the organic cropping systems mainly depend on the low *Energy Inputs* that characterise these two systems. Moreover, the presence of meadows, particularly lucerne, increased energy output due to its high DM yield. Furthermore, double crops increased energy output with a small energy input. Differences between livestock and arable organic systems also related to the higher share of energy-producing meadows and silage crops in livestock systems. The lower *Net Energy* and *Energy Use Efficiency* calculated for conventional and integrated systems related to their higher *Energy Input*. The similarity of *Energy Input* and energy output in the two systems did not permit distinction between them.

4.5 Gaseous emissions

Ammonia Emissions showed a trend like that of N fertilization, but no correlation with system grading. The EMEP/EEA methodology (EEA, 2009) explains that mineral fertiliser ammonia comes from urea that has emission values similar to manure, whereas ammonia emissions from other mineral fertilisers are lower. Livestock farms had the highest emission values due to their exclusive use of manure (high emission factor).

GHG Emissions correctly graded the cropping systems. The highest ranked system was conventional and the lowest ranked was arable organic; livestock organic systems ranked in the middle of the two. Although livestock and integrated systems had similar N fertilisation values, livestock systems yielded lower emission values due to the presence of meadows that are characterized by lower crop residue.

4.6 Radar

Radar graphs described and made evident the impact of the different cropping systems (Bockstaller et al., 1997; Sattler et al., 2010). They made it easy to understand how driving

671 forces and their consequent pressures determine the grade of the different systems. Figure 10
672 shows two radar graphs, one for driving force indicators and one for pressure indicators. The
673 indicator *Irrigation* varied highly among and within the cropping system groups because of
674 differing pedological and climatic conditions; at the same time, it is unaffected by the
675 different intensification levels. As it was unable to differentiate farms based on their intensity
676 levels, it was excluded from the graph. Among pressure indicators, *Load Index* values were
677 not represented as *Environmental Impact Quotient* better graded the different farms.

678 The two radar graphs show that the grade of the driving forces is not necessarily reflective of
679 the grade of the pressures. The graphs underscore that most indicators graded the cropping
680 systems according to their defined intensification level. However, some indicator groups
681 graded the cropping systems differently in driving force and pressure graphs.

682 Conventional systems demonstrated the worst grade of driving force indicators, while the best
683 were related to arable organic cropping systems. Analysing the graph as a whole, it is not
684 possible to clearly discriminate integrated from livestock organic systems as both presented
685 intermediate values, although they graded differently on single indicators. Conventional
686 cropping systems presented the highest impact from pressure indicators, followed by
687 integrated cropping systems. In this case, arable organic cropping systems presented higher
688 impact than did livestock systems. This suggests that pressure indicators reflected
689 intensification grading better than driving force indicators.

690 The agro-environmental indicator set analyses underlined two main correlated factors, which
691 allowed differentiation of intensification levels among the cropping systems. First, legal input
692 limits and management practices do reflect on intensification levels. Second, organic
693 production regulations that defined management practices, in particular the presence of
694 meadows and use of organic fertilisers, do influence those systems, and could similarly

influence the environmental performance of other cropping systems if practiced. This potential calls for evaluation on how to improve regulations to increase system sustainability. The goal to design and develop usable tools to assess the environmental impact of agricultural policy has grown in recent years. Improvements in agro-environmental policy evaluation standards, direct support schemes, and recommendations from the Common Monitoring and Evaluation Framework of the European Commission, which requires Member States to assess the impacts of their RDP (Schuh et al., 2011), have converged to focus on the same goal. Member States often use routine administrative data to monitor the effectiveness of agro-environmental measures, but this often does not reliably measure the environmental impacts of the policy. Adoption of agro-environmental measures does not guarantee that environmental standards will be attained (Mauchline et al., 2012).

The indicator set in this research was selected to allow comparison and grading of farm management intensities in order to assess environmental pressures and to inform decision- and policy-makers on how to manage, implement, and evaluate *ex post* agro-environmental measures and policy impacts. Following the recommendations of Bechini and Castoldi (2009), who suggested that indicators be simple, synthetic, and derived from data that can be easily obtained, input variables for the calculation of selected indicators should be collectable in farm interviews by questionnaire and/or data should be obtainable from official farm databases, thus coupling scientific soundness with cost-effectiveness of the process.

5 Conclusion

The result of this study showed that the indicator set presented was mostly able to correctly grade cropping system intensification levels, and that it could evaluate their agro-environmental sustainability. However, in some cases, the expected grade did not result. This work showed that this is not due to indicator fault, but rather that some analysed variables did

not reflect the intensification expected. This phenomenon happened mainly for driving force indicators.

The analysis also showed that higher input levels do not always reflect higher environmental pressure. Therefore, outside ethical aspects that are not in the aim of this work, regulations should be preferable based on pressure indicator thresholds instead of on system inputs.

Acknowledgments

The authors want to thank Simona Avagnina, Aurelio Del Vecchio, Giancarlo Bourlot, and Carlo Grignani for their contribution, and Joan Leonard for the English revision of the manuscript. This work has been funded by Regione Piemonte, Assessorato all'Agricoltura e Foreste e alla Caccia e Pesca through its MITANet project.

References

Alluvione, F., Moretti, B., Sacco, D., Grignani, C., 2011. EUE (energy use efficiency) of cropping systems for a sustainable agriculture. *Energy* 36, 4468-4481.

Amon, B., Hutchings, N., Piertrzak, S., Vinther, F.P., Nielsen, P.K., Poulsen, H.D., Kristensen, I.S., 2011. Analysis of methodologies for calculating greenhouse gas and ammonia emissions and nutrient balances reporting. In: Selenius, J., Baudouin, L., Kremer, A.M. (Eds.), *Eurostat methodologies and working papers*. Publications Office of the European Union, Luxembourg.

Bassanino, M., Grignani, C., Sacco, D., Allisiardi, E., 2007. Nitrogen balances at the crop and farm-gate scale in livestock farms in Italy. *Agric., Ecosyst. Environ.* 122, 282-294.

745

746 Bassanino, M., Sacco, D., Zavattaro, L., Grignani, C., 2011. Nutrient balance as a
747 sustainability indicator of different agro-environments in Italy. *Ecol. Indic.* 11, 715-723.

748

749 Basset-Mens, C., van der Werf, H.M.G., 2005. Scenario-based environmental assessment of
750 farming systems: the case of pig production in France. *Agric., Ecosyst. Environ.* 105, 127-
751 144.

752

753 Bechini, L., Castoldi, N., 2009. On-farm monitoring of economic and environmental
754 performances of cropping systems: Results of a 2-year study at the field scale in northern
755 Italy. *Ecol. Indic.* 9, 1096-1113.

756

757 Bertora, C., Zavattaro, L., Sacco, D., Monaco, S., Grignani, C., 2009. Soil organic matter
758 dynamics and losses in manured maize-based forage systems. *Eur. J. Agron.* 30, 177-186.

759

760 Bockstaller, C., Girardin, P., van der Werf, H.M.G., 1997. Use of agro-ecological indicators
761 for the evaluation of farming systems. *Eur J. Agron.* 7, 261-270.

762

763 Bosch, R., 1996. *Automotive handbook*. 4th ed. Bentley Publishers, Cambridge (MA).

764

765 Canakci, M., Topakci, M., Akinci, I., Ozmerzi, A., 2005. Energy use pattern of some field
766 crops and vegetable production: Case study for Antalya Region, Turkey. *Energy Conv.*
767 *Manag.* 46, 655-666.

768

769 Castoldi, N., Bechini L., 2006. Agro-ecological indicators of field-farming systems
 770 sustainability. I. Energy, landscape and soil management. Italian J Agrometeorol. 1, 19-31.
 771

772 Castoldi, N., Bechini, L., 2010. Integrated sustainability assessment of cropping systems with
 773 agro-ecological and economic indicators in northern Italy. Eur. J. Agron. 32, 59-72.
 774

775 CRPA (Centro Ricerche Produzioni Animali), 1993. Manuale per la gestione e l'utilizzazione
 776 agronomica dei reflui zootecnici. Assessorato agricoltura della Regione Emilia Romagna,
 777 Reggio Emilia.
 778

779 Council of the European Communities, 1992. Council Directive 92/43/EEC of 21 May 1992
 780 on the conservation of natural habitats and of wild fauna and flora. Brussels, Belgium.
 781

782 Cruse, M.J., Liebman, M., Raj Raman, D., Wiedenhoef, M., 2010. Fossil energy use in
 783 conventional and low-external-input cropping systems. Agron. J. 102, 934-941.
 784

785 Dalgaard, T., Halberg, N., Porter, J.R., 2001. A model for fossil energy use in Danish
 786 agriculture used to compare organic and conventional farming. Agric., Ecosyst. Environ 87,
 787 51–65.
 788

789 Dennis, P., Herzog, F., Jeanneret, P., 2010. Selection and field validation of candidate
 790 biodiversity indicators, including field manual. Handbook for testing candidate indicators of
 791 organic/low-input farming and biodiversity.
 792 <http://www.biobio-indicator.org> Accessed December 2012
 793

794 EC (European Commission), 2000. Directive 2000/60/EC of the European Parliament and of
795 the Council of 23 October 2000 establishing a framework for Community action in the field
796 of water policy. Brussels, Belgium.

797

798 EC (European Commission), 2006. COM(2006) 508. Development of agri-environmental
799 indicators for monitoring the integration of environmental concerns into the common
800 agricultural policy. Brussels, Belgium.

801

802 EC (European Commission), 2007. Council Regulation (EC) No 834/2007 of 28 June 2007 on
803 organic production and labelling of organic products and repealing Regulation (EEC) No
804 2092/91. Brussels, Belgium.

805

806 EC (European Commission), 2008. Commission Regulation (EC) No 889/2008 of 5
807 September 2008 laying down detailed rules for the implementation of Council Regulation
808 (EC) No 834/2007 on organic production and labelling of organic products with regard to
809 organic production, labelling and control. Brussels, Belgium.

810

811 EC (European Commission), 2009a. Council regulation (EC) No 73/2009 of 19 January 2009
812 establishing common rules for direct support schemes for farmers under the common
813 agricultural policy and establishing certain support schemes for farmers, amending
814 Regulations (EC) No 1290/2005, (EC) No 247/2006, (EC) No 378/2007 and repealing
815 Regulation (EC) No 1782/2003. Brussels, Belgium.

816

817 EC (European Commission), 2009b. Directive 2009/128/EC of the European Parliament and
818 of the Council of 21 October 2009 establishing a framework for Community action to achieve
819 the sustainable use of pesticides. Brussels, Belgium.

820

821 EC (European Commission), 2009c. Directive 2009/147/ of the European Parliament and of
822 the Council of 30 November 2009 on the conservation of wild birds. Brussels, Belgium.

823

824 EEA (European Environment Agency), 2005. Agriculture and environment in EU-15 – the
825 IRENA indicator report. EEA Report No 6/2005. European Environment Agency,
826 Copenhagen, Denmark.

827

828 EEA (European Environment Agency), 2009. EMEP/EEA air pollutant emission inventory
829 guidebook 2009. European Environment Agency, Copenhagen, Denmark.

830

831 Eltun, R., Korsæth, A., Nordheim, O., 2002. A comparison of environmental, soil fertility,
832 yield, and economical effects in six cropping systems based on an 8-year experiment in
833 Norway. Agric. Ecosyst. Environ. 90, 155-168.

834

835 FAO, 2012. FAO Agri-environmental statistics website.
836 <http://www.fao.org/economic/ess/ess-agri/en/>. Accessed January 2012.

837

838 Flessa, H., Ruser, R., Dörsch, P., Kamp, T., Jimenez, M.A., Munch, J.C., Beese, F., 2002.
839 Integrated evaluation of greenhouse gas emissions (CO₂, CH₄, N₂O) from two farming
840 systems in southern Germany. Agric. Ecosyst. Environ. 91, 175-189.

841

842 Fließbach, A., Oberholzer, H.R., Gunst, L., Mäder, P., 2007. Soil organic matter and
843 biological soil quality indicators after 21 years of organic and conventional farming. *Agric.*
844 *Ecosyst. Environ.* 118, 273-284.

845

846 Fluck, R., 1992. Energy of Agriculture Products. In: Stout, B.A. (editor-in-chief), *The Energy*
847 *in world agricultural.* Agricultural Engineering Department, Texas A & M University College
848 station, U.S.A. pp. 39.

849

850 Fumagalli, M., Acutis, M., Mazzetto, F., Vidotto, F., Sali, G., Bechini, L., 2011. An analysis
851 of agricultural sustainability of cropping systems in arable and dairy farms in an intensively
852 cultivated plain. *Eur. J. Agron.* 34, 71-82.

853

854 Gomiero, T., Pimentel, D., Paoletti, M.G., 2011. Environmental impact of different
855 agricultural management practices: conventional vs. organic agriculture. *Crit. Rev. Plant Sci.*
856 30, 95-124.

857

858 Green, M.B., 1987. Energy in pesticide manufacture, distribution and use. In: Helsen, Z.R.
859 (editor), *Energy in plant nutrition and pest control.* Elsevier, Amsterdam.

860

861 Grignani, C., Bassanino, M., Sacco, D., Zavattaro, L., 2003. Il bilancio degli elementi nutritivi
862 per la redazione del piano di concimazione. *Riv. di Agron.* 37, 155-172.

863

864 Halberg, N., van der Werf, H.M.G., Basset-Mens, C., Dalgaard, R., de Boer, I.J.M., 2005.
865 Environmental assessment tools for the evaluation and improvement of European livestock
866 production systems. *Livest. Prod. Sci.* 96, 33-50.

867

868 IPCC (Intergovernmental Panel on Climate Change), 2006. IPCC Guidelines for National
869 Greenhouse Gas Inventories. Prepared by the National Greenhouse Gas Inventories
870 Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). IGES,
871 Japan.

872

873 IPLA (Istituto per le Piante da Legno e l'Ambiente), 2012. Riduzione della potenziale
874 contaminazione di acque e suoli da input agricoli. Programma di Sviluppo Rurale 2007-2013:
875 monitoraggio degli indicatori agricoli, forestali e ambientali finalizzato alla valutazione in
876 itinere, assistenza tecnica all'autorità di gestione. Direzione 11, Agricoltura. Regione
877 Piemonte.

878

879 ISO 2006. Norma ISO 14040:2006. Environmental management. Life cycle assessment.
880 Principles and framework. ISO, Geneva, Switzerland.

881

882 ISPRA (Istituto Superiore per la Protezione e la Ricerca Ambientale), 2011. Italia Greenhouse
883 Gas Inventory 1990 – 2009. National Inventory Report. ISPRA, Rome.

884

885 Kendall, M.G., 1938. A new measure of rank correlation. *Biometrika* 30, 81-93.

886

887 Kirchmann, H., Bergström, L., 2001. Do organic farming practices reduce nitrate leaching?.
888 *Commu. Soil Sci. Plant Anal.* 32, 997-1028.

889

890 Kirchmann, H., Ryan, M.H., 2005. Nutrient exclusivity in organic farming – Does it offer
891 advantages? *Better Crops* 89, 24-27.

892

893 Kongshaug, G., 1998. Energy consumptions and greenhouse gas emissions in fertiliser
894 production. In: International Fertilizer Industry Association technical conference, Marrakech,
895 Morocco, September 28–October.

896

897 Kovach, J., Petzoldt, C., Degnil, J., Tette, J., 1992. A method to measure the environmental
898 impact of pesticides. New York's Food and Life Sciences Bulletin, 139.

899

900 Kramer, S.B., Reganold, J.P., Glover, J.D., Bohannan, B.J.M., Mooney, H.A., 2006. Reduced
901 nitrate leaching and enhanced denitrifier activity and efficiency in organically fertilized soils.
902 PNAS 103, 4522-4527.

903

904 Kristensen, P., 2004. The DPSIR Framework. Paper presented at the 27-29 September 2004
905 workshop on a comprehensive / detailed assessment of the vulnerability of water resources to
906 environmental change in Africa using river basin approach. UNEP Headquarters, Nairobi,
907 Kenya.

908

909 Lampkin, N., 2002. Organic Farming. (Revised edition). Old Pond Publishing, Suffolk, UK.

910

911 Liebman, M., Gibson, L.R., Sundberg, D.N., Heggenstaller, A.H., Westerman, P.R., Chase,
912 C.A., Hartzler, R.G., Menalled, F.D., Davis, A.S., Dixon, P.M., 2008. Agronomic and
913 economic performance characteristics of conventional and low-external-input cropping
914 systems in the central corn belt. Agron. J. 100, 600-610.

915

916 Liu, B., Tu, C., Hu, S., Gumpertz, M., Ristaino, J.B., 2007. Effect of organic, sustainable,
 917 and conventional management strategies in grower fields on soil physical, chemical, and
 918 biological factors and the incidence of Southern blight. *Appl. Soil Ecol.* 37, 202-214.

919

920 Mauchline, A.L., Mortimer, S.R., Park, J.R., Finn, J.A., Haysom, K., Westbury, D.B., Purvis,
 921 G., Louwagie, G., Northey, G., Primdahl, J., Vejre, H., Kristensen, L.S., Teilmann, K.V.,
 922 Vesterager, J.P., Knickel, K., Kasperczyk, N., Balázs, K., Podmaniczky, L., Vlahos, G.,
 923 Christopoulos, S., Kroger, L., Aakkula, J., Yli-Viikari, A., 2012. Environmental evaluation of
 924 agri-environment schemes using participatory approaches: Experiences of testing the agri-
 925 environmental footprint index. *Land Use Policy* 29, 317-328.

926

927 Meisinger, J.J., Randall, G.W., 1991. Estimating nitrogen budgets for soil–crop systems. In:
 928 Follett, R.F., Kenney, D.R., Cruse, R.M. (Eds.), *Managing Nitrogen for Groundwater Quality*
 929 *and Farm Profitability*. Soil Science Society of America, Madison, WI, pp. 85–124.

930

931 Meul, M., Nevens, F., Reheul, D., Hofman, G., 2007. Energy use efficiency of specialised
 932 dairy, arable and pig farms in Flanders. *Agric. Ecosyst. Environ.* 119, 135-144.

933

934 Michos, M.C., Mamolos, A.P., Menexes, G.C., Tsatsarelis, C.A., Tsirakoglou, V.M.,
 935 Kalburtji, K.L., 2012. Energy inputs, outputs and greenhouse gas emissions in organic,
 936 integrated and conventional peach orchards. *Ecol. Indic.* 13, 22-28.

937

938 MIPAAF (Ministero delle politiche agricole alimentari e forestali), 2000. *Metodi di analisi*
 939 *chimica del suolo* (2 version). Franco Angeli Editore, Milano.

940

941 MIPAAF (Ministero delle politiche agricole alimentari e forestali), 2012. Website Fitofarmaci
 942 e Sostanze Attive - Centro Ricerca Per la Patologia Vegetale. <http://www.sian.it/fitovis/>
 943 Accessed December 2012.

944

945 Mitchell, G., May, A., McDonald, A., 1995. PICABUE: a methodological framework for the
 946 development of indicators of sustainable development. *Int. J. Sust. Dev. World Ecol.* 2, 104-
 947 123.

948

949 Morris, C., Winter, M., 1999. Integrated farming systems: the third way for European
 950 agriculture? *Land Use Policy* 16, 193-205.

951

952 Nemecek, T., Huguenin-Elie, O., Dubois, D., Gaillard, G., Schaller, B., Chervet, A., 2011.
 953 Life cycle assessment of Swiss farming systems: II. Extensive and intensive production.
 954 *Agric. Syst.* 104, 233-245.

955

956 Nielsen, A.H., Kristensen, I.S., 2005. Nitrogen and phosphorus surpluses on Danish dairy and
 957 pig farms in relation to farm characteristics. *Livest. Prod. Sci.* 96, 97-107.

958

959 OECD (Organisation for Economic Co-operation and Development), 2004. Summary Report
 960 of the OECD Project on Pesticide Terrestrial Risk Indicators (TERI). OECD Publications,
 961 Paris, France.

962

963 OECD (Organisation for Economic Co-operation and Development), 1999. Environmental
 964 indicators for agriculture. Volume 1 Concepts and framework.

965

966 Oenema, O., Amon, B., van Beek, C., Hutchings, N., Perez-Soba, M., Procter, C., Pietrzak,
967 S., Velthof, G.L., Vinther, F., Wilson, L., 2011. Farm data needed for Agri-environmental
968 reporting. In: Selenius, J., Baudouin, L., Kremer, A.M. (Eds.), Eurostat methodologies and
969 working papers. Publications Office of the European Union, Luxembourg.

970

971 Oenema, O., Kros, H., de Vries, W., 2003. Approaches and uncertainties in nutrient budgets:
972 Implications for nutrient management and environmental policies. *Eur J. Agron.* 20, 3-16.

973

974 Pimentel, D., Hepperly, P., Hanson, J., Douds, D., Seidel, R., 2005. Environmental, energetic,
975 and economic comparisons of organic and conventional farming systems. *BioScience* 55,
976 573-582.

977

978 University of Hertfordshire, 2012. The Pesticide Properties Database (PPDB) developed by
979 the Agriculture & Environment Research Unit (AERU), University of Hertfordshire, 2006-
980 2012
981 <http://sitem.herts.ac.uk/aeru/footprint/index2.htm> Accessed December 2012

982

983 Pointereau, P., Bochu, J.L., Doublet, S., 2012. Characterization and elements for a definition
984 and an analysis of low input farming system in EU-27.
985 http://www.biobio-indicator.org/publications/Pointereau_2012.pdf Accessed October 2013.

986

987 Rees, W.E., 2000. Eco-footprint analysis: merits and brickbats. *Ecol Econ.* 32, 371-374.

988

989 Regione Piemonte, 2009. Disposizioni attuative del Regolamento Regionale 29 ottobre 2007,
 990 n. 10/R. Indicazioni operative per la redazione del Piano di utilizzazione agronomica. DGR
 991 12/10/2009, n° 30-12335. Regione Piemonte, Torino, Italy.
 992
 993 Regione Piemonte, 2010. Anagrafe Agricola Unica del Piemonte.
 994 [http://www.ruparpiemonte.it/cms/servizi-rupar/servizio/12-anagrafe-agricola-unica-del-](http://www.ruparpiemonte.it/cms/servizi-rupar/servizio/12-anagrafe-agricola-unica-del-piemonte-.html)
 995 [piemonte-.html](http://www.ruparpiemonte.it/cms/servizi-rupar/servizio/12-anagrafe-agricola-unica-del-piemonte-.html). Accessed December 2012.
 996
 997 Sacco, D., Bassanino, M., Grignani, C., 2003. Developing a regional agronomic information
 998 system for estimating nutrient balances at a larger scale. *Eur. J. Agron.* 20, 199-210.
 999
 1000 Sattler, C., Nagel, U.J., Werner, A., Zander, P., 2010. Integrated assessment of agricultural
 1001 production practices to enhance sustainable development in agricultural landscapes. *Ecol.*
 1002 *Indic.* 10, 49-61.
 1003
 1004 Schröder, J.J., Neeteson, J.J., 2008. Nutrient management regulations in The Netherlands.
 1005 *Geoderma* 144, 418-425.
 1006
 1007 Schröder, J.J., van Asperen, P., van Dongen, G.J.M., Wijnands, F.G., 1996. Nutrient surpluses
 1008 on integrated arable farms. *Eur. J. Agron.* 5, 181-191.
 1009
 1010 Schuh, B., Lukesch, R., Michalek, J., Kaufmann, P., Pufahl, A., Schiller, S., Koorberg, P.,
 1011 Beaufoy, G., Pinay, G., Moran, D., Gomann, H., Storti, D., Rossi, P., Paracchini, M.L., 2011.

1012 Approaches for assessing the impacts of the Rural Development Programmes in the context of
 1013 multiple intervening factors.
 1014 <http://ageconsearch.umn.edu/bitstream/99365/2/schuhlukeschmichalek.pdf> Accessed October
 1015 2013.
 1016
 1017 Spears, R.A., Young, A.J., Kohn, R.A., 2003. Whole-farm phosphorus balance on western
 1018 dairy farms. *Journal of Dairy Science* 86, 688-695
 1019
 1020 Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural
 1021 sustainability and intensive production practices. *Nature* 418, 671-677.
 1022
 1023 Torstensson, G., Aronsson, H., Bergström, L., 2006. Nutrient use efficiencies and leaching of
 1024 organic and conventional cropping systems in Sweden. *Agron. J.* 98, 603-615.
 1025
 1026 van der Werf, H.M.G., Tzivilakis, J., Lewis, K., Basset-Mens, C., 2007. Environmental
 1027 impacts of farm scenarios according to five different methods. *Agric. Ecosyst. Environ.* 118,
 1028 327-338.
 1029
 1030 Vereijken, P., Wijnands, F., Stol, W., 1995. Progress Report 2. Designing and Testing
 1031 Prototypes. Progress reports of the research network on integrated and ecological arable
 1032 farming systems for EU and associated countries (Concerted action AIR 3 -CT920755). AB-
 1033 DLO, Wageningen, The Netherlands.
 1034

Tables

Table 1 – Surface and soil characteristic of the eight farms.

	State indicator	CONV1	CONV2	INT1	INT2	ORG1	ORG2	LIV1	LIV2
Surface	UAA (ha)	36	84	50	83	24	19	35	54
Soil quality	Texture	silt loam	loam	silt loam	silt loam	silt loam	silt loam	loam	loam
	pH	6.6	5.9	6.3	5.6	8.3	6.1	5.9	6.9
	Organic matter (%)	3.1	1.7	1.2	1.9	2.0	2.3	4.0	3.6
	N (%)	0.21	0.10	0.08	0.14	0.14	0.13	0.23	0.24
	C/N	8.3	9.7	9.3	7.8	8.7	10.2	10.0	8.7
	P (ppm)	20.3	66.3	35.0	30.8	27.3	19.3	34.8	27.8
	K (meq 100g ⁻¹)	0.15	0.21	0.15	0.15	0.24	0.08	0.15	0.09

Table 2 – Crop DM yield (t ha⁻¹) of the eight farms studied.

Product	CONV1	CONV2	INT1	INT2	ORG2	ORG2	LIV1	LIV2
Maize grain	11.3	11.3	7.9	10.1			8.8	
Maize silage							17.3	
Maize straw							7.3	
Wheat	3.6		5.0	4.4	4.5	5.4		
Wheat straw	2.3		4.1	2.3	4.2	2.9		
Lucerne					8.1	15.0		13.7
Soybean (II) ^a		3.9		3.2		3.5 (3.3)		
Meadow							12.0	
Barley						4.8	4.1	4.4
Barley straw						2.9	3.4	5.7
Switchgrass								4.2
Bean					1.6			
Italian ryegrass					6.8			
Sorghum silage							10.2	
Grassland						3.4		
Tomato						6.5		
^a second crop								

1048 Table 3 – Coefficients used to calculate Energy Input, Net Energy and Energy Use Efficiency.

1049 Energy equivalent represents the energy embedded in each input/output product.

Input/Output		U.M.	Energy equivalent
Crop	Product		
Maize	grain ^a	MJ kg ⁻¹ DM	18.92
	silage ^b		17.9
	straw ^a		18.67
	seed ^b		113.2
Wheat	grain ^a		18.42
	straw ^a		18.17
	seed ^b		31.3
Lucerne	hay ^c		18.84
	seed ^d		31.4
Soybean	grain ^a		23.65
	seed ^b		40.6
Meadow	hay ^b		17.6
	seed ^d		31.3
Barley	grain ^b		18.4
	straw ^b		16.8
	seed ^b		31.4
Switchgrass	hay ^d		17.2
	seed ^d		31.3
Bean	grain ^d		16.74
	seed ^d		40.6
Italian ryegrass	hay ^b		17.2
	seed ^b		31.3
Sorghum	silage ^d		18
	seed ^d		31.4
Grassland	hay ^b		17.6
Tomato	fruit ^e	MJ kg ⁻¹ FM	1.3
	seedling ^f	MJ plant ⁻¹	0.28
Fertiliser			
N- NH ₄ ^{b,g}		MJ kg ⁻¹	39
N-ureic ^{b,g}			48
N-NO ₃ ^{b,g}			32
N-other ^h			50
P ₂ O ₅ ^{b,g}			4
K ₂ O ^{b,g}			5
FPT ^h for N fertilisers ⁱ			1.5
FPT ^h for P fertilisers ⁱ			9.8
FPT ^h for K fertilisers ⁱ			7.3
FPT ^h for NP fertilisers ⁱ			5.7
FPT ^h for other fertilisers ⁱ			6
Pesticide			
Herbicides ^a		MJ kg ⁻¹ a.i.	264
Fungicides ^a			168
Insecticides ^a			214
Formulation ^a			20
Packaging ^a			2
Transport ^a		MJ kg ⁻¹	1.3
Other			
Diesel ^a		MJ l ⁻¹	46.9
Lubricants ^h		MJ l ⁻¹ diesel	3.6
Machinery energy embedded			12
^a Alluvione et al., 2011	^f Canakci et al., 2005		
^b Bechini and Castoldi, 2009	^g Kongshaug, 1998		
^c Fluck, 1992	^h Dalgaard et al., 2001		
^d Estimated	ⁱ Castoldi and Bechini, 2006		
^e Meul et al., 2007			
^h FPT = Formulation, Packaging and Transportation			

1050

1051 Table 4 – Diesel consumption for the different farm practices.
1052

Operation	l ha ⁻¹ (min-max)
chisel plowing	15-32
combined ridging-fertilisation	14
combined row cultivation-fertilisation	4
combined sowing-disking	12-14
combined sowing-fertilisation	12-14
combined sowing-pesticides treatment	6
cutting	5
drying	38-210
fertilisation	2-20
harrowing	8-26
harvesting	17-35
hay spreading	4-5
irrigation	5-185
laser levelling	9-23
pesticide treatment	3-9
plowing	16-39
ridging	4
rolling	3-4
rotary hoe	46
rototilling	7-25
row cultivation	4-9
silaging	17-29
sowing	7-26
straw harvest	4-22
straw shredding	8-30
tine arrowing	3-6
transplanting	24
windrowing	4-6

1053
1054
1055 Table 5 – Coefficients used to calculate the nutrient balance crop off-take.
1056

Crop	Product	N (%DM)	P (%DM)	K (%DM)
Maize	grain ^a	1.70	0.35	0.67
	silage ^a	1.20	0.22	1.00
	straw ^a	0.70	0.13	1.50
Wheat	grain ^a	2.30	0.39	0.50
	straw ^a	0.60	0.09	1.25
Lucerne	hay ^a	2.80	0.31	1.83
Soybean	grain ^a	6.70	0.74	2.25
Meadow	hay ^a	2.20	0.35	2.58
Barley	grain ^a	2.00	0.35	0.83
	straw ^a	0.60	0.09	1.08
Switchgrass	hay ^b	1.50	0.35	2.25
Bean	grain ^a	6.20	0.96	3.42
Italian ryegrass	hay ^a	1.50	0.35	2.25
Sorghum	silage ^a	0.90	0.13	0.83
	grain ^a	1.80	0.52	1.33
Grassland	hay ^a	2.20	0.35	2.08
Tomato	fruit ^b	2.50	0.18	4.83
Generic seeds ^c		1 ^d	1 ^d	3 ^d
^a Grignani et al., 2003				^d kg ha ⁻¹
^b Estimated				
^c Nielsen and Kristensen, 2005; Schröder et al., 1996				

1057

1058 Table 6 – Driving force indicators determined for the eight farms.

1059

Indicator group	Driving force indicator	CONV1	CONV2	INT1	INT2	ORG1	ORG2	LIV1	LIV2
Land use	Number of Crops	2	2	2	3	4	6	4	3
	Tillage Practices (% UAA)	100	100	100	100	73	85	70	28
	Irrigation (% UAA)	55	100	0	68	81	37	60	0
Fertiliser use	Mineral fertilisation (kg ha ⁻¹)	771	1180	694	473	0	0	0	0
	Organic fertilisation (kg ha ⁻¹)	0	0	0	0	1977	8710	25163	33524
	N fertilisation (kg ha ⁻¹)	222.7	310.4	163.4	123.1	11.1	42.7	158.5	214.6
	P fertilisation (kg ha ⁻¹)	17.2	31.7	25.3	13.4	3.8	16.7	79.5	85.6
	K fertilisation (kg ha ⁻¹)	32.8	121.0	131.0	68.9	10.6	47.2	167.9	257.4
Pesticides use	Consumption of Pesticides (kg a.i. ha ⁻¹)	1.5	2.2	0.8	0.5	-	-	-	-
	Equivalent Treatments	3.3	2.2	1.7	2.7	-	-	-	-
Energy use	Number of Practices	10.9	11.4	11.4	10.1	13.9	16.3	17.0	22.5
	Energy Input (GJ ha ⁻¹)	30.6	40.2	25.5	31.9	9.4	20.0	13.1	15.3

1060

1061

1062

1063 Table 7 – Kendall's tau b values of correlation between indicator grading against expected
 1064 grading. Expected grading correspond to CONV = 1; INT = 2; ORG = 3; LIV = 4 for all
 1065 indicators except for pesticides use indicators where ORG and LIV = 3.

1066

Indicator group	Indicators	Kendall's tau_b	Sig.
Driving force indicators			
Land use	Number of Crops	0.60	n.s.
	Tillage Practices	-0.87	0.006
	Irrigation	-0.27	n.s.
Fertiliser use	Mineral fertilisers	-0.87	0.006
	Organic fertilisers	0.87	0.006
	N fertilisers	-0.39	n.s.
	P fertilisers	0.15	n.s.
	K fertilisers	0.31	n.s.
Pesticides use	Consumption of Pesticides	-0.95	0.004
	Equivalent Treatment	-0.86	0.009
Energy use	Number of Practices	0.69	0.022
	Energy Input	-0.69	0.022
Pressure indicators			
Land use	Soil Cover	0.69	0.022
Fertiliser use	Gross N Balance	-0.77	0.011
	Gross P Balance	0.00	n.s.
	Gross K Balance	0.39	n.s.
Pesticide use	Load Index algae	-0.95	0.004
	Load Index fishes	-0.76	0.021
	Load Index bees	-0.86	0.009
	Load Index earthworms	-0.76	0.021
	Load Index mammals	-0.86	0.009
	Load Index birds	-0.95	0.004
	Environmental Impact Quotient	-0.95	0.004
Energy use	Net Energy	0.31	n.s.
	Energy Use Efficiency	0.77	0.011
Gaseous emissions	Ammonia emissions	0.00	n.s.
	GHG emissions	-0.62	0.041

1067

Figures

Figure 1 – Relationships between driving force indicators and pressure indicators. The symbol reported for each indicator specifies the optimal value of the indicator: “+” sustainability is higher when the indicator is high; “-” sustainability is higher when the indicator is low; “0” sustainability is higher when the indicator is zero.

Figure 2 – Energy Inputs.

Figure 3 – Percentage of the year with soil cover.

1086 Figure 4 – N, P, and K inputs (left) and gross nutrient balance (right).
1087

Figure 5 – Load Index for different non-target organisms.

Figure 6 – Different components of Environmental Impact Quotient (EIQ).

Figure 7 – Net Energy (NE) and Energy Use Efficiency (EUE).

Figure 8 – Ammonia emissions.

Figure 9 – Greenhouse gases emissions.

Figure 10 – Radar graphs representing driving force indicators (left) and pressure indicators (right).

List of abbreviation - Crop: Number of Crop; EI: Energy Input; Practices: Number of Practices; EqTreat: Equivalent Treatment; Pesticides: Consumption of Pesticides; K: K fertilisers; P: P fertilisers; N: N fertilisers; Tillage: Tillage Practices; SC: Soil Cover ; NE: Net Energy; EUE: Energy Use Efficiency; GHG: GHG emission; AE: Ammonia Emission; EIQ: Environmental Impact Quotient; GKB: Gross K Balance; GPB: Gross P Balance; GNB: Gross N Balance.