

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Management of downy mildew of sweet basil (*Ocimum basilicum*) caused by *Peronospora belbahrii* by means of resistance inducers, fungicides, biocontrol agents and natural products.

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/145852> since 2016-11-12T14:10:03Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Gilardi G.; Demarchi S.; Garibaldi A.; Gullino M.L.. Management of downy mildew of sweet basil (*Ocimum basilicum*) caused by *Peronospora belbahrii* by means of resistance inducers, fungicides, biocontrol agents and natural products.. *PHYTOPARASITICA*. 41 pp: 59-72.

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/145852>

1 **Management of downy mildew of sweet basil (*Ocimum basilicum* L.) caused by *Peronospora***
2 ***belbahrii* by means of resistance inducers, fungicides and alternative control measures**

3
4 Giovanna Gilardi, Stefano Demarchi, Angelo Garibaldi and Maria Lodovica Gullino

5
6 Centre of Competence for the Innovation in the Agro-environmental Sector (AGROINNOVA),
7 University of Torino, Via L. da Vinci 44, 10095 Grugliasco (TO), Italy

8
9 Corresponding author. Tel.:+ 39 0116708540; Fax: +39 0116709307.

10 E-mail address: marialodovica.gullino@unito.it

11

12 **Abstract** In the present study a number of compounds (acibenzolar-S-methyl, mineral and organic
13 fertilizers, plant extracts), known for their capability of inducing resistance in plants to several
14 pathogens, were tested in 2011 and 2012, in comparison with registered fungicides, copper-based
15 fungicides and biocontrol agents, to control downy mildew of sweet basil (*Ocimum basilicum* L.),
16 incited by *Peronospora belbahrii*. Four experimental trials were carried out under glasshouse
17 conditions in the presence of a high disease incidence. The tested compounds were used alone or
18 applied in rotation in spray programmes. In all trials the best results, in terms of reduction of disease
19 incidence and disease severity were offered by metalaxyl-M + copper hydroxide, by the mineral
20 fertilizer Alexin, by mandipropanid and by azoxostybin, followed by the glucohumates activator
21 complex and acibenzolar-S-methyl. Such products did significantly reduce disease incidence and
22 severity also 20 days after the last treatment. Among the copper-based products, the best results
23 were provided by copper hydroxide with terpenic alcohols and copper oxychloride + copper
24 hydroxide. The mineral fertilizer Kendal and prohexadione calcium, the biocontrol agent *B. subtilis*
25 QST 713, thyme oil extract and mustard oil on the contrary, were not effective. When different
26 combination of various products used in rotation were tested, it was possible to reduce disease
27 incidence and severity with different strategies, based on the rotation of fungicides and resistance
28 inducers as well as with the rotation of resistance inducers.

29

30 **Keywords** organic and mineral fertilizers; fungicides; acibenzolar-S-methyl; *Bacillus subtilis*;
31 integrated control

32

33 **Introduction**

34

35 Sweet basil (*Ocimum basilicum* L.) is an economically important herb crop in several
36 Mediterranean regions, in the USA and in many other parts of the world. This popular herb is used
37 both as a fresh and dried food spice, and in traditional medicine (Csizinsky, 1993; Lucier, 1993). In
38 Italy, most of the basil production takes place on the Riviera Ligure, and ‘Genovese Gigante’ is the
39 most appreciated variety for fresh consumption. It is the only cultivar used for industrial production
40 of the pesto sauce and covers 90% of the total growing area (Montalti, 1995). Also in Piedmont
41 basil is an important crop, grown in particular outdoor, especially for the production of pesto sauce,
42 with an average production of 4,000-4,500 tons/year, not considering the small producers. In
43 general, basil is mostly grown under greenhouses; in this environment average night temperatures
44 and dew depositions caused by high RH favour the development of diseases such as grey mould
45 (*Botrytis cinerea*), downy mildew (*Peronospora belbahrii*) and foliar spots (*Alternaria alternata*,
46 *Colletotrichum* sp.) especially after cool nights following warm, humid days (Garibaldi *et al.*, 1997;
47 Garibaldi *et al.*, 2011).

48 In particular, downy mildew, incited by *P. belbahrii*, was observed in northern Italy in 2003
49 (Garibaldi *et al.*, 2004) and quickly spread to other Italian regions in Central and Southern Italy
50 (Minuto *et al.*, 2004) as well as in France (Garibaldi *et al.*, 2005). This pathogen was first reported
51 in Uganda, identified as *Peronospora* sp. (Hansford, 1934) and much later in Switzerland (Lefort *et*
52 *al.*, 2003). After this report in Switzerland, the pathogen spread to many basil growing areas. Its
53 spread has been probably favoured by the fact that it is seed-transmitted (Garibaldi *et al.*, 2004).
54 The causal agent of basil downy mildew has been recently identified as *Peronospora belbahrii*
55 (Belbahri *et al.*, 2005; Thines *et al.*, 2009).

56 The disease was recently reported also in Belgium (Coosemans, 2004), in USA (Roberts *et al.*,
57 2009), in Cuba (Martinez de La Parte *et al.*, 2010) and in Hungary (Nagy and Horvath, 2011).

58 One of the hypothesis of its sudden, almost simultaneous spread in so many countries, is linked to
59 the production of seeds in developing countries where the pathogen is endemic.

60 The management of downy mildew of basil is complicated by the very limited availability of
61 chemicals on the crop, due to the risk of the presence of residues at harvest as well as on the
62 difficulty to have registered fungicides on minor crops such as basil (Leadbeater and Gisi, 2010).
63 The few fungicides registered in Italy on basil (azoxystrobin, belonging to QoI, metalaxyl-M,
64 belonging to phenylamides, and mandipropamid belonging to carboxylic acid amides) must be used
65 in a very limited number of sprays for avoiding the selection of resistant strains of the pathogen.
66 The search for varieties of basil resistant to the pathogen are still at early stages (Wyenandt *et al.*,
67 2010).

68 In the present study a number of compounds, known for their capability of inducing resistance
69 in plants to several pathogens, was tested, in comparison with registered fungicides and biocontrol
70 agents to control basil downy mildew.

71

72 **Materials and methods**

73

74 *Basil growth and experimental conditions* Four experimental trials (Table 1) were carried in 2011
75 and 2012 out at Grugliasco (Torino) in a glasshouse belonging to Agroinnova. Details about the
76 trials are reported in table 1. Basil seeds (cv. Genovese selection 'Italiano classico', Pagano
77 Sementi) were sown (0.2 g/pot, corresponding to 70-80 plants/pot) in plastic pots (1.5 L vol., 12 x
78 12 cm) containing a peat:perlite (80:20 v/v) mix (Turco Silvestro, Albenga, Savona), steamed (90°C
79 for 30 minutes).

80 For each treatment, four replicates (2 pots each) with a randomized design were used. All replicates
81 of the same treatment were managed similarly in terms of fertilization.

82 During the trials temperatures ranged between 19 and 25 °C. The basil pots were maintained over
83 the benches and covered with a plastic sheet placed over all the plants by using iron supports (1.0
84 m high, 3.0 m wide and 6.0 m long). The plastic sheets were placed on the iron support immediately
85 after each artificial inoculation and maintained until the last assessment. The relative humidity was
86 maintained close to 100% by misting during the experimental trials. The dates of the different
87 operations carried out are reported under table 1.

88

89 *Artificial inoculation* The populations of *Peronospora belbahrii*, collected in Piedmont (Northern
90 Italy) from diseased plants, were maintained on basil plants. The artificial inoculation was carried
91 out throughout nebulisation with a laboratory spray bottle (10 ml of capacity) of a suspension of
92 1×10^5 sporangia/ml. One ml of suspensions/treatment (corresponding to 8 pots), was used. The
93 artificial inoculation was carried out 24 h after the second treatment in trials 1-3, while in trial 4 two
94 artificial inoculations were carried out (Table 1).

95

96 *Treatments* Several compounds, known for their capability to induce resistance in the host, salts,
97 fertilizers, copper-based fungicides as well as fungicides registered on basil were tested. In trial 1-3
98 the different compounds were tested separately, while in trial 4 several spray programmes were
99 tested.

100

101 *Plant inducers, salts and fertilizers* Acibenzolar-S-methyl (Bion 50WG, 50% a.i., Syngenta, Italy),
102 prohexadione calcium (Regalis, 10% a.i, BASF, Italy), organic-mineral fertilizer based on
103 glucohumates complex N:P (Glucoinductor + GlucoActivator, N 4%, P₂O₅ 18%, International
104 patent PCT, IB2004\001905, Fertirev, Torino, Italy), organic-mineral fertilizer N:K (Kendal,
105 soluble organic nitrogen 3.5%, soluble potassium oxide 15.5%, organic carbon 3-4% Valagro,
106 Atessa, Chieti, Italy), mineral fertilizer (Kendal TE, Cu 23%, Mn 0,5%, Zn 0,5%, Valagro, Atessa,
107 Chieti, Italy) and (Alexin 95PS, P₂O₅ 52%, K₂O 42%, Massò, Milano, Italy), were tested at the
108 dosages reported under tables 2- 5.

109

110 *Plant extracts and BCAs* Thyme oil (*Thymus vulgaris* 100% a.i. plant extract, Soave & C, Italy),
111 mustard oil (Duolif, soluble organic nitrogen 3%, soluble sulphur 15%, organic matter 80%,
112 Cerealtoscana S.p.A., Livorno, Italy), *Bacillus subtilis* QST 713 (Serenade MAX, 15,67% a.i.,
113 BASF, Italy) were tested alone or in alternation.

114

115 *Copper-based products* Copper oxychloride (Cupravit Flow, 20% a.i., Bayer, Italy), copper
116 oxychloride and copper hydroxide (Airone, 10%+10% a.i. Isagro, Italy), copper sulfate (Cuproxat
117 SDI, 15.2% a.i., Nufarm GnbH & CoKG, Austria), copper hydroxide and terpenic alcohols
118 (Heliocuvire, 26.7% a.i., Intrachem Bio Italia, Italy), peptide complex copper sulfate, tackifying
119 compound and natural inducers of systemic induced resistance (SAR) (Labimethyl, 3%+ 2% a.i.,
120 Macasa, Spain), were tested at the dosages reported under tables 2-5.

121

122 *Chemical fungicides* Azoxystrobin (Ortiva, Syngenta Crop Protection S.p.A., Milano, Italy, 23.2%
123 a.i.), mandipropamid (Pergado SC, 23.4% a.i., Syngenta Crop Protection S.p.A., Italy), metalaxyl-
124 M + copper oxychloride (Ridomil R WG 2.5% + 40% a. i., Syngenta Crop Protection S.p.A., Italy)
125 were tested. Such fungicides are registered for use on basil in Italy and routinely applied in the
126 commercial production.

127 All treatments were carried out on basil plants 16-21 days after sowing at the dates reported under
128 table 1 by using 800 L ha⁻¹.

129 In the trials 1-3, two treatments at 6 day interval were carried out, with the exception of chemical
130 fungicides tested that were applied once at the dates reported under table 1. One artificial
131 inoculation with the pathogen was made 24 hours after the last treatment applications.

132 In trial 4, three treatments were carried out at 6 days interval, by using the tested products alone and
133 in alternation. Two artificial inoculations with the pathogen were carried out respectively, 24 hours
134 after the first and third treatment (Table 1).

135

136 *Data collection and analysis* Starting at the appearance of the first symptoms, the evaluation of the
137 percent of infected leaves (disease incidence) and of diseased leaf tissue (disease severity) was
138 made by using a disease rating scale with: 1 = up to 5 % infected leaf area; 2 = up to 10% infected
139 leaf area; 3 = up to 25% infected leaf area; 4 = up to 50% infected leaf area; 5= up to 75 %
140 infected leaf area. Disease incidence and severity was estimated on 100 leaves/treatment.

141 In trials 4, plant biomass, as fresh weight of plants at the end of the trial, was also measured.

142 All data collected were statistically analysed by univariate Anova with Tukey's test using SPSS
143 software 18.

144

145 **Results**

146

147 In all the four trials the artificial inoculation with *P. belbahrii* led to a high level of disease in the
148 untreated inoculated control; at the last estimating the development of pathogen attacks, disease
149 incidence ranged respectively from 66.8 to 88.8 and disease severity from 40.7 to 66.6. The level of
150 infection reached with the artificial inoculation was consistent in the different trials (Tables 2-5). In
151 the different trials and especially in the first one, infection was present also in the non inoculated
152 control plants; this is probably due to a contamination of the used seeds as well as with the spread
153 of the pathogen among plants (Table 2).

154 In trial 1, the best results, in terms of reduction of disease incidence and disease severity were
155 offered by the mixture metaxyl-M + copper hydroxide, by the mineral fertilizer Alexin, by
156 mandipropanid and by azoxostybin, followed by the glucohumates activator complex. Such
157 products did significantly reduce disease incidence and severity also 20 days after the last treatment
158 (Table 2). Twenty days after the last treatment, disease incidence was reduced from 88.8 in the
159 inoculated control to 5.5, 13.5, 16.3, 18.0 and 30.8 in the plants treated with the above mentioned
160 products. The same treatments reduced disease severity from 66.6 to 3.8, 9.1, 9.8, 11.9 and 22.0,
161 respectively (Table 2). The copper-based treatments offered a partial control of the disease as well
162 as acibenzolar-S-methyl. Among the copper-based products, the best results were provided by
163 copper hydroxide mixed with terpenic alcohols and copper sulphate + copper gluconate.
164 Probexadone calcium and mineral fertilizer Kendal, alone or combined with copper (Kendal TE)
165 provided only a very partial activity. *Bacillus subtilis* QST 713 and thyme oil extract did not
166 control downy mildew (Table 2).

167 Similar results were obtained in trial 2, in the presence of a similar level of infection. Metalaxyl-
168 M + copper hydroxide, mandipropanid, azoxystrobin, the mineral fertilizer Alexin, followed by

169 glucohumates activator complex were very effective in reducing downy mildew incidence and
170 severity. In this trial acibenzolar-S-methyl offered very good results. Also prohexadione calcium,
171 and the mustard oil provided interesting results in terms of reducing disease incidence and disease
172 severity. Among the different copper-based products tested, copper oxychloride + copper
173 hydroxide and copper hydroxide with terpenic alcohols provided interesting results followed by
174 copper sulphate compounds and by the mineral fertilizer Kendal TE (Table 3). *Bacillus subtilis* and
175 thyme oil extract were not effective (Table 3).

176 In trial 3, a similar trend was observed, with the same products providing the best control, in
177 terms of reduction of disease incidence and severity. Among the copper based products, with the
178 exception of copper oxychloride + copper hydroxide, copper hydroxide with terpenic alcohols, and
179 copper sulphate, copper oxychloride and the mineral fertilizer Kendal TE were partially effective,
180 as well as mustard oil and prohexadione calcium. *B. subtilis* and thyme oil confirmed to be not
181 effective (Table 4).

182 In trial 4, when used alone in 3 treatments, the fungicides metalaxyl-M + copper hydroxide,
183 mandipropanid, azoxystrobin, the mineral fertilizer Alexin and glucohumates activator complex
184 confirmed their very good activity (Table 5). Very interestingly, when different combination of
185 various products used in rotation were tested, it was possible to reduce disease incidence and
186 severity with different strategies, based on the rotation of fungicides and resistance inducers as well
187 as with the rotation of resistance inducers.

188 When also the effect on biomass was evaluated, in trial 4, the mineral fertilizer Alexin provided the
189 best results, while the glucohumate activator complex caused some reduction (Table 5).

191 **Discussion**

192
193 Downy mildew incidence was consistent and relevant in all trials as a consequence of the artificial
194 inoculation and permitted to evaluate the different products and strategies of disease control under
195 severe disease pressure. The infection observed in some trials in the not inoculated control was
196 probably caused by natural infection of seeds as well as by the spread of the pathogen among plants
197 with consequent cross contamination. The results obtained in the different trials were quite
198 consistent in terms of response of the different tested products. Among the tested fungicides,
199 metalaxyl-M + copper hydroxide and azoxystyrobin confirmed its efficacy (Gullino *et al.*, 2009).
200 Also mandipropamid, admitted for use on basil in Italy with a maximum of 50% of the total number
201 of applications or in alternation with fungicides having other modes of action, was quite effective.
202 The copper-based fungicides provided a partial control of the disease, except copper oxychloride

203 which was not effective. Copper-based products were mostly effective when applied in rotation
204 with resistance inducers such as the mineral fertilizer Alexin and with the glucohumates activator
205 product.

206 The most interesting results were consistently provided by the mineral fertilizer Alexin and by
207 the glucohumates activator complex, which provided, used alone or in rotation with other products,
208 also 20 days after the last treatment, results similar to those offered by the most active fungicides.
209 Also acibenzolar-S-methyl provided a good disease control of basil downy mildew with pre-
210 infection treatment, thus confirming what already observed for this compound when applied against
211 other foliar diseases, such as grey mould (*Botrytis cinerea*) on tomato (Malolepsza, 2006),
212 anthracnose (*Colletotrichum lagenarium*) on cucumber (Ishii *et al.*, 1999) and blue mould
213 (*Peronospora tabacina*) on tobacco (LaMondia, 2009).

214 The mineral fertilizer Kendal, the biocontrol agent *B. subtilis*, thyme oil extract and mustard oil, on
215 the contrary, were not effective.

216 Resistance inducers, also frequently referred to as plant strengthening agents, are legal under
217 current understanding and are treated under the fertilizer regulation (Kappert *et al.*, 2011). In
218 horticulture, particularly on vegetables and ornamentals, they are increasingly used as a means of
219 production and disease control. Their application is particularly interesting in the case of minor
220 crops, because of the lack of registered fungicides, as well as in organic farming, where they can
221 contribute in health maintenance and production quality (Kappert *et al.*, 2011).

222 Plant resistance inducers or improvers provided good results under different conditions, for
223 instance against *Peronospora destructor*, *P. parasitica*, *Bremia lactucae* and *Pseudoperonospora*
224 *cubensis* under greenhouse conditions (Kofot and Fisher, 2007): in general they are more effective
225 in the presence of moderate infection pressure and are generally influenced by different factors,
226 such as genotype, inoculum density, climatic conditions. In the case of the previously mentioned
227 pathogens, only phosphonates confirmed their efficacy also under field conditions (Kofot and
228 Fischer, 2007). Also a positive effects of their post-infection application have been reported (Wicks
229 *et al.*, 1999). In the case of *Peronospora parasitica* of cauliflower, acibenzolar-S-methyl was
230 effective by inducing the production of pathogenesis-related proteins (Ziadi *et al.*, 2001). The same
231 compound proved effective against Phytophthora blight, caused by *P. capsici*, on squash (*Cucurbita*
232 *pepo*) (Ji *et al.*, 2011).

233 In our study, some of the resistance inducers and fertilizers provided very interesting results,
234 when applied alone as well as in rotation with fungicides. As shown in trial 4, several combination
235 of products, applied in rotation in three treatments, permit to strongly reduce downy mildew
236 incidence and severity. Of particular interest are the strategies that, besides providing a good disease

237 control, do not cause strong reduction in the biomass. Moreover, it must be pointed out that the
238 rotation of effective fungicides with resistance inducers strongly reduces the risk of presence of
239 fungicide residues in a crop such as basil characterized by a high number of harvests. In the case of
240 basil crops for pesto production, harvest is carried out every 20 days. In the mean time, the rotation
241 between fungicides with resistance inducers will reduce the selection pressure by the fungicides,
242 thus reducing the risk of development of resistance towards the few registered fungicides.

243

244 **Acknowledgements**

245

246 Work carried out in the frame of the project SAFEFOODCONTROL, funded by Piedmont Region
247 “Sviluppo di sistemi e tecnologie innovative per la produzione, conservazione, trasformazione e
248 valorizzazione dell’orto-frutticoltura piemontese di qualità”.

249

250 **References**

251

252 Belbahri, I., Calmin, G., Pawlowski J., & Lefort, F. (2005). Phylogenetic analysis and real time PCR
253 detection of a presumably undescribed *Peronospora* species on sweet basil and sage.
254 *Mycological Research*, 109, 1276-1287.

255 Cohen, Y., Rubin, A., & Gotlieb, D. (2008). Activity of carboxylic acid amide (CAA) fungicides
256 against *Bremia lactucae*. *European Journal of Plant Pathology*, 122, 169-183.

257 Coosemans, J. (2004). First report of *Peronospora lamii*, downy mildew on basil (*Ocimum*
258 *basilicum*) in Belgium. *Parasitica*, 60, 27.

259 Csizinsky, A.A. (1993). The potential for aromatic plant production with plastic mulch culture in
260 Florida. *Acta Horticulturae*, 331, 27-34.

261 EPPO (2004). EPPO Standards PP1, 2nd Edition, Vol. 2. European and Mediterranean Plant
262 Protection Organization, Paris.

263 Garibaldi, A., Gilardi, G., Bertoldo, C., & Gullino, M.L. (2011). First report of leaf spot of sweet
264 basil (*Ocimum basilicum*) caused by *Alternaria alternate* in Italy. *Journal of Plant*
265 *Pathology*, 93, S4.71.

266 Garibaldi, A., Gullino, M. L., & Minuto, G. (1997). Diseases of basil and their management. *Plant*
267 *Disease*, 81, 124-132.

268 Garibaldi, A., Minuto, A. & Gullino, M.L. (2005). First report of downy mildew of basil (*Ocimum*
269 *basilicum*) in France. *Plant Disease*, 89, 683.

270 Garibaldi, A., Minuto, A., Minuto, G. & Gullino, M.L. (2004). First report of downy mildew of basil
271 (*Ocimum basilicum*) in Italy. *Plant Disease*, 88, 312.

- 272 Garibaldi, A., Minuto, G., Bertetti, D. & Gullino, M.L. (2004). Seed transmission of *Peronospora* of
273 basil. *Journal of Plant Diseases and Protection*, 111, 465-469.
- 274 Gisi, U., & Sierotzki, H. (2008). Fungicide modes of action and resistance in downy mildews. *Eur.*
275 *Journal of Plant Pathology*, 122, 157-167.
- 276 Gullino, M.L., Gilardi, G., & Garibaldi, A. (2009). Chemical control of downy mildew on lettuce
277 and basil under greenhouse. *Communication Agricultural and Applied Biological Sciences*,
278 74 (3), 933-940.
- 279 Hansford, C.G. (1932). Annual report of the mycologist. *Rev. appl. Mycol.*, 12, 421.
- 280 Ishii, H., Tomita, Y., Horio, T., Naruska, Y., & Nakazawa, Y. (1999). Induced resistance of
281 Acibenzolar-S-methyl (CGA245704) to cucumber and Japanese pear diseases. *European*
282 *Journal of Plant Pathology*, 105, 77-85.
- 283 Ji, P., Yin, J., & Koné, D. (2011). Application of acibenzolar-S-methyl and standard fungicides for
284 control of Phytophthora blight on squash. *Crop Protection*, 30, 1601-1605.
- 285 Kappert, R., Djufri, I., & Balas, J. (2011). Testing a plant strengthening agent in horticulture.
286 *Bulletin UASVM Horticulture*, 68, 375-379.
- 287 Kofoet, A., & Fischer, K. (2007). Evaluation of plant resistance improvers to control *Peronospora*
288 *destructor*, *P. parasitica*, *Bremia lactucae* and *Pseudoperonospora cubensis*. *Journal of Plant*
289 *Diseases and Protection*, 114, 54-61.
- 290 LaMondia, J.A. (2009). Efficacy of fungicides and a systemic acquired resistance activator
291 (acibenzolar-S-methyl) against tobacco blue mould. *Crop Protection*, 28, 72-76.
- 292 Leadbeater, A., & Gisi, U. (2010). The challenges of chemical control of plant diseases. In: Recent
293 developments in management of plant diseases (Gisi U., Chet I., Gullino M.L. eds.), Springer,
294 Dordrecht, The Netherlands, 3-17.
- 295 Lefort, F., Gigon, V., & Amos, B. (2003). Le mildiou s'étend. Déjà Détecté dans des nombreux
296 pays européens, *Peronospora lamii*, responsable du mildiou de basilic, a été observé en Suisse
297 dans la région lémanique. *Réussir Fruits et Légumes*, 223, 66.
- 298 Lucier, G. (1993). Vegetables and specialties situation and outlook report. *Commod. Econ. Res.*
299 *Serv.*, U.S. Dep. Agric., Nov: TVS-261.
- 300 Malolepsza, U. (2006). Induction of disease resistance by acibenzolar-S-methyl and o-
301 hydroxyethylrutin against *Botrytis cinerea* in tomato plants. *Crop Protection*, 25, 956-962.

302

303 Martinez de la Parte, E., Pérez-Vicente, L., Bernal, B., & Garcia, D. (2010). First report of
304 *Peronospora* sp. on sweet basil (*Ocimum basilicum*) in Cuba. *Plant Pathology*, 59, 800.

305 Montalti, M. (1995). Producendo basilico in coltura protetta. *Colture Protette*, 24, 45-49.

306 Nagy, G., & Horvath, A. (2011). Occurrence of downy mildew caused by *Peronospora belbahrii* on
307 sweet basil in Hungary. *Plant Disease*, 95, 1034.

308 Roberts, P.D., Raid, R.N., Harmon, P.F., Jordan, S.A., & Palmateer, A.J. (2009). First report of
309 downy mildew caused by a *Peronospora* sp. on basil in Florida and the United States. *Plant*
310 *Disease*, 93, 199.

311 Thines, M., Telle, S., Ploch, S., & Runge, F. (2009). Identity of the downy mildew pathogens of
312 basil, coleus, and sage with implications for quarantine measures. *Mycological Research*, 113,
313 532-540.

314 Wicks, T.J., Magarey, P.A., Wachtel, H.F., & Fresham, A.B. (1991). Effect of post infection
315 application of phosphonic acid on *Plasmopara citricola* on grapevine. *Plant Disease*, 75, 40-
316 43.

317 Wyenandt, C.A, Simon, J.A., McGrath, M.T., & Ward, D.L. (2010). Susceptibility of basil
318 cultivars and breeding lines to downy mildew (*Peronospora belbahrii*). *HortScience*,
319 45,1416-1419.

320 Ziadi, S., Barbedette, S., Godard, J.F., Monot, C., Le Corre, D., & Silué, D. (2001). Production of
321 pathogenesis-related proteins in the cauliflower (*Brassica oleracea* var. *Botrytis*) - downy
322 mildew (*Peronospora parasitica*) pathosystem treated with acibenzolar-S-methyl. *Plant*
323 *Pathology*, 50, 579-586.

324

325 **Table 1** Operations carried out in the four trials

Operation	Trial number			
	1	2	3	4
Date of sowing	18/08/2011	21/09/2011	22/09/2011	27/12/2011
Dates of treatments	9/09; 15/09	7/10; 13/10	14/10; 20/10	11/01/2012; 18/01/2012; 25/01/2012
Date of artificial inoculation	16/09	14/10	21/10	12/01/2012; 26/01/2012
Dates of disease assessment	23/09; 30/09; 5/10	26/10; 2/11; 7/11	31/10; 7/11;14/11	1/02/2012; 8/02/2012;17/02/2012
Date of biomass evaluation	-	-	-	17/02/2012

Table 2 Efficacy of different treatments, expressed as disease incidence (DI, % of infected leaves) and disease severity (DS, % of infected leaf area) against downy mildew, incited by *Peronospora belbahii*, on basil cv. Genovese, selection ‘Italiano Classico’, Trial

Active ingredient	Commercial product ^x	Dosage a.i.100 L	DI at day after last treatment ^x			DS at day after last treatment ^x								
			8	15	20	8	15	20						
-	Inoculated control	-	40.4	gh ^k	66.8	e	88.8	f	22.4	f-i ^k	46.7	e	66.6	f
Copper oxychloride	Cupravit flow	50	33.3	d-h	44.9	d	64.3	de	23.3	g-i	28.7	cd	46.8	c-e
Copper oxychloride + copper hydroxide	Airone	40+40	27.5	c-g	39.0	d	56.5	de	18.8	e-h	25.8	cd	40.1	b-e
Acibenzolar-S-methyl	Bion	1	5.5	ab	21.5	bc	49.3	cd	1.0	a	12.4	ab	32.6	bc
Organic mineral fertilizer N:K	Kendal	10.5+45	35.6	e-h	46.3	d	72.5	d-f	23.3	g-i	33.2	cd	54.4	d-f
Mineral fertilizer Cu+Mn+Zn	Kendal TE	46+1.5+1.5	26.8	c-f	39.1	d	65.0	d-f	16.4	d-g	25.7	cd	47.3	c-f
Prohexadione calcium	Regalis	5	22.0	cd	42.0	d	58.8	de	14.1	c-f	29.3	cd	41.9	b-e
Thyme oil extract	Tyme oil	100	44.0	h	50.5	de	77.5	ef	30.4	i	36.5	de	57.5	ef
<i>Bacillus subtilis</i> QST713	Serenade	58.4	23.5	c-e	38.0	cd	70.0	d-f	16.1	d-g	24.8	cd	52.5	d-f
Glucosaminates activator complex ^z	Glucosinductor	400	7.8	ab	16.5	ab	30.8	bc	5.5	a-c	11.5	ab	22.2	ab
Peptide complex copper sulfate	Labimethyl	9+6	22.3	c-e	40.9	d	54.9	c-e	14.8	d-g	25.7	cd	39.9	b-e
Copper hydroxide and terpenic alcohols	Heliocouivre	60	15.5	bc	35.0	cd	51.5	cd	10.0	b-d	21.9	bc	36.1	b-d
Copper sulfate	Cuproxat	53.2	16.6	bc	40.3	d	56.3	de	11.8	b-e	26.3	cd	39.5	b-e
Mineral fertilizer P ₂ O ₅ 52%, K ₂ O 42%	Alexin	130+105	0.0	a	6.3	ab	13.5	ab	0.0	a	3.7	a	9.1	a
Metalaxyl-M + copper hydroxide	Ridomil Gold R	7.5+ 120	0.0	a	1.3	a	5.5	a	0.0	a	0.7	a	3.8	a
Mandipropamid	Pergado	11.7	4.5	ab	9.5	ab	16.3	ab	3.4	ab	5.4	a	9.8	a
Azoxystrobin	Ortiva	18.6	1.0	a	6.0	ab	18.0	ab	0.8	a	2.7	a	11.6	a
-	Not inoculated control	-	37.5	f-h	48.2	d	73.0	d-f	25.8	hi	33.6	cd	54.4	d-f

^x Two treatments were carried out on: 9/09/2011; 15/09/2011. One artificial inoculation was carried out on: 16/09/2011

^k Means of the same column, followed by the same letter do not differ according to Tukey’s test (P<0.05)

^z Dosage (ml 100 L⁻¹) of the commercial formulation

Table 3 Efficacy of different treatments, expressed as disease incidence (DI, % of infected leaves) and as disease severity (DS, % of infected leaf area) against downy mildew, incited by *Peronospora belbahii* on basil cv. Genovese, selection ‘Italiano Classico’, Trial 2

Active ingredient	Commercial product ^x	Dosage g a.i.100 L ⁻¹	DI at at day after last treatment						DS at day after last treatment ^x					
			13 ^x	20	25	13 ^x	20	25						
-	Inoculated control	-	53.5	e ^k	81.8	f	84.5	f	32.9	e ^k	58.6	f	61.8	e
Copper oxychloride	Cupravit flow	50	19.0	b-d	36.3	cd	43.8	c-e	7.1	a-c	25.0	d	28.9	b-d
Copper oxychloride + copper hydroxide	Airone	40+40	11.0	ab	10.8	ab	13.0	ab	3.6	a	6.8	a-c	7.3	a
Acibenzolar-S-methyl	Bion	1	2.0	ab	6.0	ab	7.5	ab	0.8	a	3.9	a-c	4.7	a
Organic mineral fertilizer N:K	Kendal	10.5+45	31.3	cd	26.3	bc	50.5	de	17.9	b-d	18.4	cd	36.1	cd
Mineral fertilizer Cu+Mn+Zn	Kendal TE	46+1.5+1.5	9.3	ab	25.3	bc	27.8	b-d	3.1	a	16.3	b-d	18.1	a-c
Prohexadione calcium	Regalis	5	13.3	a-c	26.0	bc	25.3	a-e	5.3	ab	17.6	b-d	18.4	a-c
Thyme oil extract	Time oil	100	36.2	de	58.5	de	61.3	ef	22.9	de	41.3	e	44.7	de
Mustard oil ^z	Duolif	1000	12.5	ab	16.5	a-c	25.5	a-e	5.1	ab	10.5	a-d	15.4	a-c
<i>Bacillus subtilis</i> QST713	Serenade	58.4	35.5	de	74.5	ef	83.8	f	20.0	c-e	55.8	ef	60.3	e
Glucohumates activator complex ^z	Glucoinductor	400	0.0	a	4.3	ab	3.8	ab	0.0	a	1.6	ab	2.1	a
Peptide complex copper sulfate	Labimethyl	9+6	6.0	ab	15.5	a-c	24.8	a-e	2.7	a	10.9	a-d	15.4	a-c
Copper hydroxide and terpenic alcohols	Heliocuvire	60	1.3	ab	7.3	ab	13.3	ab	0.4	a	4.5	a-c	7.4	a
Copper sulfate	Cuproxat	53.2	8.5	ab	15.0	a-c	18.3	a-c	2.8	a	9.9	a-d	10.3	ab
Mineral fertilizer P ₂ O ₅ 52%, K ₂ O 42%	Alexin	130+105	0.0	a	0.0	a	2.5	ab	0.0	a	0.0	a	1.8	a
Metalaxyl-M + copper hydroxide	Ridomil Gold R	7.5+ 120	0.0	a	0.0	a	0.8	a	0.0	a	0.0	a	0.2	a
Mandipropamid	Pergado	11.7	0.0	a	0.3	a	1.3	ab	0.0	a	0.1	a	0.8	a
Azoxystrobin	Ortiva	18.6	0.0	a	0.0	a	1.3	ab	0.0	a	0.0	a	0.6	a
-	Non inoculated control	-	0.0	a	0.3	a	6.8	ab	0.0	a	0.1	a	4.6	a

^x Two treatments were carried out on:7/10; 13/10. One artificial inoculation was carried out on: 14/10

^k Means of the same column, followed by the same letter do not differ according to Tukey’s test (P<0.05)

^z Dosage (ml 100 L⁻¹) of the commercial formulation

Table 4 Efficacy of different treatments, expressed as disease incidence (DI, % of infected leaves) and as disease severity (DS, % of infected leaf area) against downy mildew, incited by *Peronospora belbahii* on basil cv. Genovese, selection ‘Italiano Classico’, Trial 3

Active ingredient	Commercial product ^x	Dosage g a.i.100 L ⁻¹	DI at day after last treatment ^x						DS at day after last treatment ^x					
			11	18	25	11	18	25	11	18	25			
-	Inoculated control	-	38.3	e ^k	47.5	h	83.8	h	26.8	f ^k	35.6	h	62.8	h
Copper oxychloride	Cupravit flow	50	22.0	d	27.5	e-f	61.3	f-h	15.8	e	20.6	fg	45.9	f-h
Copper oxychloride + copper hydroxide	Airone	40+40	8.5	a-c	11.7	a-c	21.3	a-d	6.0	a-d	8.1	a-d	15.9	a-d
Acibenzolar-S-methyl	Bion	1	3.3	a	6.5	ab	14.0	a-c	2.1	a	3.6	ab	10.5	a-c
Organic mineral fertilizer N:K	Kendal	10.5+45	17.0	cd	25.0	d-f	53.8	e-g	12.0	de	18.8	e-g	40.3	e-g
Mineral fertilizer Cu+Mn+Zn	Kendal TE	46+1.5+1.5	7.0	ab	13.8	b-d	30.0	b-e	4.6	a-c	10.3	b-e	22.5	b-e
Prohexadione calcium	Regalis	5	16.0	b-d	16.3	b-e	37.5	c-f	11.3	c-e	12.2	b-f	28.1	c-f
Thyme oil extract	Tyme oil	100	21.8	d	39.8	gh	58.8	fg	14.8	e	24.9	g	44.1	fg
Mustard oil ^z	Duolif	1000	14.5	b-d	23.3	c-f	42.5	d-g	10.1	b-e	17.4	e-g	31.9	d-g
<i>Bacillus subtilis</i> QST713	Serenade	58.4	14.8	b-d	26.3	e-f	62.5	gh	10.6	b-e	19.7	fg	46.9	gh
Glucohumates activator complex ^z	Glucoinductor	400	0.0	a	5.5	ab	10.8	ab	0.0	a	3.7	ab	8.1	ab
Peptide complex copper sulfate	Labimethyl	9+6	21.0	d	27.5	e-f	53.8	e-g	14.4	e	20.6	fg	40.3	e-g
Copper hydroxide and terpenic alcohols	Heliocuire	60	4.8	a	9.0	ab	22.5	a-d	3.6	ab	6.8	a-c	16.9	a-d
Copper sulfate	Cuproxtat	53.2	6.8	ab	29.5	fg	20.0	a-d	4.6	a-c	14.4	c-f	15.0	a-d
Mineral fertilizer P ₂ O ₅ 52%, K ₂ O 42%	Alexin	130+105	2.8	a	6.3	ab	11.3	ab	2.1	a	4.6	ab	8.4	ab
Metalaxyl-M + copper hydroxide	Ridomil Gold R	7.5+ 120	0.0	a	0.0	a	5.0	a	0.0	a	0.0	a	3.8	a
Mandipropamid	Pergado	11.7	0.0	a	0.0	a	10.0	ab	0.0	a	0.0	a	7.5	ab
Azoxystrobin	Ortiva	18.6	0.0	a	0.0	a	9.3	ab	0.0	a	0.0	a	6.9	ab
-	Non inoculated control	-	15.5	b-d	22.5	c-f	56.3	fg	10.4	b-e	16.9	d-g	42.2	fg

^x Two treatments were carried out on:14/10; 20/10. One artificial inoculation was carried out on: 21/11

^k Means of the same column, followed by the same letter do not differ according to Tukey’s test (P<0.05)

^z Dosage (ml 100 L⁻¹) of the commercial formulation

Table 5 Effect of different treatments against downy mildew, incited by *Peronospora belbaharii*, expressed as disease incidence (DI, % of infected leaves) as disease severity (DS, % of infected leaf area) and on the yield of basil cv. Genovese, selection 'Italiano Classico', Trial 4.

Active ingredient (code)	Commercial product ^x	Dosage g a.i.100 L ⁻¹	DI at days after the last treatment ^x						DS at days after the last treatment						Biomass (g)	
			8	13	22	8	13	22	8	13	22					
-	Inoculated control	-	22.5	fg ^k	36.8	e	66.8	f	10.7	ef	25.8	c	40.7	f	64.0	cd
Copper oxychloride + copper hydroxide (A)	Airone	40+40	9.3	c-e	17.5	cd	24.5	d	2.6	a-c	9.8	b	14.9	d	90.7	b-d
Acibenzolar-S-methyl (B)	Bion	1	2.8	a-c	5.5	ab	12.8	a-d	0.8	ab	1.4	a	8.1	a-d	80.4	a-d
Mineral fertilizer Cu+Mn+Zn	Kendal TE	46+1.5+1.5	10.5	de	10.8	a-d	24.5	d	3.4	bc	5.3	ab	14.0	cd	69.5	a-d
Thyme oil	Tyme oil	100	24.5	g	36.0	e	58.8	f	11.4	f	23.6	c	37.8	f	68.0	d
Mustard oil ^z	Duolif	1000	21.8	fg	28.8	e	37.8	e	7.8	de	19.3	c	24.6	e	60.8	cd
Glucosaminates activator complex ^z (F)	Glucosinductor	400	0.5	a	0.5	a	6.3	ab	0.1	a	0.0	a	3.4	a	75.4	b-d
Copper hydroxide and terpenic alcohols (G)	Heliocuvivre	60	15.8	ef	17.8	d	40.3	e	5.0	cd	10.1	b	24.7	e	86.2	a-d
Mineral fertilizer P ₂ O ₅ 52%, K ₂ O 42% (H)	Alexin	130+105	0.0	a	0.0	a	6.0	ab	0.0	a	0.0	a	1.9	a	124.8	a
Metalaxyl-M + copper hydroxide (I)	Ridomil Gold R	7.5+ 120	0.0	a	7.0	a-d	9.8	ab	0.0	a	2.3	a	4.6	ab	89.4	a-c
Mandipropamid (L)	Pergado	11.7	2.3	a-c	3.0	ab	8.3	ab	0.6	ab	0.4	a	3.1	a	78.0	a-c
Azoxystrobin (M)	Ortiva	18.6	0.0	a	0.0	a	8.0	ab	0.0	a	0.0	a	3.1	a	76.6	a-d
A-H-A ^y	Airone; Alexin; Airone	40+40; 130+105; 40+40	1.5	a	3.0	ab	11.8	a-c	0.4	ab	0.9	a	4.6	ab	78.2	a-c
A-I-A	Airone; Ridomil Gold R; Airone	40+40; 7.5+ 120; 40+40	0.0	a	4.5	ab	10.5	a-c	0.0	a	1.3	a	4.9	ab	64.0	a-d
A-L-A	Airone; Pergado; Airone	40+40;11.7;40+40	5.5	a-d	9.5	a-d	16.8	b-d	1.8	ab	2.8	ab	8.3	a-d	73.6	cd
A-M-A	Airone; Ortiva;Airone	40+40;18.6;40+40	0.0	a	8.3	a-d	10.3	ab	0.0	a	2.2	a	4.0	ab	95.2	a-c
A-B-A	Airone; Bion; Airone	40+40; 1; 40+40	8.8	b-e	11.8	b-d	22.8	cd	1.2	ab	1.6	a	5.9	a-c	100.4	a-d
A-F-A	Airone; Glucosinductor; Airone	40+40; 400; 40+40	4.3	a-d	4.3	ab	10.5	a-c	1.3	ab	2.5	ab	3.2	a	84.4	a-c
B-H-B	Bion ;Alexin; Bion	1; 130+105; 1	0.5	a	0.8	a	4.0	a	0.1	a	0.2	a	1.6	a	93.4	a-c
B-I-B	Bion ;Ridomil Gold R; Bion	1; 7.5+120; 1	0.0	a	5.3	ab	7.0	ab	0.0	a	1.6	a	2.8	a	83.2	a-c
B-L-B	Bion; Pergado; Bion	1; 11.7; 1	2.0	ab	9.5	a-d	17.5	b-d	0.6	ab	2.8	ab	7.6	a-d	66.4	a-d
B-M-B	Bion; Ortiva; Bion	1; 18.6; 1	0.0	a	2.8	ab	5.8	ab	0.0	a	0.7	a	2.0	a	85.0	a-d
B-F-B	Bion; Glucosinductor; Bion	1; 400; 1	5.0	a-d	6.8	a-c	8.3	ab	1.1	ab	0.6	a	2.8	a	94.4	a-c
F-H-F	Glucosinductor; Alexin; Glucosinductor	400; 130+105;400	4.0	a-d	2.0	ab	5.5	ab	0.2	a	0.0	a	2.4	a	36.8	cd
F-I-F	Glucosinductor ; Ridomil Gold R; Glucosinductor	400; 7.5+120;400	0.8	a	0.0	a	6.8	ab	0.0	a	0.4	a	4.7	ab	79.0	a-d
F-L-F	Glucosinductor; Pergado; Glucosinductor	400; 11.7;400	0.0	a	1.8	ab	12.3	a-d	0.1	a	0.0	a	2.2	a	100.6	a-d
F-M-F	Glucosinductor; Ortiva; Glucosinductor	400; 18.6;400	0.5	a	0.0	a	4.0	a	0.5	ab	0.3	a	4.4	ab	111.6	ab
G-H-G	Heliocuvivre; Alexin; Heliocuvivre	60; 130+105;60	2.0	ab	1.3	ab	7.5	ab	1.1	ab	1.1	a	3.9	ab	104.4	a-c
G-I-G	Heliocuvivre ; Ridomil Gold R; Heliocuvivre	60; 7.5+120;60	4.0	a-d	3.8	ab	8.5	ab	1.6	ab	4.5	ab	9.0	a-d	106.2	a-c
G-L-G	Heliocuvivre; Pergado; Heliocuvivre	60; 11.7;60	5.0	a-d	8.8	a-d	17.3	b-d	0.8	ab	1.6	a	6.1	a-d	88.6	a-d
G-M-G	Heliocuvivre; Ortiva; Heliocuvivre	60; 18.6; 60	3.0	a-c	4.0	ab	12.3	a-d	2.6	abc	5.3	ab	12.4	b-d	97.6	a-c
-	Not inoculated control	-	0.0	a	1.5	ab	8.3	ab	0.0	a	0.5	a	3.1	a	93.8	a-c

^x Three treatments were carried out on: 11/1; 18/1; 25/1. Two artificial inoculations were carried out on: 12/01 and 26/01

^k Means of the same column, followed by the same letter do not differ according to Tukey's test ($P < 0.05$)

^z Dosage (ml 100 L⁻¹) of the commercial formulation

^y Spray program adopted where letter corresponding to A, Copper oxychloride + copper hydroxide (Airone); H, Mineral fertilizer P₂O₅ 52%, K₂O 42% (Alexine); I, Metalaxyl-M + copper hydroxide (Ridomil Gold R); L, Mandipropamid (Pergado); M, azoxystrobin (Ortiva); B, Acibenzolar-S-methyl (Bion); F, Glucohumates activator complex (Glucoinductor); G, Copper hydroxide and terpenic alcohols (Heliocuire)