

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Genetic variation of two local chicken populations: 'Bianca di Saluzzo' and 'Bionda Piemontese'

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/147272> since

Publisher:

DEPARTMENT OF AGRONOMY, FOOD, NATURAL RESOURCES, ANIMALS AND ENVIRONMENT

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Genetic variation of two local chicken populations: 'Bianca di Saluzzo' and 'Bionda Piemontese'

S. Sartore^{1}, M. Marco¹, S. Maione¹, D. Soglia¹, P. Sacchi¹, R. Rasero¹, M.T. Bottero¹, A. Dalmaso¹, D. Pattono¹, M. Giacobini¹, L. Bertolotti¹, P. Bajardi¹, I. Zoccarato², L. Gasco², A. Brugiapaglia², M. Tarantola¹, A. Schiavone¹, S. Sponza¹*

*¹Dipartimento di Scienze Veterinarie. ²Dipartimento di Scienze Agrarie, Forestali e Alimentari.
University of Torino. 10195 Grugliasco (TO). Italy*

In Piemonte region two autochthonous chicken breeds exist: 'Bianca di Saluzzo' and 'Bionda Piemontese'. These slow-growing meat breeds are reared in farms mainly located in Cuneo and Asti provinces. In the past decades both breeds underwent to a strong reduction in size and they were frequently replaced by fast-growing lines. Only recently, conservation plans were undertaken. The main objective of this work was to determine the genetic structure of these breeds. We used 26 microsatellites suggested by the literature. Allelic richness was 5.9 and 6.4 for the 'Bianca' and the 'Bionda' respectively, gene diversity was 0.665 in both breeds. Differentiation was measured by F_{st} (0.056) and was highly significant ($P < 0.001$). Using the Bayesian analysis (Structure software) under the hypothesis of two clusters, all runs split the dataset into samples from the 'Bianca' and from the 'Bionda' and most individuals were correctly assigned with membership > 0.9 . With three or more clusters the 'Bionda' split into two different ecotypes whereas the cluster including most of the 'Bianca' was very robust, showing that this breed was the most different but also the most homogeneous within the dataset.