

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Cytogenetic investigations in two endangered pig breeds raised in Southern-Italy.

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/150211> since

*Published version:*

DOI:10.1007/s10577-014-9435-7

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

*The final publication is available at Springer via <http://dx.doi.org/10.1007/s10577-014-9435-7>*

### Cytogenetic investigations in two endangered pig breeds raised in Southern-Italy

A. Perucatti<sup>1</sup>, V. Genuardo<sup>1</sup>, A. Iannuzzi<sup>1</sup>, L. Pucciarelli<sup>1</sup>, D. Incarnato<sup>1</sup>, A. Pauciullo<sup>1</sup>, D. Marletta<sup>2</sup>, G. Scopino<sup>1,3</sup>, B. Castiglioni<sup>4</sup>, V. Peretti<sup>3</sup>, L. Iannuzzi<sup>1</sup> ([angela.perucatti@ispaam.cnr.it](mailto:angela.perucatti@ispaam.cnr.it))

<sup>1</sup>National Research Council (CNR) of Italy, ISPAAM, Laboratory of Animal Cytogenetics and Gene Mapping, Naples, Italy; <sup>2</sup>DISPA, University of Catania, Catania, Italy; <sup>3</sup>Dept. Veterinary Medicine and Animal Production, University of Naples Federico II, Naples, Italy; <sup>4</sup>CNR-IBBA, Milan, Italy.

Pig from Casertana and Siciliana breeds are two endangered breeds raised in the southern-Italy (Campania and Sicily regions, respectively) and characterized to have a black skin. Special projects are trying to save and characterize both of them by using several approaches. In this study we report the preliminary results we obtained after a cytogenetic investigation we performed by using both C and R-banding techniques and the sister chromatid exchange (SCE) test to verify their chromosome stability under the environmental conditions. Fifty-two pigs from Casertana breed and 19 pigs from Siciliana breed were investigated. All animals from both breeds showed a normal karyotype, with the exception of two male pig from Siciliana breed which were found heterozygous carrier of rob(15;17) (2n=37, XY), probably being hybrids with the wild pig (2n=36) present in the Nebrodi mountains where this breed is raised in Sicily. SCE-test applied on 42 pigs from Casertana breed (22 males and 20 females) and 19 pigs from Siciliana breed (8 males and 11 females) revealed no statistical differences between the SCE-mean number in Casertana pig ( $7.13 \pm 3.20$ ) than that ( $6.87 \pm 3.12$ ) achieved in Siciliana pig. Statistical differences were found between males ( $7.26 \pm 3.38$ ) and females ( $6.59 \pm 2.90$ ) of Siciliana pig breed, as well as between females of Casertana ( $7.24 \pm 3.26$ ) and Siciliana ( $6.59 \pm 2.90$ ), while no statistical differences were found between males of the breeds, as well as between males and females of Casertana breed.

*Acknowledgements. The study has been supported by project "RARECA, PSR, Misura 214 e2" of Campania region and "CISIA-GenePig" project, National Research Council (CNR) of Italy.*