AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Evidence of genotoxicity in lymphocytes of non-smoking alcoholics

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1506171> since 2017-06-27T14:19:40Z

Published version:

DOI:10.1007/s11033-014-3739-8

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Evidence of genotoxicity in lymphocytes of non-smoking alcoholics

Alfredo Santovito · Piero Cervella ·
Massimiliano Delpero

Received: 1 March 2013 / Accepted: 11 September 2014
© Springer Science+Business Media Dordrecht 2014

Abstract Alcohol abuse is a significant public health issue. Epidemiological studies conducted on different populations consistently showed that consumption of alcoholic beverages is associated with cytogenetic damages and higher risk for several types of cancer. However, the interpretation of many cytogenetic studies resulted complicated because some confounding factors, such as smoking habit, are not always taken into account. In the present study, the frequency of sister chromatid exchanges (SCEs), chromosome aberrations (CAs) and micronuclei (MNs) in cultured human lymphocytes was assessed on 15 alcoholic and 15 non-alcoholic control male subjects. Moreover, considering the implication of the Glutathione S-transferases gene polymorphisms in the genetic susceptibility to alcoholic liver diseases, we considered an important issue to evaluate the relationship between these gene polymorphisms and the cytogenetic damage. In our sample we exclusively considered individuals that did not smoke nor consume drugs for a period of at least 2 years prior to the analysis. Statistically significant differences were found between alcoholics and controls in the frequency of SCEs/cell ($P = 0.001$), RI value ($P = 0.001$), CAs ($P = 0.002$) and CAB ($P = 0.002$). Vice versa, no significant differences were found between alcoholics and controls in terms of MNs frequency and CBPI value. In both samples, no statistically significant association was found between the analysed GSTs gene polymorphisms and the frequencies of MNs, SCEs and CAs. Finally, among alcoholics we found a positive correlation between

SCEs and CAs frequencies and the duration of alcohol abuse. **37**
39

Keywords Micronuclei · Sister chromatid exchanges · **40**
Chromosomal aberrations · Genotoxicity · Alcohol **41**

Introduction **42**

Alcohol abuse is a significant social and public health concern. **43**
In 1988, the International Agency for Research on Cancer **44**
(IARC) classified alcoholic drinks as group I carcinogens in **45**
humans for the upper airways, digestive tract and liver [20]. **46**
Excessive chronic ingestion of ethanol was also associated **47**
with serious damages to most of the major organ systems such **48**
as the liver, pancreas, thyroid, pituitary glands, and adrenal **49**
gland [1, 8, 21, 32, 33]. From a cytogenetic point of view, **50**
ethanol was shown to induce sister chromatid exchanges **51**
(SCEs), micronuclei (MNs), and aneuploidy in mouse eggs **52**
[38]; mis-segregation and/or nondisjunction in *Aspergillus* **53**
nidulans [9, 23], *Drosophila melanogaster* [41], and rodents **54**
[25–28], as well as MNs in *Zea* and *Tradescantia* [38]. **55**

In vivo cytogenetic studies among humans showed **56**
increased frequencies of chromosomal aberrations (CAs) **57**
[17, 38], SCEs [38], and MNs [19, 32] in peripheral blood **58**
lymphocytes of alcoholics. Nevertheless, the interpretation **59**
of some cytogenetic studies could be problematic because **60**
some confounding factors, such as smoking, were not **61**
always taken into account. It is known that alcohol **62**
dependent is usually associated with smoking habit. This **63**
association could result in further increased levels of CAs **64**
and MNs [37, 48] and cancer incidence [22, 30, 50]. **65**

In this study, we analysed the MNs, SCEs and CAs **66**
frequencies in cultured lymphocytes from a sample of non- **67**
smoker alcoholic and non-alcoholic control subjects. MN **68**

A1 A. Santovito (✉) · P. Cervella · M. Delpero
A2 Department of Life Sciences and Systems Biology, University of
A3 Turin, Via Accademia Albertina n. 13, 10123 Turin, Italy
A4 e-mail: alfredo.santovito@unito.it

assay detects both clastogenicity (chromosome breakage) and aneugenicity (chromosome lagging due to dysfunction of mitotic apparatus) [12, 14, 39], while the SCE analysis reveals alterations in the chromosome structure only. Moreover, increased levels of CAs, in particular of chromosome breaks, were found closely related to cancer development [18].

Metabolism of alcohol results in the generation of several classes of DNA-adducts and/or DNA-damaging molecules, including reactive oxygen species (ROS), lipid peroxidation products and acetaldehyde [32]. These compounds were found to affect the DNA-repair systems and can contribute to the hepatocarcinogenesis process [47]. Detoxification cellular systems protect the cells from DNA damage caused by various reactive substances. Glutathione S-transferases (GSTs) represent one of the major groups of phase II detoxifying enzymes, found in virtually all eukaryotes and evolved providing protection against reactive oxygen metabolites and toxic substances present in the food and the environment [36]. In particular, the GST enzymes are able to detoxify harmful ethanol metabolites in the liver by conjugating acetaldehyde and ROS to reduced glutathione [31]. *GSTT1* and *GSTM1* genes are known to be polymorphic in humans for a deletion of a segment of DNA, that results in the absence of protein synthesis and consequent reduced detoxification of xenobiotics in homozygous individuals [42].

These genes have been also considered as potential candidates for alcoholic liver disease (ALD) susceptibility [34]. Savolainen et al. [46], found a statistically significant association between the occurrence of alcoholic liver cirrhosis and *GSTM1* “null” genotype, suggesting that persons with homozygous deletion of the *GSTM1* gene could be genetically more prone to develop alcoholic liver fibrosis. Nevertheless, no data were found in literature about the possible association of *GSTs* gene polymorphism and cytogenetic damage in samples of alcoholics.

In this study we aimed to perform a cytogenetic analysis on lymphocytes from a selected sample of alcoholic beverage dependent individuals, in order to evaluate possible risks of genomic damage due alcohol genotoxicity. Moreover, considering the implication of the *GSTs* gene polymorphisms in the genetic susceptibility to alcoholic liver diseases and cancer, we considered an important issue to evaluate the possible relationship between these gene polymorphisms and the cytogenetic damage in chronic alcoholics.

Materials and methods

Groups studied

The study was conducted on 15 alcoholics and 15 non-drinker controls, all males. Alcoholic subjects declared to drink

Table 1 Demographic characteristics and alcohol consumption habits of the studied groups

Characteristics	Alcoholics	Controls
Subjects (n)	15	15
Age		
Mean years (SD)	50.73 (8.89) ^a	49.33 (10.83) ^a
Range (years)	36–70	38–70
Alcohol habit		
Years (SD)	7.67 (2.74)	–
Range (years)	4–12	–
Daily assumption		
Mean litres of alcoholic beverages (SD)	3.13 (2.20)	–
Mean grams of ethanol (SD)	284.50 (196.50)	–

SD Standard deviation

^a $P = 0.944$

regularly both wine and distilled beverages, also during the period while the study was conducted. All participants were extensively interviewed by a specialized physician with a detailed questionnaire in order to collect meaningful information for the study. Age, smoking habits, work-related exposure to hazardous agents, dietary habits, use of therapeutic drugs and alcohol consumption were all recorded.

All subjects were selected avoiding individuals occupationally exposure to known or suspected mutagens/genotoxic agents. In our sample we exclusively considered individuals that have not consumed drugs and have not been subjected to invasive diagnostic examinations for a period of at least 2 years prior to the analysis. Among controls, 9 subjects declared to be teetotalers, and 6 occasional drinkers who declared to drink no more than one glass of wine during week-end.

All the subjects were volunteers, received information about the study and gave their written informed consent. The procedures followed in this work were in accordance with the ethical standards of the local responsible committee on human experimentation and with the Helsinki Declaration of 1975, as revised in 1983.

Blood sample collection

Blood samples were obtained by venipuncture (5–10 mL) and collected into heparinised tubes, for genotoxicity testing. All blood samples were coded, cooled (4 °C), and processed within 2 h after collection.

Cytokinesis-block micronucleus assay

MNs assay was conducted following the protocol described in [43]. Micronuclei were scored in 1,000 bi-nucleated

lymphocytes per subject, following the established criteria for the MN evaluation [15]. The cytokinesis-block proliferation index (CBPI) was calculated according to the formula: $[1 \times N_1] + [2 \times N_2] + [3 \times (N_3 + N_4)]/N$, where N_1 – N_4 represent the number of cells with 1–4 nuclei, respectively, and N is the total number of cells scored.

Sister chromatid exchanges assay

SCEs assay was performed according to [45]. In order to determine the number of SCE/cell for each subject, we scored 50 well-spread second-division metaphases containing 46 (± 1) chromosomes. A total of 100 cells from each donor were scored for the determination of the replication index (RI), calculated according to the formula: $RI = (M_1 + 2M_2 + 3M_3)/N$, where M_1 , M_2 and M_3 represent the number of cells undergoing first, second, and third mitosis and N is the total number of metaphase scored.

Chromosomal aberrations assay

CAs assay was performed according to [44]. For each subject, a total of 200 well-spread first-division complete metaphases were analyzed for the following categories of CAs: chromatid breaks (B'), chromosome breaks (B''), dicentric (Dic), acentric fragments (AF), and tri- or tetra-

radials (RAD). Cells containing any type of chromosomal aberrations were scored as "cells with aberrations" (CAB).

DNA Extraction and Genotyping

Genomic DNA was extracted using the Chelex solution protocol [51]. *GSTM1* and *GSTT1* genotypes were determined by polymerase chain reaction (PCR) using primers described in [53] and [40], respectively. In addition, as internal control, a fragment of the β -globin gene was co-amplified using the primers 5'-CAACTCATCCACGTT-CACC-3' and 5'-ACACAAC-TGTGTTCACTAGC-3'. PCR reactions were carried out following the procedure described in [44]. Genotypes with homozygous deletion of the *GST* genes are identified as "*GST*-null", whereas genotypes having at least one copy of the gene are "*GST*-positive".

Statistical analysis

Statistical analysis was assessed using the SYSTAT software statistical package (version 10.0, Inc., Chicago, Illinois, USA). The non-parametric Wilcoxon test was used to compare the mean frequencies of SCEs, MNs and CAs between alcoholics and controls. Multiple regression analysis was used to evaluate the influence of age and years of employment on SCEs, CAs and MNs frequencies of both groups. All P values were two tailed and the level of statistical significance was set at $P < 0.05$ for all tests.

Table 2 Frequency of SCEs and RI values in metaphases of lymphocytes from alcoholics and controls

Groups	N	NSM	SCEs	SCEs/NSM (SE)	M_1	M_2	M_3	RI (SE)
<i>Alcoholics</i>	15	750	6,831	9.11 (0.33)^a	559	353	288	1.75 (0.04)^b
<i>GSTT1</i> -positive	10	500	4,581	9.16 (0.38) ^c	419	269	212	1.77 (0.04) ^f
<i>GSTT1</i> -null	5	250	2,250	9.00 (0.71) ^c	140	84	76	1.71 (0.06) ^f
<i>GSTM1</i> -positive	10	500	4,528	9.06 (0.48) ^d	416	278	206	1.76 (0.05) ^g
<i>GSTM1</i> -null	5	250	2,303	9.21 (0.36) ^d	143	75	82	1.72 (0.05) ^g
<i>GSTs</i> double positives	8	400	3,709	9.27 (0.47) ^e	374	245	181	1.76 (0.05) ^h
<i>GSTs</i> double nulls	3	150	1,431	9.54 (0.51) ^e	159	81	60	1.67 (0.07) ^h
<i>Controls</i>	15	750	4,019	5.36 (0.30)^a	398	486	316	1.93 (0.02)^b
<i>GSTT1</i> -positive	9	450	2,523	5.61 (0.33) ⁱ	249	278	173	1.91 (0.02) ⁿ
<i>GSTT1</i> -null	6	300	1,496	4.99 (0.56) ⁱ	149	208	143	1.96 (0.04) ⁿ
<i>GSTM1</i> -positive	8	400	2,118	5.30 (0.44) [*]	207	242	151	1.91 (0.03) ^o
<i>GSTM1</i> -null	7	350	1,901	5.43 (0.42) [*]	191	244	165	1.96 (0.04) ^o
<i>GSTs</i> double positives	7	350	1,941	5.55 (0.42) ^m	247	280	173	1.89 (0.03) ^p
<i>GSTs</i> double nulls	5	250	1,319	5.28 (0.49) ^m	154	201	145	1.98 (0.05) ^p

Superscript letters indicate each comparison (a–p) performed between groups. In bold are highlighted the statistically significant differences (a and b)

SE Standard error; NSM Number of scored metaphases; SCEs Sister chromatid exchanges; RI Replication index = $(M_1 + 2M_2 + 3M_3)/N$, where M_1 , M_2 and M_3 represent the number of cells undergoing first second and third mitosis and N is the total number of metaphase scored

^{a,b} $P = 0.001$; ^c $P = 0.500$; ^d $P = 0.686$; ^{e,h} $P = 1$; ^f $P = 0.500$; ^g $P = 0.892$; ⁱ $P = 0.753$; ^m $P = 0.893$; ⁿ $P = 0.143$; ^o $P = 0.345$; ^p $P = 0.068$

^{*} $P = 0.612$

Table 3 Frequency of chromosomal aberrations in metaphases of lymphocytes from alcoholics and controls

Groups	N	NSM	CAs							CAs/NSM %	CAB/NSM %
			B'	B''	Dic	AF	RAD	Totals CAs	Totals CAB		
<i>Alcoholics</i>	15	3,000	17	8	1	2	1	29	29	0.97 (0.11)^a	0.97 (0.11)^b
<i>GSTT1</i> -positive	10	2,000	12	4	1	1	1	19	19	0.95 (0.11) ^c	0.95 (0.11) ^f
<i>GSTT1</i> -null	5	1,000	5	4	0	1	0	10	10	1.00 (0.13) ^c	1.00 (0.13) ^f
<i>GSTM1</i> -positive	10	2,000	13	3	1	0	1	18	18	0.90 (0.12) ^d	0.90 (0.12) ^g
<i>GSTM1</i> -null	5	1,000	4	5	0	2	0	11	11	1.10 (0.11) ^d	1.10 (0.11) ^g
<i>GSTs</i> double positives	8	1,600	10	2	1	0	1	14	14	0.88 (0.16) ^e	0.88 (0.16) ^h
<i>GSTs</i> double nulls	3	600	2	3	0	1	0	6	6	1.00 (0.29) ^e	1.00 (0.29) ^h
<i>Controls</i>	15	3,000	0	1	0	2	1	4	4	0.13 (0.08)^a	0.13 (0.08)^b
<i>GSTT1</i> -positive	9	1,800	0	1	0	1	1	3	3	0.17 (0.09) ⁱ	0.17 (0.09) ⁿ
<i>GSTT1</i> -null	6	1,200	0	0	0	1	0	1	1	0.08 (0.05) ⁱ	0.08 (0.05) ⁿ
<i>GSTM1</i> -positive	8	1,600	0	1	0	1	1	3	3	0.19 (0.10) [*]	0.19 (0.10) ^o
<i>GSTM1</i> -null	7	1,400	0	0	0	1	0	1	1	0.07 (0.05) [*]	0.07 (0.05) ^o
<i>GSTs</i> double positives	7	1,400	0	1	0	1	1	3	2	0.21 (0.15) ^m	0.14 (0.09) ^p
<i>GSTs</i> double nulls	5	1,000	0	0	0	1	0	1	1	0.10 (0.10) ^m	0.10 (0.10) ^p

Superscript letters indicate each comparison (a–p) performed between groups. In bold are highlighted the statistically significant differences (a and b)

^{a,b} $P = 0.002$; ^c $P = 1$; ^{d,g} $P = 0.257$; ^{e,h} $P = 0.157$; ^f $P = 1$; ⁱ $P = 0.317$; ^{*,o} $P = 0.414$; ^{m,n,p} $P = 0.317$

N Number of tested individuals; NSM Number of scored metaphases; CAs Chromosome aberrations; B' Chromatid breaks; B'' Chromosome breaks; Dic Dicentric chromosome; AF Acentric fragments; RAD Tri- or tetra-radials; CAB Cells with aberrations; AB.C % percentage of cells with aberrations

194 Results

195 Table 1 summarizes the main characteristics of the two studied
196 groups. Alcoholic subjects had a mean age of $50.73 \pm$
197 8.89 years (range: 36–70 years) and an average duration of
198 alcohol abuse of 7.67 ± 2.74 years (range: 4–12 years). The
199 mean age of controls was 49.33 ± 10.83 years (range: 38–70).
200 No significant differences were found between groups in terms
201 of mean age ($P = 0.944$).

202 Table 2 reports the SCEs frequencies recorded in
203 peripheral blood lymphocytes of alcoholics and control
204 subjects. A total of 750 metaphases for each group were
205 analysed. Statistically significant differences were found
206 between alcoholics and controls in both SCEs/cell fre-
207 quency ($P = 0.001$) and RI value ($P = 0.001$).

208 Table 3 shows the CAs frequencies in the two subject
209 groups. A total of 3000 metaphases for each group were
210 analysed. Statistically significant differences were found
211 between alcoholics and controls in terms of CAs and CAB
212 frequencies ($P = 0.002$).

213 Table 4 shows the frequencies of MNs and cells with
214 MNs in the studied groups. A total of 15,000 bi-nucleated
215 cells for each group were scored. No statistical significant
216 differences were found between alcoholics and controls in
217 the frequency of MNs ($P = 0.509$), cells with MNs
218 ($P = 0.394$) and CBPI values ($P = 0.691$).

219 Among both the alcoholics and control groups, we did
220 not find statistically significant associations between any
221 GST genes polymorphisms and the frequencies of SCEs
222 (Table 2), CAs (Table 3) or MNs (Table 4).

223 Results of multiple regression analysis are summarized
224 in Table 5. The duration of alcohol exposure (expressed as
225 years of alcohol abuse) was seen to have effects on the
226 frequencies of SCEs ($P = 0.015$), CAs ($P = 0.45$), but not
227 on MNs frequency ($P = 0.396$). Finally, in both groups, no
228 significant correlations were found between the age of
229 subjects and the frequencies of the analysed cytogenetic
230 biomarkers.

231 Discussion

232 Epidemiological studies conducted on different human
233 populations consistently showed that consumption of
234 alcoholic beverages is associated with a higher risk for
235 several types of cancer, such as oral, pharyngeal, colorectal
236 and liver cancers [21]. Nevertheless, the interpretation of
237 some cytogenetic studies is complicated by factors, such as
238 smoking or other potential confounders, not always taken
239 into account. Specifically, the effects of smoking and
240 alcohol consumption appear to be multiplicative. It seems
241 plausible that the synergism between tobacco and alcohol

Table 4 Frequencies of MNs, Cells with MNs and CBPI values in bi-nucleated lymphocytes from alcoholics and controls

Groups	N	NSCs	Distribution of BNCs according to the number of MNs						% MNs/NSCs (SE)	% CMNs/NSCs (SE)	CBPI (SE)
			1	2	3	4	Total MNs	Total CMNs			
<i>Alcoholics</i>	15	15,000	163	26	11	0	248	205	1.65 (0.23) ^a	1.37 (0.20) ^b	1.91 (0.06) ^c
<i>GSTT1</i> -positive	10	10,000	111	20	6	0	169	142	1.69 (0.23) ^d	1.42 (0.16) ^g	1.97 (0.05) ^l
<i>GSTT1</i> -null	5	5,000	52	6	5	0	79	163	1.58 (0.58) ^d	1.26 (0.54) ^g	1.78 (0.14) ^l
<i>GSTM1</i> -positive	10	10,000	93	17	5	0	142	120	1.42 (0.24) ^e	1.20 (0.19) ^h	1.97 (0.07) ^m
<i>GSTM1</i> -null	5	5,000	70	9	6	0	106	85	2.12 (0.48) ^e	1.70 (0.44) ^h	1.80 (0.25) ^m
<i>GSTs</i> double positives	8	8,000	85	16	4	0	129	110	1.61 (0.25) ^f	1.38 (0.19) ⁱ	1.99 (0.07) ⁿ
<i>GSTs</i> double nulls	3	3,000	44	5	4	0	66	53	2.20 (0.78) ^f	1.77 (0.44) ⁱ	1.73 (0.19) ⁿ
<i>Controls</i>	15	3,000	146	10	14	0	208	170	1.39 (0.28) ^a	1.13 (0.21) ^b	1.95 (0.07) ^c
<i>GSTT1</i> -positive	9	9,000	99	8	12	0	151	119	1.68 (0.43) ^o	1.32 (0.32) ^r	1.96 (0.10) ^u
<i>GSTT1</i> -null	6	6,000	47	2	2	0	57	51	0.95 (0.23) ^o	0.85 (0.18) ^r	1.94 (0.08) ^u
<i>GSTM1</i> -positive	8	8,000	75	8	7	0	112	90	1.40 (0.43) ^p	1.13 (0.31) ^s	1.85 (0.08) ^v
<i>GSTM1</i> -null	7	7,000	71	2	7	0	96	80	1.37 (0.40) ^p	1.14 (0.31) ^s	2.07 (0.10) ^v
<i>GSTs</i> double positives	7	7,000	66	7	7	0	101	80	1.44 (0.49) ^q	1.14 (0.36) ^t	1.86 (0.10) ^w
<i>GSTs</i> double nulls	5	5,000	38	1	2	0	46	41	0.92 (0.27) ^q	0.82 (0.22) ^t	1.96 (0.09) ^w

Superscript letters indicate each comparison (a–w) performed between groups
^a *P* = 0.509; ^b *P* = 0.394; ^c *P* = 0.691; ^d *P* = 0.686; ^e *P* = 0.225; ^f *P* = 0.593; ^g *P* = 0.786; ^h *P* = 0.343; ^{m,q,t} *P* = 0.500; ⁿ *P* = 0.285, ^o *P* = 0.462; ^p *P* = 0.499; ^{r,u} *P* = 0.600; ^s *P* = 0.553; ^v *P* = 0.173; ^w *P* = 0.498
NSCs Number of scored metaphases; BNCs Bi-nucleated cells; MNs Micronuclei; CMNs Cells with micronuclei; SE Standard error; CBPI (Cytokinesis-block proliferation index) = $[1 \times N_1] + [2 \times N_2] + [3 \times (N_3 + N_4)]/N$, where *N*₁–*N*₄ represent the number of cells with 1–4 nuclei, respectively, and *N* is the total number of cells scored

in the causation of cancer is due to the enhancement of the effects of tobacco carcinogens by ethanol [11]. While several previous studies have addressed this topic, this is the first study about non-smoker alcoholics.

We found a significantly high frequency of SCEs among alcoholics compared to healthy controls (Table 2). Moreover, the alcohol assumption appeared to influence the lymphocyte replication capacity, as shown by RI values significantly different between alcoholics and controls (*P* = 0.001). These results are concordant with a previously published study of [Butler et al. [7]] who observed an increase of SCEs frequency in alcoholics, that was not related to age, sex, cigarette smoking and duration in years of alcohol abuse. Similarly, [29] and [24] reported an increase in SCEs rates among alcoholics, although in those studies the higher SCEs frequency was associated to smoking habit and age. Considering that SCEs seem to reflect the repair efficiency of DNA lesions by homologous recombination, these results provide further support to the hypothesis that high alcohol consumption could be associated with impaired DNA repair mechanisms, as also suggested by other authors [16, 52].

Accordingly to other published studies [6, 10], among alcoholics we also found a significantly higher frequency of CAs compared to controls (Table 3). This finding could

suggest a possible clastogenic effect of ethanol in chronic alcoholics and a higher risk of cancer incidence among them. Indeed, it is known that cancer incidence among healthy individuals of a population increases with increased levels of CAs in their circulating lymphocytes [4, 5].

Although a possible aneugenic effect of ethanol has been evidenced by different authors [8, 19, 32, 49], in our study the MNs test did not reveal any significant difference between alcoholics and controls in terms of MNs frequency formation, as well as the number of cells with MNs and CBPI value (Table 4). Nevertheless, results reported in the above studies are referred to subjects that were both alcoholics and smokers, and thus the effects of smoking as a confounder factor should be taken into account. Vice versa, our selected sample included only non-smoking alcoholics, and thus the observed cytogenetic damage appeared to be exclusively due to the effects of alcohol consumption.

Overall, our data suggest that alcoholism may cause chromosome damage in humans, in terms of increased levels of SCEs and CAs. However, it cannot be concluded that the direct action of ethanol on chromosomes is responsible for these effects. Indeed, ethanol is quickly metabolized and significant levels of ROS and acetaldehyde accumulate in the blood during ethanol oxidation

Table 5 Multiple regression analysis of confounding factors on SCEs and CAs frequencies in peripheral lymphocytes of the study groups

CF	SCEs frequency			CAs frequency			MNs frequency		
	β -co	P value	95 % CI Lower-upper	β -co	P value	95 % CI Lower-upper	β -co	P value	95 % CI Lower-upper
Alcoholics									
Age	0.33	0.15	−0.94–5.70	0.21	0.40	−0.11–0.26	0.21	0.45	−0.39–0.82
YA	0.60	0.02	4.16–30.99	0.53	0.05	0.020–1.52	0.24	0.40	−1.46–3.44
Controls									
Age	0.37	0.18	−1.01–4.87	−0.05	0.87	−0.040–0.03	0.23	0.40	−0.35–0.83

CF Confounding factor; β -co β -coefficient; YA Years of alcohol abuse

[35]. This last compound cause many toxic effects associated with ethanol excess and may be responsible for its mutagenic activity [6].

Ethanol metabolizing process requires different types of enzymes, including GST enzymes, able to detoxify harmful ethanol metabolites in the liver by conjugating acetaldehyde and ROS to reduced glutathione [31]. Some *GST* alleles, such as *GSTM1* null allele, were found associated with liver diseases in alcoholics [34, 46]. However, among heavy and chronic ethanol consumers, no data are present in literature about a possible direct association between *GST* alleles and genomic damage measured in terms of SCEs, CAs and MNs. We investigated this possible association and observed in both groups, no statistically significant association between *GST* genes polymorphisms and the frequencies of SCEs, CAs, and MNs (Tables 2, 3, 4). However, the reduced sample size here considered requires further investigations with a larger number of subjects to provide convincing evidences for the absence of such correlation.

Finally, the increased levels of SCEs and CAs recorded among alcoholics appear to be associated with the duration of alcohol abuse, but not with the age of the subjects (Table 5). An age related increase of spontaneous chromosome instability was showed [2, 3, 13]. The lack of a similar age-related pattern among alcoholics could be probably due to an increased incidence of the chromosomal damage induced by alcohol among younger individuals. In this scenario the effect of alcohol abuse could obscure this relationship among alcoholics.

Acknowledgments This work was supported by Grants from the Italian Ministry of University and Scientific Research. We are grateful to Natascha Rogge for her valuable help and criticism and to all volunteers who participated in this study.

References

- Adler RA (1992) Clinical review 33: clinically important effects of alcohol on endocrine function. *J Clin Endocrinol Metab* 74:957–960
- Barnett YA, King CM (1995) An investigation of antioxidants status, DNA repair capacity and mutation as a function of age in humans. *Mutat Res* 338:115–128
- Bolognesi C, Abbondandolo A, Barale R et al (1997) Age-related increase of baseline frequencies of sister chromatid exchanges, chromosome aberrations, and micronuclei in human lymphocytes. *Cancer Epidemiol Biomark Prev* 6(4):249–256
- Bonassi S, Hagmar L, Strömberg U et al (2000) Chromosomal aberrations in lymphocytes predict human cancer independently of exposure to carcinogens. *Cancer Res* 60:1619–1625
- Bonassi S, Znaor A, Norppa H, Hagmar L (2004) Chromosomal aberrations and risk of cancer in humans: an epidemiologic perspective. *Cytogenet Genome Res* 104:376–382
- Burim RV, Canalle R, Takahashi CS et al (2004) Clastogenic effect of ethanol in chronic and abstinent alcoholics. *Mutat Res* 560(2):187–198
- Butler MG, Sanger WG, Veonett GE (1981) Increased frequency of sister-chromatid exchanges in alcoholics. *Mutat Res* 85(2):71–76
- Castelli E, Hrelia P, Maffei F, Fimognari C, Foschi FG, Caputo F, Cantelli-Forti G, Stefanini GF, Gasbarrini G (1999) Indicators of genetic damage in alcoholics: reversibility after alcohol abstinence. *Hepatogastroenterology* 46(27):1664–1668
- Crebelli R, Conti G, Conti L, Carere A (1989) A comparative study on ethanol and acetaldehyde as inducers of chromosome malsegregation in *Aspergillus nidulans*. *Mutat Res* 215:187–195
- Demirhan O, Tastemir D (2008) Cytogenetic effects of ethanol on chronic alcohol users. *Alcohol Alcohol* 43(2):127–136
- Du X, Squier CA, Kremer MJ, Wertz PW (2000) Penetration of N-nitrosornicotine (NNN) across oral mucosa in the presence of ethanol and nicotine. *J Oral Pathol Med* 29:80–85
- Elhajouji A, Cunha M, Kirsch-Volders M (1998) Spindle poisons can induce polyploidy by mitotic slippage and micronucleate mononucleates in the cytokinesis-block assay. *Mutagenesis* 13:193–198
- Fenech M (1998) Important variables that influence base-line micronucleus frequency in cytokinesis-blocked lymphocytes as biomarker for DNA damage in human populations. *Mutat Res* 404:155–165
- Fenech M (2000) The in vitro micronucleus technique. *Mutat Res* 455:81–95
- Fenech M, Chang WP, Kirsch-Volders M, Holland N, Bonassi S, Zeiger E (2003) Human Micronucleus project (HUMN project): detailed description of the scoring criteria for the cytokinesis-block micronucleus assay using isolated human lymphocyte cultures. *Mutat Res* 534(1–2):65–75
- Garcia-Sagredo JM (2008) Fifty years of cytogenetics: a parallel view of the evolution of cytogenetics and genotoxicology. *Biochim Biophys Acta* 1779(6–7):363–375
- Gattás GJ, Saldanha PH (1997) Chromosomal aberrations in peripheral lymphocytes of abstinent alcoholics. *Alcohol Clin Exp Res* 21:238–243

18. Hagmar L, Bonassi S, Stromberg U, Brogger A, Knudsen LE, Norppa H, Reuterwall C (1998) Chromosomal aberrations in lymphocytes predict human cancer: a report from the European study group on cytogenetic biomarkers and health (ESCH). *Cancer Res* 58:4117–4121
19. Iarmarcovai G, Bonassi S, Sari-Minodier I, Baciuchka-Palmaro M, Botta A, Orsiere T (2007) Exposure to genotoxic agents, host factors, and lifestyle influence the number of centromeric signals in micronuclei: a pooled re-analysis. *Mutat Res* 615:18–27
20. IARC (1988) Alcohol drinking. international agency for research on cancer. Monographs on the evaluation of carcinogenic risk to human, vol 54. International Agency for Research on Cancer, Lyon, pp 251–321
21. IARC (2010) Alcohol consumption and Ethyl Carbamate. Monographs on the evaluation of carcinogenic risk to human, vol 96. International Agency for Research on Cancer, Lyon, pp 173–1001
22. Inoue M, Tajima K, Hirose K, Kuroishi T, Gao CM, Kitoh T (1994) Life-style and subsite of gastric cancer-joint effect of smoking and drinking habits. *Int J Cancer* 56:494–499
23. Kafer E (1984) Disruptive effects of ethyl alcohol on mitotic chromosomes segregation in diploid and haploid strains of *Aspergillus nidulans*. *Mutat Res* 135:53–75
24. Karaoguz MY, Coşar B, Arıkan Z, Başaran F, Menevşe A, Menevşe S (2005) Increased frequency of sister chromatid exchanges in peripheral lymphocytes of alcoholics and cigarette smokers. *Cell Biol Int* 29(2):165–168
25. Kaufman MH (1983) Ethanol-induced chromosomal abnormalities at conception. *Nature* 302:258–260
26. Kaufman MH (1985) An hypothesis regarding the origin of aneuploidy in man: indirect evidence from an experimental model. *J Med Genet* 22:171–178
27. Kaufman MH (1997) The teratogenic effects of alcohol following exposure during pregnancy, and its influence on the chromosome constitution of the pre-ovulatory egg. *Alcohol Alcohol* 32:113–128
28. Kaufman MH, Bain IM (1984) Influence of ethanol on chromosome segregation during the first and second meiotic divisions in the mouse egg. *J Exp Zool* 230:315–320
29. Lazutka JR, Dedon V, Lekevicius RK (1992) Sister chromatid exchanges in lymphocytes of normal and alcoholic subjects. *Experientia* 48(5):508–512
30. Lee KW, Kuo WR, Tsai SM et al (2005) Different impact from betel quid, alcohol and cigarette: risk factors for pharyngeal and laryngeal cancer. *Int J Cancer* 117:831–836
31. Lieber CS (1994) Alcohol and the liver: update. *Gastroenterology* 106:1085–1105
32. Maffei F, Fimognari C, Castelli E, Stefanini GF, Forti GC, Hrelia P (2000) Increased cytogenetic damage detected by FISH analysis on micronuclei in peripheral lymphocytes from alcoholics. *Mutagenesis* 15(6):517–523
33. Mandayam S, Jamal MM, Morgan TR (2004) Epidemiology of alcoholic liver disease. *Semin Liver Dis* 24:217–232
34. Marcos M, Pastor I, Chamorro AJ, Ciria-Abad S, Gonzalez-Sarmiento R, Laso FJ (2011) Meta-analysis: glutathione-S-transferase allelic variants are associated with alcoholic liver disease. *Aliment Pharmacol Ther* 34:1159–1172
35. Nakamura K, Iwalashi K, Itoh M, Ameno K, Ijiri I, Takeuchi Y, Suwaki H (2000) Immunohistochemical study on acetaldehyde adducts in alcohol-fed mice. *Alcohol Clin Exp Res* 24(4):93S–96S
36. Nebert DW, McKinnon RA, Puga A (1996) Human drug metabolising enzyme polymorphisms effects on risk of toxicity and cancer DNA. *Cell Biol* 15:273–280
37. Nersesyan A, Muradyan R, Kundi M, Knasmueller S (2011) Impact of smoking on the frequencies of micronuclei and other nuclear abnormalities in exfoliated oral cells: a comparative study with different cigarette types. *Mutagenesis* 26(2):295–301
38. Obe G, Anderson D (1987) International commission for protection against environmental mutagens and carcinogens. IC-PEMC Working Paper No. 15/1. Genetic effects of ethanol. *Mutat Res* 186(3):177–200
39. Parry EM, Parry JM, Corso C et al (2002) Detection and characterization of mechanisms of action of aneugenic chemicals. *Mutagenesis* 17(6):509–521
40. Pemble S, Schroeder KR, Spencer SR, Meyer DJ, Hallier E, Bolt HM, Ketterer B, Taylor JB (1994) Human glutathione S-transferase theta (GSTT1): cDNA cloning and the characterization of a genetic polymorphism. *Biochem J* 300:271–276
41. Rey M, Palermo AM, Muñoz ER (1992) Nondisjunction induced by ethanol in *Drosophila melanogaster* females. *Mutat Res* 268(1):95–104
42. Roy B, Majumder PP, Dey B, Chakraborty M, Banerjee S, Roy M, Mukherjee N, Sil SK (2001) Ethnic differences in distributions of GSTM1 and GSTT1 homozygous “Null” genotypes in India. *Hum Biol* 73(3):443–450
43. Santovito S, Cervella P, DelPero M (2011) In vitro aneugenic effects of the fungicide thiabendazole evaluated in human lymphocytes by the micronucleus assay. *Arch Toxicol* 85:689–693
44. Santovito A, Schilirò T, Castellano S, Cervella P, Bigatti MP, Gilli G, Bono R, DelPero M (2011) Combined analysis of chromosomal aberrations and glutathione S-transferase M1 and T1 polymorphisms in pathologists occupationally exposed to formaldehyde. *Arch Toxicol* 85:1295–1302
45. Santovito S, Cervella P, DelPero M (2014) Increased frequency of chromosomal aberrations and sister chromatid exchanges in peripheral lymphocytes of radiology technicians chronically exposed to low levels of ionizing radiations. *Environ Toxicol Phar* 37:396–403
46. Savolainen VT, Pajarinen J, Perola M, Penttilä A, Karhunen PJ (1996) Glutathione-S-Transferase GST M1 “Null” genotype and the risk of alcoholic liver disease. *Alcohol Clin Exp Res* 20(8):1340–1345
47. Seitz HK, Stickel F (2006) Risk factors and mechanisms of hepatocarcinogenesis with special emphasis on alcohol and oxidative stress. *Biol Chem* 387(4):349–360
48. Sierra-Torres MS, Arboleda-Moreno YY, Hoyos LS, Sierra-Torres CH (2004) Chromosome aberrations among cigarette smokers in Colombia. *Mutat Res* 562(1–2):67–75
49. Stich HF, Rosin MP (1983) Quantitating the synergistic effect of smoking and alcohol consumption with the micronucleus test on human buccal mucosa cells. *Int J Cancer* 31(3):305–308
50. Vioque J, Barber X, Bolumar F, Porta M, Santibáñez M, de la Hera MG, Moreno-Osset E, PANESOS Study Group (2008) Oesophageal cancer risk by type of alcohol drinking and smoking: a case-control study in Spain. *BMC Cancer* 8(1):221
51. Walsh P, Metzger DA, Higuchi R (1991) Chelex 100 as a Medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques* 10:506
52. Wilson DM 3rd, Thompson LH (2007) Molecular mechanisms of sister-chromatid exchange. *Mutat Res* 616(1–2):11–23
53. Zhong S, Wyllie AH, Barnes D, Wolf CR, Spurr NK (1993) Relationship between GSTM1 genetic polymorphism and susceptibility to bladder, breast and colon cancer. *Carcinogenesis* 14:1821–1824