

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Frequency of aneuploidy in in vitro matured MII oocytes and corresponding first polar bodies in two dairy cattle (*Bos taurus*) breeds as determined by dual color fluorescent in situ hybridization (FISH)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1507155> since 2018-03-16T18:28:04Z

Published version:

DOI:10.1016/j.theriogenology.2009.10.007

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

D. Nicodemo, A. Pauciullo, G. Cosenza, V. Peretti, A. Perucatti, G.P. Di Meo, L. Ramunno, L. Iannuzzi, J. Rubes, D. Di Berardino. Frequency of aneuploidy in in vitro-matured MII oocytes and corresponding first polar bodies in two dairy cattle (*Bos taurus*) breeds as determined by dual-color fluorescent in situ hybridization. *Theriogenology*, 73 vol.4, year 2010, 523–529
DOI: 10.1016/j.theriogenology.2009.10.007

The publisher's version is available at:

<https://www.sciencedirect.com/science/article/pii/S0093691X09004671?via%3Dihub>

When citing, please refer to the published version.**Link to this full text:**

<https://www.sciencedirect.com/science/article/pii/S0093691X09004671?via%3Dihub>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

1 **Frequency of aneuploidy in in vitro–matured MII oocytes and corresponding first polar bodies**
2 **in two dairy cattle (*Bos taurus*) breeds as determined by dual-color fluorescent in situ**
3 **hybridization**

4

5 D. Nicodemo ^a, A. Pauciullo ^a, G. Cosenza ^a, V. Peretti ^b, A. Perucatti ^c, G.P. Di Meo ^c, L.

6 Ramunno ^a, L. Iannuzzi ^c, J. Rubes ^d, D. Di Berardino ^{a,*}

7

8

9 ^a Department of Soil, Plant, Environment and Animal Production Sciences, University of Naples

10 “Federico II,” Portici, Italy

11 ^b Department of Animal Science and Food Inspection, University of Naples “Federico II,” Naples,

12 Italy

13 ^c Research National Council (CNR), ISPAAM, Laboratory of Animal Cytogenetics and Gene

14 Mapping, Naples, Italy

15 ^d Veterinary Research Institute, Brno, Czech Republic

16

1 **Abstract**

2 The current study was undertaken to investigate the aneuploidy rates in in vitro–matured meiosis II
3 (MII) oocytes and corresponding first polar bodies in two dairy cattle (*Bos taurus*) breeds by using
4 dual-color fluorescent in situ hybridization (FISH). A total of 159 and 144 in vitro–matured MII
5 oocytes of the Italian Friesian and Italian Brown breeds, respectively, were obtained according to the
6 standard methods and analyzed by FISH using “Xcen” and “5” chromosome-specific painting
7 probes, produced by chromosome microdissection and Degenerate Oligonucleotide Primer-
8 Polymerase Chain Reaction (DOP-PCR). Oocytes with unreduced chromosome number were 10.1%
9 and 16.7% in the two breeds, respectively. To avoid bias due to possible artifacts, the aneuploidy
10 rates were determined by analyzing only oocytes with the corresponding polar bodies. In the Italian
11 Friesian, 100 of 143 (69.9%) secondary MII oocytes showed clear MII plates with corresponding first
12 polar bodies and were scored for aneuploidy detection; one oocyte was “nullisomic” for
13 chromosome X (1.0%) and one “disomic” for chromosome 5 (1.0%). In the Italian Brown, 100 of
14 120 (83.3%) MII oocytes with corresponding first polar bodies were analyzed; one oocyte was
15 nullisomic (1.0%) and one was disomic (1.0%), both for chromosome 5. Totally, 303 oocytes were
16 analyzed, 40 of which showed an unreduced chromosome complement (13.2%); of 200 MII oocytes
17 with the corresponding first polar bodies, the aneuploidy rate (nullisomy + disomy) for the two
18 chromosomes scored was 2%. Assuming that each chromosome is equally involved in aneuploidy, it
19 results that in cattle oocytes matured in vitro, at least 30% of the oocytes (1 x 30 haploid
20 chromosomes) should be aneuploid. Premature separation of sister chromatids (PSSC) was also
21 observed in 2% of the oocytes in the Italian Friesian breed involving chromosome 5 and in 1% of the
22 Italian Brown breed involving the X chromosome. Estimation of the “baseline” level of aneuploidy
23 in the in vitro–matured oocytes of the various domestic animal species and breeds is, to our opinion,
24 a useful reference for improving the in vitro production of embryos as well as for monitoring future
25 trends of the reproductive health of the species/breeds engaged in zootechnical productions,
26 especially in relation to management errors and environmental hazards.

1 **Keywords:** Aneuploidy; Bovine oocytes; Cattle breeds; Diploidy; FISH analysis; Polar bodies

2

3 **1. Introduction**

4 Bovine (*Bos taurus*) oocytes have mainly been investigated in the past for studying meiotic
5 progression and maturation under in vitro conditions [1–7]. Most of the authors analyzed only the
6 ploidy (reduced/unreduced) status of the oocytes, whereas only two articles [8,9] reported on the
7 aneuploidy rates. All these works used the conventional air-drying method [10], which, as is known,
8 allows visualization and counting of metaphase chromosomes at the MII stage. This method,
9 however, has serious limitations, due partly to the quality of the spreading, which is not always
10 optimal (overlapping chromosomes, presence of cytoplasmic residual, compacted metaphases) and
11 partly to the fact that it induces chromosomal loss due to spreading.

12 Under these circumstances, the aneuploidy rate can be underestimated or overestimated. In addition,
13 the chromosomal material of the corresponding first polar body cannot be analyzed because usually
14 the chromosomes are too condensed and overlapped. The possibility to detect aneuploidy also in
15 interphase cells (i.e., without the need to display metaphase chromosomes) came along with the
16 fluorescence in situ hybridization (FISH) technique [11] by using chromosome-specific “painting”
17 probes or Bacterial Artificial Chromosomes (BACs). This technique, if applied to MII oocytes with
18 the corresponding first polar body, can provide a more precise estimation of aneuploidy, because the
19 lack of any chromosome in the MII metaphase (nullisomy) should have its counterpart in the
20 corresponding polar body, which should therefore result disomic, and vice versa. The limiting factor
21 of this technique, however, is the scarce availability of the chromosome-specific probes. The PRINS
22 technique [12] is another interesting way to analyze aneuploidy, but because there are no specific
23 reports on domestic animals, we preferred to use the FISH approach. Because all the information we
24 have to date on the aneuploidy rates in cattle oocytes has been collected by using the conventional
25 air-drying method, we decided to review the available data in cattle aneuploidy by using the more
26 resolutive FISH technique applied on in vitro-matured MII oocytes plus first polar bodies by using

1 painting probes from chromosome X and chromosome 5 prepared by chromosome microdissection
2 and DOP-PCR. These probes were chosen because they both provided strong and specific signals.
3 Furthermore, while the Xcen probe was previously used for sperm FISH analysis [13], the probe for
4 chromosome 5 was specifically prepared for this purpose.

5 The current contribution analyzes the rate of aneuploidy in the Italian Friesian and Italian Brown
6 cattle breeds to estimate the “baseline” level of aneuploidy in the in vitro–matured oocytes of the
7 species *Bos taurus*. These breeds have been chosen because of their worldwide importance for milk
8 production. Once defined, this baseline level could be used as a reference for improving the in vitro
9 production of embryos as well as for monitoring future trends of the reproductive health of the
10 species/breeds engaged in zootechnical productions, especially in relation to management errors and
11 environmental hazards.

12

13 **2. Materials and methods**

14 **2.1. Karyotyping of donor cows**

15 Cows ready for slaughtering were previously karyotyped according to standard methods [14]. The
16 donor cows used in this study were all karyologically normal.

17

18 **2.2. Collection of ovaries and oocytes and in vitro maturation**

19 Ovaries were collected from slaughtered cows of the Italian Friesian and Italian Brown breeds, aging
20 between 12 and 36 mo and transported to the laboratory within 2 h. Cumulus-oocyte complexes
21 (COCs) were collected through aspiration with 21-gauge needles, washed in TC-199 medium (No.
22 M2154; Sigma, St. Louis, MO, USA), and examined on Petri dishes under a stereomicroscope. Only
23 oocytes with compact-intact cumulus cell layers and good morphology were selected for the study.
24 Groups of oocytes selected from each donor were transferred into 50- μ L droplets of maturation
25 medium consisting of TC-199 medium + 10% fetal bovine serum (No. 10106-151; Gibco, Invitrogen,
26 Carlsbad, CA, USA), supplemented with 0.5 mg/mL follicle-stimulating hormone (FSH; No. F8174;

1 Sigma), 5 mg/mL luteinizing hormone (LH; No. L5269; Sigma), and 1 mg/mL estradiol (No. E2257;
2 Sigma), covered with sterile mineral oil (No. M5310; Sigma) and allocated in a humidified
3 atmosphere containing 5% CO₂ in air at 39 °C for 24 h.

4

5 **2.3. Oocyte fixation**

6 After 24 h maturation, the COCs were incubated for a few minutes in a hyaluronidase solution (1
7 mg/mL; No. H4272; Sigma) to remove the cumulus cells, washed in Phosphate Buffered Saline
8 (PBS), and exposed to a hypotonic sodium citrate solution (0.8% wt/vol) for 3 min, followed by KCl
9 (75 mM) treatment for 3 min. The fixation was carried out using cold methanol/glacial acetic acid
10 (3:1) solution. Oocytes were individually fixed at the center of a precleaned slide, air-dried, and kept
11 at -20°C until analysis.

12

13 **2.4. Chromosome microdissection and probes preparations**

14 Metaphase cells for the production of probes via microdissection were prepared according to the
15 standard cytogenetic techniques [14]. For microdissection, the fixed lymphocyte suspension was
16 spread onto a precleaned 24 x 60 mm coverslip, which was then airdried and treated for GTG-banding.
17 The Xcen probe was produced by isolating the pericentromeric region, corresponding with the
18 centromere and with the region Xp11-14 of the standardized GTG-banded karyotype [13]; the probe
19 for chromosome 5 was produced by scraping the entire chromosome. Microdissected chromosomes
20 were amplified following the protocol of Engelen et al. [15]. Probes were labeled with dUTP-11-
21 digoxigenin (chr. Xcen) and dUTP-16-biotin (chr. 5) (No. 11558706910 and No. 11093070910,
22 respectively; Roche, Mannheim, Germany) in a second DOP-PCR reaction using 2 mL of products
23 from the first reaction as template.

24

25 **2.5. In situ hybridization**

1 The Xcen and 5 probes were hybridized simultaneously on metaphase plates for validation and
2 subsequently used for oocytes analysis. Probes were precipitated in the presence of 10 mg salmon
3 sperm DNA (No. D7656; Sigma) and 10 mg of calf thymus DNA (No. D8661; Sigma) dissolved in
4 15 mL hybridization solution (50%formamide in 2X SSC + 10% dextran sulfate; No. F7503 and No.
5 D8906, respectively; Sigma) (SSC – Standard Saline Citrate), denatured at 72 8C for 10 min, and
6 incubated at 37 8C for 90 min. Fixed oocytes were denatured for 2 min in a solution of 70%
7 formamide in 2X SSC (pH 7.0) at 72 8C. The hybridization mixture containing the Xcen and 5 probes
8 was applied on the slides and covered with 24 _ 24 mm coverslips. The slides were hybridized in a
9 moist chamber at 37 8C overnight. After hybridization and slide washing, the biotin-labeled probe
10 was revealed using a green Alexa 488 fluorochrome conjugated to streptavidin (No. S-11223;
11 Invitrogen, Carlsbad, CA, USA), and the digoxigenin-labeled probe was revealed using a red
12 rhodamine fluorochrome conjugated to an antidigoxigenin antibody from sheep (No. 11207750910;
13 Roche). Slides were counterstained with DAPI (40,60-diamidino-2-phenylindole, 0.24 mg/mL) (No.
14 D9542; Sigma) in Antifade (No. H1000; Vector Laboratories, Burlingame, CA, USA).

15

16 **2.6. Fluorescence analysis and scoring**

17 The slides were observed at _ 100 magnification with a Leica (Wetzlar, Germany) DMRA
18 fluorescence microscope equipped with DAPI, Fluorescein isothiocyanate (FITC), and Texas Red
19 (TXRD) specific filters, the DAPI/FITC/TXRD triple filter, and phase-contrast optics. Digital images
20 were captured using the Leica Q4000 software. To avoid possible bias, reduced secondary oocytes
21 without the corresponding first polar bodies were excluded from the analysis. An oocyte was defined
22 as “nullisomic” when one of the two signals (either X or 5) was lacking from the MII plate but
23 present twice in the corresponding polar body; vice versa, an oocyte was defined as “disomic” when
24 one extra signal (either X or 5) was present in the MII plate but absent from the polar body. Chi-
25 square analysis was used for statistical analysis of data.

26

1 **3. Results and discussion**

2 The Xcen and 5 probes were hybridized simultaneously on cattle metaphase preparations for
3 validation of the probes; Fig. 1 shows the Xcen and 5 painting probes, the DAPI staining, and their
4 diagrammatic representation. Fig. 2 shows normal and abnormal in vitro–matured MII oocytes after
5 FISH by using Xcen and 5 chromosome-specific painting probes. The detailed results are shown in
6 Table 1. Animals with less than four reduced secondary oocytes with analyzable polar bodies (PBs)
7 were grouped. Totally, 549 COCs were collected through aspiration (295 and 254 in the Italian
8 Friesian and Italian Brown, respectively). On the average, 13 COCs were recovered from each subject
9 (range, 6 to 34). The percentage of COCs selected for in vitro maturation was around 70% in both
10 breeds, but it was different among the subjects (range of selected COCs varied between 54% and
11 82%). Of 204 and 179 COCs selected for maturation, respectively, in the Italian Friesian and Italian
12 Brown, 159 and 144 reached the MII stage; the efficiency of the in vitro maturation process was
13 thus 78% and 80% in the two breeds, respectively. Individually, the maturation rate varied between
14 57% and 100% among the subjects. Significant interindividual differences were found in the yield of
15 in vitro–matured MII oocytes in the Italian Friesian as well as in the Italian Brown breed. As shown,
16 these differences are mainly due to the number of oocytes obtained from individual donors and
17 selected for in vitro maturation and partly due to the different efficiency of the in vitro maturation
18 process in the different subjects. In the Italian Friesian, 16 oocytes of 159 showed an unreduced
19 chromosome number (10.1%); of 100 MII + PB analyzed, one oocyte was nullisomic for the X
20 chromosome and one disomic for chromosome 5, with an overall frequency of aneuploidy (nullisomy
21 + disomy) of 2%; other two oocytes (2%) were affected by PSSC involving chromosome 5. In the
22 Italian Brown, 24 oocytes of 144 showed an unreduced chromosome number (16.7%); of 100 MII +
23 PB analyzed, one oocyte was nullisomic for chromosome 5, and one was disomic for the same
24 chromosome, with an overall frequency of aneuploidy of 2%; only one oocyte (1%) showed PSSC
25 involving the X chromosome.

1 By considering the two breeds together, on the average, 40 oocytes of 303 (13.2%) were found to be
2 unreduced; of 200 MII + PB analyzed, 2 oocytes were nullisomic (1%), 2 oocytes were disomic (1%),
3 with an overall aneuploidy rate, for these two chromosomes, of 2%, nullisomy and disomy being
4 equivalent (i.e., 1% each). Totally, chromosome 5 showed a higher frequency of aneuploidy
5 compared with that of the X chromosome (1.5% vs. 0.5%, respectively), but this difference was not
6 statistically significant. The results of the current study demonstrated that in bovine oocytes matured
7 in vitro, the mean rate of unreduction was on the average 13.2%. This value is “within” the interval
8 already reported in the literature (around 8% to 12%) by use of conventional methods [3,5–8].
9 Considering the animal individually, 54% of the subjects showed one or more unreduced oocytes.
10 Again, this value is within the range of other studies [4,7]. In the current contribution, the Brown
11 breed showed a higher frequency compared with the Friesian (16.7% vs. 10.1%), but the difference
12 was not statistically significant.

13 Interestingly, chromosome 5 was found to be three times more involved in aneuploidy compared with
14 the X chromosome (1.5% vs. 0.5%, respectively); even though this difference was not statistically
15 significant, it is indicative of the existence of interchromosomal differences in the rate of
16 nondisjunction. Whereas in humans the interchromosomal differences have already been
17 demonstrated [16,17], in domestic animals to date they have only been suggested.

18 By assuming an average rate of aneuploidy of 1% per chromosome and an equal probability of each
19 chromosome being involved in aneuploidy, it results that in cattle, nearly 30% (1 x 30 haploid
20 chromosomes) of the in vitro–matured oocytes might be aneuploid.

21 This estimate is higher compared with 5.8% and 7.1% previously reported in cattle by Yadav et al. [8]
22 and by Lechniak and Switonski [9], respectively, by using conventional cytogenetic methods but very
23 close to that obtained in the pig by using the FISH technique. In fact, Vozdova´ et al. [18] estimated
24 that about 27% of the in vitro–matured pig oocytes are aneuploid, whereas conventional methods
25 indicated an average frequency of 4% to 5% [19,20]. In humans, the aneuploidy rates vary among
26 different laboratories, with the highest value reaching 47% [17]. By considering the results of this

1 study in cattle and those reported by Vozdova et al. in the pig [18], it seems that the FISH method
2 overestimates -compared with the classic method- the incidence of aneuploidy, but this is simply due
3 to the extrapolation from only two or three chromosomes, under the general assumption of no
4 interchromosomal differences (i.e., equal likelihood for each haploid chromosome). This assumption,
5 however, may not be true, and future studies should focus on this topic by using FISH with probes
6 for all the chromosomes. Balanced PSSC was also observed in 2% of the oocytes in the Italian
7 Friesian breed involving chromosome 5 and in 1% of the Italian Brown breed involving the X
8 chromosome. A balanced PSSC is characterized by balanced chromosomal sets with separation of
9 sister chromatids. This anomaly is not considered responsible for aneuploidies, although it may
10 indicate a predisposition to nondisjunction. On the contrary, unbalanced PSSC can lead to embryonic
11 aneuploidy in 50% of the cases, depending upon the behavior of the extra chromatid during the second
12 meiotic division. No oocytes with unbalanced PSSC were observed in this study.

13 As known, aneuploidy in in vitro-matured oocytes is strongly dependent upon the culture system
14 [21]; furthermore, it has been demonstrated that the percentage of embryos with chromosomally
15 abnormal cells increases from 25% to 72% in in vivo-produced and in vitro-produced bovine
16 embryos, respectively [22]. The whole in vitro embryo production system in cattle, starting from
17 immature oocytes, has an efficiency of 30% to 40% [23]; this low efficiency is mainly due to the high
18 rate of aneuploidy in the in vitro-matured oocytes used for fertilization. Further studies should,
19 therefore, concern the optimization of the current in vitro culture system for oocyte maturation.
20 Fertility in mammals is strongly affected by chromosomal abnormalities, which represent one of the
21 major causes of embryonic mortality, as it has been demonstrated in humans [24–27] and in domestic
22 animals [28–30]. Meiotic errors may originate spontaneously or may be due to the genotoxic effects
23 of a variety of environmental mutagens, hormonal unbalancements, nutritional and dietetical
24 mistakes, and so forth [17,31]. In any case, estimation of the baseline level of aneuploidy in the
25 oocytes of the various domestic species and breeds is an essential step for improving the in vitro
26 production of embryos destined to the embryo transfer industry as well as for monitoring future trends

1 of the reproductive health of domestic animals in relation to management errors and/or environmental
2 hazards.

3

4 **Acknowledgment**

5 This work was supported by the financial contribution of the Ministry of Agricultural and Forestry
6 Politics (MiPAAF) of Rome (SpermovoFISH project no. 291/7303/06), which is gratefully
7 acknowledged.

8

9 **References**

- 10 [1] King WA, Xu KP, Sirard MA, Greve T, Leclerc P, Lambert RD, Jacques P. Cytogenetic study of
11 parthenogenetically activated bovine oocytes matured in vivo and in vitro. *Gamete Res* 1988;20:265–
12 74.
- 13 [2] Suss U, Wuthrich K, Stranzinger G. Chromosome configuration and time sequence of the first
14 meiotic division in bovine oocytes matured in vitro. *Biol Reprod* 1988;38:871–80.
- 15 [3] Ectors FJ, Konlischer L, Jamar M, Herens C, Verloes A, Remy B, Beckers JF. Cytogenetic study
16 of bovine oocytes matured in vitro. *Theriogenology* 1995;44:445–50.
- 17 [4] Sosnowski J, Switonski M, Lechniak D, Molinski K. Cytogenetic evaluation of in vitro matured
18 bovine oocytes collected from ovaries of individual donors. *Theriogenology* 1996;45:865–72.
- 19 [5] Lechniak D, Switonski M, Sosnowski J. The incidence of bovine diploiid oocytes matured in
20 vitro. *Theriogenology* 1996;46:267– 77.
- 21 [6] Ocana-Quero JM, Pinedo-Merlin M, Moreno-Millan M. Influence of follicle size, medium,
22 temperature and time on the incidence of diploid bovine oocytes matured in vitro. *Theriogenology*
23 1999;51:867–72.
- 24 [7] Lechniak D, Kaczmarek D, Stanislavski D, Adamowicz T. The ploidy of in vitro matured bovine
25 oocytes is related to the diameter. *Theriogenology* 2002;57:1303–8.

- 1 [8] Yadav B, King W, Xu K, Pollard J, Plante L. Chromosome analysis of bovine oocytes cultured in
2 vitro. *Genet Sel Evol* 1991;23:191–6.
- 3 [9] Lechniak D, Switonski M. Aneuploidy in bovine oocytes matured in vitro. *Chromosome Res*
4 1998;6:504–1504.
- 5 [10] Tarkowski AK. An air-drying method for chromosome preparations from mouse eggs.
6 *Cytogenetics* 1966;5:394–400.
- 7 [11] Langer PR, Waldrop AA, Ward DC. Enzymatic synthesis of biotin-labelled polynucleotides:
8 novel nucleic acid affinity probes. *Proc Natl Acad Sci U S A* 1981;78:6633–7.
- 9 [12] Pellestor F, Girardet A, Andre´o B, Charlieu JP. A polymorphic alpha satellite sequence specific
10 for human chromosome 13 detected by oligonucleotide primed in situ labelling (PRINS). *Hum Genet*
11 1994;94:346–8.
- 12 [13] Nicodemo D, Pauciullo A, Castello A, Roldan E, Gomendio M, Cosenza G, et al. X-Y sperm
13 aneuploidy in two cattle (*Bos taurus*) breeds as determined by dual colour fluorescent in situ
14 hybridization (FISH). *Cytogenet Genome Res* 2009;126: doi:10.1159/000245923, in press.
- 15 [14] Iannuzzi L, Di Berardino D. Tools of the trade: diagnostic and research applied to domestic
16 animal cytogenetics. *J Appl Genet* 2008;49:357–66.
- 17 [15] Engelen JJ, Albrechts JC, Hamers GJ, Geraedts JP. A simple and efficient method for
18 microdissection and microFISH. *J Med Genet* 1998;35:265–8.
- 19 [16] Kuliev A, Cieslak J, Verlinskly Y. Frequency and distribution of chromosome abnormalities in
20 human oocytes. *Cytogenet Genome Res* 2005;111:193–8.
- 21 [17] Pacchierotti F, Adler I, Eichenlaub-ritter U, Mailhes JB. Gender effects on the incidence of
22 aneuploidy in mammalian germ cells. *Environ Res* 2007;104:46–69.
- 23 [18] Vozdova´ M, Machatkova´ M, Kubikova´ S, Zudova´ D, Jokesova´ E, Rubes J. Frequency of
24 aneuploidy in pig oocytes matured in vitro and of the corresponding first polar bodies detected by
25 fluorescent in situ hybridization. *Theriogenology* 2001;56: 771–6.

- 1 [19] Mcgaughey RW, Polge C. Cytogenetic analysis of pig oocytes matured in vitro. *J Exp Zool*
2 1971;176:383–95.
- 3 [20] Sosnowski J, Waroczyk M, Switonski M. Chromosome abnormalities in secondary pig oocytes
4 matured in vitro. *Theriogenology* 2003;60:571–81.
- 5 [21] A' Arabi SY, Roussel JD, Chandler JE. Chromosomal analysis of mammalian oocytes matured
6 in vitro with various culture systems. *Theriogenology* 1997;48:1173–83.
- 7 [22] King WA. Chromosome variation in the embryos of domestic animals. *Cytogenet Genome Res*
8 2008;120:81–90.
- 9 [23] Lonergan P. State-of-the-art embryo technologies in cattle. *Soc Reprod Fertil Suppl*
10 2007;64:315–25.
- 11 [24] Hecht F, Hecht BK. Environmental chromosome damage. *Am J Med Genet* 1987;27:399–400.
- 12 [25] Martin HR, Ko E, Rademaker A. Distribution of aneuploidy in human gametes: comparison
13 between human sperm and oocytes. *Am J Med Genet* 1991;39:321–31.
- 14 [26] Pellestor F. The cytogenetic analysis of human zygotes and preimplantation embryos. *Hum*
15 *Reprod Update* 1995;1:581–5.
- 16 [27] Hassold TJ. Nondisjunction in the human male. In: Handel MA, editor. *Meiosis and*
17 *Gametogenesis*. Academic Press; 1998. p.383–406.
- 18 [28] King WA. Chromosome abnormalities and pregnancy failure in domestic animals. *Adv Vet Sci*
19 *Comp Med* 1990;23: 229–50.
- 20 [29] Kawarski SJ, Basrur PK, Stubbings RB, Hansen PJ, King WA. Chromosomal abnormalities in
21 bovine embryos and their influence on development. *Biol Reprod* 1996;54:53–9.
- 22 [30] Vanroose G, de Kruif A, Van Soom A. Embryonic mortality and embryo-pathogen interactions.
23 *Anim Reprod Sci* 2000;60–61: 131–43.
- 24 [31] Maihes JB, Marchetti F. Mechanisms and chemical induction of aneuploidy in rodent germ cells.
25 *Cytogenet Genome Res* 2005; 111:384–91.