AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Nesting strategies affect altitudinal distribution and habitat use in Alpine dung beetle communities

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1508339> since 2016-03-09T15:29:47Z

Published version:

DOI:10.1111/een.12195

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Ecological Entomology 40: 372-380. 10.1111/een.12195

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://onlinelibrary.wiley.com/doi/10.1111/een.12195/abstract;jsessionid=A3A45192C935E4667347375E4A9923AA.f02t04>

**The influence of altitude, habitat and nesting strategy on key
parameters of ecosystem service provision in Alpine dung beetles**

Dan Chamberlain, Claudia Tocco, Alice Longoni, Stefano Mammola, Claudia Palestini & Antonio
Rolando*

Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università di Torino, Via Accademia
Albertina 13, Turin, Italy

*Corresponding author.

Email: dan.chamberlain99@gmail.com; Tel. +39 0116704534

Running head: Dung beetles along altitudinal gradients

Abstract

1. Dung beetles are key contributors to a suite of ecosystem services. Understanding the factors that dictate their diversity is a necessary step towards preventing negative impacts of biodiversity loss.
2. We analysed Alpine dung beetle communities along altitudinal gradients to assess how different components of the community, defined in terms of nesting strategy (dung-ovipositing Aphodidae [DOAs], soil-ovipositing Aphodidae [SOAs] and two paracoprid [PAR] groups, Geotrupidae and Onthophagini) and parameters relevant to ecosystem functioning (species richness, total biomass and functional diversity relative to dung removal), are distributed, and to identify to which environmental factors they respond.
3. Species richness declined with altitude. There was no significant variation in functional diversity, or total biomass in relation to altitude. There were significant variations when considered by nesting group: DOA species richness and biomass decreased, SOA biomass increased, and Geotrupidae biomass showed a non-linear trend, as altitude increased.
4. Functional diversity and total species richness were positively related to vegetation cover. DOA species richness was highest in forest and scrub, SOA species richness was highest in grassland and PAR species richness was lowest in rocky areas.
5. Dung beetle species show different trends in species richness and biomass depending on nesting strategy. Management to promote the dung beetle community should include maintenance of a mosaic of habitat types. Given the importance of species richness and biomass to ecosystem functioning, and the complimentary effect of different dung beetle groups, such a strategy will protect and enhance the ecosystem services that Alpine dung beetles provide.

Key words. Aphodidae, dung removal, ecosystem functioning, endocoprid, functional diversity, Geotrupidae, Onthophagini, paracoprid.

60 **Introduction**

61

62 Biodiversity is facing many pressures globally, leading to population declines, range contractions and
63 species extinctions (Butchart *et al.*, 2010), and consequent negative impacts on ecosystem services, causing
64 a loss of ecosystem functioning (Naeem *et al.*, 2012). Understanding the factors that dictate the diversity
65 of key providers of ecosystem services is therefore a necessary step towards halting or even reversing
66 damaging impacts of biodiversity loss.

67 Dung beetles contribute to a suite of important ecosystem services, including dung removal and
68 decomposition, nutrient cycling, plant growth enhancement and seed dispersal (e.g. Nichols *et al.* 2008,
69 Beynon *et al.*, 2012; Braga *et al.*, 2013). Dung beetle species show different nesting strategies, e.g.
70 paracoprids dig tunnels below the dung mass in which they bury brood balls, and endocoprids brood their
71 young inside the dung-mass itself, or at the soil-dung interface. The action of these different groups has
72 been shown to have a complementary effect on dung removal rates (Beynon *et al.*, 2012), although the
73 contribution of individual species is not equal, larger species having a disproportionate effect (Rosenlew &
74 Roslin, 2008). More generally, greater biomass has been shown to enhance decomposition rates for
75 decomposer communities (O’Hea *et al.*, 2010). A diverse and healthy dung beetle community is therefore
76 likely to enhance ecosystem services, and this effect may be particularly important in perturbed
77 ecosystems (Beynon *et al.*, 2012).

78 Mountainous areas are likely to be particularly sensitive to environmental change. In the European
79 Alps, negative impacts on biodiversity are expected due to increased disturbance (e.g. from leisure
80 activities, Negro *et al.*, 2009), changes in pastoral management (e.g. Marini *et al.*, 2009) and elevational
81 shifts in vegetation zones due to climate change (e.g. Cannone *et al.*, 2007). Dung beetles are likely to be
82 sensitive to such factors, in particular changes in livestock management (e.g. Tocco *et al.*, 2013) and climate
83 change (e.g. Menéndez *et al.*, 2014). Most studies of dung beetles in European mountains have taken place
84 in pastures at altitudes at or below the natural treeline (e.g. Jay-Robert *et al.*, 2008; Negro *et al.*, 2011,
85 Tocco *et al.*, 2013). However, there have been studies considering variations in overall abundance and
86 diversity along altitudinal gradients up to 3000m (e.g. Jay-Robert *et al.*, 1997).

87 In this paper, we advance the study of dung beetle ecosystem service provision by considering how
88 key parameters, shown experimentally to be associated with dung removal, respond to environmental
89 variations in the field, specifically by analyzing functional diversity, biomass and species richness along
90 altitudinal gradients (c. 1900-2800m) in the European Alps. Functional diversity of dung beetles has been
91 shown to be sensitive to land use changes (Barragàn *et al.*, 2011) and specifically to grazing management in
92 the Alps (Tocco *et al.*, 2013), whilst biomass and species richness have been shown to be linked to dung
93 removal rates, one of the key ecosystem functions provided by dung beetles (Beynon *et al.*, 2012; Nervo *et*

al., 2014). Given the differing contributions to ecosystem functioning between dung beetle nesting groups (e.g. Beynon *et al.*, 2012), we also consider the effect of dung beetle nesting strategy on these parameters.

The altitudinal gradient approach can prove very useful in understanding the distributions of species, as inferences can be drawn about larger-scale and longer-term processes from data collected at a relatively small spatial scale (Hodkinson 2005). We are unaware of any studies that have considered dung beetle functional diversity or biomass variations along altitudinal gradients in alpine (i.e. above the treeline) habitat, although species richness gradients have been considered (e.g. Jay-Robert *et al.* 1997). For all parameters, we consider both altitudinal gradients and habitat gradients, the latter being particularly important in terms of informing management strategies in a landscape subject to several environmental pressures. Furthermore, we control statistically for variations in trapping effort, variations in exposure time, and variations in dung resources in the surrounding area of each sampling point, a potentially important variable which has yet to be taken into account in studies of Alpine dung beetles. Our over-arching goals are therefore to clarify how the different components of the assemblage, defined in terms of nesting strategy and parameters relevant to ecosystem functioning, are distributed, and to identify to which environmental factors they respond.

Methods

The study was carried out in 2010 and 2011 at five sites in the mountains of the western Italian Alps in Val Gressoney, Val Grande, Val Chisone, Val Troncea and Val Argentera (Fig. 1). At each site, four sampling points were identified along gradients ranging from c. 1700m to c. 2800m in altitude, each one in a different habitat type: larch forest, scrub, alpine meadow and grass/rock mosaics. Between four and eight hang-bait traps (Palestrini *et al.*, 2008), baited with c. 200g of fresh cow dung, were placed at each point at a minimum distance of 20m. Traps were first set between mid- June and mid-July and were left for variable periods between visits (range 9 – 26 days). Note was made of any traps destroyed or removed. Dung beetles trapped were stored in 70% alcohol before later identification in the laboratory.

At each point, the percentage cover of different broadly-defined habitat types was estimated within 100m radius of the point: canopy, shrubs, field-layer vegetation, bare rock. The presence of dung on the first visit (i.e. dung of the previous year) and the presence of livestock within the current year at a point were noted. The topographic variables altitude, slope, easting and northing were extracted for each point from a Digital Terrain Model in GIS. Easting and northing were expressed as an index between 1 (facing directly south or east) and -1 (facing directly north or west) following Bradbury *et al.*, (2011). Site descriptions and precise locations are given in the Supplementary Material (Table S1).

Dung beetle nesting group

129

130 Dung beetles were defined into groups according to nesting strategy and taxonomy, based on Beynon *et al.*
131 (2012): dung-ovipositing Aphodidae (DOAs; species that live in the dung pad itself, and which either lay
132 eggs freely in the dung, or fashion brood balls into the dung pad), soil-ovipositing Aphodidae (SOAs; species
133 that construct the nest immediately below the dung pad in the soil-dung surface) and paracoprids (PARs;
134 species that deposit dung containing their eggs at the end of tunnels beneath the dung pad). Within
135 paracoprids, there is also considerable variation in ecology and morphology (Jay-Robert *et al.* 2008)
136 according to phylogenetic group, so for most analyses we also further divide paracoprids into Geotrupidae
137 (GEO) and Scarabaeidae, which in our case is represented only by the Onthophagini (ONT).

138

139 *Dung beetle parameters*

140

141 We considered three measures to describe the dung beetle community following Negro *et al.* (2011) and
142 Tocco *et al.* (2013): functional diversity (FD), species richness, and biomass. FD is a component of
143 biodiversity that refers to the range and value of organismal traits that influence ecosystem properties
144 (Hooper *et al.* 2002), and can be calculated by combining a matrix of species occurrence with data on
145 species traits. We focused on dung removal capability by taking into account four functional traits
146 following Tocco *et al.* (2013), using an ecomorphological approach in order to calculate the functional
147 diversity index (Petchey & Gaston, 2006). Functional traits used were nesting behaviour (as above), weight,
148 and two prothoracal leg allometries, all expected to be linked to dung removal efficiency. Dry weight,
149 body size (maximum width of the pronotum) and linear measurements of the prothoracal tibia for ten
150 randomly selected specimens within each species were recorded. These traits were then used in
151 conjunction with abundance data to calculate FD per trap, using the vegan package in R 3.03 (Oksanen *et*
152 *al.*, 2013). This translates species traits into pairwise distances among species from which clustering trees
153 of the traits are derived (Oksanen *et al.*, 2013). FD is the total branch length in a tree connecting all
154 species, and high values of FD indicate communities with a high diversity of functional traits.

155 The total biomass of dung beetles per trap was estimated by multiplying the abundance of each
156 species by its mean mass, and then calculating the sum over all species. As nesting group is intrinsic to the
157 expression of FD, this parameter was calculated only at the whole community level, and not specifically by
158 nesting group. Biomass and species richness were calculated for all groups combined, and for each nesting
159 group separately.

160

161 *Statistical Methods*

162

163 The data were analysed primarily through GLMMs which accounted for multiple observations from the
164 same locations by specifying site, point and (for FD and species richness) trap as random factors (see
165 below). Many dung beetle species show seasonal variations in abundance (e.g. Tocco *et al.*, 2013). Sites
166 had variable sampling periods (total trapping days varying from 54 to 95 days per site), so ‘month’ was
167 fitted as a three-level factor (i.e. June/July, August and September) in each model in addition to altitude,
168 habitat and/or group category variables (see below) in order to account for seasonal effects. All analyses
169 were initially carried out on all species combined. For biomass and species richness, the analytical
170 approaches were then repeated including the different nesting groups, and the interaction between
171 nesting group and the environmental variable of interest. Where there was a significant interaction, the
172 analysis was carried out on the groups separately.

173 Species richness is typically related non-linearly to sampling effort (Magurran, 2004), which varied
174 between visits due to trap loss. Therefore, species richness was analysed at the level of the individual trap
175 for each visit. Species accumulation curves were derived for each site using the vegan package in R in order
176 to assess sampling completeness. Species richness contributes to FD, and it is expected that FD will be
177 similarly sensitive to variation in trapping effort, therefore it was also analysed at the level of the individual
178 trap. Species richness and FD were analysed in relation to environmental variables using respectively the
179 glmer command (Poisson-distributed data) and the lmer command (normally distributed data) in the
180 package lme4 (Bates *et al.*, 2013) in R 3.0.1 (R Development Core Team, 2013). Non-independence of traps
181 within points and points within sites was accounted for by specifying each as a nested random factor,
182 specifically trap nested within point nested within site. Note FD can only be calculated where a species is
183 present (a single species equates to FD = 0 for a given trap), so this analysis had a lower sample size (there
184 were 38 trap/visits where no dung beetles were captured, leaving a total of 338 traps from 69 point/visits)
185 compared to other analyses. There was a significant positive linear effect of trapping effort on the total
186 biomass per point ($z = 2.59$, $P = 0.015$), so mean biomass was expressed per trap for each point and visit for
187 the analysis, allowing a simplified model structure, with point nested within site defined as random factors.
188 The lmer command was used for analysis, after applying a square-root transformation to normalise the
189 data.

190 For all analyses, a statistical hypothesis testing framework was adopted, whereby model reduction
191 was carried out on the full model by sequentially deleting non-significant terms until a minimum adequate
192 model of significant fixed effects remained. For normally distributed data (FD and biomass), P-values were
193 obtained from *F*-tests using the lmerTest package in R. For Poisson models (species richness), P-values
194 were based on *z*-tests.

195 The period between visits varied from 9 to 26 days. The first part of the analysis was therefore to
196 determine if the number of days for which a trap was left between visits (termed exposure time) was
197 related to a given parameter, as longer exposure times might result in more individuals and more species,

198 which would then have to be taken into account in subsequent models. The next goal was then to describe
199 the altitudinal variation in abundance, richness and diversity, hence the model contained altitude, altitude²
200 (to assess non-linear distributions), month, and exposure time if appropriate.

201 The next step was to identify other environmental correlates of species parameters. There was a
202 high level of inter-correlation between most of the habitat variables considered. PCA was therefore used to
203 describe community variation in relation to continuous environmental variables across the gradient
204 (canopy, shrubs, grass, rock, easting, northing, slope and altitude). The princomp command from the vegan
205 package (Oksanen *et al.*, 2013) was used, specifying the scale option. The relationships between habitat
206 gradients, as measured by the derived PCA axes, and dung beetle parameters were analysed using the
207 modelling approaches described above. These models also included the categorical fixed factors presence
208 of livestock and presence of dung. Nesting group and the interaction between nesting group and each PCA
209 axis was also included to assess whether different groups showed different responses to the habitat
210 gradient.

211 Altitude may be a proxy for a multitude of effects operating at various scales (Hodkinson, 2005). If
212 such factors are important, it may be expected that altitude models will perform better than habitat
213 models. If habitat models are better, this suggests that dung beetle communities may be significantly
214 influenced by habitat management. This was assessed by comparing model performance of both full
215 models and reduced models between those for altitude and those for habitat using Akaike's Information
216 Criterion corrected for small sample size (AICc; Burnham & Anderson, 2002).

217

218 **Results**

219

220 One point at the highest altitude in Val Grande was abandoned due to repeated losses of all traps, leaving a
221 total of 19 points sampled. For the five sites, altitudinal range was between 297m and 813m. There were a
222 total of 510 trap/visits placed (i.e. the sum of the number of times an individual trap was set across all sites
223 and visits), but 104 traps (20.4%) were destroyed or removed (due mostly to suspected large wild and
224 domestic mammals), leaving a total sample size of 406 trap/visits and 76 point/visits. Rates of trap loss
225 varied between sites from 11.3% ($n = 88$ traps set) in Val Argentera to 45.6% ($n = 68$) in Val Grande.

226 A total of 24 species and 12218 individuals of three families (Aphodiidae, Scarabaeidae and
227 Geotrupidae) were captured (a summary of the numbers and occurrence of each species caught, and their
228 nesting group, is given in Supplementary Material Table S2). The total number of species recorded in the
229 different sites was: Val Gressoney = 17, Val Grande = 9, Val Chisone = 19, Val Troncea = 20, Val Argentera =
230 11. Accumulation curves showed adequate sampling in all except Val Grande, which experienced the
231 highest rate of trap loss (Supplementary Material Fig. S1). This site was omitted from subsequent analyses,
232 leaving a total of 369 trap/visits and 69 point/visits, although results were generally very similar when

including these sites (full model details for models derived from four sites, and from all five sites, are given in Supplementary Material Table S3).

There was no significant effect of exposure time on species richness ($z = 1.54$, $P = 0.12$) or biomass ($F_{1,55.2} = 0.02$, $P = 0.93$), but there was a significant negative effect on FD ($F_{1,193.7} = 4.03$, $P = 0.032$). The general weak effect of exposure was probably because the dung bait only has a finite period during which it is attractive (Lumaret & Stienet, 1991). Exposure time was included in all subsequent models for FD, but was not considered further for biomass or species richness.

Distribution along the altitudinal gradient

There was a decrease in species richness with increasing altitude ($z = -2.21$, $P = 0.027$; Fig. 2A). There was no significant relationship between altitude and either FD or biomass. For species richness, there were only two species in ONT, therefore this was combined with GEO for the analyses. There was a significant interaction between altitude and nesting group indicated by significant differences between levels (test relative to PARs: AltA*DOAs, $z = -8.00$, $P < 0.001$; AltA*SOAs, $z = 6.404$, $P < 0.001$), so the three groups were subsequently analysed separately. There was a significant decrease in species richness of DOAs with increasing altitude ($z = -5.58$, $P < 0.001$; Fig. 2B), but no significant variation in the richness of SOAs. PARs showed a non-linear trend (Fig. 2C), with significant effects of both altitude ($z = 2.75$, $P = 0.006$) and altitude² ($z = -2.81$, $P = 0.005$). When considering only GEO, results were very similar in that there was also an overall interaction, and there were significant effects of both altitude ($z = 3.96$, $P < 0.001$) and altitude² ($z = -4.00$, $P < 0.001$). FD and most measures of species richness showed a decrease from July to September, with the exception of DOA richness (highest in August) and richness of paracoprid groups (PAR, GEO and ONT), none of which showed significant seasonal variation (Table S3).

There was a significant interaction between altitude and nesting group on biomass ($F_{3,253.7} = 10.85$, $P < 0.001$). SOAs showed a significant increase in biomass with altitude ($F_{1,13.7} = 8.72$, $P = 0.011$; Fig. 3A), whereas there was a significant decrease in DOAs ($F_{1,14.3} = 29.67$, $P < 0.001$; Fig. 3B), but no significant effect in GEO or ONT. There was no significant effect of altitude when considering biomass of all paracoprids combined ($F_{1,15.3} = 1.24$, $P = 0.28$). Seasonal trends in biomass broadly followed those of species richness, although both PAR and ONT biomass showed significantly higher values in July (Table S3).

Associations with habitat

The first two axes of the PCA explained 85.5% of variation in the data. The first axis described a gradient from open to closed habitat, being positively correlated with canopy (biplot score = 0.66) and negatively with grass (-0.63). The second axis was related to vegetation cover in general, being positively correlated

with grass cover (0.53) and to a lesser extent canopy (0.23), but negatively with rock cover (-0.80). The biplot of habitat scores for the first two axes is shown in Fig. 4. The topographic variables had little influence on these axes.

FD was significantly positively related to PCA2 ($F_{1,11.6} = 11.58$, $P = 0.003$; Fig. 5A), indicating lower values in habitat with greater rock cover. For species richness, there was a positive relationship with PCA2 ($z = 5.01$, $P < 0.001$; Fig. 5B). There were significant interactions between nesting group and both PCA1 (test relative to PARs: PCA1*DOA, $z = 9.63$, $P < 0.001$; PCA2*SOA, $z = -8.18$, $P < 0.001$) and PCA2 (PCA2*DOA, $z = 3.39$, $P < 0.001$; PCA2*SOA, $z = -4.80$, $P < 0.001$), so groups were analysed separately (figures are given in the Supplementary Material). DOAs had higher species richness at higher values of PCA1, i.e. more closed habitats ($z = 5.24$, $P < 0.001$), and at higher values of PCA2 i.e. well vegetated habitat ($z = 5.96$, $P < 0.001$), although species richness was lower when dung was present from the previous year ($z = -2.42$, $P = 0.016$, parameter estimate = -0.197 ± 0.081). SOAs conversely showed a negative association with PCA1 ($z = -2.68$, $P = 0.007$). They also showed a positive association with PCA2 ($z = 1.98$, $P = 0.047$) and a higher species richness when livestock were present ($z = 2.52$, $P = 0.012$, parameter estimate = 0.342 ± 0.136). PAR species richness was significantly higher at higher values of PCA2 ($z = 5.23$, $P = 0.001$), significantly higher when livestock were present ($z = 2.71$, $P = 0.007$, parameter estimate = 0.352 ± 0.137), but significantly lower when dung from the previous year was present ($z = -2.35$, $P = 0.019$ parameter estimate = -0.311 ± 0.133). However, when only GEO was considered, there was no significant effect of any habitat variable.

There was no overall effect of habitat on biomass, but there was a significant interaction between PCA1 and nesting group ($F_{2,253.2} = 12.67$, $P < 0.001$). Separate analyses showed that the two endocoprid groups had differing responses to the habitat gradient (see Supplementary Material Fig. S3), DOAs increasing in biomass with PCA1 ($F_{1,11.1} = 15.00$, $P = 0.003$) and PCA2 ($F_{1,16.9} = 4.45$, $P = 0.050$), but SOAs decreasing in biomass with PCA1 ($F_{1,10.3} = 14.87$, $P = 0.003$). There was no significant effect of habitat in either GEO or ONT, nor when both paracoprid groups were combined.

293

294 *Model comparison*

295

In the majority of cases, the habitat model was better (i.e. lower AICc) than the altitude model for species richness, the single exception being for the Geotrupidae full model (Table 1), whereas the altitude models clearly performed better for biomass (when environmental variables were included in the minimum adequate models, which was not always the case). FD was the only parameter to show inconsistencies between $\Delta AICc$ for the full model and reduced model, the altitude model being better for the former and the habitat model for the latter.

302

Discussion

There was a significant decrease in species richness, but no evidence of variation in FD or biomass of dung beetles, in relation to altitude. There were, however, significant variations in biomass and species richness in relation to altitude when considered by nesting group, but these patterns were often opposing. We are unaware of previous studies that have considered variations in the FD or biomass of dung beetle communities along altitudinal gradients. There is no consistent pattern across studies that have considered species richness in relation to altitude, with evidence of decreases (e.g. Jay-Robert *et al.*, 1997; Lobo *et al.*, 2007), increases (Negro *et al.*, 2011), non-linear effects (Escobar *et al.*, 2005) or no effect (Romero-Alcaraz & Ávila, 2000). The lack of stronger effects of altitude in our study seems likely to have been due to differing responses of the nesting groups, where DOAs showed a decreasing trend in species richness and biomass with altitude, and SOAs showed no effect on species richness but an increase in biomass with altitude. Paracopids showed a non-linear trend in biomass with altitude, a result largely driven by variations in Geotrupidae biomass. Jay-Robert *et al.* (1997) found a decrease in paracoprids, but an increase in all endocoprids, with altitude. Some of the variability in the findings of our and other studies is likely to result both from the length of the gradient and the maximum altitude considered, but geographic (Errouissi *et al.*, 2004), year-to-year (Jay-Robert *et al.*, 1997) and sampling period (see below) variation in the response to the altitudinal gradient may also be important.

Both FD and species richness were lower when rock cover was higher, demonstrating that dung beetles are linked to well-vegetated areas. Species richness of SOA and PAR was also higher when livestock were present, suggesting that resources in the area local to the sampling location will influence positively the number of species caught and will not therefore have a competition effect on the pitfall traps. There was a negative effect of presence of dung from the previous year on the species richness of DOAs. Given the role of dung beetles in dung decomposition, this result may indicate a slower rate of dung decomposition in sites where dung beetles are scarce – for example, in their absence, dung has been shown to take up to four years to fully decompose (Walters, 2008). A scarcity could be caused by particular local meteorological conditions, or could also arise if livestock in the area had been treated with pesticides such as avermectins, which may be passed into the dung which then becomes less attractive to the dung beetles (e.g. Webb *et al.* 2010).

Different groups also responded differently to habitat gradients, both in terms of species richness and biomass, in particular DOAs and SOAs showed opposing trends: both the richness and biomass of SOAs was higher in open grassland, but for DOAs they were higher in closed habitats. The different responses to habitat and altitude may be influenced by the physical factors affecting the dung. In particular, the opposing trends of DOAs and SOAs are probably driven by the different rates of dung dessication in closed and open habitats. DOAs require largely intact dung pads for a high rate of breeding success, and they are

likely to be more sensitive to dung dessication, hence their apparent preference for closed habitats and lower elevations. On the other hand, the burying activity of SOAs may mean they are less sensitive to higher rates of dung drying in open habitats. We suggest that the location of eggs and larvae on or under the soil surface is an adaptation to the harsh, high altitude environmental conditions, which enables SOAs to occur in great numbers in dung pads at high altitude (in keeping with the increase of SOA biomass with increasing altitude). Moreover, since dung drying makes the microhabitat structurally more suitable for predators to invade (Slansky & Rodriguez, 1987), the DOA strategy might result in greater exposure to predators in open habitats.

All paracoprids combined (PAR) showed higher species richness in vegetated habitats (i.e. PCA2), but otherwise, they did not show significant variation in relation to habitat gradients. Paracoprids excavate relatively deep tunnels beneath the dung pad, and hence they avoid rocky areas with little or no soil. More broadly, models for paracoprids, including those separately for Geotrupidae and Onthophagini, tended to explain less variation in the data compared to DOAs and SOAs (see pseudo- R^2 values in Table S3). This may be because paracoprid distribution is more closely linked to the ability to excavate deep tunnels, and therefore to soil structure (e.g. Sowig 1995), rather than broader measures of habitat variation.

Habitat models were better for describing patterns in FD and species richness, whereas altitude models were better for describing biomass (Table 1). This suggests that habitat management alone is likely to have major effects on species richness of Alpine dung beetle communities. However, the biomass, which will be in part dictated by abundance and in part by the presence of larger species, is likely also to be strongly linked to other factors along the altitudinal gradient (Hodkinson, 2005) – in other words, the broad-scale habitat measures considered here may have been too coarse to identify key small-scale habitat elements for certain components of the community. Further research into effects of micro-habitat and micro-climate on dung beetles at high altitude would help resolve these issues.

Caveats on interpretation

There are a number of caveats that need to be made when interpreting the results. First, the sampling for the majority of points was carried out between July and mid-September. Other comparable studies have often also included samples from October (e.g. Lumaret & Stienet, 1991), or early June (e.g. Jay-Robert *et al.*, 1997), but there is much variation in sampling periods from published Alpine studies (e.g. May to October, Jay-Robert *et al.*, 2008; June and July only, Jay-Robert *et al.*, 1997). Our study may have missed some early and late species, and indeed for lower altitude points, species richness was towards the lower end of the range recorded in other studies that considered different periods. Conversely, species richness at the higher altitudes was higher than reported in some comparable studies (e.g. Lumaret & Stienet, 1991; Palestini *et al.*, 2008). The conclusions here must therefore be considered as relevant to the summer

373 period. Second, there was a high rate of trap loss in some sites, which was probably mostly due to
374 disturbance by livestock. For example, at Val Grande, the highest point suffered 100% trap loss due to the
375 continued presence of sheep. This issue seems to have been particularly severe in our area compared to
376 other similar studies, and may reflect a generally higher level of grazing. We were, however, able to
377 account for the variation in trapping effort between sites statistically using nested random factors. Third,
378 higher altitude species may be more adapted to, and hence more attracted to, dung of wild animals such as
379 Alpine Ibex *Capra ibex* L. and Chamois *Rupicapra rupicapra* L., so sheep dung may have been a better
380 alternative than the cow dung used as bait in the traps. Indeed, it has been shown to be preferred by a
381 number of species recorded in our study (Dormont *et al.*, 2007). Nevertheless, the vast majority of studies
382 cited here also used cow dung. Fourth, although we took a measure of dung resources from domestic
383 animals into account, we did not measure dung of wild animals. Both the amount and diversity of dung
384 could have affected local dung beetle communities. Further research on altitudinal patterns using different
385 types of bait, and accounting for all dung resources at a given sampling point, would be interesting
386 methodological topics to pursue.

387

388 *Management implications*

389

390 There are two general findings relevant to habitat management arising from this work. First, vegetated
391 habitats are important. Whilst this may seem self-evident, such a simple relationship may become more
392 important in the future if elevational treeline shifts continue at their current rate (Harsch *et al.*, 2009),
393 because rocky habitats are generally predominant at higher altitudes, but development of vegetation in
394 these areas is likely to be a slow process (e.g. Cannone *et al.*, 2007), resulting in asymmetric shifts in
395 vegetation zones which are likely to have negative consequences for the biodiversity of Alpine grasslands
396 (e.g. Chamberlain *et al.*, 2013). Second, different dung beetle species may show different trends in terms of
397 richness and biomass depending on their nesting strategy. In terms of management to promote the dung
398 beetle community, maintenance of a mosaic of habitat types would therefore seem necessary.
399 Management such as targeted grazing (Tocco *et al.*, 2013) has the potential to maintain a landscape mosaic
400 of open and closed habitat areas, particularly where treelines are advancing due to climate change and
401 changes in grazing practices. Given the likely importance of species richness and biomass to ecosystem
402 functioning, and given the likely complimentary effect of different dung beetle groups, such practices
403 ultimately will protect and enhance the ecosystem services that Alpine dung beetles provide.

404

405 **Acknowledgements**

406

407 We would like to thank Lindsay Hernandez-Nova and Andrea Battisti for help in the field, and Matteo Negro
408 for production of Fig. 1. DEC was funded by the People Programme (Marie Curie Actions) of the European
409 Union's Seventh Framework Programme FP7/2007-2013/. CT was funded by a grant from Turin University.
410 Part of the field work was supported by the Regione Autonoma Valle d'Aosta Phenoalps project.

411

412 **Author contributions**

413 The project was developed by DC and AR. Data were collected in the field by DC, CT, AL and SM. CT
414 identified specimens and took trait measurements in the lab. CP advised on dung beetle sampling methods
415 and identification. DC analysed the data and took the lead role in writing the paper. All co-authors
416 contributed significantly to the submitted manuscript through discussions and additions to the text.

417

418 **References**

419

420 Barragán, F., Moreno, C.E., Escobar, F., Halffter, G. & Navarrete, D. (2011) Negative impacts of human land
421 use on dung beetle functional diversity. *PLOS ONE*, **6**, e17976.

422

423 Bates, D., Maechler, M., Bolker, B. & Walker, S. (2013) *lme4: Linear mixed-effects models using Eigen and*
424 *S4*. R package version 1.0-5, <http://CRAN.R-project.org/package=lme4>

425

426 Beynon, S.A., Mann, D.J., Slade, E.M. & Lewis, O.T. (2012) Species-rich dung beetle communities buffer
427 ecosystem services in perturbed agro-ecosystems. *Journal of Applied Ecology*, **49**, 1365-1372.

428

429 Bradbury, R.B., Pearce-Higgins, J.W., Wotton, S., Conway, G.J. & Grice, P.V. (2011) The influence of climate
430 and topography in patterns of territory establishment in a range-expanding bird. *Ibis*, **153**, 336-344.

431

432 Braga, R.F., Korasaki, V., Andresen, E. & Louzada, J. (2013) Dung beetle community and functions along a
433 habitat-disturbance gradient in the Amazon: A rapid assessment of ecological functions associated to
434 biodiversity. *PLOS ONE*, **8**, e57786.

435

436 Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference. A practical information-*
437 *theoretic approach*, 2nd edn. Springer-Verlag, New York, USA.

438

439 Butchart, S. *et al.* (2010). Global biodiversity: indicators of recent declines. *Science*, **328**, 1164-1168.

440

441 Cannone, N., Sgorbati, S. & Guglielmin, M. (2007) Unexpected impacts of climate change on alpine
 442 vegetation. *Frontiers in Ecology and Environment*, **5**, 360-364.

443

444 Chamberlain, D.E., Negro, M., Caprio, E. & Rolando, A. (2013) Assessing the sensitivity of alpine birds to
 445 potential future changes in habitat and climate to inform management strategies. *Biological Conservation*,
 446 **167**, 127-135.

447

448 Dormont, L., Rapior, S., McKey, D. B. & Lumaret, J. P. (2007) Influence of dung volatiles on the process of
 449 resource selection by coprophagous beetles. *Chemoecology*, **17**, 23-30.

450

451 Errouissi, F., Jay-Robert, P., Lumaret, J. & Piau, O. (2004) Composition and structure of dung beetle
 452 (Coleoptera: Aphodiidae, Geotrupidae, Scarabaeidae) assemblages in mountain grasslands of the Southern
 453 Alps. *Annals of the Entomological Society of America*, **97**, 701-709.

454

455 Escobar, F., Lobo, J.M. & Halffter, G. (2005). Altitudinal variation of dung beetle (Scarabaeidae:
 456 Scarabaeinae) assemblages in the Colombia Andes. *Global Ecology and Biogeography*, **14**, 327-337.

457

458 Finn, J.A. & Gittings, T. (2003) A review of competition in north temperate dung beetle communities.
 459 *Ecological Entomology*, **28**, 1-13.

460

461 Harsch, M.A., Hulme, P.E., McGlone, M.S. & Duncan, R.P. (2009) Are treelines advancing? A global meta-
 462 analysis of treeline response to climate warming. *Ecology Letters*, **12**, 1040-1049.

463

464 Hodkinson, I.D. 2005. Terrestrial insects along elevational gradients: species and community responses to
 465 altitude. *Biology Reviews*, **80**, 489-513.

466

467 Hooper, D.U., Solan, N., Symstad, A., Diaz, S., Gassner, M.O., Buchmann, N. Degrange, V., Grime, P., Hulot,
 468 P., Mermillod-Blondin, F., Roy, J., Spehn, E. & van Peer, L. (2002) Species diversity, functional diversity and
 469 ecosystem functioning. *Biodiversity and Ecosystem Functioning. Synthesis and Perspectives* (ed. M.
 470 Loreau,, S. Naeem & P. Inchausti) pp 195-208. Oxford University Press, Oxford.

471

472 Jay-Robert, P., Lobo, J.M. & Lumaret, J.P. (1997) Altitudinal turnover and species richness variation in
 473 European montane dung beetle assemblages. *Arctic Antarctic and Alpine Research*, **29**, 196-205.

474

475 Jay-Robert, P., Lumaret, J.-P., & Lebreton, J.-D. (2008) Spatial and temporal variation of mountain dung
 476 beetle assemblages and their relationships with environmental factors (Aphodiinae: Geotrupinae:
 477 Scarabaeinae). *Annals of the Entomological Society of America*, **101**, 58-69.

478

479 Lobo, J.M., Chehlarov, E. & Guèorguiev, B. (2007) Variation in dung beetle (Coleoptera: scarabaeoidea)
 480 assemblages with altitude in the Bulgarian Rhodopes Mountains: A comparison. *European Journal of*
 481 *Entomology*, **104**, 489–495.

482

483 Lumaret, J.-P. & Stiernet, N. (1991) Montane dung beetles. *Dung Beetle Ecology* (ed. I. Hanski & Y.
 484 Cambefort) pp. 242-254. Princeton University Press, Princeton.

485

486 Magurran, A.E. (2004) *Measuring Biological Diversity*. Blackwell, Oxford.

487

488 Marini, L., Fontana, P., Klimek, S., Battisti, A. & Gaston, K.J. (2009) Impact of farm size and topography on
 489 plant and insect diversity of managed pastures in the Alps. *Biological Conservation*, **142**, 394-403.

490

491 Menéndez, R., González-Megías, A., Jay-Robert, P. & Marquèz-Ferrando, R. 2014. Climate change and
 492 elevational range shifts : evidence from dung beetles in two European mountain ranges. *Global Ecology*
 493 *and Biogeography*, **23**, 646-657.

494

495 Naeem, S. *et al.* 2012. The Functions of Biological Diversity in an Age of Extinction. *Science*, **336**, 1401–1406.

496

497 Negro, M., Isaia, M., Palestini, C. & Rolando, A. (2009) The impact of forest ski-pistes on diversity of
 498 ground-dwelling arthropods and small mammals in the Alps. *Biodiversity and Conservation*, **18**, 2799–2821.

499

500 Negro, M., Palestini, C., Giraudo, M.T. & Rolando, A. (2011) The effect of local environmental
 501 heterogeneity on the diversity of species of alpine dung beetles. *European Journal of Entomology*, **108**, 91-
 502 98

503

504 Nervo, B., Tocco, C., Caprio, E., Palestini, C. & Rolando, A. (2014) The effects of body mass on dung removal
 505 efficiency in dung beetles. *PLoS ONE*, **9**, e107699.

506

507 Nichols, E., Spector, S., Louzada, J., Larsen, T., Amequita, S. & Favila, M.E. (2008) Ecological functions and
 508 ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation*, **141**, 1461-1474.

509

510 O’Hea, N.M., Kirwan, L. & Finn, J.A. (2010) Experimental mixtures of dung fauna affect dung decomposition
 511 through complex effects of species interactions. *Oikos*, **119**,1081-1088.
 512

513 Oksanen, J., Guillaume Blanchet, F., Kindt, R., Legendre, P., Minchin, P.R., O’Hara, R.B., Simpson, G.L.,
 514 Solymos, P., Henry, M., Stevens, H. & Wagner, H. (2013) vegan: Community Ecology Package. R package
 515 version 2.0-8.
 516

517 Palestrini, C., Roggero, A., Negro, M., Quaglia, E., Rovei, R. & Barbero E. (2008). A study on dung beetle
 518 fauna (Coleoptera: Scarabaeoidea) in some areas of Mont Avic Regional Natural Park (Valle d’Aosta, Italy).
 519 *Revue Valdôtaine Histoire Naturelle*, **61/62**, 189-217.
 520

521 Petchey, O.L. & Gaston, K.J. (2006) Functional diversity: back to basics and looking forward. *Ecology Letters*,
 522 **9**, 741–758.
 523

524 R Development Core Team, 2013 - *R: A language and environment for statistical computing*. R Foundation
 525 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
 526

527 Romero-Alcaraz, E. & Ávila, J.M. (2000) Effect of elevation and type of habitat on the abundance and
 528 diversity of Scarabaeoid dung beetles (Scarabaeoidea) assemblages in a Mediterranean area from southern
 529 Iberian Peninsula. *Zoological Studies*, **39**, 351-359.
 530

531 Rosenlew, H. & Roslin, T. (2008) Habitat fragmentation and the functional efficiency of temperate dung
 532 beetles. *Oikos*, **117**, 1659-1666.
 533

534 Slansky, Jr. F, F. & Rodriguez, J. G. (1987). *Nutritional ecology of insects, mites, spiders, and related*
 535 *invertebrates*. John Wiley, New York.
 536

537 Sowig, P. (1995) Habitat selection and offspring survival rate in three Paracoprid dung beetles: The
 538 influence of soil type and soil moisture. *Ecography*, **18**, 147-154.
 539

540 Tocco, C., Probo, M., Lonati, M., Lombardi, G., Negro, M., Nervo, B., & Rolando, A. & Palestrini, C. (2013)
 541 Pastoral practices to reverse shrub encroachment of sub-alpine grasslands: dung beetles (Coleoptera,
 542 Scarabaeoidea) respond more quickly than vegetation. *PLoS ONE*, **8**, e83344.
 543

544 Walters, C. (2008) *Dung Beetles: The greatest invention and a cowman’s profits*. Acres U.S.A. Austin, Texas.

545

546 Webb, L., Beaumont, D.G., Nager, R.G. & McCracken, D.I. (2010) Field-scale dispersal of *Aphodius* dung
547 beetles (Coleoptera: Scarabaeidae) in response to avermectin treatments on pastured cattle. *Bulletin of*
548 *Entomological Research*, **100**, 175-183.

549

550 **Table 1.** A comparison of performance of altitude and habitat models based on AICc. Full models were
551 considered which included either altitude and altitude² (altitude models), or PCA1, PCA2, presence of cattle
552 and presence of dung (habitat models). Month was in all models. ΔAICc is the difference in AICc between
553 habitat and altitude models, where $\Delta\text{AICc} > 2$ suggests altitude is the 'best' model and $\Delta\text{AICc} < -2$ suggests
554 habitat is the 'best' model. ΔAICc MAM is the same comparison but considering the final reduced models
555 (see Table S3). In some cases, no environmental variables were significant, in which case ΔAICc MAM is
556 zero (e.g. when only month was significant). Note that species richness was not analysed for Onthophagini.
557

Group	Parameter	ΔAICc	ΔAICc MAM
All	FD	3.7	-2.3
	Species richness	-13.4	-12.4
	Biomass	31.7	0
Dung-ovipositing Aphodidae	Species richness	-12.7	-10.7
	Biomass	35.8	26.6
Soil-ovipositing Aphodidae	Species richness	-7.2	-6.3
	Biomass	32.9	7.9
Paracoprids (all)	Species richness	-2.7	-9.4
	Biomass	35.3	0
Geotrupidae	Species richness	5.88	0
	Biomass	35.4	0
Onthophagini	Biomass	36.4	0

558

559

Fig. 1. Map of study area showing location of each sampling site. Shading indicates altitudes above 1700m. From north to south the sites were Val di Gressoney, Valli di Lanzo, Val Chisone, Val Troncea, Val Argentera (see text for precise coordinates).

Fig. 2. Variation in species richness of dung beetles in relation to altitude. A. All species. B. Dung-ovipositing Aphodidae (DOA). C. Paracoprids (PAR). Lines fitted from Poisson GLMMs with site, point and trap fitted as random effects ($n = 376$). Symbol sizes represent the number of observations for a given point (range: A = 1 to 11; B = 1 to 26; C = 1 to 31).

562
563
564

Fig. 3. Variation in the biomass of different groups of dung beetles across an altitudinal gradient, defined according to reproductive strategy. A. Dung-ovipositing Aphodidae (DOA). B. Soil-ovipositing Aphodidae (SOA). Filled circles show mean biomass (g) per trap for each point and each visit, square-root transformed. The lines were fitted from linear mixed models which included site and sampling point as random effects. N = 69 point/visits. Larger symbols indicate 2 observations for a given point.

565
566
567

Fig. 4. Bi-plot of PCA scores for the first two axes based on habitat data collected at 16 sampling points along altitudinal gradients at four sites. Each point is classified from 1 to 4 according to the rank in altitude at each site, and corresponds approximately to forest (circles), shrubs (diamonds), continuous grassland (squares) and high altitude grassland, rock and scree (triangles). Arrows show the loadings for each environmental variable. Note that the variables easting, northing and altitude had negligible scores (<5) on each axis and are not shown in the figure.

Fig. 5. Variation in dung beetle functional diversity and species richness in relation to a habitat gradient as measured by PCA score (PCA2 which represents a gradient from high rock cover to high vegetation cover. – see Fig. 4). A. Functional diversity per trap. B. Species richness (all species) per trap. Lines fitted from GLMMs with site, point and trap fitted as random effects ($n = 376$ trap). Symbol sizes represent the number of observations for a given point (range: A = 1 to 2; B = 1 to 4).