AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Aero-dispersed mutagenicity attributed to particulate and semi volatile phase in an urban environment

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/150871> since 2016-11-30T12:56:33Z

Published version:

DOI:10.1016/j.chemosphere.2014.12.033

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in CHEMOSPHERE, 124 (-), 2015, 10.1016/j.chemosphere.2014.12.033.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.chemosphere.2014.12.033

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0045653514014611>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/150871>

TITLE page:

Aero-dispersed mutagenicity attributed to particulate and semi volatile phase in an urban environment.

Short running title:

Particulate and non-particulate mutagenicity in an urban environment

AUTHORS: Deborah Traversi^{1*}, Festa Evelina¹, Cristina Pignata¹, Giorgio Gilli¹

AFFILIATIONS:

¹ Department of Public Health and Pediatrics, University of Torino, piazza Polonia 94, 10126 Torino, Italy

***CORRESPONDING AUTHOR:**

Deborah Traversi

tel +390116705822

fax +390116705874

Department of Public Health and Pediatrics, University of Torino, piazza Polonia 94, 10126 Torino, Italy

e-mail: deborah.traversi@unito.it

KEY WORDS: particulate matter, mutagenicity, , urban air pollution, gas phase pollution

ABSTRACT

Commonly the atmospheric pollution research is focussed on particulate indicators especially when mutagenicity was studied. On the other hand the volatile and semi-volatile compounds not adsorbed on to the particles can be genotoxic and mutagenic. Moreover some mutagenic compounds, such as polycyclic aromatic hydrocarbons, are present both in the particulate and in the gas-phase in according to chemical conditions. This work is focussed on the assessing of the total mutagenicity shifting the gas-phase and

particulate phase, during two seasons, in Turin. Two sampling sessions are conducted for total particulate matter and gas-phase pollutants. Moreover meteorological and usual air pollution monitoring data were collected at the same sampling station. The Salmonella assay using the strains TA98 and YG1021 was conducted on each organic extract. The mean level of total suspended particles, PM₁₀ and PM_{2.5} were 73.63 ± 26.94 , 42.85 ± 26.75 and $31.55 \pm 26.35 \mu\text{g}/\text{m}^3$. The observed mutagenicity was PM induced YG1021 > PM induced TA98 > PM induced TA98+S9 >> non-particle induced YG1021 > non-particle induced TA98 > non-particle induced TA98+S9. The multivariate regression is significant when we consider air pollution and meteorological indicators and chemical conditions as predictors.

HIGHLIGHTS

- Both chemicals and meteo-chemical parameters can influence the mutagenicity of air pollution.
- The gas phase and particulate phase mutagenicity can be different and affected by season.
- The gas phase accounted for only 1% of the observed mutagenicity.
- The particulate mutagenicity is approximately 5-fold higher during winter.
- The contribution of the nitro-derived compounds seems to be crucial.

1. Introduction

Air pollution is one of the most important worldwide health concerns (WHO-Europe, 2013). Particularly in the last 10 years, in both the US and Europe, new directives and regulations supporting more restrictive pollution limits were published (Krzyzanowski, 2008). However, the early effects of air pollution cannot be avoided, especially for the urban population (EEA, 2012). A recent Eurobarometer survey showed that European citizens are deeply concerned about the impact of air pollution and that more than 70% of the European population is worried that air pollution and air quality is worsening over time (EU, 2013). The decision to designate 2013 as the Year of Air reflects both the economic seriousness of the problem but also the impacts on humans. Approximately 3 % of cardiopulmonary and 5 % of lung cancer deaths are attributable to particulate matter (PM) pollution worldwide (HEI, 2013), while the disease burden related specifically to PM_{2.5} pollution accounts for approximately 3.1% of the global disability-adjusted life years

55 (Lim et al., 2012).

56 The total suspended particulate (TSP) air pollution is widespread and consists of a mixture of solid and
57 liquid particles suspended in the air. The physical and chemical characteristics of TSP vary by site. Common
58 chemical constituents of PM include sulphates, nitrates, ammonium and other inorganic ions, but also
59 include organic carbon, crustal material, particle-bound water, metals, aromatic hydrocarbons such as
60 polycyclic hydrocarbons and their nitrated, oxidised, sulphated forms (Claxton et al., 2004; Breyse et al.,
61 2013). Especially in urban polluted locations, the secondary particulates formed from precursor gases are
62 the prevalent toxic agents. Particle accumulation and coagulation reactions in the atmosphere produce a
63 fine fraction of particulate matter (PM_{2.5}) that often constitutes more than fifty percent of the TSP
64 (Dimitriou and Kassomenos, 2013). The emitted chemicals, the dispersion conditions, and physical
65 parameters such as humidity and temperature (Zhang et al., 2012) can all influence particle formation.
66 A large number of studies provide evidence of a correlation between both for short term and long-term
67 exposure to PM pollution and health effects such as morbidity and mortality from cardiovascular and
68 respiratory diseases, as well as from lung cancer (Krzyzanowski, 2008). At the end of 2013, outdoor air
69 pollution and its major component, outdoor particulate matter were classified as carcinogenic for humans
70 (1 Group) (Loomis et al., 2013).

71 Many mutagenic and genotoxic compounds are present in air pollution, and the effects are widely known
72 and reviewed (de Kok et al., 2006; Claxton and Woodall, 2007a; Valavanidis et al., 2008; DeMarini, 2013).
73 The finest air pollution fractions, PM₁₀ (particles with a diameter of less than 10 µm) and PM_{2.5} (particles
74 with a diameter of less than 2.5 µm) show greater genotoxicity (Claxton et al., 2004) , while the ultrafine
75 particles (particles having a diameter of less than 0.1 µm) are the subject of in-depth analyses (Hoek et al.,
76 2010; Kovats et al., 2013). The studies conducted using *in vivo* and *in vitro* models show the induction of
77 mutations and genotoxic effects. However, non-genotoxic effects also occur and various studies focused on
78 the epigenetic effects of the ambient particles (Ji and Hershey, 2012).

79 Among the typical air pollution chemicals, Polycyclic Aromatic Hydrocarbons PAHs have a relevant role in air
80 pollution toxicity. These compounds are reactive in the atmosphere and primarily form oxidised products,
81 the most notable being oxy-derivatives (mostly quinones) and nitrated compounds (Kim et al., 2013). Some

of these compounds, such as benzo(a)pyrene and 6-ditrochrysene and the 7,12-dimethylbenz(a)anthracene, are also present in primary emissions. Benzo(a)pyrene is the reference compound for the carcinogenic relative potency factor, while others previously cite PAHs as having a carcinogenic factor of 10 and 64, respectively. Also among the secondary PAHs are compounds with high carcinogenic relative potency factors such as benz(j)aceanthrylene (60) and 1,6-dinitropyrene (10) (ATSDR, 1995). The historic list of 16 USEPA priority PAHs is an important source of information, but was developed when knowledge of the relative toxicity of PAH congeners was more limited than at present. As such, it is useful as reference for monitoring but limited for the assessment of human health risks attributable to air PAH mixture exposition (Yang et al., 2007).

Vapor–particle partitioning of mutagens can be quantified using the gas–particle-partitioning coefficient for each compound. This coefficient is influenced by both the adsorption and absorption processes and is strongly temperature dependent (Albinet et al., 2008). Moreover, volatile and semi-volatile organic compounds associated with particulate matter can be influenced by heterogeneous photochemical reactions in the atmosphere (Fraser et al., 2000; Xie et al., 2013). Our typical samplings were conducted using standard methods that are affected by relevant limits (Liu et al., 2007; Forbes et al., 2012).

The aim of this work is to assess the mutagenicity of particulate and not-particulate air pollution and to determine the effects of seasonality and the contribution of nitro-compounds to the mutagenic effects in an urban environment.

100

101 **2. Materials and methods**

102 **2.1 Sampling**

103 Sampling was performed from 20 November to 22 December 2009 for the winter period and from 4 May to
104 4 June 2010 for the spring period at a meteorological–chemical station of the Environmental Protection
105 Regional Agency (Piedmont A.R.P.A.) located at Torino, in the northwest of the Padana Plain, Italy. The
106 sampling site, called Lingotto, was located outdoors in a small green area within an enclosed zone classified
107 as urban background (ARPA Piemonte, 2010). Turin has 872,367 inhabitants and a population density of
108 approximately 7,000 inhabitants per km²; thus, the pressure on the territory that is associated with human

activity is very high (ISTAT, 2012). Moreover, the climate and topographical characteristics of the area contribute to critical air pollution (Poncino et al., 2009; Eeftens et al., 2012). The Total Suspended Particles (TSP) were collected on glass micro-fibre filters (Type Fiberfilm T60A20, 150 mm, SKC, 863 Valley View Road Eighty Four, PA 15330, USA) and micro-pollutants were collected in Polyurethane Foam (PUF) Sorbent Tubes (SKC, 226-131 Valley View Road Eighty Four, PA 15330, USA) using an AirFlowPuf Sampler and conforming with the US EPA methods TO-4A, TO-9A, TO13A, ASTM D-6209 and ISO-12884, ISO-16362 (Analitica Strumenti Samplers, via degli Abeti 144 61100 Pesaro, Italy).

The TSP were collected on glass fibre filters, and the polyurethane foam (PUF) cartridge was placed in series after the glass fibre filter. The volatile compounds, which were not trapped on the filter, were retained in the PUF cartridge.

The sampling flow was electronically controlled to be 250 L/min. Each sampling duration was controlled by a timer that was accurate to ± 15 min over a 48-hour sampling period. The exact flow rate was calculated daily and corrected for variations in atmospheric pressure and actual differential pressure across the filter. The filters were conditioned for 48 h and were weighed using an analytical balance (± 10 μ g) before and after sampling to calculate the mass of the TSP trapped on the filter. The procedures were conducted according to the European Committee for Standardization. Additionally, , the PUF had been pre-cleaned by 24 h Soxhlet extractions using acetone.

126

127 **2.2 Extraction and mutagenicity assays**

Each sample was extracted with acetone in a Soxhlet apparatus for a minimum of 80 cycles. The samples were dried in a Rotavapor instrument, and suspended in dimethyl sulfoxide (DMSO) to obtain an equivalent concentration of 0.1 m³ of sampled air per μ l of solution. The mutagenicity assay was conducted as previously described (Maron and Ames, 1983; Traversi et al., 2009). Defined amounts of organic extract were tested to generate a dose–response curve (2, 5, 10, 20, 30 air equivalent m³ for the TSP extracts and 10, 20,30, 50, 100 air equivalent m³ for the PUF extracts). The slope of the dose–response curve (revertants/m³) was calculated by the least squares linear regression beginning at the first linear portion of the dose–response curve (Traversi et al., 2011). All experiments were performed in triplicate using at least

three doses. The results are expressed as net revertants per cubic metre (rev/m³) (the total revertants minus the spontaneous revertants) and were calculated using the dose–response curve (Cassoni et al., 2004; Claxton et al., 2004). The mutagenic activity of the airborne particulate extracts was determined using the *Salmonella typhimurium* strain TA98, with and without S9 mix, as well as the YG1021 strain. YG1021 is a ‘classical’ nitroreductase-overproducing strain obtained by cloning the nitroreductase gene of *S. typhimurium* TA1538 into the pBR322 vector and introducing the recombinant plasmid into TA98 (Josephy et al., 1997). YG1021 has a nitrofurazone reductase activity more than 50 times higher than the original TA98 strain, permitting the efficient detection of mutagenic nitroarenes. The mean number of spontaneous revertants, obtained during a 10 bioassay series, one every two samplings, was 16 ± 4 for TA98, 21 ± 1 for TA98+S9 and 23 ± 5 for YG1021. The genotype of each tester strain was routinely confirmed. In each assay session, positive and negative controls were included. Moreover, the known mutagen 2-nitrofluorene (1 µg/plate) was tested in each assay as a positive control for the strains TA98 and YG1021 while aminofluorene (1 µg/plate) was used as a positive control for the TA98 strain in presence of S9 mix.

150

151 **2.3 Chemicals and inhalable particles data**

152

Chemical data and inhalable particles data (PM₁₀ and PM_{2.5}) were extracted from a specialised database provided by the Regional System for the real-time monitoring of Air Quality, AriaWeb (ARPA Piemonte, 2014). The data were obtained for the same day as our sampling and for the same sampling station. For example, the NO_x data represent a monthly mean of hourly data collected using the standard monitoring method EN 14211:2005 (2008/50/EC, annex VI, section B). All the adopted methods conform to the directive and were validated before being published in the AriaWeb database (ARPA Piemonte, 2014).

159

160 **2.4 Statistics**

The seasons were designated as winter for the first sampling session (November and December) and as spring for the second sampling session (May and June). The statistical analyses were performed using the

163 SPSS Package, version 21.0. In particular we applied: (1) a log transformation of non-normally distributed
164 data, (2) the Spearman rank-order correlation coefficient to assess relationships between variables, (3) a
165 Wilcoxon-Mann-Whitney test to compare means. The mean differences and correlations were considered
166 significant if $p < 0.05$.

167

168 3. Results

169 3.1 Gravimetric analysis

170 The descriptive analysis of the collected data is shown in **Table 1**. The gravimetric data showed that, on
171 average, **meanly** the TSP proportion in the samples was 58% PM₁₀ and 43% PM_{2.5}. Moreover, during the
172 high pollution period in winter, these proportions increased up to 70% and 61%, with a PM_{2.5}/PM₁₀ ratio
173 equal to 0.87. Moreover, **Figure 1** highlights the marked seasonal differences for all three particulate
174 indicators, with the **mean comparison** between winter and spring means for TSP, PM₁₀ and PM_{2.5} being
175 significant ($p < 0.01$). The mean reduction in TSP in the spring with respect to winter was 30%, with mean
176 reductions of 68% for PM₁₀ and 82% for PM_{2.5}. The mean temperature differences between sampling
177 seasons was significantly different by ($p < 0.01$) with the mean winter temperature being 2.95 ± 4.09 °C and
178 the mean spring temperature being 16.55 ± 4.20 °C. However neither the average humidity nor the wind
179 speed were significantly different, with 0.73% humidity during the winter vs. 0.64% spring ($p > 0.05$) and ,an
180 average wind speed in both seasons of approximately 10 m/s (**Table 1**).

181

182 3.2 Mutagenicity

183 The mutagenicity tests show **a significantly elevated** of net revertants per unit of exposure (air equivalent
184 m³) **respect to the negative control. An elevated number of net revertants (250) was recorded at the**
185 **highest test doses for the winter TSP extracts in the YG1021 strain, while the mutagenicity of the PUF**
186 **extracts was markedly lower (Table 1)**. The PUF extracts contribute only about 2% to the total
187 mutagenicity. As **figure 2** also shows, the mutagenicity of the samples, expressed as net revertants, from
188 higher to the lower was PM induced YG1021 > PM induced TA98 > PM induced TA98+S9 >> non-particle
189 induced YG1021 > non-particle induced TA98 > non-particle induced TA98+S9. Moreover, the seasonal

190 trend is clearly evident and significant only for particulate-induced mutagenicity (YG1021 $p < 0.01$; TA98
191 $p < 0.01$; TA98+S9 $p < 0.01$). The mutagenicity of the spring TSP samples is less than 10% of the mutagenicity
192 recorded for the winter samples in all the strains.

193 The higher mutagenicity of the winter particles was confirmed also adjusting the data for particles mass
194 unit (Figure 3), highlighting the worse quality of the particles- in terms of mutagen presence - and not only
195 the higher level of aero-dispersed pollution for each volume unit.

196 Among the chemicals variability we observed a not so great changeability during the year for PAHs and
197 metals, observing a difference due to seasonality. More variability is instead observable for NO_x and ozone,
198 however also in this case the levels are clearly affected by seasonality (with highest value recoding in winter
199 with the ozone exception)(table 1). Table 2 showed the correlations between variables. Only the variables
200 for which at least one correlation with mutagenicity is significant was included, the not particles induced
201 mutagenicity was however included for its experimental origin, favoring the mutagenicity results
202 comparison.

203 As presented, the mutagenicity attributed to the non-particle phase was not influenced by the
204 environmental temperature or wind speed and, furthermore, does not correlate with the mutagenicity of
205 the particle phase. Additionally, the chemical parameters did not correlate with the minimal mutagenic
206 effects of the non-particle phase (table 2).

207 In contrast, the temperature and wind speed significantly inversely correlated with the TSP levels and to
208 mutagenicity of this mixture. The TSP level correlated with mutagenicity and, in particular, this correlation
209 showed a higher Spearman's rho for TA98 strain, with and without the addition of the S9 mixture. The
210 results of the mutagenicity assays conducted using the TSP extracts all correlate with each other (**Table 2**).

211 Among the chemical parameters, the TSP mutagenicity correlates with the presence of nitrogen oxides and,
212 in particular, this relationship is more marked for the nitrogen monoxides. The ozone levels inversely and
213 significantly correlate with the TSP mutagenicity. The cadmium and nickel levels significantly correlate with
214 direct mutagenicity (i.e., without the introduction of the metabolic activation). The TSP mutagenicity
215 correlates to the concentration of the finest fraction of the particulate matter and, in particular, there is a
216 better correlation with the PM₁₀ fraction in the TA98 strain with and without metabolic activation. A

217 significant correlation is not observed between benzo(a)pyrene or benzo(a)anthracene and mutagenicity but
218 there was a high correlation between PAHs and with metals (0.929 $p < 0.01$) due probably mainly to the
219 same seasonality.
220 Among the meteo climatic variables the temperature showed the high influence to the particulate pollution
221 and associated mutagenicity, moreover this physical parameter is not significantly correlated to the wind
222 that also showed an influence on the particulate pollution dispersion but not on the NO_x and ozone levels.
223 The humidity during the sampling showed a quite constant level so in this study we can't observe an
224 influence on the pollution level.
225 The NO_x, in particular NO, among the chemicals correlated with particulate pollution and associated
226 mutagenicity, moreover with PAHs, cadmium and nickel. This result was similar to those previously
227 observed in other studies (Du Four et al., 2004; Du Four et al., 2005).

228

229 4. Discussion

230 In our study, the inhalable fraction and the high-risk inhalable fraction represented a very high proportion
231 of the TSP, highlighting a human health hazard comparable to that estimated for urban sites. The observed
232 pollution levels are significantly higher than both the WHO guidelines (Krzyzanowski, 2008) and the EU
233 regulations 2008/50/CE. In addition, critical particle concentrations are present particularly during the
234 winter and especially for PM_{2.5}. Recently, the IARC classified outdoor pollution and particulate matter, as
235 its major component, as carcinogenic for humans (Loomis et al., 2013). Consequently, reducing air pollution
236 and particle matter to the lowest amount possible is becoming a marked priority.
237 Particulate matter clearly contributed to the overall mutagenicity (**Figure 2**). This observation confirmed
238 the evidence of other studies where PM total air toxicity and genotoxicity was higher than the gas phase
239 fraction. In particular, PM₁ was responsible for approximately 80% of the observed effects at various
240 sampling localities (Novak et al., 2014), and the fine particles generally showed higher mutagenicity
241 (Claxton et al., 2004; Claxton and Woodall, 2007b; Lemos et al., 2012). The gas phase mutagenicity was very
242 low and often indeterminable, with the exception of particular sampling sites such as industrial sites (Du
243 Four et al., 2005) and exhaust emissions from gasoline- and diesel-powered passengers cars (Pohjola et al.,

2003a; Pohjola et al., 2003b). The contribution of the gas phase with respect to the particulate phase seems to be higher during summer and related to the major PAHs content (Du Four et al., 2004; Kennedy et al., 2010). In the present study, the contribution of the gas phase with respect to the particulate phase is relative; in summer the particulate phase mutagenicity is reduced while the gas phase mutagenicity remains quite constant. It is supposable that this level of mutagenicity is not imputable to climatic or chemical stress condition and it indicates probably a background mutagenicity level hardly to avoid.

The benzo(a)pyrene concentration was higher during the winter than summer and higher than the WHO guide line value of 0.12 ng/m³ (Krzyzanowski, 2008; WHO-Europe, 2013). As widely observed, the PAHs are generally higher in the gas phase (Lemos et al., 2012), however, this fraction is less genotoxic and mutagenic, and thus PAH concentration explains only a small part of air pollution toxicity. Moreover, PAHs can react with nitrogen oxides, generating more genotoxic and mutagenic compounds (Albinet et al., 2008). The contribution of the nitro-derivate compounds to the overall mutagenicity, as assessed by comparing the number of Salmonella YG1021 net revertants to the strain without the modified nitro-reductase activity, was marked. The ratio of the net revertants observed in the TA98 and YG1021 strains is approximately 1:2, during both summer and winter. This observation is widely confirmed by other studies (Ramos de Rainho et al.; Traversi et al., 2009). Moreover the direct mutagens action is higher than indirect mutagens as highlighted by the ratio of the net revertants observed in the TA98 and TA98+S9 that is approximately of 1:1.7.

Air pollution and its major components have a marked seasonality, and the toxic content in the gas phase and particulate phase can vary based upon the meteo-climatic conditions (Albinet et al., 2008). In particular, more nitro-derived compounds can be present in the particulates during winter, thus enhancing the genotoxic and mutagenic properties.

266

267 5. Conclusions

268 By combining data on meteo-climatic conditions, various air pollution indicators and mutagenicity assays
269 we produced an evaluation of particulate and non-particulate air pollution in Turin during different season..
270 We present the following results:

- In the present study, the mutagenicity of the gas phase sampled by PUF method is practically negligible with respect to the mutagenicity of the particulate phase. The gas phase accounted for only 1% of the observed mutagenicity.
- The mutagenicity of the non-particulate phase remained constant during the summer and winter, while the particulate mutagenicity is approximately 5-fold higher during winter when the finest fraction of the PM increases.
- The contribution of the nitro-derived compounds seems to be crucial in Turin, in both winter and summer.
- Both chemicals (such as NO_x, metals and PAHs) and meteo-chemical parameters (such as temperature, wind speed and humidity) can influence the mutagenicity of particulate matter. Moreover, the total mutagenicity recorded in winter most likely results from the combination of not only additive but also synergistic effects among the components of the air pollution, conducting both to higher particulate level and to a higher content of mutagens in each unit particulate mass;
- Although PUF sampling is a common approach used in gas phase studies, there were relevant uncertainties regarding the applicability to biological *in vitro* models. A crucial point is the necessary extraction procedure between the sampling and the *in vitro* test. It is not presumably able to avoid a partial loss of the volatile and semi-volatile compounds. This more research is necessary to understand this problem.

Finally, the biological assays are relevant tools for the evaluation of the environmental and human health impact of air pollution.

6. Conflict of interest statement

The authors have nothing to declare. Funding source: this study was co-funded by the University of Turin (local funds ex-60% 2012) and the Piedmont Region (Italy) in the field of health projects.

7. Acknowledgements

297 The authors thank the Environmental Protection Agency of Piedmont, especially dott. F. Lollobrigida and
298 dott.ssa M. Maringo, for their collaboration in the sample collection and Dr. T. Nohmi of the National
299 Institute of Hygienic Sciences of Tokyo for the *S. typhimurium* YG1021 and TA98NR strains.

300

301 **8 List of abbreviations:**

302	PAHs	Polycyclic Aromatic Hydrocarbons
303	PCR	Polymerase Chain Reaction
304	TSP	Total Suspended Particles
305	PM	Particulate matter
306	PM10	Particulate matter with an aerodynamic diameter < 10 µm
307	PM2.5	Particulate matter with an aerodynamic diameter < 2.5 µm

308

309 **Table legends:**

310 **Table 1** -Descriptive analysis on 20 total measurements for each parameter are showed median and first
311 and third quartiles .

312

313 **Table 2** -Spearman's correlation between the mutagenicity, gravimetric, chemical and meteorological
314 variables ¹ rho = -0.436, p=0.054

315

316 **Figure legends:**

317 **Figure 1** - Mean and standard deviation of TSP, PM10 and PM2.5 levels recorded during the winter and
318 spring sampling sessions.

319 **Figure 2** - Total mutagenicity, subdivided into gas phase and particulate phase, recorded for the winter and
320 summer samples with metabolically different strains.

321 **Figure 3** - Net revertants expressed as unit mass of total suspended particulate for the different strain and
322 the different seasons.

323

324 **Bibliography**

- 325 Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., Jaffrezo, J.L., 2008. Nitrated and oxygenated
326 derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys - Part 1:
327 Concentrations, sources and gas/particle partitioning. *Atmospheric Environment* 42, 43-54.
- 328 ARPA Piemonte, P.d.T., 2010. Uno sguardo all'aria 2010. Relazione annuale sui dati rilevati dalla rete
329 provinciale di monitoraggio della qualità dell'aria - Anno 2010. Provincia di Torino, Torino.
- 330 ARPA Piemonte, S.P., CSI Piemonte, 2014. ARIAWEB - Sistema Regionale di Rilevamento della Qualità
331 dell'Aria della Regione Piemonte.
- 332 ATSDR, 1995. Toxicological Profile for Polycyclic Aromatic Hydrocarbons (PAHs). In: profile, A.t. (Ed.).
- 333 Breyse, P.N., Delfino, R.J., Dominici, F., Elder, A.C.P., Frampton, M.W., Froines, J.R., Geyh, A.S., Godleski,
334 J.J., Gold, D.R., Hopke, P.K., Koutrakis, P., Li, N., Oberdorster, G., Pinkerton, K.E., Samet, J.M., Utell, M.J.,
335 Wexler, A.S., 2013. US EPA particulate matter research centers: summary of research results for 2005-2011.
336 *Air Quality Atmosphere and Health* 6, 333-355.
- 337 Cassoni, F., Bocchi, C., Martino, A., Pinto, G., Fontana, F., Buschini, A., 2004. The Salmonella mutagenicity of
338 urban airborne particulate matter (PM_{2.5}) from eight sites of the Emilia-Romagna regional monitoring
339 network (Italy). *Sci Total Environ* 324, 79-90.
- 340 Claxton, L.D., Matthews, P.P., Warren, S.H., 2004. The genotoxicity of ambient outdoor air, a review:
341 Salmonella mutagenicity. *Mutat Res* 567, 347-399.
- 342 Claxton, L.D., Woodall, G.M., 2007a. A review of the mutagenicity and rodent carcinogenicity of ambient
343 air. *Mutation Research-Reviews in Mutation Research* 636, 36-94.
- 344 Claxton, L.D., Woodall, G.M., Jr., 2007b. A review of the mutagenicity and rodent carcinogenicity of ambient
345 air. *Mutat Res* 636, 36-94.
- 346 de Kok, T.M.C.M., Drieste, H.A.L., Hogervorst, J.G.F., Briede, J.J., 2006. Toxicological assessment of ambient
347 and traffic-related particulate matter: A review of recent studies. *Mutation Research-Reviews in Mutation*
348 *Research* 613, 103-122.
- 349 DeMarini, D.M., 2013. Genotoxicity biomarkers associated with exposure to traffic and near-road
350 atmospheres: a review. *Mutagenesis* 28, 485-505.
- 351 Dimitriou, K., Kassomenos, P., 2013. The fine and coarse particulate matter at four major Mediterranean
352 cities: local and regional sources. *Theoretical and Applied Climatology* 114, 375-391.
- 353 Du Four, V.A., Janssen, C.R., Brits, E., Van Larebeke, N., 2005. Genotoxic and mutagenic activity of
354 environmental air samples from different rural, urban and industrial sites in Flanders, Belgium. *Mutation*
355 *Research-Genetic Toxicology and Environmental Mutagenesis* 588, 106-117.
- 356 Du Four, V.A., Van Larebeke, N., Janssen, C.R., 2004. Genotoxic and mutagenic activity of environmental air
357 samples in Flanders, Belgium. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis* 558,
358 155-167.
- 359 EEA, 2012. European Environmental Agency - Air quality in Europe - 2012 report Office for Official
360 Publications of the European Union, Copenhagen.
- 361 Eeftens, M., Tsai, M.Y., Ampe, C., Anwander, B., Beelen, R., Bellander, T., Cesaroni, G., Cirach, M., Cyrys, J.,
362 de Hoogh, K., De Nazelle, A., de Vocht, F., Declercq, C., Dedele, A., Eriksen, K., Galassi, C., Grazuleviciene, R.,
363 Grivas, G., Heinrich, J., Hoffmann, B., Iakovides, M., Ineichen, A., Katsouyanni, K., Korek, M., Kramer, U.,
364 Kuhlbusch, T., Lanki, T., Madsen, C., Meliefste, K., Molter, A., Mosler, G., Nieuwenhuijsen, M., Oldenwening,
365 M., Pennanen, A., Probst-Hensch, N., Quass, U., Raaschou-Nielsen, O., Ranzi, A., Stephanou, E., Sugiri, D.,
366 Udvardy, O., Vaskoevi, E., Weinmayr, G., Brunekreef, B., Hoek, G., 2012. Spatial variation of PM_{2.5}, PM₁₀,

367 PM2.5 absorbance and PMcoarse concentrations between and within 20 European study areas and the
 368 relationship with NO₂ - Results of the ESCAPE project. *Atmospheric Environment* 62, 303-317.

369 EU, E.C., 2013. Attitudes of Europeans towards air quality. In: Environment, D.-G.f.t. (Ed.). *Flash*
 370 *Eurobarometer* 360

371 Forbes, P.B.C., Karg, E.W., Zimmermann, R., Rohwer, E.R., 2012. The use of multi-channel silicone rubber
 372 traps as denuders for polycyclic aromatic hydrocarbons. *Analytica Chimica Acta* 730, 71-79.

373 Fraser, M.P., Kleeman, M.J., Schauer, J.J., Cass, G.R., 2000. Modeling the atmospheric concentrations of
 374 individual gas-phase and particle-phase organic compounds. *Environmental Science & Technology* 34,
 375 1302-1312.

376 HEI, E.C.E., 2013. Understanding the Health Effects of Air Pollution: Recent Advances to Inform EU Policies
 377 Brussels, Belgium Auditorium, Madou Tower, Place Madou

378 Hoek, G., Boogaard, H., Knol, A., De Hartog, J., Slottje, P., Ayres, J.G., Borm, P., Brunekreef, B., Donaldson,
 379 K., Forastiere, F., Holgate, S., Kreyling, W.G., Nemery, B., Pekkanen, J., Stone, V., Wichmann, H.E., Van der
 380 Sluijs, J., 2010. Concentration Response Functions for Ultrafine Particles and All-Cause Mortality and
 381 Hospital Admissions: Results of a European Expert Panel Elicitation. *Environmental Science & Technology*
 382 44, 476-482.

383 ISTAT, 2012. 15° Censimento generale della popolazione e delle abitazioni

384 Ji, H., Hershey, G.K.K., 2012. Genetic and epigenetic influence on the response to environmental particulate
 385 matter. *Journal of Allergy and Clinical Immunology* 129, 33-41.

386 Josephy, P.D., Gruz, P., Nohmi, T., 1997. Recent advances in the construction of bacterial genotoxicity
 387 assays. *Mutat Res* 386, 1-23.

388 Kennedy, K., Macova, M., Bartkow, M.E., Hawker, D.W., Zhao, B., Denison, M.S., Mueller, J.F., 2010. Effect
 389 based monitoring of seasonal ambient air exposures in Australia sampled by PUF passive air samplers.
 390 *Atmospheric Pollution Research* 1, 50-58.

391 Kim, K.H., Jahan, S.A., Kabir, E., Brown, R.J., 2013. A review of airborne polycyclic aromatic hydrocarbons
 392 (PAHs) and their human health effects. *Environ Int* 60, 71-80.

393 Kovats, N., Acs, A., Ferincz, A., Kovacs, A., Horvath, E., Kakasi, B., Jancsek-Turoczi, B., Gelencser, A., 2013.
 394 Ecotoxicity and genotoxicity assessment of exhaust particulates from diesel-powered buses. *Environmental*
 395 *Monitoring and Assessment* 185, 8707-8713.

396 Krzyzanowski, M., 2008. WHO Air Quality Guidelines for Europe. *J Toxicol Environ Health A* 71, 47-50.

397 Lemos, A.T., Coronas, M.V., Rocha, J.A.V., Vargas, V.M.F., 2012. Mutagenicity of particulate matter fractions
 398 in areas under the impact of urban and industrial activities. *Chemosphere* 89, 1126-1134.

399 Lim, S.S., Vos, T., Flaxman, A.D., Danaei, G., Shibuya, K., Adair-Rohani, H., Amann, M., Anderson, H.R.,
 400 Andrews, K.G., Aryee, M., Atkinson, C., Bacchus, L.J., Bahalim, A.N., Balakrishnan, K., Balmes, J., Barker-
 401 Collo, S., Baxter, A., Bell, M.L., Blore, J.D., Blyth, F., Bonner, C., Borges, G., Bourne, R., Boussinesq, M.,
 402 Brauer, M., Brooks, P., Bruce, N.G., Brunekreef, B., Bryan-Hancock, C., Bucello, C., Buchbinder, R., Bull, F.,
 403 Burnett, R.T., Byers, T.E., Calabria, B., Carapetis, J., Carnahan, E., Chafe, Z., Charlson, F., Chen, H.L., Chen,
 404 J.S., Cheng, A.T.A., Child, J.C., Cohen, A., Colson, K.E., Cowie, B.C., Darby, S., Darling, S., Davis, A.,
 405 Degenhardt, L., Dentener, F., Des Jarlais, D.C., Devries, K., Dherani, M., Ding, E.L., Dorsey, E.R., Driscoll, T.,
 406 Edmond, K., Ali, S.E., Engell, R.E., Erwin, P.J., Fahimi, S., Falder, G., Farzadfar, F., Ferrari, A., Finucane, M.M.,
 407 Flaxman, S., Fowkes, F.G.R., Freedman, G., Freeman, M.K., Gakidou, E., Ghosh, S., Giovannucci, E., Gmel, G.,
 408 Graham, K., Grainger, R., Grant, B., Gunnell, D., Gutierrez, H.R., Hall, W., Hoek, H.W., Hogan, A., Hosgood,
 409 H.D., Hoy, D., Hu, H., Hubbell, B.J., Hutchings, S.J., Ibeanusi, S.E., Jacklyn, G.L., Jasrasaria, R., Jonas, J.B., Kan,
 410 H.D., Kanis, J.A., Kassebaum, N., Kawakami, N., Khang, Y.H., Khatibzadeh, S., Khoo, J.P., Kok, C., Laden, F.,
 411 Lalloo, R., Lan, Q., Lathlean, T., Leasher, J.L., Leigh, J., Li, Y., Lin, J.K., Lipshultz, S.E., London, S., Lozano, R.,
 412 Lu, Y., Mak, J., Malekzadeh, R., Mallinger, L., Marcenes, W., March, L., Marks, R., Martin, R., McGale, P.,

McGrath, J., Mehta, S., Mensah, G.A., Merriman, T.R., Micha, R., Michaud, C., Mishra, V., Hanafiah, K.M.,
 Mokdad, A.A., Morawska, L., Mozaffarian, D., Murphy, T., Naghavi, M., Neal, B., Nelson, P.K., Nolla, J.M.,
 Norman, R., Olives, C., Omer, S.B., Orchard, J., Osborne, R., Ostro, B., Page, A., Pandey, K.D., Parry, C.D.H.,
 Passmore, E., Patra, J., Pearce, N., Pelizzari, P.M., Petzold, M., Phillips, M.R., Pope, D., Pope, C.A., Powles, J.,
 Rao, M., Razavi, H., Rehfuess, E.A., Rehm, J.T., Ritz, B., Rivara, F.P., Roberts, T., Robinson, C., Rodriguez-
 Portales, J.A., Romieu, I., Room, R., Rosenfeld, L.C., Roy, A., Rushton, L., Salomon, J.A., Sampson, U.,
 Sanchez-Riera, L., Sanman, E., Sapkota, A., Seedat, S., Shi, P.L., Shield, K., Shivakoti, R., Singh, G.M., Sleet,
 D.A., Smith, E., Smith, K.R., Stapelberg, N.J.C., Steenland, K., Stockl, H., Stovner, L.J., Straif, K., Straney, L.,
 Thurston, G.D., Tran, J.H., Van Dingenen, R., van Donkelaar, A., Veerman, J.L., Vijayakumar, L., Weintraub,
 R., Weissman, M.M., White, R.A., Whiteford, H., Wiersma, S.T., Wilkinson, J.D., Williams, H.C., Williams, W.,
 Wilson, N., Woolf, A.D., Yip, P., Zielinski, J.M., Lopez, A.D., Murray, C.J.L., Ezzati, M., 2012. A comparative
 risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21
 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380, 2224-
 2260.

Liu, S.Z., Tao, S., Liu, W.X., Liu, Y.N., Dou, H., Zhao, J.Y., Wang, L.G., Wang, J.F., Tian, Z.F., Gao, Y., 2007.
 Atmospheric polycyclic aromatic hydrocarbons in north China: A winter-time study. *Environmental Science
 & Technology* 41, 8256-8261.

Loomis, D., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Baan,
 R., Mattock, H., Straif, K., Iarc, 2013. The carcinogenicity of outdoor air pollution. *Lancet Oncology* 14, 1262-
 1263.

Maron, D.M., Ames, B.N., 1983. Revised methods for the Salmonella mutagenicity test. *Mutat Res* 113, 173-
 215.

Novak, J., Hilscherova, K., Landlova, L., Cupr, P., Kohut, L., Giesy, J.P., Klanova, J., 2014. Composition and
 effects of inhalable size fractions of atmospheric aerosols in the polluted atmosphere. Part II. In vitro
 biological potencies. *Environment International* 63, 64-70.

Pohjola, S.K., Lappi, M., Honkanen, M., Rantanen, L., Savela, K., 2003a. DNA binding of polycyclic aromatic
 hydrocarbons in a human bronchial epithelial cell line treated with diesel and gasoline particulate extracts
 and benzo[a]pyrene. *Mutagenesis* 18, 429-438.

Pohjola, S.K., Lappi, M., Honkanen, M., Savela, K., 2003b. Comparison of mutagenicity and calf thymus DNA
 adducts formed by the particulate and semivolatile fractions of vehicle exhausts. *Environmental and
 Molecular Mutagenesis* 42, 26-36.

Poncino, S., Bande, S., Muraro, M., 2009. Meteo-diffusive analysis: a case study of Turin. *Epidemiologia &
 Prevenzione* 33, 27-33.

Ramos de Rainho, C., Machado Correa, S., Luiz Mazzei, J., Alessandra Fortes Aiub, C., Felzenszwalb, I.,
 Genotoxicity of polycyclic aromatic hydrocarbons and nitro-derived in respirable airborne particulate
 matter collected from urban areas of Rio de Janeiro (Brazil). *Biomed Res Int* 2013, 765352.

Traversi, D., Degan, R., De Marco, R., Gilli, G., Pignata, C., Villani, S., Bono, R., 2009. Mutagenic properties of
 PM2.5 urban pollution in the Northern Italy: The nitro-compounds contribution. *Environment International*
 35, 905-910.

Traversi, D., Schiliro, T., Degan, R., Pignata, C., Alessandria, L., Gilli, G., 2011. Involvement of nitro-
 compounds in the mutagenicity of urban Pm2.5 and Pm10 in Turin. *Mutation Research-Genetic Toxicology
 and Environmental Mutagenesis* 726, 54-59.

Valavanidis, A., Fiotakis, K., Vlachogianni, T., 2008. Airborne Particulate Matter and Human Health:
 Toxicological Assessment and Importance of Size and Composition of Particles for Oxidative Damage and
 Carcinogenic Mechanisms. *Journal of Environmental Science and Health Part C-Environmental
 Carcinogenesis & Ecotoxicology Reviews* 26, 339-362.

459 WHO-Europe, 2013. Review of evidence on health aspects of air pollution - Technical Report - REVIHAAP
460 Project

461 Xie, M., Barsanti, K.C., Hannigan, M.P., Dutton, S.J., Vedal, S., 2013. Positive matrix factorization of PM2.5 -
462 eliminating the effects of gas/particle partitioning of semivolatile organic compounds. Atmospheric
463 Chemistry and Physics 13, 7381-7393.

464 Yang, H.H., Chien, S.M., Cheng, M.T., Peng, C.Y., 2007. Comparative study of regulated and unregulated air
465 pollutant emissions before and after conversion of automobiles from gasoline power to liquefied
466 petroleum gas/gasoline dual-fuel retrofits. Environmental Science & Technology 41, 8471-8476.

467 Zhang, Y.W., Gu, Z.L., Cheng, Y., Shen, Z.X., Dong, J.G., Lee, S.C., 2012. Measurement of Diurnal Variations of
468 PM2.5 Mass Concentrations and Factors Affecting Pollutant Dispersion in Urban Street Canyons under
469 Weak-Wind Conditions in Xi'an. Aerosol and Air Quality Research 12, 1261-1268.

470

471