

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Lepiota sanguineofracta (Basidiomycota, Agaricales), a new species with a hymeniform pileus covering from Italy

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/151918> since 2016-08-10T12:04:22Z

Published version:

DOI:10.1007/s11557-013-0950-2

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

A. Vizzini;E. Ercole;S. Voyron. *Lepiota sanguineofracta* (Basidiomycota, Agaricales), a new species with a hymeniform pileus covering from Italy. MYCOLOGICAL PROGRESS. 13 pp: 683-690.
DOI: 10.1007/s11557-013-0950-2

The publisher's version is available at:

<http://link.springer.com/content/pdf/10.1007/s11557-013-0950-2>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/151918>

Lepiota sanguineofracta (*Basidiomycota*, *Agaricales*), a new species with a hymeniform pileus covering from Italy

- Alfredo Vizzini,
- Enrico Ercole,
- Samuele Voyron

DOI: 10.1007/s11557-013-0950-2

Cite this article as:

Vizzini, A et al. Mycol Progress (2014) 13: 683. doi:10.1007/s11557-013-0950-2

Abstract

Lepiota sanguineofracta sp. nov., a taxon with a hymeniform pileus covering, from Italy, is here described. A full description, colour pictures of basidiomata, line drawings of microscopic features, and ITS phylogenetic analysis are provided. It is morphologically and molecularly close (sister) to *Lepiota coloratipes* from which it differs mainly by olivaceous tinges on the pileus surface, basidiome surfaces and context strongly reddening on handling, a sweetish smell of withered rose and binucleate, not metachromatic spores.

Keywords

Agaricomycetes *Agaricaceae* *Lepiota* Hymeniform pileus covering ITS sequences Taxonomy

Introduction

Species of *Lepiota* (Pers.) Gray with a hymeniform pileus covering correspond to the morphology-based section *Cristatae* (Kühner ex Wasser) Bon as delimited by Kühner (1936) and Singer (1986) or section *Lilaceae sensu* Vellinga and Huijser (1999) and Vellinga (2001). Bon (1993) distributed these species over sections *Cristatae* (subg. *Lepiotula* (Mre.) Locquin ex Horak), *Integrellae* (Kühner ex Bon) Bon and *Lilaceae* Bon (subg. *Paralepiotula* M. Bon) as a whole, mainly based on different spore shapes (either ellipsoid or spurred) and spore nuclear number (mononucleate vs binucleate). According to recent molecular analyses, the species with a hymeniform pileus covering do not form a monophyletic lineage (Vellinga 2001, 2003, 2010a; Vizzini et al. 2013), even though most of them (with different spore shapes and nuclear number) fall in a clade (named clade 3 in Vellinga 2003) which includes also taxa (*L. cystophoroides* Joss., *L. luteophylla* Sundb. and *L. scaberula* Vellinga) with a hymeniform pileus covering giving rise to loose globose elements (a transition between hymeniderm and epithelium, see Vellinga 1998).

During a survey of macrofungi in the Botanical Garden of Turin (Piedmont, western Italy), collections of a remarkable *Lepiota* with a hymeniform pileus covering were recorded in flowerbeds.

A thorough literature search including monographic treatments and papers on world lepiotoid fungi (e.g., Morgan [1906](#); Murrill [1914](#); Kauffman [1924](#); Beeli [1932](#), [1936](#); Dennis [1952](#), [1970](#); Smith [1954](#), [1966](#); Aberdeen [1962](#); Pegler [1968](#), [1972](#), [1975](#), [1977](#), [1983](#), [1986](#), [1987a](#), [b](#), [1990](#); Pegler and Rayner [1969](#); Wasser [1980](#); Natarajan and Manjula [1982](#); Enderle and Krieglsteiner [1989](#); Sundberg [1989](#); Candusso and Lanzoni [1990](#); Guzmán and Guzmán-Dávalos [1992](#); Bizio et al. [1993](#); Bon [1993](#); Akers [1997](#); Vellinga and Huijser [1999](#); Vellinga [2001](#), [2010a](#); Montoya and Bandala [2005](#); Wang and Yang [2005](#); Zelený [2006](#); Kosakyan et al. [2008](#); Kumar and Manimohan [2009](#); Albuquerque et al. [2010](#); Gierczyk et al. [2011](#); Liang and Yang [2011](#); Ferreira and Cortez [2012](#); Razaq et al. [2012](#); Kaur et al. [2013](#); and Nawaz et al. [2013](#)), highlighted the unique nature of this taxon: its features do not fit with the description of any published species. The aim of the present contribution is to describe this *Lepiota* as a new species, providing full morphological and molecular data (ITS sequences analysis) and a comparison with allied taxa.

Materials and methods

Morphology

Macroscopic description was based from detailed field notes of fresh basidiomes. Colour terms in capital letters (e.g., Mars Orange, Plate II) are those of Ridgway ([1912](#)). Micromorphologic features were observed on dried material; sections were rehydrated in 5 % KOH, and then mounted in ammoniacal Congo Red, Cotton Blue, Cresyl Blue, Methyl Blue and Melzer's reagent, separately. Measurements were made at $\times 1000$ magnification with a calibrated ocular micrometer (Wild M11 optical light microscope). The width of each basidium was measured at the widest part, and the length was measured from the apex (sterigmata excluded) to the basal septum. The DNA fluorescent dye 4',6-diamidino-2-phenyl-indoldihydrochloride (DAPI) was used to stain nuclei in spores following Horton ([2006](#)). The number of nuclei in spores were then determined using a Leica TCS-SP2 confocal microscope. Samples were excited with 405 nm light and fluorescence was recorded at 440–500 nm.

The following abbreviations are used: L = number of lamellae reaching the stipe, l = number of lamellulae between each pair of lamellae; the notation [X, Y, Z] indicates that measurements were made on X spores, in Y samples from Z collections; Q = the spore quotient (length/width ratio); Q_{av} = the average spore quotient. Terminology for descriptive terms is according to Vellinga ([2001](#)). Herbarium abbreviations follow Thiers ([2013](#), continuously updated). The holotype collection is kept in TO. Author citations follow the Index Fungorum –Authors of Fungal Names (<http://www.indexfungorum.org/authorsoffungalnames.htm>).

DNA extraction, PCR amplification and DNA sequencing

Genomic DNA was isolated from 25 mg of two herbarium specimens (TO-HG2916 and TO-HG2917) using the DNeasy Plant Mini Kit (Qiagen, Milan Italy). Universal primers ITS1F/ITS4 were used for the ITS region amplification (White et al. [1990](#); Gardes and Bruns [1993](#)). Amplification reactions were performed in a PE9700 thermal cycler (Perkin-Elmer, Applied Biosystems) in a 25 μ l reaction mixture using the following final concentrations or total amounts: 10 ng DNA, 1 \times PCR buffer (20 mM Tris/HCl pH 8.4, 50 mM KCl), 0.4 μ M of each primer, 2.5 mM MgCl₂, 0.25 mM of each dNTP, 0.5 units of Taq polymerase (Promega). The PCR

studies focused on *Lepiota* with a hymeniform pileus covering (Vellinga [2003](#), [2010a](#); Vizzini et al. [2013](#)) sequences were retrieved from GenBank for the comparative phylogenetic analysis. Besides *Lepiota* species with a hymeniform pileus covering, eight species (indicated by an asterisk in Fig. [1](#)) representative of the major clades in *Lepiota* as delimited by Vellinga ([2003](#)) were chosen for comparison.

Alignments were generated using MAFFT (Kato et al. [2002](#)) with default conditions for gap openings and gap extension penalties. The sequence alignments were then imported into MEGA 5.10 (Tamura et al. [2011](#)) for manual adjustment. *Chamaemyces fracidus* (Fr.) Donk (AY176343) was used as an outgroup taxon because it is basal in the *Agaricaceae* (Vellinga [2004](#), [2010a](#)). Best-fit models were estimated by both the Akaike information criterion (AIC) and the Bayesian information criterion (BIC) with jModelTest 0.1.1 (Posada [2008](#)) to provide a substitution model for the alignment. Phylogenetic analyses were performed using the Bayesian Inference (BI) and Maximum likelihood (ML) approaches. The BI was performed with MrBayes 3.1.2 (Huelsenbeck and Ronquist [2001](#)) with four incrementally heated simultaneous Monte Carlo Markov Chains (MCMC) run over 10 million generations, under the GTR + Γ evolutionary model. Trees were sampled every 1,000 generations resulting in an overall sampling of 10,001 trees; the first 2,500 trees were discarded as “burn-in” (25 %). For the remaining trees, a majority rule consensus tree showing all compatible partitions was computed to obtain estimates for Bayesian Posterior Probabilities (BPP).

ML estimation was performed through RAxML v.7.3.2 (Stamatakis [2006](#)) with 1,000 bootstrap replicates (Felsenstein [1985](#)) using the GTRGAMMA algorithm to perform a tree inference and search for a good topology. Support values from bootstrapping runs (MLB) were mapped on the globally best tree using the “-f a” option of RAxML and “-x 12345” as a random seed to invoke the novel rapid bootstrapping algorithm.

BI and ML analyses were run on the CIPRES Science Gateway web server (Miller et al. [2010](#)). Only BPP values over 0.70 and MLB over 50 % are reported in the resulting tree (Fig. [1](#)). Branch lengths were estimated as mean values over the sampled trees.

Pairwise % identity values of ITS sequences were calculated using MEGA 5.10 (Tamura et al. [2011](#)).

Results

Molecular analysis

Both Bayesian and Maximum likelihood analyses produced the same topology; therefore, only the Bayesian tree with both BPP and MLB values is shown (Fig. [1](#)). The ITS data matrix comprised a total of 57 sequences (including 55 from GenBank). The alignment comprised 803 characters, and contains 395 variable sites.

In the obtained Bayesian phylogram (Fig. [1](#)), our two *Lepiota sanguineofracta* sequences clustered sister (BPP = 1; MLB = 94) to seven sequences of *L. coloratipes*. The pairwise % identity value between the *L. sanguineofracta* and *L. coloratipes* clades is 94.1.

Taxonomy

Lepiota sanguineofracta Vizzini, sp. nov. (Figs. [2](#), [3](#), and [4](#))



Fig. 2

Lepiota sanguineofracta. **a** Basidiomata; **b** pilei with green tinges; **c** reddening of context and basidiome surfaces; **d, e** details of veil remnants on stipe and pileus margins; **f** detail of the reddish stipe base. Photos by Alfredo Vizzini (all from the holotype). Scale bars **a–d, f** = 10 mm; **e** = 5 mm




Fig. 3

Lepiota sanguineofracta. Microscopic characters (in Cresyl blue, from the holotype). **a** Spores; **b** pileus covering elements. Scale bars **a** = 10 μm ; **b** = 40 μm (Photos by Alfredo Vizzini)




Fig. 4

Lepiota sanguineofracta. Microscopic characters (from the holotype). **a** Spores; **b** basidia; **c** cheilocystidia; **d** pileus covering elements. Scale bars **a** = 10 μm ; **b-c** = 20 μm ; **d** = 40 μm (Drawings by Enrico Bizio)

Mycobank MB 807073

Etymology the specific epithet from the Latin words *sanguineus* (meaning blood red coloured) and *fractus* (broken) refers to the basidiome surfaces and context which turn blood-red on bruising.

Macrocharacters Pileus 4–22 mm wide, at first slightly obtusely campanulate, hemispherical or broadly conical, later plano-convex to applanate-expanded, subumbonate, with a shallow umbo; not hygrophanous; margin not striated when wet and dry, slightly exceeding the lamellae when young, sinuous-undulate, lobate, entire or slightly fringed with age; surface dry, at first smooth, micaceous, later rugulose-wrinkled, sometimes irregularly cracking in concentric zones towards the margin; variable in colour, the pileus is at first completely white, soon becoming brownish-orange (Mars Orange, Orange Rufous, Plate II; Xathine Orange, Mars Yellow, Plate III; Ferruginous, Cinnamon-Rufous, Plate XIV; Vinaceous-Tawny, Plate XXVIII) spotted in the centre, when mature turns light brown (5D4, 5D5), pale ivory (Pale Pinkish Buff, Plate XXIX), remaining paler towards the margin; often with large green spots (Oriental Green, Rinnemann's Green, Plate XVIII) in the centre and near margin. Stipe 15–35 \times 1.0–1.8(–2.4) mm, central, cylindrical, usually regular, but sometimes also slightly flexuous, hollow; shiny, at first white, soon becoming reddish-brown (Acajou Red, Pompeian Red, Madder Brown, Plate XIII) overall starting from the base and progressing upward; minutely silky fibrillose along all length; stipe and pileus margin connected initially with whitish fibrils of veil, forming an indistinct ring zone on the upper part of the stipe and soon disappearing. Lamellae 2–3(4) mm wide, L = 33–42, l = 1–3(4), subfree to adnexed, at first white, soon with evident pinkish tints (Pale Salmon Color, Seashell Pink, Plate XIV); edge finely

granulose. Context elastic, in pileus dull white, in stipe concolorous with its surface, but shiny white in central cavity; without specific taste, smell strong, sweetish, reminiscent of dried rose petals as in *Lepiota helveola*/*L. josserandii*. Pileus and stipe surface, lamellae edge and context strongly reddening (Scarlet-Red, Plate I; Acajou Red, Dragon's-blood Red, Plate XIII) on bruising or with age. Spore print pale cream.

Microcharacters Spores [60, 2, 2] (3–)3.5–4.5(–5) × (2.5–)2.9–3.3(–3.5) μm, on average 4.23 × 2.94 μm, Q = (1.29–)1.3–1.55(–1.66), Q_{av} = 1.43, ellipsoid, rarely subcylindrical, hyaline, thin-walled, smooth, not verruculose in Melzer's reagent, not metachromatic in Cresyl Blue, non-amyloid, non-dextrinoid, cyanophilic in Cotton Blue and Methyl blue, congophilic, binucleate. Basidia 4-spored, 15–22 × 4–6 μm, clavate, hyaline, thin-walled; sterigmata up to 3–4 μm long. Cheilocystidia (10–)15–20 × 5–8(–9) μm, numerous and crowded, hyaline, thin-walled, various in shape, mostly clavate to subutriform, occasionally subfusiform, subcapitulate, lecythiform. Pleurocystidia absent. Pileus covering (pileipellis) a hymeniderm with transition to an epithelium, with up to 3(4) colourless elements on top of each other; terminal elements tightly packed, (12–)18–30(–50) × (10–)15–20(–25) μm, vesiculose, sphaeropedunculate to clavate-pyriform, sometimes with a short and obtuse apical mucro; slightly thick-walled (walls ca. 0.5 μm), with walls embedded in gelatinous matter; olivaceous tinges, when present, as intracellular pigment; subpellis composed of densely arranged and branching cylindrical hyphae, up to 5 μm wide. Hymenophoral trama subregular, consisting of hyphae 3–15 μm wide; trama of pileus and stipe similar to that in lamellae. Caulocystidia present only at apex of stipe, absent towards the base, very various in shape, 10–20 × 5–8 μm, hyaline, thin-walled, cylindrical, clavate, subglobose, sometimes sinuose, moniliform; occasionally in the form of simply lateral projections (diverticula) of the stipe hyphae. Clamp connections present and abundant in all tissues. Extracellular oil droplets absent in all tissues and on spore surface.

Habitat and distribution in small groups, terrestrial, on loamy or sandy humus-rich soils, growing at the edges of and in flowerbeds in Turin Botanical Garden. Fruiting in September. So far it is known only from the type locality.

Collections examined ITALY, Piedmont, Turin, Turin Botanical Garden, in flowerbeds of *Lonicera* spp. and *Viburnum* spp., 07 Sept 2011, leg. A. Vizzini (TO-HG2916, Holotype; GenBank acc. n. KF879620); ibidem, 13 Sept 2012, leg. A. Vizzini (TO-HG2917; GenBank acc. n. KF879621).

Discussion

Among all the described species of *Lepiota* with a hymeniform pileus covering, *L. sanguineofracta* is clearly circumscribed on the basis of a distinctive set of features such as a micaceous and not squamulose pileus surface with evident green tinges when mature, a fugacious partial veil not forming an annulus but leaving fibrillose remnants on the stipe surface, a stipe with reddish tinges towards the base, context smelling of dried rose petals, basidiome surfaces and context strongly reddening on handling, with the hymeniform pileus covering consisting of a tightly arranged palisade of clavate to sphaeropedunculate elements, binucleate spores, versiform cheilocystidia, and a unique ITS sequence. The reddening of the context is striking and quick, reminding one of *L. maculans* Peck (sect. *Lepiota*, Birkebak et al. [2011](#)) and several species of *Leucoagaricus* Locq. ex Singer sect. *Piloselli* Singer (Vellinga [2010b](#)).

According to both morphological and molecular data (Fig. [1](#)), *Lepiota coloratipes* is the most closely related species to *L. sanguineofracta*. This recently described species (Vizzini et al. [2013](#)), only so far known from Europe, China and North America, was established for the taxon usually referred to as *L. rufipes* in Europe, non-ss. orig. (= *Cystolepiota seminuda* according to Vellinga

[2010a](#)). It shares with *L. sanguineofracta* the reddish stipe base, the usually not diffracted pileus surface, the micaceous pileus surface due to the tightly packed claviform elements of pileipellis, the versiform cheilocystidia and the absence of a well-formed annulus, but differs in having an unchanging (not reddening) context and surfaces, a not distinctive smell, lamellae without cream-pink hues, slightly smaller (on average $3.4 \times 2.5 \mu\text{m}$) and mononucleate spores, that are metachromatic when mature and usually verruculose when observed in Melzer, and minute oil droplets in all tissues (Kühner and Maire [1937](#); Smith ([1954](#)); Josserand ([1955](#)); Lanzoni [1986](#); Candusso and Lanzoni [1990](#); Vellinga and Huijser [1999](#); Vellinga [2001](#), [2010a](#); Mohr [2008](#); Lange [2012](#); Vizzini et al. [2013](#)). Greenish tinges may be present on its pileus surface, but very faint and limited (Vizzini et al. [2013](#)).

The other species of *Lepiota* with hymeniform pileus covering and not squamulose-diffracted pileus surface, all without colour changes when bruised or with age, are also clearly distinguished from the new species: *L. neophana* Morgan (including var. *europaea* Bizio & Migl. and f. *papillata* Migl. & L. Perrone) shows a buff to dark brown and umbonate pileus centre, a well-formed annulus, very rare clamp-connections in the pileus trama and no cheilocystidia (Anonymous [1992](#); Bizio et al. [1993](#); Vellinga and Huijser [1999](#); Vellinga [2010a](#)). *L. psalion* Huijser & Vellinga is characterised by a distinct annulus, a looser pileipellis structure and clavate cheilocystidia (Vellinga and Huijser [1999](#); Vellinga [2001](#)) and, finally, *L. pyrochroa* Malenç. has an orange-red pileus and stipe, oil droplets in tissues, rough spores when studied in Melzer's reagent and clavate cheilocystidia (Malençon and Bertault [1970](#); Rioussset and Josserand [1976](#); Bon and Rioussset [1992](#); Bizio et al. [1993](#); Antonín and Vágner [1998](#); Vellinga and Huijser [1999](#)).

Acknowledgments

Enrico Bizio (Venice, Italy) is acknowledged for providing the microscopic line drawings. We are also grateful to Andrea Genre (University of Turin, Italy) for his technical assistance in confocal/fluorescence microscopy.

References

1. Aberdeen JEC (1962) Notes on Australian *Lepiota* in the Kew Herbarium. Kew Bull 16(1):129–137
2. Akers BP (1997) The family *Lepiotaceae* (*Agaricales*, *Basidiomycetes*) in Florida. M.Sc. Thesis. Southern Illinois University, Carbondale, IL
3. Albuquerque MP, Pereira AB, Carvalho AA Jr (2010) A família *Agaricaceae* Chevall. em trechos de Mata Atlântica da Reserva Biológica do Tinguá, Nova Iguaçu, Rio de Janeiro, Brasil: gêneros *Agaricus*, *Cystolepiota* e *Lepiota*. Acta bot bras 24(2):497–509
4. Anonymous (1992) Nova taxa in *Lepiota* s.l. Boll Ass micol ecol Romana 27:44–45
5. Antonín V, Vágner A (1998) [1997] New, rare and less known macromycetes in Moravia (Czech Republic) – IV. Acta Mus Moraviae Sci Biol 82:29–38
6. Beeli M (1932) Fungi Goossensiani: IX Genre *Lepiota*. B Soc Roy Bot Belg 64(2):206–222
7. Beeli M (1936) Genre *Lepiota* (Pers.) Fr. Flore iconographique des champignons du Congo, 2. Bruxelles
8. Birkebak JM, Vellinga EC, Franco-Molano AE, Wood MG, Matheny PB (2011) *Lepiota maculans*, an unusual mushroom rediscovered after 105 years. Southeast Nat 10(2):267–274
9. Bizio E, Migliozzi V, Zecchin G (1993) La sezione *Integrellae* (Kühner ex M. Bon) M. Bon del genere *Lepiota* (Persoon) Gray. Riv Micol 36:223–244
10. Bon M (1993) Flore mycologique d'Europe, 3. Les Lépiotes. *Lepiotaceae* Roze. Doc Mycol Mémoire hors série no. 3. L'Association d'Ecologie et Mycologie, Lille

11. Bon M, Rioussset G (1992) *Lepiotes* méridionales ou thermophiles, nouvelles ou intéressantes (1 – Tribu *Lepioteae*, genre *Lepiota*). *Doc Mycol* 22(85):63–73
12. Candusso M, Lanzoni G (1990) *Lepiota* s.l. *Fungi Europaei* 4. G. Biella, Saronno
13. Dennis RWG (1952) *Lepiota* and allied genera in Trinidad, British West Indies. *Kew Bull* 7(4):459–499
14. Dennis RWG (1970) Fungus flora of Venezuela and adjacent countries. *Kew Bull Addit Ser* 3:1–530
15. Drummond AJ, Ashton B, Buxton S, Cheung M, Cooper A, Duran C, Field M, Heled J, Kearse M, Markowitz S, Moir R, Stones-Havas S, Sturrock S, Thierer T, Wilson A (2010) Geneious v5.3. Available from <http://www.geneious.com/>
16. Enderle M, Krieglsteiner GJ (1989) Die Gattung *Lepiota* (Pers.) S.F. Gray emend. Pat. in der Bundesrepublik Deutschland (Mitteleuropa). *Z Mykol* 55(1):43–104
17. Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783–791
18. Ferreira AJ, Cortez VG (2012) Lepiotoid *Agaricaceae* (*Basidiomycota*) from São Camilo State Park, Paraná State, Brazil. *Mycosphere* 3(6):962–976
19. Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Mol Ecol* 2:113–118
20. Gierczyk B, Kujawa A, Szczepkowski A, Chachula P (2011) Rare species of *Lepiota* and related genera. *Acta Mycol* 46:137–178
21. Guzmán G, Guzmán-Dávalos L (1992) A checklist of the lepiotaceous fungi. Koeltz Scientific Books USA/Germany
22. Horton TR (2006) The number of nuclei in basidiospores of 63 species of ectomycorrhizal *Homobasidiomycetes*. *Mycologia* 98:233–238
23. Huelsenbeck JP, Ronquist F (2001) MrBayes: Bayesian inference of phylogeny. *Bioinformatics* 17:754–755
24. Jossierand M (1955) Notes critiques sur quelques champignons de la région Lyonnaise, 5. *Bull Soc Myc Fr* 71:65–125
25. Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucl Acids Res* 30:3059–3066
26. Kauffman CH (1924) The genus *Lepiota* in the United States. *Papers Mich Acad Sci, Arts Letters* 4:319–344
27. Kaur NJ, Saini MK, Kaur H (2013) Two new species of Agarics from India. *Mycosphere* 4(5):856–863
28. Kosakyan A, Ur Y, Wasser SP, Nevo E (2008) Rare and noteworthy lepiotaceous species (*Basidiomycota*, *Agaricales*, *Agaricaceae*) from Israel. *Mycotaxon* 103:59–74
29. Kühner R (1936) Recherches sur le genre *Lepiota*. *Bull Soc Myc Fr* 52:177–238
30. Kühner R, Maire R (1937) Trois *Lépiotes* peu connues. *Bull Soc Hist nat Afr N* 28:108–112
31. Kumar TKA, Manimohan P (2009) The genus *Lepiota* (*Agaricales*, *Basidiomycota*) in Kerala State, India. *Mycotaxon* 107:105–138
32. Lange C (2012) *Lepiota* (Pers. : Fr.) Gray. In: Knudsen H, Vesterholt J (eds), *Funga Nordica. Agaricoid, boletoid, clavarioid, cyphelloid and gastroid genera*. Nordsvamp, Copenhagen, pp 627–637
33. Lanzoni G (1986) Raccolte interessanti del 1984. *Boll Gruppo Micol Bres* 29:81–90
34. Liang JF, Yang ZL (2011) Two new taxa close to *Lepiota cristata* from China. *Mycotaxon* 116:387–394
35. Malençon G, Bertault R (1970) Flore des champignons supérieurs du Maroc, Tome I. Faculté des Sciences de Rabat, Rabat
36. Miller MA, Pfeiffer W, Schwartz T (2010) “Creating the CIPRES Science Gateway for inference of large phylogenetic trees” in *Proceedings of the Gateway Computing Environments Workshop (GCE)*, 14 Nov. 2010, New Orleans, LA, pp 1–8

37. Mohr P (2008) Interessante Schirmlinge aus Berlin und der Mark Brandenburg. *Boletus* 30(2):47–79
38. Montoya L, Bandala VM (2005) A new species and a new record of *Lepiota* occurring in the Gulf of Mexico area. *Mycotaxon* 94:111–125
39. Morgan AP (1906) North American species of *Lepiota*. *J Mycol* 12:154–159; 195–203; 242–248
40. Murrill WA (1914) *Agaricaceae*. *North American Flora* 10(1):41–65
41. Natarajan K, Manjula B (1982) South Indian *Agaricales*: XI. *Mycologia* 74(1):130–137
42. Nawaz R, Khalid AN, Hanif M, Razaq A (2013) *Lepiota vellingana* sp. nov. (*Basidiomycota*, *Agaricales*) a new species from Lahore, Pakistan. *Mycol Prog*. doi:10.1007/s11557-012-0884-0
43. Pegler DN (1968) Studies on African *Agaricales* I. *Kew Bull* 21:499–538
44. Pegler DN (1972) A revision of the genus *Lepiota* from Ceylon. *Kew Bull* 27(1):155–202
45. Pegler DN (1975) A revision of the Zanzibar *Agaricales* described by Berkeley. *Kew Bull* 30:429–442
46. Pegler DN (1977) A preliminary agaric flora of East Africa. *Kew Bull Addit Ser* 6:1–615
47. Pegler DN (1983) Agaric flora of the Lesser Antilles. *Kew Bull Addit Ser* 9:1–668
48. Pegler DN (1986) Agaric flora of Sri Lanka. *Kew Bull Addit Ser* 12:1–519
49. Pegler DN (1987a) A revision of the *Agaricales* of Cuba 1. Species described by Berkeley & Curtis. *Kew Bull* 42:501–585
50. Pegler DN (1987b) A revision of the *Agaricales* of Cuba 2. Species described by Earle and Murrill. *Kew Bull* 42:855–88
51. Pegler DN (1990) *Agaricales* of Brazil described by J.P.F.C. Montagne. *Kew Bull* 45:161–177
52. Pegler DN, Rayner RW (1969) A contribution to the *Agaric* flora of Kenya. *Kew Bull* 23(3):347–412
53. Posada D (2008) jModeltest: phylogenetic model averaging. *Mol Biol Evol* 25:1253–1256
54. Razaq A, Khalid AN, Vellinga EC (2012) *Lepiota himalayensis* (*Basidiomycota*, *Agaricales*), a new species from Pakistan. *Mycotaxon* 121:319–325
55. Ridgway R (1912) Color standards and color nomenclature. Self-published, Washington D.C
56. Rioussset G, Josserand M (1976) Récolte en France d'un basidiomycète marocain. "*Lepiota pyrochroa* Mal. *Bull mens Soc Linn Lyon* 45:198–200
57. Singer R (1986) The *Agaricales* in modern taxonomy, 4th edn. Koeltz Scientific Books, Koenigstein
58. Smith HV (1954) A revision of the Michigan species of *Lepiota*. *Lloydia* 17:307–328
59. Smith HV (1966) Contributions toward a monograph on the genus *Lepiota*, I. Type studies in the genus *Lepiota*. *Mycopath & Mycol appl* 29:97–117
60. Stamatakis A (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688–2690
61. Sundberg WJ (1989) *Lepiota* sensu lato in California III. Species with a hymeniform pileipellis. *Mycotaxon* 34:239–248
62. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731–2739
63. Thiers B (2013) [continuously updated]. Index herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>
64. Vellinga EC (1998) Glossary. In: Bas C, Kuyper TW, Noordeloos ME, Vellinga EC (eds) *Flora Agaricina Neerlandica*. Critical monographs on families of agarics and boleti occurring in the Netherlands 1. Balkema, Rotterdam, pp 54–64

65. Vellinga EC (2001) *Lepiota* (Pers. : Fr.) S.F. Gray. In: Noordeloos ME, Kuyper TW, Vellinga EC (eds) *Flora Agaricina Neerlandica. Critical monographs on families of agarics and boleti occurring in the Netherlands 5*. A.A. Balkema Publishers, Lisse, pp 109–151
66. Vellinga EC (2003) Phylogeny of *Lepiota* (*Agaricaceae*) – evidence from nrITS and nrLSU sequences. *Mycol Prog* 2:305–322
67. Vellinga EC (2004) Genera in the family *Agaricaceae* – evidence from nrITS and nrLSU sequences. *Mycol Res* 108:354–377
68. Vellinga EC (2010a) *Lepiota* in California: species with a hymeniform pileus covering. *Mycologia* 102:664–674
69. Vellinga EC (2010b) Lepiotaceous fungi in California, U.S.A. *Leucoagaricus* sect. *Piloselli*. *Mycotaxon* 112:393–444
70. Vellinga EC, Huijser HA (1999) [1998] Studies in *Lepiota* I. Species with a hymeniform pileus covering. *Belg J Bot* 131:191–210
71. Vizzini A, Liang JF, Jančovičová S, Adamčík S, Ercole E, Contu M, Yang ZL, Vellinga EC (2013) *Lepiota coloratipes*, a new species for *Lepiota rufipes* ss. *Auct europ non ss orig Mycol Progress*. doi:10.1007/s11557-013-0905-7
72. Wang HC, Yang ZL (2005) Notes on *Lepiota shixingensis* and an allied new species (Basidiomycota). *Nova Hedwigia* 81:463–469
73. Wasser SP (1980) Flora gribov Ukrainy, Agarikovye griby. [Fungal Flora of the Ukraine: Agaricoid Fungi]. Naukova Dumka, Kiev
74. White TJ, Bruns TD, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand D, Sninsky J, White T (eds) *PCR Protocols, a guide to methods and applications*. Academic, Orlando, pp 315–322
75. Zelený L (2006) Taxonomic literature on the genus *Lepiota* s. l. in the Czech Republic. *Czech Mycol* 58(3–4):225–265

About this article

- **Print ISSN**

1617-416X

- **Online ISSN**

1861-8952

Not logged in Universita' degli Studi di Torino (3000088961) - INDEST AICTE Consortium Indian Institute of Technology (3000185589) - National Institute of Technology Rourkela (3000194080) - Consortia Annali dell'Università di Ferrara (3002076494) 130.192.101.166