


 CrossMark
 click for updates

 Cite this: *RSC Adv.*, 2015, 5, 74734

Polyhydroxylated GdDTPA-derivatives as high relaxivity magnetic resonance imaging contrast agents†

 Lorenzo Tei,^{*a} Alessandro Barge,^b Matteo Galli,^c Roberta Pinalli,^c Luciano Lattuada,^{*c} Eliana Gianolio^d and Silvio Aime^d

The search for new MRI agents endowed with high relaxivity at the magnetic field strength of clinical scanners (1.5–3 T) is still receiving great attention from researchers involved in the development of new probes. Such Gd(III) complexes should combine a fast inner-sphere water exchange rate (k_{ex}) with an enhanced contribution from water molecules in the second and outer coordination spheres. In the present work, dendrimeric-like structures were synthesized by coupling diethylenetriaminepentaacetic acid (DTPA) and its mono-methylphosphonic derivative (P-DTPA) with two differently branched, highly hydrophilic, gluconyl moieties. A ^1H and ^{17}O NMR relaxometric study on the corresponding Gd(III) complexes reveals that the Gd-P-DTPA-polyol complex displays very high relaxivities (around $20 \text{ mM}^{-1} \text{ s}^{-1}$ at 298 K) over the 0.5–3 T range of field strengths as a result of a fast k_{ex} and the presence of a strong second sphere contribution.

 Received 29th July 2015
 Accepted 28th August 2015

DOI: 10.1039/c5ra15071j

www.rsc.org/advances

Introduction

The use of Gd(III)-chelates as contrast enhancing agents (CA) for Magnetic Resonance Imaging (MRI) has developed considerable research activity aimed at the design of effective, specific and safe CAs.¹ The efficacy of a contrast agent is expressed by its relaxivity, r_1 , which is the enhancement of the longitudinal proton relaxation rate induced by the paramagnetic agent at 1 mM concentration. Theory predicts high relaxivity for a Gd(III) complex when the rate of water exchange between the inner sphere and the bulk solvent ($k_{\text{ex}} = 1/\tau_{\text{M}}$) is around 10^6 – 10^7 s^{-1} and when the molecular reorientational time and electron spin relaxation time are similar and long at the selected field strength.^{1,2} Nowadays, clinical MR imaging is increasingly moving to higher fields since the number of installed 3 Tesla scanners is steadily growing worldwide.³ Higher fields improve the signal-to-noise ratio (SNR) and provide higher spatial resolution and/or reduced acquisition times.⁴ Considering this trend, there is the need of contrast agents endowed with a high

relaxivity over an extended range of magnetic field strength.⁵ High-relaxivity CAs permit the use of lower doses in routine clinical applications and are the candidates of choice in molecular imaging procedures aimed at detecting low-concentration targets. Optimizing the three contributions that determine the observed relaxation rates of water protons, namely inner-, second- and outer-coordination sphere, could lead to highly efficient CAs.

One route followed in the search for higher relaxivities relied on the lengthening of the molecular reorientational time through the formation of polymers or of covalent and non-covalent conjugates between the paramagnetic chelate and slowly moving substrates⁶ (dendrimers,⁷ proteins,⁸ carbohydrates⁹) and the formation of supramolecular adducts like micelles or liposomes.¹⁰ In most cases, these systems display maximum r_1 values at 20–40 MHz (*i.e.* 0.5–1 T) but their relaxation enhancement ability decreases quickly at higher fields. Therefore, medium size molecular weight Gd-based compounds have been proposed in order to obtain good relaxivity values over the 0.5–3 T magnetic field range. Examples of these systems are based either on oligomeric (2–8 units) Gd(III) complexes^{4,11} or on monomeric Gd-complexes with the Gd³⁺ ion at the baricentre of the macromolecule with a well-structured second sphere of hydration.¹² Similar symmetric structures based on DOTA (DOTA = 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid) but with higher molecular weights (5–7 kDa) were reported to have a relaxivity drop at magnetic fields higher than 1 T (40 MHz).¹³ Moreover, their dimensions allow only a slow diffusion in the extravascular space.

^aDipartimento di Scienze e Innovazione Tecnologica, Università del Piemonte Orientale “Amedeo Avogadro”, Viale T. Michel 11, 15121 Alessandria, Italy. E-mail: lorenzo.tei@uniupo.it

^bDipartimento di Scienza e Tecnologia del Farmaco, Università di Torino, via P. Giuria 9, 10125 Torino, Italy

^cBracco Imaging SpA, Bracco Research Centre, via Ribes 5, 10010 Colletterto Giacosa, TO, Italy. E-mail: luciano.lattuada@bracco.com

^dDepartment of Molecular Biotechnology and Health Sciences, Molecular Imaging Center, Università di Torino, Via Nizza 52, 10126, Torino, Italy

† Electronic supplementary information (ESI) available: ^1H and ^{13}C NMR spectra of novel compounds. See DOI: 10.1039/c5ra15071j


Due to their pharmacokinetic profile, the diagnostic applications of this type of contrast agents are different not only from the already marketed non-specific agents that freely and quickly distribute into the extracellular space, but also from blood pool agents that mainly distribute into the blood system. In light of these considerations, there still is the need for improved non-specific contrast agents able to combine high relaxivity with rapid extravasation, as required for angiographic- and perfusion-based diagnostic applications.

We have faced this problem by synthesizing new Gd(III) complexes based on the structure of DTPA functionalized with highly hydrophilic substituents in order to remarkably increase the second sphere contribution to the relaxivity by creating a structured network of water molecules localised around the polyhydroxyl groups. Ligands **L1** and **L2** are DTPA-bisamides bearing two differently branched gluconyl moieties and **L3** is based on a DTPA derivative in which the central arm has been replaced by a methyl phosphonate moiety (P-DTPA) and two outer acetate moieties bear each the triply gluconyl substituted polyol. The choice of a P-DTPA derivative relies on the published observation that its Gd(III) complex shows an optimal value of τ_M (about 88 ns).¹⁴ Moreover, phosphonate moieties have been shown to yield good second sphere contributions to the observed relaxivity due to hydrogen bonded water molecules localised between the phosphonate group and the Gd.¹⁵ In the present work, we report on the synthesis of **L1–L3** ligands and on a ¹H and ¹⁷O NMR relaxometric study on the corresponding Gd(III) complexes.

Results and discussion

Ligands design


The introduction of polyhydroxylic or glycosylic groups at the periphery of a Gd(III) complex has been considered previously with different purposes. In some reports, CAs structures that




Scheme 1 Synthesis of the DTPA-bisamide **L1**.

incorporate glucose and galactose/mannose moieties were developed with the intent to enhance targeting *in vivo*.¹⁶ Others groups engineered complexes in which the Gd(III) ion lied at the barycentre of the macromolecular structure surrounded by a well-defined second hydration sphere to obtain higher relaxivities.^{12,13} Furthermore, a water-solubilizing polyhydroxylic dendron was incorporated to improve the poor water solubility of the parent complex.¹⁷ Our work started with the synthesis of two diamides of DTPA (**L1** and **L2**) bearing differently branched polygluconyl groups on the outer surface. The assessment of the relaxometric properties of their Gd-complexes allowed us to determine the best hydrophilic pendant group for the attainment of a large second sphere contribution to relaxivity. Two examples of DTPA-bisamides bearing two acetylglucose units and a dendrimeric glycosidic cluster with six acetylgluconamides at the periphery were already reported with the aim to create new blood pool CAs more than to evaluate the effect of the second sphere water molecules surrounding the Gd(III) centre.¹⁸ As we were aware that DTPA-bisamides are characterized by a very slow water exchange rate (k_{ex}) that “quenches” any putative inner sphere relaxivity gain, our next task dealt with the introduction of two polyhydroxy-containing moieties on P-DTPA ligand.

Bifunctional derivatives of DTPA have been reported either with the remote functional group on the central acetate arm or on the diethylenetriamine backbone.¹⁹ Only one example of



Scheme 2 Synthesis of the DTPA-bisamide **L2**.



Scheme 3 Synthesis of the fully acetylated aminopolyol **9**; (i) Boc_2O , TMAOH, DMF, 48h; (ii) Ac_2O , Py, 90 °C, 24 h.

functionalization of a lateral acetate pendant arm was described²⁰ and to the best of our knowledge a bis-functionalized DTPA-like agent has never been reported. Thus, the bifunctional chelating ligand **15** (Scheme 4) was designed having four carboxylic and the phosphonic groups *tert*-butyl protected and two free carboxylic groups able to react with amino functionalized moieties. Then, the ligand **L3** could be prepared by reacting the bifunctional agent **15** with two dendronized polygluconyl groups with the aim to increase the amount of hydroxyl groups and to create a symmetric Gd(III) complex with lateral hydrophilic moieties and a central phosphonate group.


Synthesis

The polyol functionalised DTPA bisamide **L1** (Scheme 1) was synthesized in one step with an overall 61% yield starting from DTPA bis-anhydride **4**²¹ and *N,N'*-(iminodi-2,1-ethanediyl)bis- δ -gluconamide **3**, previously obtained by amidation of diethylenetriamine **1** with δ -gluconolactone **2** (Scheme 1). Similarly, the DTPA bisamide **L2** was synthesised by reaction of DTPA bis-anhydride **4** with the tris-gluconoyl amide derivative **6** obtained by reaction between pentaerythritol tetramine **5**²² and δ -gluconolactone (Scheme 2). Aiming to use this amino-functionalised polyol for further syntheses, we decided to prepare the fully acetylated derivative **9** (Scheme 3). Thus, the amino group of **6** was protected with $(\text{Boc})_2\text{O}$ and tetramethylammonium hydroxide in DMF to give **7**. Then, all the hydroxyl groups were acetylated with Ac_2O in pyridine to give **8** and, finally, **9** was obtained by removing the Boc protection with TFA in CH_2Cl_2 .


The synthetic procedure to obtain **L3**, depicted in Scheme 4, relies on the strategy published by Williams and Rapoport who reported the *N*-bisalkylation of *p*-nitrophenylalanine benzyl ester with different bromoethylamines as a direct method for the construction of protected DTPA analogues.²³ Moreover, following this procedure, modified amino acids have been bisalkylated with [*N*-(bromoethyl)amino]diacetic acid *t*-butyl ester in order to obtain functionalized DTPA pentaesters as versatile intermediates for the preparation of conjugates of metal complexes.^{19,24} We exploited this idea in a different way by synthesising a bromoethyl derivative of protected aspartic acid and reacting it with an aminomethyl phosphonate. By this

route, after orthogonal deprotection of two carboxy benzylester groups, we were able to attach an amino-polyol molecule on both sides of the P-DTPA derivative. In particular, *L*-aspartic acid 4-benzyl ester **10** was esterified, using *t*-butyl acetate in the presence of catalytic perchloric acid, into *L*-aspartic acid 1-*tert*-butyl 4-benzyl diester **11**, which was directly reacted with one equivalent of *t*-butyl bromoacetate under Rapoport's two-phase conditions (*i.e.*: MeCN/pH 8 phosphate buffer) to yield, after chromatographic purification, the mono-alkylated compound **12** in 76.5% yield. Triester **12** was then further alkylated with 2-bromoethyl-trifluoromethanesulfonate in order to obtain in good yield the bromoethyl component **13** to be used for the construction of the P-DTPA derivative (Scheme 4).

The other component was the aminomethylphosphonic acid di-*tert*-butyl ester **19** (Scheme 5) obtained in two steps by Mannich reaction of dibenzylamine with paraformaldehyde and *tert*-butyl phosphite, followed by hydrogenolysis of the benzyl group at atmospheric pressure with 10% Pd/C catalyst. The Rapoport's conditions were also used for the alkylation of **19** with two equivalents of **13** (no excess was used) to get the P-DTPA derivative **14** in 42.4% yield after column purification. Cleavage of the benzyl ester protections of **14** by hydrogenolysis yielded the diacid pentaester **15**. This compound was then coupled to the amino-protected polyol **9** in DMF at room temperature for 48 h by using HATU (*O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluroniumhexafluoro phosphate) as coupling reagent and



Scheme 4 Synthesis of **L3**.



Scheme 5 Synthesis of aminomethylphosphonic acid di-*tert*-butyl ester **19**.

N,N-diisopropyl ethylamine as base. Product **16**, together with a small amount of the mono-substituted derivative, was collected by precipitation in cold water. Due to difficulties met during the purification procedures, it was decided to use this mixture without separating the mono-functionalized impurity. Deacetylation of the polyalcohol was afforded by bubbling ammonia in a refrigerated methanol solution of **16**. This reaction was completed only after 9 days and after bubbling more ammonia every two days. In order to obtain the final ligand **L3**, the last step was the cleavage of the *t*-butyl esters by reaction with a 50% solution of trifluoroacetic acid in CH_2Cl_2 and precipitation by addition of diethyl ether. In order to get pure **L3** a simple procedure was followed: the trifluoroacetate salt obtained after *t*-butyl ester deprotection was redissolved in water, brought to pH 7, lyophilized and recovered with methanol. Ligand **L3** was thus obtained as a white solid in 30% yield by filtration of the MeOH solution. On the other hand, evaporation of the solvent give rise to an oil which resulted to be the mono-polyol derivative present in about 28% of the total raw material.

Relaxivity of the Gd(III) complexes

The Gd(III) complexes Gd**L1**, Gd**L2** and Gd**L3** were prepared by ^1H NMR relaxometric titration with a stock solution of GdCl_3 at pH 6.5 monitoring the change in the longitudinal water proton relaxation rate (R_1) at 20 MHz and 298 K as a function of the concentration of Gd^{3+} . The slope of the straight line obtained corresponds to the relaxivity (r_1) of the complex. The residual




Fig. 1 Relaxivities (r_1) values, measured at 298 K and 20 MHz, of Gd**L1**, Gd**L2** and Gd**L3** as a function of their molecular weight compared to those reported for other $q = 1$ complexes. Values for Gd-DOTA, Gd-DTPA and Gd-HPDO3A come from ref. 2, for Gd-4 and Gd-5 from ref. 12a, for Gd-6b, 6c, 6d and 6e from ref. 12b and for P760 and P792 from ref. 13.

free Gd^{3+} ion was quantified by Orange Xylenol UV method as lower than 0.2% in all complexes.²⁵ The exact concentration of Gd^{3+} ions was determined by measurement of the bulk magnetic-susceptibility shifts of the *t*BuOH signal. The relaxivity (r_1) values at 20 MHz and 298 K are $5.0 \text{ mM}^{-1} \text{ s}^{-1}$ for Gd**L1**, $4.3 \text{ mM}^{-1} \text{ s}^{-1}$ for Gd**L2** and $19.2 \text{ mM}^{-1} \text{ s}^{-1}$ for Gd**L3**. In Fig. 1, the relaxivities of the three Gd-chelates are plotted as a function of their molecular weight and compared to those reported in the literature for monomeric Gd-complexes with one coordinated water molecule at $T = 298 \text{ K}$ and $B_0 = 0.5 \text{ T}$. While the r_1 values for the two GdDTPA-bisamides Gd**L1** and Gd**L2** are significantly lower than the values expected on the basis of their molecular weight, the r_1 of the Gd(III) complex with the monophosphonic DTPA derivative **L3**, is markedly higher than expected for a molecule of such molecular weight. Noteworthy, the relaxivity of Gd**L3** is more than four times higher than that of the parent GdDTPA ($r_1 = 4.7 \text{ mM}^{-1} \text{ s}^{-1}$). A more detailed relaxometric characterization was carried out through the registration of the ^{17}O NMR- R_{2p} profiles as a function of temperature (for an accurate determination of the exchange lifetime (τ_M) of the coordinated water molecule – Fig. 2) and of the ^1H -NMRD profiles over the range of frequencies from 0.01 to 120 MHz (Fig. 3A and B).

The analysis of the ^{17}O NMR experimental data yielded the τ_M values reported in Table 1. The value found for Gd**L3** (93 ns) is very similar to that already reported for related GdP-DTPA derivatives¹⁴ and very close to the optimal value for the achievement of high relaxivity.^{1,2} On the contrary, very long τ_M values ($5.6 \mu\text{s}$ and $7.6 \mu\text{s}$, for Gd**L1** and Gd**L2** respectively) were found for the two bisamides derivatives. This slow water exchange rate, which is invariably found in GdDTPA-bisamides,²⁶ is responsible for the low relaxivity shown by Gd**L1** and Gd**L2**. The proton relaxivity data measured as a function of the magnetic field strength at 298 K and neutral pH (Fig. 3A and B) were well interpolated with the set of values calculated on the basis of the classical Solomon-Bloembergen-Morgan theory by fixing the τ_M values to those obtained by ^{17}O NMR- R_{2p} analysis. The main parameters obtained from the fitting of experimental to calculated data are reported in Table 1. As expected, a




Fig. 2 Temperature dependence of the paramagnetic contribution to the ^{17}O transverse relaxation rate of water for Gd**L1**, Gd**L2** and Gd**L3** (10 mM) measured at 2.1 T and pH 7.




Fig. 3 $1/T_1$ NMRD profiles of GdL1, GdL2 (A) and GdL3 (B) recorded at 298 K and pH 7. The data refer to a 1 mM concentration of the paramagnetic complexes. The solid curves through the data points were calculated with the parameters reported in Table 1, whereas the dotted lines in (B) refer to the different contributions to the overall relaxivity.

Table 1 Best-fit parameters obtained from ^1H -NMRD at 298 K and ^{17}O - R_{2p} versus T analysis shown in Fig. 2 and 3^a

	Δ^2 ($\times 10^{19} \text{ s}^{-2}$)	τ_V (ps)	τ_R (ps)	τ_M (μs)	q	q^{ss}	r^{ss} (\AA)	τ^{ss} (ps)
Gd-L1	2.9 ± 0.57	22.1 ± 1.8	305 ± 12.8	5.6 ± 1.2	1	0	—	—
Gd-L2	2.1 ± 0.57	25.0 ± 1.5	470 ± 68.1	7.6 ± 0.6	1	0	—	—
Gd-L3	2.2 ± 0.13	34.2 ± 0.10	560 ± 10.8	0.093 ± 0.003	1	3.5 ± 0.4	3.7	93 ± 0.15

^a On carrying out the fitting procedure, some parameters were fixed to reasonable values: $r_{\text{Gd-H}}$ (distance between Gd and protons of the innersphere water molecule) = 3.1 \AA ; a (distance of minimum approach of solvent water molecules to Gd ion) = 4 \AA ; D (solvent diffusion coefficient) = $2.24 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$; r^{ss} (distance between Gd and protons of the second sphere water molecules) = 3.7 \AA . [Δ^2] Squared mean transient zero-field splitting (ZFS) energy. [τ_V] Correlation time for the collision-related modulation of the ZFS Hamiltonian. [τ_R] Reorientational correlation time. [τ_M] Exchange lifetime of the coordinated water molecule. [q] Number of inner-sphere water molecules. [q^{ss}] Number of second-sphere water molecules. [τ^{ss}] correlation time associated with the motions (reorientation or exchange) of the second-sphere water molecules.

progressive increase in the reorientational correlation time (τ_R) occurs upon increasing the molecular weight.

In the case of GdL3 (Fig. 3B), a good fitting of the experimental data was possible only by taking into account a contribution from second sphere water molecules.²⁷ 3.5 second sphere water molecules (q^{ss}) were estimated to be located at an average distance of 3.7 \AA from the Gd^{3+} ion with an overall correlation time (τ^{ss}) of 93 ps (Table 1). The increase in the correlation time associated with the collision-related modulation of the zero-field splitting Hamiltonian (τ_V) observed in the case of GdL2 and GdL3, is likely related to the occurrence of a reduced rate of solvent collisions between the coordinated and bulk water molecules. Tentatively, one may relate the observed behavior to the enhanced shielding of the poly-hydroxylic containing moieties with respect to GdL1.

In summary, one may conclude that the high relaxivity shown by GdL3 is the result of the occurrence of three positive conditions: (i) an array of second sphere water molecules linked through hydrogen bond to the phosphonate moiety, (ii) a fast water exchange rate of the coordinated water molecule and (iii) the increased molecular weight ($\tau_R = 560 \text{ ns}$) determined by the introduction of the poly-hydroxylic substituents on the surface of the metal complex. Noteworthy, the relaxivity values for GdL3, measured at the Larmor frequencies more used on clinical scanners (40, 60, 120 MHz corresponding to 1, 1.5 and 3 T, respectively), are somehow constant, being $20.8 \text{ mM}^{-1} \text{ s}^{-1}$ at 40 and 60 MHz and $19.1 \text{ mM}^{-1} \text{ s}^{-1}$ at 120 MHz.

Experimental

General

All reagents and solvents were obtained from commercial suppliers and directly used without further purification. N,N' -Bis-[2-(2,6-dioxo-4-morpholinyl)ethyl]glycine (DTPA-bis-anhydride) 4,²¹ pentaerythrityl tetramine tetrahydrochloride 5,²² *tert*-butyl phosphite²⁸ and 2-bromoethyl trifluoromethanesulfonate²⁹ were synthesized as reported in literature. TLC was performed on Merck silica gel 60 TLC plates F254 and visualized by using UV (254 nm) or 1% KMnO_4 in 1 N NaOH. Column chromatography was performed by using silica gel 60 (70–230 mesh) while flash chromatography was carried out on silica gel 60 (230–400 mesh). The ^1H , ^{13}C and ^{31}P spectra were recorded on a Bruker Avance 400 instrument. Chemical shifts are reported in δ relative to an internal standard of residual chloroform (δ 7.27 for ^1H NMR and 77.16 for ^{13}C NMR). ESI MS spectra on novel compounds were acquired on a high resolving power mass spectrometer LTQ Orbitrap (Thermo Scientific, Rodano, Italy), equipped with an atmospheric pressure interface and an ESI ion source. Other mass spectra were recorded with a ThermoFinnigan TSQ700 triple-quadrupole instrument equipped with an electrospray ionization source. Analytical HPLC was performed on a Merck KGaA apparatus with the following method: stationary phase: Lichrospher RP-Select B 5 μm , $250 \times 4 \text{ mm}$ column packed by Merck KGaA; mobile phase: eluent A = 0.01 M KH_2PO_4 and 0.017 M H_3PO_4 in H_2O , eluent B = MeCN, gradient elution: $t =$

0 min (5% B), $t = 30$ min (80%B); $T = 45$ °C; flow rate: 1 mL min⁻¹; UV detection: 210 nm.

Synthesis of *N,N'*-(iminodi-2,1-ethanediy)bis-D-gluconamide (3). Diethylenetriamine **1** (1 g, 9.7 mmol) was added dropwise to a suspension of δ -gluconolactone **2** (3.6 g, 20.3 mmol) in DMF (10 mL). The reaction mixture was stirred at room temperature for 1.5 h. The precipitate was filtered, washed with CH₂Cl₂ (500 mL), to remove DMF and the excess of δ -gluconolactone, and dried to obtain **3** as a white solid (4.06 g, 8.83 mmol), yield 91%; mp 142 °C (dec.); R_f 0.12 (eluent CHCl₃/MeOH/25% NH₄OH 30 : 60 : 20); ¹H NMR (600 MHz, D₂O) δ 4.24 (d, $J = 5.0$ Hz, 2H), 4.00 (d, $J = 5.0$ Hz, 2H), 3.72 (d, $J = 16.9$ Hz, 2H), 3.66 (m, 4H), 3.55 (dd, $J = 11.0, 4.4$ Hz, 3H), 3.31 (dt, $J = 12.1, 9.2$ Hz, 4H), 2.71 (t, $J = 9.2$ Hz, 4H); ¹³C NMR (151 MHz, D₂O) δ 175.08 (CO), 73.81 (C-2), 72.50 (C-3), 71.45 (C-4), 70.70 (C-5), 62.96 (C-6), 47.33 (CH₂N), 38.45 (CH₂N).

Synthesis of *N,N'*-bis[2-[[carboxymethyl][2-bis[2-[[D-gluconoyl]amino]ethyl]amino]-2-oxoethyl]amino] ethyl]glycine (L1). *N,N'*-Bis[2-(2,6-dioxo-4-morpholinyl)ethyl]glycine (DTPA-bis-anhydride)²¹ **4** (0.19 g; 0.54 mmol) was added to a solution of **3** (0.5 g; 1.09 mmol) in DMSO (5 mL) and the yellow solution was stirred at 50 °C for 8 h. After cooling at room temperature, the reaction mixture was diluted with CH₂Cl₂ (50 mL); the precipitate was filtered, washed with CH₂Cl₂ (200 mL) then lyophilized to remove the residual DMSO. **L1** was obtained as white solid (0.42 g; 0.33 mmol), yield 61%; mp 128–130 °C R_f 0.30 (eluent MeOH/25% NH₄OH 7 : 3); HPLC: retention time 2.1 min, 98% (area%); ¹H NMR (600 MHz, D₂O, 333 K) δ 4.70–4.58 (m, 10H), 4.37 (s, 4H), 4.18 (s, 4H), 4.12–4.05 (m, 12H), 3.98–3.85 (m, 12H), 3.78 (m, 12H), 3.49 (m, 4H); ¹³C NMR (151 MHz, D₂O, 333 K) δ 175.68, 175.36, 174.90, 171.11, 167.40 (CO), 74.21 (C-2), 74.15 (C-2'), 73.97 (C-3), 73.07 (C-3'), 72.72 (C-4), 72.19 (C-4'), 71.30 (C-5), 62.56 (C-6), 58.16, 56.80, 56.02 (CH₂CO), 53.92, 50.42 (CH₂N), 47.41, 46.55 (CH₂N), 37.49, 36.50 (CH₂NH). ESI-HRMS (m/z): 1276.4563 (M + H⁺) (calcd 1276.5379).

Synthesis of *N*-[[2,2-bis[[D-gluconoylamino)methyl]-3-amino]propyl]-D-gluconamide (6). Pentaerythrityl tetramine tetrahydrochloride²² **5** (14 g; 0.051 mol) was suspended in a mixture of DMF (170 mL) and Et₃N (25.4 mL; 0.184 mmol). A solution of δ -gluconolactone **2** (32.7 g; 0.184 mol) in DMF (195 mL) was slowly added dropwise (1.5 h) at room temperature. The mixture was stirred at room temperature for 4 days. The reaction mixture was filtered and concentrated at reduce pressure. The residue was treated with MeCN to give a precipitate which was filtered and washed with MeCN. The solid was dried to give **6** as a white solid and as hydrochloride salt (30.5 g; 0.046 mmol), yield 90%; R_f 0.44 (eluent MeOH/25% NH₄OH 7 : 3); ¹H NMR (600 MHz, D₂O) δ 4.35 (m, 3H), 4.06 (dt, $J = 9.5, 3.2$ Hz, 3H), 3.75 (m, 6H), 3.69 (m, 3H), 3.60 (dd, $J = 11.8, 6.3$ Hz, 3H), 3.21–3.10 (m, 6H), 3.06–2.96 (m, 2H); ¹³C NMR (151 MHz, D₂O) δ 176.55 (CO), 175.99 (CO), 74.04 (C-2), 73.91 (C-2'), 72.95 (C-3), 72.74 (C-3'), 71.52 (C-4), 70.83 (C-5), 70.68 (C-5'), 62.96 (C-6), 62.93 (C-6'), 44.86 (C), 38.59 (CH₂N), 38.47 (CH₂N). ESI-HRMS (m/z): 667.2175 (M + H⁺) (calcd 667.2880).

Synthesis of *N,N'*-bis[[[2,2-bis[[D-gluconoylamino)methyl]-3-amino]propyl]-D-gluconamide]-2-oxoethyl]amino] ethyl]glycine (L2). *N,N'*-Bis[2-(2,6-dioxo-4-morpholinyl)ethyl]glycine (DTPA-

bis-anhydride)²¹ **4** (0.19 g; 0.54 mmol) was added to a solution of **6** (0.8 g; 1.20 mmol) in DMSO (5 mL) and stirred at 50 °C for 8 h. After cooling at room temperature, the reaction mixture was diluted with CH₂Cl₂ (50 mL); the precipitate was filtered, washed with CH₂Cl₂ (200 mL) then lyophilized to remove the residual DMSO. **L2** was obtained as white solid (0.64 g; 0.38 mmol), yield 70%. ¹H NMR (300 MHz, D₂O) δ 4.18 (m, 4H), 3.94 (d, $J = 3.2$ Hz, 2H), 3.86–3.73 (m, 6H), 3.65–3.54 (m, 8H), 3.51 (m, 8H), 3.45–3.32 (m, 8H), 3.06–2.77 (m, 6H), 2.51 (s, 18H), 2.29 (m, 6H), 0.78 (t, $J = 7.2$ Hz, 4H). ¹³C NMR (75 MHz, D₂O, pH = 10) δ 179.4, 173.7, 168.1, 74.8, 74.7, 73.9, 71.9, 71.4, 70.9, 70.3, 63.0, 45.2, 38.5, 9.9. ESI-HRMS (m/z): 1690.6211 (M + H⁺) (calcd 1690.6865).

Synthesis of (*N*-[[2,2-bis[D-gluconoylamino)methyl]-3-*tert*-butoxycarbonylamino]propyl]-D-gluconamide (7). Compound **6** (30.0 g, 45.0 mmol) was dissolved in DMF (400 mL) and a solution of (Boc)₂O (17.2 g; 65.2 mmol) in DMF (100 mL) was slowly added dropwise. Then, tetramethylammonium hydroxide (8.15 g, 45.0 mmol) was added and the mixture was stirred at room temperature for 2 days. The solution was concentrated at reduced pressure and the residue was treated with CH₂Cl₂ to give a precipitate which was filtered and washed with H₂O (100 mL) and CH₃OH (100 mL). The solid so obtained was dried to give **7** (25 g, 32.6 mmol), yield: 72%. ¹H NMR (600 MHz, D₂O) δ 4.35 (d, $J = 2.7$ Hz, 3H), 4.07 (t, $J = 3.1$ Hz, 3H), 3.75 (m, 6H), 3.69 (m, 3H), 3.61 (dd, $J = 11.8, 6.4$ Hz, 3H), 3.12 (s, 15H), 3.02 (m, 6H), 2.85 (m, 2H), 1.41 (s, 9H); ¹³C NMR (151 MHz, D₂O) δ 175.79 (CO), 158.81 (CO), 81.88 (C), 74.06 (C-2), 72.94 (C-3), 71.52 (C-4), 70.69 (C-5), 62.97 (C-6), 55.67, 45.52 (C), 39.94 (CH₂N), 38.72 (CH₂N), 28.07 (CH₃). ESI-HRMS (m/z): 789.3003 (M + Na⁺) (calcd 789.3229).

Synthesis of (*N*-[[2,2-bis[2,3,4,5,6-penta-*O*-acetyl-D-gluconoylamino)methyl]-3-*tert*-butoxycarbonylamino]propyl]-2,3,4,5,6-penta-*O*-acetyl-D-gluconamide (8). To a suspension of compound **7** (10.0 g; 13.0 mmol) in Ac₂O (186 mL; 1.97 mol), pyridine (25.7 g; 326 mmol) was added at room temperature. The mixture was then heated to 90 °C to dissolve the reagents, then cooled at room temperature. After 1 day at room temperature, the mixture was concentrated at reduce pressure and the residue was dissolved in EtOAc (200 mL) and washed with water (200 mL) and with 10% aqueous NaHCO₃ (200 mL). The organic phase was dried (Na₂SO₄) and evaporated to give **8** (15 g; 10.7 mmol), yield: 82%, ¹H NMR (600 MHz, CDCl₃) δ 8.21 (s, 1H), 7.81 (s, 2H), 6.00 (s, 1H), 5.65 (m, 3H), 5.45 (m, 3H), 5.26 (m, 3H), 5.07 (m, 3H), 4.31 (m, 3H), 4.17 (br m, 3H), 3.37 (m, 4H), 3.22 (m, 2H), 3.04 (m, 2H), 2.29, 2.10, 2.06 (m, 45H), 1.42 (s, 9H). ¹³C NMR (151 MHz, CDCl₃) δ 170.84 (CO), 170.11 (CO), 170.04 (CO), 158.25 (CO), 80.26 (C), 71.50 (C-2), 70.43 (C-3), 70.28 (C-4), 69.23 (C-5), 61.68 (C-6), 39.65, 28.66 (CH₃), 21.09 (CH₃), 21.04 (CH₃), 20.92 (CH₃), 20.64 (CH₃); ESI-HRMS (m/z): 1419.4123 (M + Na⁺) (calcd 1419.4814).

Synthesis of (*N*-[[2,2-bis[2,3,4,5,6-penta-*O*-acetyl-D-gluconoylamino)methyl]-3-amino]propyl]-2,3,4,5,6-penta-*O*-acetyl-D-gluconamide (9). Trifluoroacetic acid (3.67 g; 32.0 mmol) was added to a solution of compound **8** (4.50 g; 3.20 mmol) in CH₂Cl₂ (60 mL). After 3 days at room temperature, further TFA (1.82 g; 16.0 mmol) was added to the reaction mixture. After

further 4 days at room temperature, the solvents were removed under reduced pressure and the residue was dissolved in CH_2Cl_2 (7 mL) and TFA (5.49 g; 48.0 mmol). The mixture was stirred at room temperature for 1 day, then evaporated under reduced pressure. The residue was dissolved in CH_2Cl_2 (10 mL) and evaporated several times affording **9** as a white solid and as trifluoroacetic salt (3.00 g; 2.13 mmol), yield: 67%; R_f 0.59 (eluent EtOAc/*n*-hexane 8 : 2); ^1H NMR (400 MHz, CDCl_3) δ 8.27 (s, 1H), 7.78 (s, 2H), 7.34 (s, 2H), 5.60 (m, 3H), 5.46 (m, 3H), 5.24 (d, $J = 3.3$ Hz, 3H), 5.09 (td, $J = 6.0, 3.2$ Hz, 3H), 4.34 (m, 3H), 4.16 (dd, $J = 12.4, 5.8$ Hz, 3H), 3.02 (m, 6H), 2.70 (s, 2H), 2.24, 2.13, 2.09, 2.07 (s, 45H). ^{13}C NMR (101 MHz, CDCl_3) δ 170.99 (CO), 170.61 (CO), 170.47 (CO), 72.46 (C-2), 70.07 (C-3), 69.37 (C-4), 69.07 (C-5), 62.00 (C-6), 45.20 (C), 38.96 (CH_2N), 38.49 (CH_2N), 21.01 (CH_3), 20.96 (CH_3), 20.72 (CH_3), 20.59 (CH_3); ESI-HRMS (m/z): 1297.3996 ($\text{M} + \text{H}^+$) (calcd 1297.4470).

Synthesis of (dibenzylamino)methylphosphonic acid di-*tert*-butyl ester (18). Paraformaldehyde (2.14 g; 71.2 mmol) was added to a solution of dibenzylamine (12.7 g; 64.5 mmol) in MeCN (130 mL). The suspension was heated to 80 °C for 1 h, then the resulting solution was cooled to rt. A solution of *tert*-butyl phosphite (71.2% of purity, 25 g; 71.2 mmol) in MeCN was added dropwise over 25 min to the reaction mixture and the solution was stirred at room temperature for 22 h. The solvent was evaporated under vacuum and 0.1 N HCl (320 mL) was added to the residue. The suspension was extracted with CH_2Cl_2 (3 × 150 mL) and the combined organic phases washed with water (3 × 150 mL), dried (Na_2SO_4) and evaporated *in vacuo*. The crude (27.3 g) was purified by chromatography on silica gel (15 : 85 EtOAc/petroleum ether; R_f 0.2) and the desired product was obtained (12.4 g, 30.8 mmol) as a colourless oil. Yield: 47.7%. ^1H -NMR (CDCl_3) 400 MHz $\delta = 7.43$ (d, 4H, $J = 7.2$ Hz, H_{ar}), 7.32 (t, 4H, $J = 7.2$ Hz, H_{ar}), 7.24 (m, 2H, H_{ar}), 3.94 (s, 4H, CH_2Ph), 2.79 (d, 2H, $J = 10.2$ Hz, CH_2P), 1.49 (s, 18H, CCH_3). ^{13}C -NMR (CDCl_3) 100 MHz $\delta = 139.6$ (C_{ar}), 129.6–128.6–127.3 (CH_{ar}), 82.4 (CCH_3), 59.5 (CH_2Ph), 52.5 (CH_2P , $J = 163$ Hz), 30.9 (CCH_3); ^{31}P -NMR (CDCl_3) 162 MHz $\delta = 19.4$. MS (ESI+; MeOH): m/z 404 ($\text{M} + \text{H}^+$).

Synthesis of aminomethylphosphonic acid di-*tert*-butyl ester (19). (Dibenzylamino)methylphosphonic acid di-*tert*-butyl ester (33.3 g; 82.7 mmol) was dissolved in MeOH (500 mL) and hydrogenated at atmospheric pressure on Pd/C 10% (3.3 g). After 2 h the mixture was filtered through Millipore® apparatus (0.5 μm) and the solution evaporated under vacuum. The crude (18.4 g) was used without any further purification. Quantitative yield. ^1H -NMR (CDCl_3) 400 MHz $\delta = 2.81$ (d, 2H, $J = 10.2$ Hz, CH_2P), 1.49 (s, 18H, CCH_3). ^{13}C -NMR (CDCl_3) 100 MHz $\delta = 83.2$ (CCH_3), 40.6 (CH_2P , $J = 153$ Hz), 29.5 (CCH_3); ^{31}P -NMR (CDCl_3) 162 MHz $\delta = 21.0$. MS (ESI+; MeOH): m/z 224 ($\text{M} + \text{H}^+$).

Synthesis of L-aspartic acid, 1-*tert*-butyl-4-benzylester (11). Perchloric acid (70% aq.; 19.28 g; 0.134 mol) was dropped into a suspension of L-aspartic acid 4-(phenylmethyl) ester (25 g; 0.112 mol) in *tert*-butyl acetate (515 mL; 3822 mol) stirred at room temperature. After 18 h at room temperature the obtained clear solution was diluted with H_2O (470 mL) and the phases were separated; the aqueous phase was extracted with EtOAc (2 × 235 mL). The organic phases were collected and washed

with 5% aq. NaHCO_3 (2 × 250 mL) and H_2O (2 × 200 mL); the new aqueous phases were collected and extracted with EtOAc (3 × 100 mL). All organic phases were collected, dried over Na_2SO_4 and evaporated to obtain **11** (27 g; 0.097 mol) as a pale yellow oil. Yield 86.6%. TLC (silica gel; eluent: EtOAc; R_f 0,57). ^1H -NMR (CDCl_3) 400 MHz $\delta = 7.31$ (m, 5H, H_{ar}), 5.14 (s, 2H, CH_2Ph), 3.68 (t, 1H, $J = 6.3$ Hz, CHNH_2), 2.77 and 2.66 (dd, 2H, $J_1 = 16.3$ Hz, $J_2 = 4.9$ Hz, $\text{CH}_2\text{CO}_2\text{Bn}$), 1.76 (bs, 2H, NH_2), 1.40 (s, 9H, CCH_3). ^{13}C -NMR (CDCl_3) 100 MHz $\delta = 173.7$ (CO_2Bn), 171.5 (CO_2tBu), 136.1 (C_{ar}), 128.7 (CH_{ar}), 81.8 (CCH_3), 66.8 (CH_2Ph), 52.2 (CHNH_2), 39.6 ($\text{CH}_2\text{CO}_2\text{Bn}$), 28.3 (CCH_3). MS (ESI+; MeOH): m/z 280.2 ($\text{M} + \text{H}^+$); 302.2 ($\text{M} + \text{Na}^+$).

Synthesis of N-2-(*t*-butoxycarbonylmethyl)-L-aspartic acid, 1-*tert*-butyl-4-benzylester (12). A mixture of L-aspartic acid, 1-*tert*-butyl-4-benzylester **11** (27 g; 0.0967 mol), *t*-butyl bromoacetate (20.00 g; 0.102 mol), acetonitrile (160 mL) and 2 M phosphate buffer pH 8 (80 mL) was vigorously stirred at room temperature; after 18 h the phases was separated and the organic phase was evaporated. The residue thus obtained was dissolved in EtOAc (300 mL) and washed with H_2O (2 × 150 mL) and brine (2 × 150 mL). After drying over Na_2SO_4 , the organic solution was evaporated to give a crude (34 g) that was purified by flash-chromatography [silica gel; eluent: 4 : 1 *n*-hexane/EtOAc; R_f 0,34] to obtain 29.1 g of **1**. (Yield: 76.5%) ^1H -NMR (CDCl_3) 400 MHz $\delta = 7.32$ (m, 5H, H_{ar}), 5.14 (s, 2H, CH_2Ph), 3.59 (t, 1H, $J = 6.3$ Hz, CHNH), 3.36 (dd, 2H, $J_1 = 17.0$ Hz, $\text{CH}_2\text{CO}_2\text{tBu}$), 2.78 and 2.71 (dd, 2H, $J_1 = 16.0$ Hz, $J_2 = 6.4$ Hz, $\text{CH}_2\text{CO}_2\text{Bn}$), 1.48 and 1.44 (s, 18H, CCH_3). ^{13}C -NMR (CDCl_3) 100 MHz $\delta = 172.3$ (CO_2Bn), 171.3, 171.1 (CO_2tBu , 2C), 136.1 (C_{ar}), 128.9, 128.6 (CH_{ar}), 82.1, 81.6 (CCH_3), 66.9 (CH_2Ph), 58.0 (CHNH), 50.3 ($\text{CH}_2\text{CO}_2\text{tBu}$), 38.5 ($\text{CH}_2\text{CO}_2\text{Bn}$), 28.6, 28.3 (CCH_3). ESI-HRMS (m/z): 394.2167 ($\text{M} + \text{H}^+$) (calcd 394.2230).

Synthesis of N-2-bromoethyl-N-2-(*t*-butoxycarbonylmethyl)-L-aspartic acid, 1-*tert*-butyl-4-benzylester (13). 2-Bromoethyl trifluoromethanesulfonate²⁹ (34.93 g; 0.136 mol, 1.7 eq.) was slowly dropped into a solution of compound **12** (31.5 g; 0.08 mol) and 2,6-lutidine (27 g; 0.25 mol, 3.1 eq.) in toluene (450 mL) stirred at −15 °C under nitrogen atmosphere. The reaction mixture was stirred at room temperature for 22 h then diluted with H_2O (200 mL) and extracted with EtOAc (200 mL). The organic solution was dried over Na_2SO_4 and evaporated to give a crude (33 g) that was purified by chromatography [silica gel, eluent: 7 : 3 *n*-hexane/ i Pr₂O, R_f 0,44] to obtain **13** (24,9 g; 0,05 mol) as a pale yellow oil. Yield 62.4%. ^1H -NMR (CDCl_3) 400 MHz $\delta = 7.37$ (m, 5H, H_{ar}), 5.15 (s, 2H, CH_2Ph), 3.87 (t, 1H, $J = 7.4$ Hz, CHN), 3.40, 3.15 (m, 6H, $\text{CH}_2\text{CH}_2\text{Br}$ and $\text{CH}_2\text{CO}_2\text{tBu}$), 2.85 and 2.69 (dd, 2H, $J_1 = 16.0$ Hz, $J_2 = 7.4$ Hz, $\text{CH}_2\text{CO}_2\text{Bn}$), 1.46 and 1.44 (s, 18H, CCH_3). ^{13}C -NMR (CDCl_3) 100 MHz $\delta = 171.0$ (CO_2Bn and CO_2tBu , 3C), 136.1 (C_{ar}), 128.9, 128.7 (CH_{ar}), 82.4, 81.5 (CCH_3), 67.0 (CH_2Ph), 62.9 (CHN), 56.1, 54.9 (NCH_2 , 2C), 36.7 ($\text{CH}_2\text{CO}_2\text{Bn}$), 30.8 (CH_2Br), 28.5 (CCH_3). ESI-HRMS (m/z): 500.1206 ($\text{M} + \text{H}^+$) (calcd 500.1648).

Synthesis of N,N'-[[[bis(1,1-dimethylethoxy)phosphinyl]methyl]imino]di-2,1-ethanediyl]bis[N-2-(1,1-dimethylethoxy)-2-oxoethyl]-L-aspartic acid 1,1'-bis(1,1-dimethylethyl) 4,4'-bis(phenylmethyl) ester (14). A solution of bromo-derivative **13** (4.11 g; 8.22 mmol) in acetonitrile (20 mL) was slowly dropped

into an emulsion of aminomethylphosphonic acid di-*tert*-butyl ester **19** (916 mg; 4.11 mmol) in acetonitrile (10 mL) and 2 M phosphate buffer pH 8 (20 mL) over 8 hours under vigorous stirring. After 22 h, the phases were separated and the aqueous layer was extracted with EtOAc (2 × 30 mL). The organic layer was evaporated. The oily residue was dissolved with EtOAc (20 mL), and the solution was washed with 1 : 1 water/brine (2 × 30 mL) and dried over Na₂SO₄. The crude (4.49 g) was purified by silica gel chromatography (silica gel; EtOAc/petroleum ether 4 : 6; *R_f* 0.32). Compound **14** was obtained (1.86 g; 1.75 mmol) as a yellow oil. Yield 42.4%. ¹H-NMR (CDCl₃) 400 MHz δ = 7.32 (m, 10H, H_{ar}), 5.09 (s, 4H, CH₂Ph), 3.80 (dd, 2H, *J* = 3.6 Hz, CHN), 3.35 (s, 4H, CH₂CO₂*t*Bu), 2.82–2.62 (m, 14H, NCH₂CH₂N, NCH₂P and CH₂CO₂Bn), 1.46 and 1.41 (s, 54H, CCH₃). ¹³C-NMR (CDCl₃) 100 MHz δ = 171.3 (CO₂Bn and CO₂*t*Bu, 6C), 136.3 (C_{ar}), 128.9, 128.6 (CH_{ar}), 82.5, 81.8, 81.0 (CCH₃), 66.7 (CH₂Ph), 62.2 (NCH), 55.0 (CH₂CO₂*t*Bu), 54.7, 51.2 (NCH₂CH₂N), 52.2 (CH₂P), 36.5 (CH₂CO₂Bn), 30.9 and 28.5 (CCH₃). ³¹P-NMR (CDCl₃) 162 MHz δ = 18.6. ESI-HRMS (*m/z*): 1062.5503 (M + H⁺) (calcd 1062.6031).

Synthesis of *N,N'*-[[[bis(1,1-dimethylethoxy)phosphinyl]methyl]imino]di-2,1-ethanediyl]bis[*N*-[2-(1,1-dimethylethoxy)-2-oxoethyl]-L-aspartic acid 1,1'-bis(1,1-dimethylethyl) ester, (15**).** Compound **14** (973 mg; 1.10 mmol) was dissolved in methanol (50 mL) and hydrogenated at atmospheric pressure on 10% Pd/C (50 mg). After 30 min the reaction mixture was filtered through Millipore apparatus® (0.5 m filter) and the solution evaporated under vacuum. The crude (725 mg) was used in the next step without any further purification. ¹H-NMR (CDCl₃) 400 MHz δδ = 3.89 (m, 2H, CHN), 3.49–3.36 (m, 4H, CH₂CO₂*t*Bu), 3.25–2.95 (m, 10H, NCH₂CH₂N, NCH₂P) 2.75 (CH₂CO₂H), 1.50 and 1.46 (s, 54H, CCH₃). ¹³C-NMR (CDCl₃) 100 MHz δδ = 173.0 (COOH), 170.8, 170.2 (CO₂*t*Bu, 4C), 84.2, 82.7, 82.2 (CCH₃), 61.7 (NCH), 54.5 (CH₂CO₂*t*Bu), 54.0, 50.1 (NCH₂CH₂N), 51.8 (CH₂P, *J* = 155 Hz), 35.6 (CH₂CO₂H), 30.9 and 28.5 (CCH₃). ³¹P-NMR (CDCl₃) 162 MHz δ = 15.0. ESI-HRMS (*m/z*): 882.4855 (M + H⁺) (calcd 882.5092).

Synthesis of compound (16). A solution of diacid **15** (4.91 g; 5.58 mmol), polyalcohol **9** (17.40 g; 13.38 mmol, 2.4 equivalents) and HATU (5.08 g; 13.38 mmol) in 50 mL of DMF was brought to 0 °C with an ice bath in nitrogen atmosphere. To this solution, DIPEA (9.32 mL, 53.52 mmol, 9.6 eq.) was added keeping the temperature below 5 °C. The mixture was kept under stirring at room temperature for 48 h monitoring the reaction by TLC (silica gel; CHCl₃/MeOH 9 : 1), then the reaction solution was poured in ice, stirred for 30 min and filtered. The solid was washed with water, dissolved in chloroform and dried over Na₂SO₄. After filtration and solvent evaporation, the desired product was obtained as pale yellow gummy solid (15.3 g, 80% yield). (*R_f* 0.52, silica, CHCl₃/MeOH 9 : 1). ¹H NMR (400 MHz, CDCl₃) δ 7.86 (sb, 8H), 5.61 (m, 6H), 5.46 (m, 6H), 5.31 (m, 6H), 4.97 (m, 6H), 4.29 (m, 6H), 4.16 (m, 6H), 3.83 (m, 2H, CHN), 3.60–2.75 (m, 14H, CH₂CO₂*t*Bu, NCH₂CH₂N, NCH₂P), 2.28 (s, 16H, CCH₂N), 2.07 and 2.02 (m, 90H, CH₃COO), 1.54 and 1.46 (s, 54H, CCH₃). ¹³C NMR (101 MHz, CDCl₃) δ 172.8, 170.9, 170.5, 169.8, 169.6, 168.7, 81.4, 69.9, 68.9, 61.4, 45.8, 38.4, 30.7, 30.0, 28.3, 28.2, 20.7. ESI-HRMS (*m/z*): 3439.2986 (M + H⁺) (calcd 3439.3665).

Synthesis of compound (17). Ammonia was bubbled for 45 min into a solution of the fully protected ligand **16** (19.48 g) in methanol (400 mL) refrigerated with an ice bath. After 1 h the ice bath was removed, CaCl₂ valve was applied and the solution was allowed to stir at room temperature for 9 days. Every two days more ammonia was bubbled for 10–15 min into the solution. The reaction was followed by TLC (CHCl₃/MeOH 9 : 1) and by MS until disappearance of any peaks related to partially acetylated product. Then, the solvent was evaporated in vacuum obtaining 18.1 g of the crude which was used without any further purification in the next deprotection step. ¹H NMR (400 MHz, D₂O) δ 4.34 (bs, 6H), 4.06 (bs, 6), 3.82–3.71 (m, 8H), 3.68 (t, *J* = 6.8 Hz, 4H), 3.59 (m, 4H), 2.94 (bs, 8H), 2.78 (s, 4H), 2.73 (s, 2H), 1.42 (m, 56H). ¹³C NMR (101 MHz, CDCl₃) δ 171.9, 171.5, 170.8, 170.7, 169.7, 82.4, 78.0, 70.9, 70.0, 62.4, 46.7, 39.2, 31.7, 31.0, 29.2. ESI-HRMS (*m/z*): 2179.1102 (M + H⁺) (calcd 2179.0495).

Synthesis of compound (L3). A round-bottomed flask containing the phosphonic proligand **17** (18.1 g, 8.3 mmol) was cooled with an ice bath; then a mixture of CH₂Cl₂/TFA 1 : 1 (100 mL) was added dropwise giving a yellow-brown solution. After 20 min, the ice bath was removed and the solution was kept under stirring at room temperature for 16 h. The solvent was removed and the residue was dissolved in dichloromethane (5 mL) and evaporated; this procedure was repeated four times. The crude was treated with Et₂O to obtain the precipitation of a white solid which was filtered, dissolved in water and lyophilized. The slightly yellow solid was then re-dissolved in water, brought to pH 7 by adding 2 M NaOH and lyophilized again. The solid obtained was recovered with MeOH and kept under stirring for 1 h. The white solid formed was filtered and dried in vacuum to obtain 4.6 g (2.50 mmol, 30% yield) of product. The methanolic solution was dried at the rotary evaporator and in vacuum to obtain 2.9 g of the mono-derivative as a yellow oil. ¹H NMR (400 MHz, D₂O) δ 4.39 (bs, 6H), 4.10 (bs, 6H), 3.82–3.76 (m, 8H), 3.76–3.70 (m, 4H), 3.64 (dd, *J* = 11.6, 6.3 Hz, 4H), 3.32 (s, 4H), 3.07 (s, 8H), 2.96 (s, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 171.9, 171.5, 170.8, 170.7, 169.7, 82.4, 78.0, 70.9, 70.0, 62.4, 46.7, 39.2, 31.7, 31.0, 29.2. ESI-HRMS (*m/z*): 1842.6012 (M + H⁺) (calcd 1842.6739).

Water proton relaxation measurements

The longitudinal water proton relaxation rates were measured at 25 °C by using a Stelar Spinmaster (Stelar, Mede, Pavia, Italy) spectrometer operating at 0.5 T (21.5 MHz Proton Larmor Frequency), by mean of the standard inversion-recovery technique. The temperature was controlled with a Stelar VTC-91 air-flow heater equipped with a copper constantan thermocouple (uncertainty 0.1 °C). The proton 1/*T*₁ NMRD profiles were measured at 25 °C on a fast field-cycling Stelar relaxometer over a continuum of magnetic field strengths from 0.00024 to 0.47 T (corresponding to 0.01–20 MHz proton Larmor frequencies). The relaxometer operates under computer control with an absolute uncertainty in 1/*T*₁ of ±1%. Additional data points in the range 21.5–70 MHz were obtained on the Stelar Spinmaster spectrometer. The concentration of the solutions used for the

relaxometric characterization was determined by measuring the bulk magnetic-susceptibility shifts of the *t*BuOH signal.

¹⁷O-NMR measurements

Variable temperature ¹⁷O-NMR measurements were recorded at 2.1 T on a JEOL90 spectrometer, equipped with a 5 mm probe, by using a D₂O external lock. The experimental settings were: spectral width 9000 Hz, 90° pulse (12 μs), acquisition time 10 ms, 1024 scans and without sample spinning. Aqueous solutions containing 2.6% of ¹⁷O isotope (Yeda, Israel) were used. The observed transverse relaxation rates ($R_{2\text{obs}}^{\text{O}}$) were calculated from the signal width at half-height ($\Delta\nu_{1/2}$): $R_{2\text{obs}}^{\text{O}} = \pi\Delta\nu_{1/2}$. Paramagnetic contributions to the observed transversal relaxation rate ($R_{2\text{p}}$) were calculated by subtracting from $R_{2\text{obs}}^{\text{O}}$ the diamagnetic contribution measured at each temperature value on pure water enriched with 2.6% ¹⁷O isotope.

Conclusions

The synthesis of three novel DTPA derivatives bearing polyhydroxylated pendant arms is reported. While the two DTPA bisamides **L1** and **L2** with differently branched polygluconyl groups were easily synthesised from DTPA bis-anhydride, the monophosphonic P-DTPA ligand **L3** was prepared by a multi-step procedure which included the synthesis of a bis-functionalized DTPA bifunctional ligand *via* the Rapoport reaction. The relaxivity of Gd**L1** and Gd**L2** were very low because DTPA-bisamides are characterized by slow water exchange rates of the coordinated water molecule. On the other hand, the relaxivity values measured for Gd**L3** over a wide range of imaging fields (0.5–3 Tesla) are among the highest till now reported for monomeric Gd(III) complexes with medium sized molecular weight (~2000 Da). This finding makes Gd**L3** a very promising system for applications at the currently used magnetic fields of the clinical MRI scanners.

Notes and references

- 1 *The Chemistry of Contrast Agents in Medical Magnetic Resonance Imaging*, ed. A. Merbach, L. Helm and É. Tóth, John Wiley & Sons Ltd, Chichester, 2nd edn, 2013.
- 2 (a) C. F. G. C. Geraldes and S. Laurent, *Contrast Media Mol. Imaging*, 2009, **4**, 1; (b) P. Hermann, J. Kotek, V. Kubiček and I. Lukeš, *Dalton Trans.*, 2008, 3027; (c) P. Caravan, J. J. Ellison, T. J. McMurry and R. B. Lauffer, *Chem. Rev.*, 1999, **99**, 2293.
- 3 (a) Y. Lvovsky, E. W. Stautner and T. Zhang, *Supercond. Sci. Technol.*, 2013, **26**, 1; (b) T. C. Cosmos and M. Parizh, *IEEE Trans. Appl. Supercond.*, 2011, **21**, 2104.
- 4 (a) P. R. Luijten and D. W. J. Klomp, *Drug Discovery Today: Technol.*, 2011, **8**, e103; (b) C. K. Khul, F. Träber and H. H. Schild, *Radiology*, 2008, **246**, 675.
- 5 (a) L. Helm, *Future Med. Chem.*, 2010, **2**, 385; (b) P. Caravan, C. T. Farrara, L. Frullano and U. Ritika, *Contrast Media Mol. Imaging*, 2009, **4**, 89.
- 6 M. Botta and L. Tei, *Eur. J. Inorg. Chem.*, 2012, 1945.
- 7 (a) M. R. Longmire, M. Ogawa, P. L. Choyke and H. Kobayashi, *Wiley Interdiscip. Rev.: Nanomed. Nanobiotechnol.*, 2014, **6**, 155; (b) A. Bumb, M. W. Brechbiel and P. Choyke, *Acta Radiol.*, 2010, **51**, 751.
- 8 P. Caravan, *Acc. Chem. Res.*, 2009, **42**, 851.
- 9 J. Martinelli, M. Fekete, L. Tei and M. Botta, *Chem. Commun.*, 2011, **47**, 3144.
- 10 (a) W. J. M. Mulder, G. J. Strijkers, G. A. F. Van Tilborg, D. P. Cormode, Z. A. Fayad and K. Nicolay, *Acc. Chem. Res.*, 2009, **42**, 904; (b) E. Terreno, D. Delli Castelli, C. Cabella, W. Dastrù, A. Sanino, J. Stancanello, L. Tei and S. Aime, *Chem. Biodiversity*, 2008, **5**, 1901.
- 11 (a) Y. Song, E. K. Kohlmeier and T. J. Meade, *J. Am. Chem. Soc.*, 2008, **130**, 6662; (b) J. B. Livramento, E. Toth, A. Sour, A. Borel, A. E. Merbach and R. Ruloff, *Angew. Chem., Int. Ed. Engl.*, 2005, **44**, 1480.
- 12 (a) D. A. Fulton, E. Elemento, S. Aime, L. Chaabane, M. Botta and D. Parker, *Chem. Commun.*, 2006, 1064; (b) D. A. Fulton, M. O'Halloran, D. Parker, K. Senanayake, M. Botta and S. Aime, *Chem. Commun.*, 2005, 474.
- 13 (a) L. Van Der Elst, Y. Raynal, M. Port, P. Tisnes and R. N. Muller, *Eur. J. Inorg. Chem.*, 2005, **6**, 1142; (b) M. Port, C. Corot, O. Rousseaux, I. Raynal, L. Devoldere, J. M. Idée and A. Dencausse, *Magn. Reson. Mater. Phys., Biol. Med.*, 2001, **12**, 121; (c) M. Port, D. Meyer, B. Bonnemain, C. Corot, M. Schaefer, O. Rousseaux, C. Simonot, P. Bourrinet, S. Benderbous, A. Dencausse and L. Devoldere, *Magn. Reson. Mater. Phys., Biol. Med.*, 1999, **8**, 172.
- 14 J. Kotek, P. Lebduskova, P. Hermann, L. Vander Elst, R. N. Muller, C. F. G. C. Geraldes, T. Maschmeyer, I. Lukes and J. A. Peters, *Chem.-Eur. J.*, 2003, **9**, 5899.
- 15 M. Botta, *Eur. J. Inorg. Chem.*, 2000, 399.
- 16 (a) P. Baia, J. P. Andre, C. F. G. C. Geraldes, J. A. Martins, A. E. Merbach and E. Toth, *Eur. J. Inorg. Chem.*, 2005, 2110; (b) J. P. Andre, C. F. G. C. Geraldes, J. A. Martins, A. E. Merbach, M. I. M. Prata, A. C. Santos, J. J. P. de Lima and E. Toth, *Chem.-Eur. J.*, 2004, **10**, 5804; (c) K. Luo, G. Liu, X. Zhang, W. She, B. He, Y. Nie, L. Li, Y. Wu, Z. Zhang, Q. Gong, F. Gao, B. Song, H. Ai and Z. Gu, *Macromol. Biosci.*, 2009, **9**, 1227.
- 17 V. C. Pierre, M. Botta and K. N. Raymond, *J. Am. Chem. Soc.*, 2005, **127**, 504.
- 18 (a) G. Yu, M. Yamashita, K. Aoshima, M. Takahashi, T. Oshikawa, H. Takayanagi, S. Laurent, C. Burtea, L. Vander Elst and R. N. Muller, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 2246; (b) M. Takahashi, Y. Hara, K. Aoshima, H. Kurihara, T. Oshikawa and M. Yamashita, *Tetrahedron Lett.*, 2000, **41**, 8485.
- 19 L. Lattuada, A. Barge, G. Cravotto, G. B. Giovenzana and L. Tei, *Chem. Soc. Rev.*, 2011, **40**, 3019.
- 20 J. F. W. Keana and J. S. Mann, *J. Org. Chem.*, 1990, **55**, 2868.
- 21 (a) C. F. G. C. Geraldes, A. M. Urbano, M. C. Alpoim, A. D. Sherry, K.-T. Kuan, R. Rajagopalan, F. Maton and R. N. Muller, *Magn. Reson. Imaging*, 1995, **13**, 401; (b) W. C. Eckelman, S. M. Karesh and R. C. Reba, *J. Pharm. Sci.*, 1975, **64**, 704.

- 22 (a) S. Dutta, P. Biswas, U. Florke and K. Nag, *Inorg. Chem. Commun.*, 2010, **13**, 1074; (b) W. Hayes, H. M. I. Osborn, S. D. Osborne, R. A. Rastall and B. Romagnoli, *Tetrahedron*, 2003, **59**, 7983; (c) A. McAuley, S. Subramanian and T. W. Whitcombe, *Can. J. Chem.*, 1989, **67**, 1650.
- 23 M. A. Williams and H. Rapoport, *J. Org. Chem.*, 1993, **58**, 1151.
- 24 P. L. Anelli, F. Fedeli, O. Gazzotti, L. Lattuada, G. Lux and F. Rebasti, *Bioconjugate Chem.*, 1999, **10**, 137.
- 25 A. Barge, G. Cravotto, E. Gianolio and F. Fedeli, *Contrast Media Mol. Imaging*, 2006, **1**, 184.
- 26 (a) S. Laurent, L. Vander Elst, F. Botteman and R. N. Muller, *Eur. J. Inorg. Chem.*, 2008, 4369; (b) E. Toth, F. Connac, L. Helm, K. Adzamli and A. E. Merbach, *Eur. J. Inorg. Chem.*, 1998, 2017.
- 27 S. Aime, M. Botta, F. Fedeli, E. Gianolio, E. Terreno and P. Anelli, *Chem.–Eur. J.*, 2001, **7**, 5261.
- 28 (a) J. R. Cox Jr and M. G. Newton, *J. Org. Chem.*, 1969, **34**, 2600; (b) H. C. Manning, M. Bai, B. M. Anderson, R. Lisiak, L. E. Samuelson and D. J. Bornhop, *Tetrahedron Lett.*, 2005, **46**, 4707.
- 29 M. Yar, E. M. McGarrigle and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2008, **47**, 3784.