

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

“Ormilo disease” a disorder of zebu cattle in Tanzania: bovine cerebral theileriosis or new protozoan disease?

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1531313> since 2015-12-04T11:06:42Z

Published version:

DOI:10.1007/s11250-015-0805-8

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **“Ormilo disease” a disorder of Zebu cattle in Tanzania: bovine cerebral theileriosis or new protozoan disease?**
2
3 Deborah Catalano · Elena Biasibetti · Godelieve Lynen · Giuseppe Di Giulio · Daniele De Meneghi · Laura Tomassone ·
4 Federico Valenza · Maria Teresa Capucchio.
5
6 D. Catalano · E. Biasibetti · D. De Meneghi · L. Tomassone · F. Valenza · MT. Capucchio
7 Department of Veterinary Sciences, University of Turin, Largo Paolo Braccini 2, 10095 Grugliasco, Italy
8
9 G. Lynen · G. Di Giulio. Former affiliation: Integrated Tick and Tick-borne Disease Control Project, Arusha, United
10 Republic of Tanzania (URT). Present address: Merlino Veterinary Clinic, P.O. Box 13188, Arusha, Tanzania (URT)
11
12 Author for correspondence, **Maria Teresa Capucchio**: e-mail: mariateresa.capucchio@unito.it, telephone:
13 00390116709035 and fax numbers: 00390116709031
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

Abstract

“Ormilo” disease is a neurological disorder of cattle described by Maasai herders in Tanzania. It is attributed to infection by *Theileria* species, although no detailed data are available in the literature.

The authors describe the macroscopical and histological changes observed in 30 brains of indigenous short-horn zebu cattle from Northern Tanzania, aged 2-9 years, with the characteristic neurological signs of “Ormilo”. Moreover the ultrastructural details observed in 14 selected brains samples were reported. Areas of congestion and hemorrhages, associated with the obstruction of the cerebral vessels with large numbers of parasitized lymphoid cells, were observed. Electron microscopy showed the presence of intralymphocytic parasites morphologically comparable to flagellated protozoa, not previously described in the lymphoid cells of cattle, but only reported during the sexual stages within the vector. *Theileria taurotragi* was detected by Polymerase Chain Reaction and Reverse Line Blot in 9 samples. The authors hypothesize that the parasite detected by electron microscopy could be a strain of a *Theileria* endemic to this region till now not investigated, having an intralymphocytic phase and being associated with other *Theileria* spp. infestation. Further studies are needed to better understand the etiology of “Ormilo” disease and to characterize the morphology of the observed parasite, clarifying its role in the disease in Tanzania.

Key words: Bovine Cerebral Theileriosis, Ormilo, pathology, short-horn zebu cattle, Tanzania, *Theileria*.

Introduction

Theileriae are tick-transmitted intracellular apicomplexan parasites, which infect a wide range of mammals worldwide (Norval et al. 1992; Dobbelaere and McKeever 2002). After the tick bite, *Theileria* sporozoites invade lymphoid cells of the vertebrate host and induce them to proliferate in an unregulated manner. The cells develop into a multinucleate syncytial schizont (macroschizont), later in microschizonts that differentiate into uninucleate merozoites, that are liberated into the blood where they invade the erythrocytes and become piroplasms (Fig. 1).

Several species of *Theileria* are known. *Theileria parva* and *Theileria annulata* are considered the most important pathogens of domestic livestock in the Old World tropical and subtropical regions. *Theileria taurotragi* was first described in 1960 as a benign parasite of the eland (Martin and Brocklesby 1960). Subsequently, in cattle it was demonstrated that this parasite causes a benign parasitosis or fatal infections (De Vos et al. 1981), frequently associated with concurrent presence of *Theileria mutans* and/or *T. parva* (Young et al. 1977; Uilenberg et al. 1982).

T. parva, *T. taurotragi*, and more rarely *T. annulata* are considered causing agents of the Bovine Cerebral Theileriosis (BCT) (De Vos et al.1981; Saville 2002; Sudan et al. 2012). BCT is described in tropical and subtropical African

62 regions (Mettam et al.1936; Flanagan et al.1957), but more recently two cases caused by *T. annulata* were reported in
63 Turkey (Dabak et al. 2004) and in India (Sudan et al. 2012).

64 BCT, or Turning sickness, is a fatal disease of indigenous African short-horn cattle of cattle. It usually occurs in young
65 animals and results in a mild febrile reaction of 1-14 days duration, after an incubation period of 10-21 days. Slight
66 enlargement of the superficial lymph nodes is present, but no other clinical signs are reported. The nervous form of
67 BCT is characterized by uncontrollable movements, sometimes unilateral-bilateral blindness (corneal opacity), circling,
68 head pressing, ataxia, opisthotonus and paralysis. Usually the animals collapse after 2-21 days, but occasionally the
69 cases become chronic and they may live for up to 6 months (Van Amstel 1982; Lawrence et al. 2004). The
70 neuropathological changes include congestion and hemorrhages in the meninges and in the brain, and subacute-chronic
71 areas of malacia. Microscopically, the obstruction of the arteries with a large numbers of parasitized lymphoblasts is the
72 most prominent finding (Bader et al. 1986; Lawrence et al. 2004).

73 Cases of disease in cattle similar to BCT have been reported from Maasai herders in northern Tanzania (Arusha,
74 Manyara and Tanga regions) since the mid-1980's (Field et al. 1988; Nsengwa 1993). Although its incidence has been
75 increasing, especially in northern Maasai land, detailed pathological studies have never been documented. Local Maasai
76 livestock keepers call the disease "Ormilo" and, from information gathered during a rapid rural in 2001, they consider it
77 as the highest disease priority and, thus, as a severe constraint to livestock production (Lynen, unpublished data).

78 This paper reports the macroscopical and histological changes observed in 30 brains of indigenous short-horn zebu
79 cattle, ranging from to 2 to 9 years of age, living in Northern Tanzania (Arusha Region, Ngorongoro District, Endulen
80 ward), that presented neurological signs characteristic of "Ormilo". Ultrastructural features and Polymerase Chain
81 Reaction (PCR) investigations are also described.

82

83 **Materials and methods**

84 The brains of 30 East-African short-horn Zebu cattle aging from 2 to 9 years old, were collected between 2001 and
85 2003 in Endulen, Ngorongoro Conservation Area, Arusha region, north-eastern Tanzania. In this area, mixed herds of
86 small ruminants and cattle are kept under a traditional pastoralist livestock system.

87 These animals were selected according to the clinical description by Maasai herders of Endulen ward confirmed by the
88 veterinarians of the Integrated Tick and Tick-borne Disease Control Project. The most frequently reported clinical signs
89 was circling, associated with para-paresis and paralysis, generally followed by recumbency. In one animal, difficulties
90 in chewing and walking were also present. Except in one case, found dead, all the other animals were slaughtered due to
91 the worsening of their clinical signs.

92 Immediately after death/slaughter, the brains were fixed in 10% neutral buffered formalin and sent to the Department of
93 Veterinary Science of Torino University to perform pathological investigations. Not all the collected samples were
94 suitable for ultrastructural investigations (TEM) due to conservation problems. Therefore, meningeal blood vessels,
95 choroid plexuses and thrombosed brain vessels of 14 animals were fixed in 2.5% glutaraldehyde for TEM.

96 Tissues were postfixed in 1% osmium tetroxide, dehydrated in acetone and embedded in Spurr epoxy resin. Semithin
97 sections (0.90 μm) were cut and stained with toluidine blue. Ultrathin sections of 70 nm were then collected onto 200
98 mesh copper grids and contrasted by uranyl acetate and lead citrate. The grids were evaluated by Electron Microscopy
99 109 JD.

100 Brain tissues from the same 14 animals were also subjected to DNA extraction with the DNeasy tissue kit (QIAGEN,
101 Hilden, Germany). PCR was carried out to amplify the V4 hyper variable region of the 18S rRNA gene of *Theileria* and
102 *Babesia* spp. Species identification was carried out by reverse line blot (RLB) hybridization of the amplified products
103 on a membrane containing oligonucleotides specific for six bovine *Theileria* species and two *Babesia* spp. and a catch-
104 all *Theileria/Babesia* control probe, according to the protocol previously described (Nijhof et al. 2003).

105

106 **Results**

107 Macroscopically, moderate to severe congestion of the leptomeningeal blood vessels, disseminated sub-meningeal
108 hemorrhages and/or multifocal to diffuse hemorrhages into the brain tissue (Fig. 2a, b, c), disseminated or focal areas of
109 sub-acute/chronic malacia were the most important observed features (Fig. 2d). Histological investigations revealed
110 congestions/hemorrhages of the meningeal vessels, frequently thrombosed and necrotic, hemorrhages and
111 plasmorrhages around the ventricles and especially in the choroid plexuses (Fig. 2e), and focal to multifocal necrotic
112 areas. Severe accumulation of mononuclear cells in the cerebral and meningeal blood vessels was the most significant
113 lesion observed (Fig. 2f). Morphologically large lymphoblastic cells predominated, including sometimes small and
114 medium sized lymphocytes, rare plasma cells and macrophages. Most lymphocytic cells showed a variable number of
115 the schizonts in the cytoplasm. The main involved areas were the cortical meninges and the rostral brainstem, while the
116 pons and medulla were rarely affected by meningeal and parenchymal hemorrhages only.

117 Ultrastructurally, all the analyzed blood vessels revealed the presence of cytoplasmatic irregular structures,
118 morphologically similar to parasitic forms, in lymphoid cells. The structures observed were considered to be free
119 schizonts in the cytoplasm of the host cells whose nuclei were arranged at their periphery. Moreover, numerous free
120 merozoites were present in the cytoplasm of lymphoid cells and in the lumen of the blood vessel. In all cases, such
121 merozoites showed a flagellum and one cytoplasmatic vacuole containing a small spherical electron-dense structure.

122 These morphological data allowed the authors to propose a developing cycle of the parasite in the lymphoid cells of the
123 cattle as represented in the Fig. 3 (a - f). The first stage observed was a young schizont - spherical in shape, electron
124 dense, with diameters of about 7-9 μ m characterized by spherical dividing nuclei, numerous mitochondria and many
125 spherical organelles such as vacuoles (Fig 3a). Later the schizont appears as a spherical vesicle with 2-3 nuclei located
126 at the periphery of the cytoplasm, numerous vacuoles and some merozoites. The merozoites are spherical or pear-
127 shaped, measure about 1.6 μ m in diameter and show some micronemes, vacuoles and numerous microtubules (Fig. 3b).
128 In some cases the microtubules are observed in cross section in both poles of the merozoites, in other cases they are
129 longitudinally sectioned and organized to form a flagellum which is located at the apical pole or sometimes crossing the
130 whole merozoite from one pole to the other. In the following stage the schizont appears as an electron dense vesicle
131 containing only few mitochondria and vacuoles, whereas the merozoites, measuring about 2.1-2.2 μ m in diameter,
132 invade the cytoplasm of the host cell (Fig. 3c). Subsequently the schizont seems to degenerate and the merozoites
133 escape from the host cell and are observed free into the lumen of the blood vessel. The merozoites free in the blood
134 measure from about 1.8x2.5 to 3.5x3.7 μ m and show numerous microtubules and a clear flagellum-like structure. The
135 medium length of these merozoites is of about 4 μ m (Fig. 3d, e, f).
136 Nine DNA extracts from brain tissues were positive to the PCR and specifically hybridised with the *T. taurotragi*
137 oligonucleotide on the RLB membrane (Fig. 4, samples 10 to 14, 34 and 37 to 39). The remaining five samples were
138 negative (samples 9,15,16,35,36).

139

140 **Discussion**

141 Despite the limited information available, the clinical signs reported by Maasai herders in the cattle of the present study
142 are similar to those described in the literature in cases of BCT (Van Amstel 1982; Lawrence et al. 2004). Contrary to
143 what has generally been observed in BCT, several cases of the present study (n=13 animals) were older than 5 years,
144 reaching 8 and 9 years of age in 6 cases.

145 Macroscopical and histological lesions resemble those reported by other authors in BCT caused by *Theileria* species
146 (Giles et al.1978; Bader et al. 1986; Lawrence et al. 2004), and particularly the accumulation of parasitized lymphoid
147 cells in the cerebral and meningeal blood vessels represents the pathognomonic feature of these neurological cases as
148 reported in the cerebral form of theileriosis (Giles et al.1978; Bader et al. 1986; Lawrence et al. 2004).

149 Ultrastructurally, most of the knowledge on developmental stages in the tick and mammalian host cells of theileriae
150 species comes from studies on *T. parva* and, to a lesser extent, *T. annulata*. In the authors' opinion, the morphology of
151 developmental stages of *T. taurotragi* in lymphoid cells is still unknown. Shein and colleagues accurately described by
152 means of electron microscopy, the morphology of the four most important *Theileriae* spp. affecting bovine species (*T.*

153 *parva*, *T. annulata*, *T. mutans*, and *T. lawrencei*) during the schizogony (Schein et al. 1978). These authors evaluated
154 the development of a young schizont, spherical in shape with diameters of about 2µm converting into a multinucleate
155 mature schizont (20-50 nuclei) that produces merozoites. The latter, pear-shaped, show a length of about 1 µm and a
156 diameter of 0.6 µm; they show an apical pole where is present a polar ring system at which microtubules are anchored,
157 3-4 rhoptries, some micronemes and numerous mitochondria. This development destroys the host cell, the merozoites
158 become free and in a position to infect the next host cell, the erythrocyte. Inside the erythrocyte the piroplasms are
159 spherical and variable in size; they may measure from 1.0 to 1.5 µm in *T. parva*, they can reach up to 2.5 µm in *T.*
160 *mutans* and *T. annulata*, and are even larger in *T. taurotragi* and *T. orientalis* (Schein et al. 1978). When infected
161 erythrocytes are ingested by the tick, lysis of the erythrocytes occurs and many free piroplasms can be seen in gut
162 smears of the ticks. A variable proportion of piroplasms undergo development into sexual stages. In the tick gut, ray
163 bodies are formed by development of ovoid or spherical intra-erythrocytic stages (Mehlhorn and Schein 1984). The ray
164 bodies are spindle-shaped, measuring 8-12 µm in length with a diameter of 0.8 µm. These parasites, in cross section,
165 could show up to four nuclei and up to four flagella-like protrusions with up to 14 tubules within such protrusions.
166 The samples described in the present paper show mature schizonts in the host cells comparable to those described in the
167 literature in the *Theileria* species cycle. However, they measure 7-9 µm in diameter compared to 2-10 µm reported by
168 Schein et al. (1978). The merozoites observed in the schizont are relatively bigger (about 1.6µm in diameter) compared
169 to the same described by Schein and coll (Schein et al. 1978) (1 x 0.6 µm). The dimensions of the merozoites free in the
170 cytoplasm of the lymphoid cells and in the blood vessels are not reported in the literature and are thus not comparable.
171 The final structure of the merozoites observed in the present paper is very similar to that described in the literature.
172 They are pear-shaped and limited by a cell membrane. At the apical pole they show two inner membranes which form a
173 polar ring system at which subpellicular microtubules are anchored. Several rhoptries and few micronemes and
174 vacuoles are present. Particularly one cytoplasmatic vacuole is always present containing a spherical structure strongly
175 electrondense (with a diameter of about 0.40µm). Moreover, it is interesting to point out that the merozoites showed in
176 the schizonts or free in the vessels, always show an evident flagellum-like protrusion so far not described in the
177 literature in these phase of the life cycle. This flagellum seems to start from the pole opposite to the apical one;
178 sometimes the microtubules forming the flagellum seem to cross the whole merozoite. Some merozoites show clear
179 microtubules in cross sections in both the poles. A comparable morphology, characterized by the presence of numerous
180 microtubules and flagella-like structures was reported in details only during the sexual stages within the tick gut (Young
181 et al. 1980; Norval et al. 1992), but nobody described it in the final host such as the lymphoid cells of the cattle. These
182 flagellated protozoa were observed in all samples.

183 Out of 14 tissue samples tested by molecular assays, nine samples were positive and *T. taurotragi* was detected as the
184 unique infectious agent by RLB. Our samples belong to a larger sample of 78 tissues (brain, lymph nodes, spleen) from
185 suspected Ormilo cases from Arusha region tested at the Division of Parasitology and Tropical Veterinary Medicine,
186 Utrecht University, by PCR/RLB. Overall, 59% of the processed samples hybridized exclusively with the *T. taurotragi*
187 oligonucleotide. None hybridized with the *T. parva* oligonucleotide, thereby convincingly excluding *T. parva* as causal
188 agent of the cerebral theileriosis in the study area (Lynen et al., unpublished data).

189 The neurological signs observed and the neuropathological lesions characteristic of cerebral theileriosis, led the authors
190 to speculate that *Theileria* spp. could be considered the possible etiologic agent in these “Ormilo” disease cases.

191 Based on the electron microscopy observations, the morphology of the schizonts and the merozoites observed during
192 schizogony is partially different from the morphology of the *Theileria* species described in the literature. Moreover the
193 absence of intra-erythrocytic parasites, not even when numerous merozoites are demonstrated free in the blood, leads the
194 authors to assume that the whole cycle of this parasite is different from the one of the *Theileria* species reported in the
195 literature (*T. parva*, *T. annulata*, *T. mutans* and *T. lawrencei*) (Schein et al., 1978). On the basis of the PCR/RLB results
196 *T. taurotragi* could be hypothesized the etiological agent of "Ormilo" cases.

197 It is also possible that the observed parasites belong to a different strain of *Theileria* genus endemic to this region till
198 now not investigated, with a development cycle different from what is supposed for other species characterized by a
199 distinct morphology. In this case, it is to be assumed that additional factors could determine the simultaneous
200 involvement of the usually benign *T. taurotragi* (Martin and Brocklesby 1960; Young et al. 1977; Grootenhuis 1979;
201 Uilenberg et al. 1982) in the cerebral theileriosis pathogenesis, such as the impairment of the immunity by viruses or
202 other concomitant tick-borne infections or heavy tick infestations. Simultaneous occurrence of different tick-borne
203 pathogens within the same host is in fact a frequent feature, but little is known about their possible interactions. Other
204 contributing factors considered for BCT are parasitoses like trypanosomiasis in Kenya (Moll et al. 1985), or
205 *Amblyomma variegatum* tick infestation (Lloyd and Walker 1993).

206 As the role of the observed parasite in the “Ormilo” syndrome is at the moment unknown, further studies are needed to
207 better understand its morphology and its potential pathogenic importance.

208

209

210

211

212

213

214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244

Acknowledgements

The authors would like to thank the Maasai herders in Endulen district, who kindly collaborated to “Ormilo” cases identification and Federico Frosini for collaborating in sample collection. Research activities were funded by DFID - UK Department for International Development, Valle d'Aosta Region-development cooperation office, through the Italian NGO "AVEC PVS", Aosta (Italy) and University of Torino.

They wish to thank Prof. F. Jongejan and his research group for the support provided on PCR/RLB testing, which was carried out at the Division of Parasitology and Tropical Veterinary Medicine, Faculty of Veterinary Medicine, Utrecht University. Many thanks to Prof. B. Summers for the constructive discussion of the text.

Conflict of interest statement

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277

References

- Bader, R., Moll G., Lohding A., 1986. Morphological findings in bovine cerebral theileriosis (BCT). *Journal of Veterinary Medicine Series A*, 33, 266-285.
- Dabak, M., Dabak, D.O., Aktas, M., 2004. Cerebral theileriosis in a Holstein calf. *Veterinary Record*, 154,533-534.
- De Vos, A.J., Bessenger, R., Banting, L.F., 1981. *Theileria ? taurotragi*: a probable agent of bovine cerebral theileriosis. *Onderstepoort Journal of Veterinary Research*, 48,177-178.
- Dobbelaere, D.A.E. and McKeever, D.J., 2002. *Theileria*. In: Black SJ, Seed JR (Eds.) *World Class Parasites.*, Kluwer Acad Publishers, Boston.
- Field, C.R., Moll, G., OleSonkoi, C., 1988. Ngorongoro conservation and development project. *Livestock Development Technical Report No 1*.
- Flanagan HO, Le Roux JMW. Bovine cerebral theileriosis - A report on two cases occurring in the Union. *Onderstepoort Journal of Veterinary Research*, 27,453-461,1957.
- Giles, N., Davies, F.G., Duffus, W.P.H., Heinonen, R., 1978. Bovine cerebral theileriosis. *Veterinary Record*, 102,313.
- Grootenhuis, J.G., 1979. Characteristics of *Theileria* species (eland) infections in eland and cattle. *Research in Veterinary Science*, 27,59-68.
- Lawrence, J.A., de Vos, A.J., Irvin, A.D., 2004. Turning sickness. In: Coetzer JAW, Thomson GR, Tustin RC (Eds.) *Infectious diseases of livestock*. Oxford University Press, Oxford. 331-333.
- Lloyd, C.M., Walker, A.R., 1993. The effect of inflammatory and hypersensitivity reactions, in response to the feeding of the tick *Amblyomma variegatum*, on the progression of the experimental dermatophilosis infections. *Experimental and Applied Acarology*, 17,345-356.
- Martin, H., Brocklesby, D.W., 1960. A new parasite of eland. *Veterinary Record*, 72,331-332.
- Mehlhorn, H. and Schein, E., 1984. The Piroplasms: life cycle and sexual stages. In: Dawes B (Ed.) *Advances in parasitology*, vol 3, Academic Press, London. 38-97.
- Mettam, R.W.M. and Carmichael, J., 1936. Turning sickness : A protozoan encephalitis of cattle in Uganda. Its relationship with East Coast fever. *The Journal Parasitology Research*, 28,254-283.
- Moll, G., Agan, L., Lohding, A., 1985. Bovine cerebral theileriosis in pure Boran and sahiwalcross cattle immunized against East Coast fever and kept under continuous field challenge. In: Irvin AD (Ed.) *Immunization against theileriosis in Africa*. International Laboratory for Research on Animal Diseases, Nairobi.
- Nijhof, A.M., Penzhorn, B.L., Lynen, G., Moll, J.O., Morkel, P., Bekker, C.P., Jongejan, F., 2003. *Babesia bicornis* sp. nov. and *Theileria bicornis* sp. nov. Tick-Borne Parasites Associated with Mortality in the Black Rhinoceros (*Diceros bicornis*). *Journal of Clinical Microbiology*, 41,2249-2254.
- Norval, R.A.I., Perry, B.D., Young, A.S., 1992. The epidemiology of Theileriosis in Africa. *Academic Press London*. 130-154.

278 Nsengwa, G.R.M., 1993. An epizootic of theileriosis in cattle characterized by a nervous syndrome, in Ngorongoro
279 District, Tanzania. Tanzania Veterinary Journal, 13, 41-44.

280 Saville, W.J.A., 2002. Cerebral theileriasis. In: Bradford PS (Ed.) Large Animal Internal Medicine, 3rd ed. Mosby, St
281 Louis, Missouri. 917-918.

282 Schein, E., Mehlhorn, H., Warnecke, M., 1978. Electron microscopic studies on the schizogony of four Theileria species
283 of cattle (*T. parva*, *T. lawrencei*, *T. annulata* and *T. mutans*). Protistologica, 24,337-348.

284 Sudan, V., Sharma, R.L., Yadav, R., Borah, M.K., 2012. Turning sickness in a cross bred cow naturally infected with
285 *Theileria annulata*. Journal of parasitic diseases, 36,226-229.

286 Uilenberg, G., Perié, N.M., Lawrence, J.A., de Vos, A.J., Paling, R.W., Spanjer, A.A., 1982. Causal agents of bovine
287 theileriosis in southern Africa. Tropical Animal Health and Production, 14,127-140.

288 Van Amstel, S.R., 1982. Bovine cerebral theileriosis, some aspects on its clinical diagnosis. Proceedings 12th World
289 Congress of Diseases in Cattle. Vol II. Amsterdam, The Netherlands, September, 981-985.

290 Young, A.S., Grootenhuis, J.G., Kimber, C.D., Kanhai, G.K., Stagg, D.A., 1977. Isolation of a *Theileria* species from
291 eland (*Taurotragus oryx*) infective for cattle. Tropical Medicine and Parasitology, 28,184-194.

292 Young, A.S., Grootenhuis, J.G., Leitch, B.L., Schein, E., 1980. The development of *Theileria* = *Cytauxzoon taurotragi*
293 (*Martin and Brocklesby*, 1960) from eland in its tick vector *Rhipicephalus appendiculatus*. Parasitology, 81,129-
294 144.

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310 **Figure captions**

311 **Fig. 1** Generalized representation of the typical life cycle of the genus *Theileria*. Sporozoites are inoculated into a
312 mammalian host when an infected tick takes a meal. The sporozoites invade lymphoid cells, and develop into a
313 multinucleate syncytial schizont or macroschizont (**a**). At the same time, the parasite induces host cell transformation
314 and proliferation (**b**). Then the schizonts differentiate into merozoites and invade erythrocytes (**c**). Ticks become
315 infected by ingesting infected erythrocytes, and gametogenesis and fertilization takes place in the gut lumen (**d**). The
316 resulting zygote invades a gut epithelial cell where it remains during the tick moult cycle and develops into a single
317 motile kinete (**e**). The motile kinete egresses the gut cell and invades the salivary glands (**f**). Tick initiates rapid
318 sporozoite development in the salivary glands, and infective sporozoites that survive in the gut epithelium are
319 transmitted to another mammalian host when the resulting post-moult nymphs or adults feed (**g**) Modified from
320 <http://www.theileria.org/ahdw/background.htm> [accessed 27-02-2015]

321
322 **Fig. 2 a** Multifocal congestion of the meningeal blood vessels covering cerebral hemispheres and cerebellum.
323 Hemorrhages surrounding the brainstem and the cervical spinal cord. **b** Severe hemorrhages involving the
324 leptomeninges, the IV ventricle, the central canal and the surrounding areas. **c** Punctiform hemorrhages in the cortical
325 white matter. **d** Focus of chronic malacia in the capsula interna area. **e** Cerebellum: hemorrhages involving the
326 leptomeninges and severe congestion of the meningeal vessels. Multifocal hemorrhages and plasmorrhages in the
327 nervous tissue around the IV ventricle (H&E). **f** Severe accumulation of mononuclear cells in the meningeal blood
328 vessels many of them showing irregular round to ovoidal bodies in the cytoplasm corresponding to parasitic schizonts
329 (H&E). (e bar=100 µm, f bar=50 µm)

330
331 **Fig. 3** Electron microscopy of bovine lymphoid cells with endocellular parasites (**a – f**). **a** Lymphoid cells containing a
332 young schizont (arrow) spherical in shape, electrondense, characterized by spherical dividing nuclei, numerous
333 mitochondria and many spherical organelles such as vacuoles. **b** Lymphoid cells with the cytoplasm occupied by a
334 voluminous schizont (arrow) with two evident nuclei located at the periphery (#), numerous vacuoles and some
335 spherical or pear-shaped merozoites (asterisks). **c** In this following stage the schizont (arrow) appears as an electron
336 dense vesicle containing only some mitochondria—and few vacuoles whereas the merozoites (asterisks) invade the
337 cytoplasm of the host cell. **d** Merozoites free into the lumen of the blood vessel. **e, f** High magnification of the
338 merozoites free into the vessels. They measure from about 1.8x2.5 to 3.5x3.7µm. At the apical pole the merozoites
339 show a polar ring system at which subpellicular microtubules are anchored to form a clear flagellum-like structure

340 (asterisk). Few rhoptries (long arrows) and micronemes (short arrows) and vacuoles are present. Scale bar 1µm
341 (a,b,c,d); 0.2µm (e,f)

342

343 **Fig. 4** Reverse Line Blot results showing species-specific oligonucleotides of the 18S rRNA gene of *Theileria* spp. in
344 the horizontal lanes and PCR products in the vertical lanes. Fourteen brain tissue samples from Ormilo cases from
345 Endulen were analyzed (vertical lanes: 9 to 16 and 34 to 39). Positive *Theileria* spp. and *Babesia* spp. controls (line 1:
346 *T. parva*, 2: *T. taurotragi*, 3: *T. buffeli*, 4: *T. mutans*, 5: *B. bovis*, 6: *B. bigemina*, 7: *T. annulata*) as well as a negative
347 control (water, line 8) are included. Other vertical lanes (17 to 33 and 40) refer to amplified DNA of tissue samples
348 collected by the authors in other study areas in Arusha region.

349

350