AperTO - Archivio Istituzionale Open Access dell'Università di Torino

VvPIP2;4N aquaporin involvement in controlling leaf hydraulic capacitance and resistance in grapevine

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1558379> since 2021-12-29T14:53:38Z

Published version:

DOI:10.1111/ppl.12463

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **VvPIP2;4N aquaporin involvement in controlling leaf hydraulic capacitance and resistance in**
2 **grapevine**

3 Marco Vitali^{1*}, Hervé Cochard^{2,3}, Giorgio Gambino⁴, Alexandre Ponomarenko², Irene Perrone⁴,
4 Claudio Lovisolo^{1,4}.

5 ¹ Department of Agricultural, Forest and Food Sciences (DISAFA), University of Turin, Largo
6 Paolo Braccini 2, 10095 Grugliasco, TO, Italy

7 ² INRA, UMR 547 PIAF, F-63100 Clermont-Ferrand, France;

8 ³ University Blaise Pascal, UMR 547 PIAF, F-63177 Aubière, France

9 ⁴ Institute for Sustainable Plant Protection, National Research Council (IPSP-CNR), Grugliasco
10 unit, Largo Paolo Braccini, 10095 Grugliasco, TO, Italy

11

12 *Corresponding author:*

13 *Marco Vitali

14 Department of Agricultural, Forest and Food Sciences (DISAFA), University of Turin, Largo Paolo
15 Braccini 2, 10095 Grugliasco, TO, Italy

16 Tel: +39.011.6708885

17 Fax: +39.011.6708658

18 e-mail: marco.vitali@unito.it

19

20 **e-mail addresses:**

21 Marco Vitali: marco.vitali@unito.it

22 Hervé Cochard: herve.cochard@clermont.inra.fr

23 Giorgio Gambino: giorgio.gambino@ipspp.cnr.it

24 Alexandre Ponomarenko: alexpono@gmail.com

25 Irene Perrone: irene.perrone@ipspp.cnr.it

26 Claudio Lovisolo: claudio.lovisolo@unito.it

Abstract

Hydraulic capacitance in a plant tissue (C), buffers the xylem tension storing and releasing water and has been highlighted in recent years as an important factor that affects water relations such as drought tolerance and embolism formation. Aquaporins are well known to control leaf hydraulic resistance (Rh) but their role in the control of C is unknown. Here, we assess Rh and C on detached grapevines leaves (cv. Brachetto) wild type (WT) and over-expressing the aquaporin gene *VvPIP2;4N* (OE). For this purpose, we developed a new method inspired from the pressure-volume curve technique and the rehydration kinetic method, which allowed us to monitor the dynamics of dehydration and rehydration in the same leaf. The recovery after dehydration was measured in the dark, in light non-transpirative conditions, light-transpirative conditions and transpirative condition adding abscisic acid.

Pressurizing to dehydrate leaves in the OE line, the recorded Rh and C were respectively lower and higher than those in the WT. The same results were obtained in the dark recovery by rehydration treatment. In the presence of light, either when leaves transpired or not (by depressing vapour pressure deficit), the described effects disappeared. The change in Rh and C did not affect the kinetics of desiccation of detached leaves in the dark in air, in OE plants compared to WT ones. Our study highlighted that both Rh and C were influenced by the constitutive over-expression of *VvPIP2;4N*. The effect of aquaporins on C is reported here for the first time and may involve a modulation of cell reflexion coefficient.

Key words: *Vitis vinifera* (L.), transgenic plant, leaf water potential, pressure-volume curve, isohydric, anisohydric, osmoregulation.

Abbreviations:

AQPs, aquaporins; OE, over-expressing; WT, wild type; C, hydraulic capacitance; Rh, hydraulic resistance; RWC, relative water content; Ψ , leaf water potential; **+0.5**, dehydration treatment with pressure applied of 0,5 MPa; **+1**, dehydration treatment with pressure applied of 1 MPa; **dark**, rehydration treatment in dark condition; **light VPD $\approx$ 0**, rehydration treatment in light condition without transpiration; **light transp**, rehydration treatment in light and transpirative condition; **light transp ABA**, rehydration treatment in light transpirative condition supplying ABA in the rehydrating water.

60 **Introduction**

61 Water is the most limiting resource for plant life and yield (Lange et al. 1982). Although the
62 majority of a plant's fresh weight consists of water, the amount of water retained by the plant in the
63 biomass is less than one percent of the total water transpired via stomata. Consequently, a huge
64 quantity of water is required to enable photosynthesis and plant growth. Therefore, water uptake
65 from the soil, its transport, storage and usage are mediated through a system that has evolved to
66 fully exploit the chemical and physical properties of water.

67 The cohesion-tension theory, formulated by Dixon (1914), explains the transport of water in the
68 soil-plant-atmosphere continuum. In this system, water moves from high to low water potential (Ψ),
69 and thus, towards transpiring leaves. Transpiration itself drives the rise of the xylem sap and
70 submits water to a considerable tension. Such tension is balanced by the hydrogen bonds among
71 water molecules, which prevent the breaking of the water column. However, under different
72 conditions (e.g., water shortage, freezing, high evaporative demand), this tension can increase and
73 cavitation can occur by air seeding mechanisms (Angeles et al. 2004).

74 In Dixon's theory, water transport in plants behaves like an electrical circuit, and follows Ohm's
75 law as described by van den Honert (1948), hence, the flow is due to the water potential gradient
76 and is hindered by the hydraulic resistance (R_h). Moreover, the pathway can be split into the water
77 transport of individual organs (root, stem, leaf), each with its own R_h that affects the water flux
78 (Tyree and Ewers 1991; Sperry et al. 1998). However, in this equation, the hydraulic capacitance
79 (C) has to be considered as an important variable that affects the output. Analogously to an
80 electrical circuit, C has the function of a capacitor (or condenser) used to store charge temporarily,
81 thus buffering a power surge. Therefore, in plants, C represents the ability to store water and to
82 buffer the system reducing the degree of tension in the xylem in transient water status conditions.
83 Capacitance corresponds to the ratio between the change in water content and the change in water
84 potential ($C = \Delta RWC / \Delta \Psi$; Tyree and Ewers 1991; Sperry et al. 2008); its effect is to make the
85 amount of water entering a region different from the amount of water leaving it, whenever $\Delta \Psi$
86 changes.

87 Leaves are the final component of the water transport system and via their stomata, they balance
88 carbon nutrition and water loss by transpiration, thereby playing a key role in the regulation of the
89 water status and the strategy of responses to drought stress. To prevent deleterious dehydration,
90 stomatal conductance is controlled by a complex regulation of guard cells, involving chemical and
91 hydraulic signals (Comstock 2002). The relevance of leaf C was recently investigated in relation to
92 various physiological traits, such as leaf thickness (Sack et al. 2003), leaf water content per unit dry
93 weight, leaf mass per unit area and lignin content (Blackman and Brodribb 2011). In addition, the

latter study described the ‘dynamic C’ (computed as the volume of flowing water measured by a flowmeter) to be highly coordinated with leaf hydraulic conductance (Blackman and Brodribb 2011). Aquaporins (AQPs) exercise a strategic function in the leaf water pathway by controlling symplastic water movements (Kaldenhoff et al. 2008), and being the main link between the symplastic and apoplastic pathways, e.g., bundle sheath cells. These water channels, without changing the flux direction, can enormously increase the water movement across membranes, and therefore, decrease the Rh. Aquaporins can be modulated at several levels, via transcription, translation, trafficking and gating (opening and closing of the pore) and by environmental and developmental factors (Chaumont and Tyerman 2014), such as: irradiation (Prado et al. 2013; Lopez et al. 2013), transpiration (Sakurai-Ishikawa et al. 2011; Laur and Hacke 2013), circadian rhythms (Hachez et al. 2008), abscisic acid (ABA) feeding (Shatil-Cohen et al. 2011; Pantin et al. 2013), auxin feeding (Péret et al. 2012) and shoot wounding (Sakurai-Ishikawa et al. 2011; Vandeleur et al. 2014). Several experiments using transgenic plants overexpressing or silencing AQP genes have been performed (reviewed by Martínez-Ballesta and Carvajal 2014) and have demonstrated that the transcriptional modulation of AQPs generally modifies the Rh, however, to date, no results exist concerning the effects of AQPs on C. In this study, grapevine plants over-expressing *VvPIP2;4N* (an aquaporin previously described by Perrone et al. 2012, extremely efficient in facilitating cell-to-cell water pathways), were used to assess the role of this AQP isoform on leaf Rh and C during leaf dehydration and recovery. The hydraulic parameters were evaluated by a new method derived from the pressure-volume curve (Tyree and Hammel 1972) and the rehydration kinetic technique explained by Blackman and Brodribb (2011).

Materials and Methods

Plant material

The experiments were performed on leaves of potted ‘Brachetto’ grapevines; 10 wild-type (WT) and 10 transgenic plants from line 16, which overexpressed *VvPIP2;4N* (OE), previously described by Perrone et al. (2012). The 4-year-old plants (two buds pruned with bud-break in March, non-grafted) were grown in a greenhouse on a mixture of peat–loam, under natural light and CO₂ concentration conditions. Plants were irrigated regularly according to their needs. In this experiment, fully expanded, mature leaves were used.

127 *Assessment of Rh and C in the dehydration and rehydration processes*

128 To assess leaf Rh and C during dehydration, a method similar to the pressure-volume curve
129 technique was used (PV curve, Tyree and Hammel 1972), whereas for the rehydration phase, a
130 modified rehydration kinetic method (see C_{dyn} measurements, Blackman and Brodribb 2011) was
131 applied.

132 The new method proposed required the use of a high-precision balance (Mettler Toledo AT261
133 deltarange; Greifensee, CH) and a modified Scholander pressure bomb, inverted over the balance,
134 and controlled by an external manometer (Bourdon, FR; class 0.1). The cut surface of the petiole,
135 which passed through the sealing system of the pressure chamber, was immersed in a cylinder (50
136 mL), filled with deionised water, placed on the balance plate (Fig. 1). The balance plate was
137 isolated from the laboratory atmosphere and the relative humidity inside the balance chamber was
138 kept close to 100% using wet paper. By applying and releasing the pressure in the chamber, the
139 flow out/in of the leaf was measured by an increase or decrease in weight measured by the balance,
140 as explained below.

141

142 Dehydration phase (Fig. 1a):

143 Leaves were collected at 18.00. The petiole extremity, cut under water, was submerged in deionised
144 water in non-transpirative conditions (dark, sealed bag) overnight, to allow full leaf hydration.
145 During the following day (at 9.00; 12.00; 15.00), the leaves were removed from water and were
146 immediately placed in the pressure chamber. After measurement of the native water potential
147 (Ψ_{leaf}), the pressure chamber was upturned on the balance, placing the petiole in the 50-mL
148 cylinder. Starting from this steady state, the pressure was increased to a value of +0.5 (noted **+0.5**
149 thereafter) and +1.0 MPa (**+1**) and was kept constant; the pressure rose by 0.05 MPa per second
150 regulated by a needle valve. The mean native Ψ_{leaf} , after one night hydration in water, was -0.01
151 MPa for both WT and OE leaves.

152 This kind of measurement was also possible with water-stressed leaves, taking care to maintain the
153 pressure inside the chamber slightly lower than that balancing the leaf water potential. This was
154 necessary to avoid water uptake by the leaf and thus, to maintain a stable weight on the balance to
155 begin measurements.

156

157 Rehydration phase (Fig. 1b):

158 The pressure was applied to the dehydrated leaves until $\Psi_{\text{leaf}} = -1$ MPa (dehydration, **+1**). After
159 reaching this level of dehydration, the pressure was released and the chamber was removed. The
160 petioles remained suspended by the lid of the pressure chamber and immersed in the 50-mL

cylinder. The water uptake by the leaves began immediately after depressurisation and was monitored for 1 h in different conditions:

- 1) dark and non-transpirative condition (**Dark**, leaf in a dark glass bell);
- 2) light and low transpirative condition (**Light VPD $\approx$ 0**, a 2L glass baker was placed over the leaf with artificial light set at 500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$, wet paper was previously used to reduce the VPD to 0 with formation of condensed water on the glass surface);
- 3) light and transpirative condition (**Light transp**, artificial light set at 500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$, leaf in laboratory atmosphere);
- 4) light, transpirative condition and ABA (**Light transp ABA**, conditions as in 3), and ABA solution (100 μmol final concentration) supplied in the cylinder.

In the transpiring treatments, the air temperature was between 19°C and 24°C and the relative humidity was between 45 and 60% (max VPD 10 Pa/KPa).

The Rh and capacitance C were obtained from the equation describing the pressure-volume curves and was computed using the formula:

$$f(x) = P * C * 1^{(-x/(Rh*C))} \quad (\text{Tyree and Hammel 1972})$$

where: x is the cumulative water in/out of the leaf and P is the pressure applied. Using this equation Rh correspond to the slope of the first part of the curve and C to the *plateau* phase, as highlighted in Fig. 2.

To avoid errors in the calculation due to the transpiration, in the rehydration phase, C was computed subtracting mathematically the transpiration rate from the water uptake weight measurements above 600 seconds. Until this time, transpiration marginally affected the water uptake (see Fig. 4). Data were normalised to the dry weight (70°C, 12 h) and the leaf area (measured by the area-meter Li3000, Lincoln NE, USA) of the single leaves. SigmaPlot 12.3 software (Systat Software, San Jose CA, USA) was used for data elaboration and statistical analysis by *t*-test and one-way ANOVA (after passed Shapiro-Wilk test). To perform ANOVA analysis, data were normalised when the homogeneity of variance test (Bartlett's test) failed.

Aquaporin expression profile

Leaves for the aquaporin expression analysis were collected in light and dark conditions following the same time-course and sampling protocol as for leaves used in the physiological tests (collected at 18.00 and left rehydrated overnight in a dark, sealed bag). Leaves in the dark treatment were harvested in liquid nitrogen at 9.00, whereas leaves in the light treatment were submitted to artificial irradiation for 1 h before harvesting. The real-time RT-PCR (qRT-PCR) quantification of transgenic *VvPIP2;4N*, endogenous *PIP2* genes and *PIP1*-type aquaporins were carried out as

195 previously reported (Perrone et al., 2012) on two biological replicates (three technical replicates
196 each).

197

198 *Leaf dehydration dynamics in dark conditions*

199 For each line, 22 leaves were sampled at 18.00; petioles were cut under water to avoid embolism
200 formation. Leaves were left free to rehydrate through the petiole in deionised water overnight, as
201 above. The following day at 8.00, water was removed and leaves were left dehydrated on the bench
202 in the dark in the laboratory atmosphere. The fresh weight of fully hydrated leaves was measured
203 with a balance (Denver Instruments Company TR603D; Arvada CO, USA), and then for each line,
204 the weight and Ψ_{leaf} were measured every hour during dehydration. The dry weight was recorded
205 after drying the leaves at 70°C for 12 h as above, and was used to calculate the relative water
206 content (RWC). The C was newly computed according to Koide et al. (2000), as:

$$C = \frac{\text{RWC}/\Psi_{\text{leaf}}}{\text{dry weight}}$$

207

208 **Results**

209 *Dehydration phase*

210 During dehydration, water was forced to exit through the petioles, and a higher flow out through the
211 petiole of the OE leaves was observed in comparison to the WT, as shown in Fig. 2, when leaves
212 were pressurised to +0.5 MPa (+0.5). These differences were also observed if the data were
213 normalised by the leaf dry weight or leaf area (Fig. 2).

214 By collecting the data during three different daily time-points (Fig. S1); morning, noon and
215 afternoon, we observed that for both lines, the amount of water that exited from the petioles
216 followed an increasing trend and reached a maximum at noon and a minimum in the morning.
217 The mean Rh and C, obtained by pressurising leaves (dehydration phase), are shown in Fig. 3 (and
218 Fig. S3). When leaves were pressurised to +0.5 MPa (+0.5), Rh was very low in both lines with no
219 significant differences, whereas at +1 MPa (+1), the Rh increased drastically and a significant
220 difference was observed in the OE line ($P < 0.01$), where the Rh was lower than in WT (Fig. 3a).
221 Conversely, C was significantly higher (+38%, $P < 0.05$) in the OE line in the first treatment (+0.5),
222 but this difference disappeared when the pressure was increased to +1 MPa (+1) (Fig. 3b).

223

224 *Rehydration phase*

225 Subsequent to dehydration to $\Psi_{\text{leaf}} = -1.0$ MPa (+1 treatment), leaves were left to rehydrate for 1 h
226 (rehydration phase) by subjecting them to different stimuli. The total time course of the amount of
227 water flowing into the petiole after the pressure release is shown in Fig. 4. Moreover, the first 600

seconds of the experiment, which was used to compute Rh and C, are highlighted in Fig. 4 frame **b**. These figures illustrate that for all lines and conditions, the recovery from stress occurred via a slow rise in the volume of water absorbed, and that transpiration began at about 600 s following depressurisation, as suggested by the divergence among treatments with (**light transp**, **light transp ABA**) or without transpiration (**dark**, **light VPD≈0**). Finally, although the standard errors overlapped for all treatments, the leaves of the WT line in dark conditions appeared to behave differently from those in the other treatments, which uptook more water.

The recovery behaviour following dehydration can be analysed by Rh, C (calculated from the dynamics shown in Fig. 5) and the Ψ_{leaf} reached after 1 h of rehydration (Table 1). Major differences in Rh between the two lines were observed in dark conditions (Fig. 5a), where in WT leaves, the Rh was significantly higher than in OE leaves ($P < 0.01$). This result agrees with the difference observed in the +1 treatment (where dark conditions were ensured by the pressure chamber) between WT and OE, although at a higher order of magnitude. The switch from **dark** to **light VPD≈0** conditions decreased the Rh in WT to levels similar to those in the OE line, whereas Rh was not affected by the transition between dark and light in OE leaves. Transpiration (**‘light transp’** treatment) did not have any effect on the Rh in WT lines, whereas a slight increase was observed in the transgenic line. Finally, after the addition of ABA to the solution absorbed through the petiole (**light transp ABA**), the Rh in WT leaves increased, but not significantly, compared to other rehydration conditions in the light and from the transgenic line. However, in general, we observed a reduction in Rh in WT following the transfer from the dark to the light conditions adopted in the rehydration experiments, whereas the Rh in the OE line tended to increase with increased transpiration.

The C computed from the same dataset did not differ between WT and transgenic leaves; only the dark condition strongly affected this parameter, causing it to be significantly lower ($P < 0.001$) in the WT than in all other treatments.

The Ψ_{leaf} recorded in the pressurisation experiment and its recovery are reported in Table 1. During dehydration, there were no differences in the native Ψ_{leaf} and consequently in the final Ψ_{leaf} reached. However, in contrast, during rehydration, the leaves of the two lines revealed a different ability to recover the Ψ_{leaf} within 1 h after de-pressurisation. In particular, both lines in dark conditions showed a higher recovery rate, reaching a Ψ_{leaf} close to 0. On the contrary, Ψ_{leaf} decreased when leaves were subjected to artificial light and transpiration, whereas ABA treatment facilitated the recovery of Ψ_{leaf} . The statistical analysis showed differences in the Ψ_{leaf} between the two lines in the **‘light VPD≈0’** and **‘light transp’** treatments.

Based on these observations, the expression levels of transgenic *VvPIP2;4N*, together with those of other known *PIP2* genes and a PIP1-type aquaporin were quantified by qRT-PCR in dark and light-transpiration conditions in both lines. The WT showed the same AQP expression profile in dark and light-transpiration conditions, suggesting a light-independent expression of these *PIP* genes (Fig. 6). Furthermore, in the OE line, the expression profile of AQPs and transgenic *VvPIP2;4N* was generally not affected by light; only *VvPIP2;2* was slightly more highly expressed in the dark.

Leaf dehydration dynamics in dark conditions

The dynamics of the dehydration of detached leaves in darkness was observed from the relationship between Ψ_{leaf} and relative water content (RWC). In addition, C was calculated as $\Delta\text{RWC}/\Delta\Psi \cdot \text{dry weight}$. The linear regression indicated a slightly higher RWC coupled to the decrease in Ψ_{leaf} in OE compared to WT leaves (Fig. 7a). The hyperbole describing Ψ_{leaf} versus C was similar in both lines, showing a reduction of C that was related to the decrease in Ψ_{leaf} (Fig. 7b). Overall, the mean values of C for the two lines confirmed the higher C in OE lines (156 ± 26 for the WT, 261 ± 52 for OE; $P < 0.05$).

Finally, to evaluate whether the observed differences were attributable to anatomical or morphological traits, the pairwise relationships between leaf area, dry weight and fresh weight were assessed, without identifying any significant differences between WT and OE samples (Fig. S2).

Discussion

In this study, transgenic grapevines that constitutively over-expressed *VvPIP2;4N* under the *Cauliflower mosaic virus* 35S promoter (Perrone et al. 2012) were used to assess the role of this AQP on leaf Rh and C during leaf dehydration and recovery. Many studies in several transgenic plants had previously shown that overexpression of aquaporin genes decreased the Rh (Ding et al., 2004; Lee et al. 2012; Perrone et al. 2012), whereas the silencing of AQPs resulted in an increase in Rh (Siefritz et al. 2004, Sade et al. 2014). However, no information is available concerning the relationship between AQP and C.

The effect of aquaporins on hydraulic resistance (Rh)

As expected, the Rh was lower in OE leaves than WT leaves when a high over-pressure was applied to the leaves (+1) and when recovery was performed in **dark** conditions. These two results can be ascribed to a direct effect of transgenic *PIP2;4N*, since an increase in PIP2;4N protein in the membranes improves the membrane permeability to water.

Several studies have demonstrated that AQPs expression and activity are regulated in leaves by circadian rhythms (Siefritz et al. 2002; Nardini et al. 2005; Hachez et al. 2008). In this study, the dynamics of the cumulative water outflow from leaves showed an influence of circadian rhythms both in OE and WT leaves (Fig. S1). These differences might have affected the computation of the hydraulic traits (Fig. S3); however, to limit the impact of the biological clock, the experiments were performed at distinct times during the day and averaged together in both genotypes. AQP expression has been assessed just during the morning. However, the impact of circadian rhythms on the extremely high expression of the transgene (meanly 7 times higher than endogenous aquaporins) can be reasonably neglected. In addition, it is known that the 35S gene promoter, controlling expression of our transgene, shows low or no sensitivity to the biological clock (Millar et al. 1992; Xu and Johnson 2001).

The leaf Rh increase with increased dehydration as observed in Fig. 3, where Rh drastically increased from the +0.5 to the +1 treatment, which was reported previously (Sack & Holbrook 2006; Scoffoni et al. 2014). However, this might represent a physical artefact. One hypothesis might be that a pressure of 1 MPa leads to a massive flow of water in the leaf hydraulic system in a short time interval. Probably, the anatomy of the leaf itself (e.g., connectivity between cells, bundle-sheath permeation, petiole conductivity) hinders the runoff of a large amount of water in very short period, leading to an overestimation of the Rh. This phenomenon might explain the different magnitude of the Rh values in the dehydration +1 and recovery treatments.

In the recovery trial (rehydration, Fig. 6), the impact of various stimuli, such as i) light, ii) ABA and iii) transpiration on the aquaporin activity was studied.

i) The light effect cancelled the differences in Rh observed in the **dark** between WT and OE. Indeed, in WT, the hydraulic resistance decreased from dark to light conditions, whereas this parameter was not affected in OE leaves. This change in the Rh in WT leaves agrees with the increase in leaf conductivity under irradiation previously reported by several authors (Nardini et al. 2010, Sellin et al. 2010; Guyot et al. 2012; Lopez et al. 2013; Prado et al. 2013). Cochard et al. (2007) indicated two potential light-modulated mechanisms of water movement in leaves: activated AQPs in light conditions allows water to move freely in the symplast and apoplast, whereas at low irradiance, deactivated AQPs force the water to move apoplastically, limited by the bundle sheath. Voicu et al. (2009) highlighted that the light-dependent change in leaf hydraulic conductance in bur oak (*Quercus macro-carpa*) was not linked to any AQP transcriptional changes. Similarly, in this study, we observed only slight differences in the expression profile between WT and OE following changes in the light conditions (Fig. 6). The qRT-PCR data showed clearly that the major difference between WT and OE lines derived exclusively from the high and constitutive expression of

transgenic *VvPIP2;4N* in all conditions (Figure 6). Thus, for the OE line, the low levels of Rh in dark conditions were probably linked to the constitutive over-expression of *VvPIP2;4N*. Some type of contrasting regulation can be hypothesised between the light-mediated activation of leaf AQPs (as in WT) and *VvPIP2;4N*. In WT plants, light activates AQPs depressing the Rh recorded upon dark condition, whereas in OE plants the effect of *VvPIP2;4N* (a root-specific AQP isoform, presumably insensitive to light modulation) is masked from the light-activation of the other leaf aquaporins.

ii) ABA modulates AQP activity, having opposite effects on root and leaf AQPs: downregulating the bundle-sheath AQPs and thus limiting hydraulic conductivity in leaf (Pantin et al. 2013; Shatil-Cohen et al. 2011), and upregulating the AQP isoforms and increasing the hydraulic conductivity in the root (Jang et al. 2004; Hose et al. 2000; Thompson et al. 2007; Parent et al. 2009). In this study, no significant differences in Rh between OE and WT were observed in the presence of ABA. However, comparing the **light transp** and **light transp ABA** treatments, Rh values were twice as high in WT, even if this difference was not statistically significant, whereas no changes were observed in OE leaves. We can speculate that ABA caused a closed conformation of PIP2;4N or inhibited the expression of all AQP isoforms in WT leaves. This downregulation might also occur in OE leaves, but additionally and in contrast, ABA might promote the expression or open conformation of PIP2;4N. The PIP2;4N protein is a root-specific AQP, and therefore, is putatively upregulated by ABA; its presence might have led to a lack of increase in Rh. The first ABA signalling transduction pathway that mediates water transport in roots has been recently demonstrated in maize (Fan et al. 2015). Notably, the post-translational regulation of ZmPIP through ABA signalling appears to be particularly important to regulate root hydraulic conductivity. This might also be the case for transgenic *VvPIP2;4N* aquaporin in leaf: since it is under the control of a constitutive promoter (35s), ABA might promote its activity in a phosphorylation-dependent manner. Moreover, Chitarra et al. (2014) demonstrated that ABA promoted *VvPIP2;4N* expression in vessel-associated cells (VACs), but not in whole petiole tissue. Similarly to the VACs, the leaf bundle sheath cells regulate exchange between the xylem and other parenchyma cells. The ABA-induced upregulation of *VvPIP2;4* at this level might easily explain the lack of rise of Rh level in the transgenic leaves.

A secondary effect of ABA was observed in the recovery of the final Ψ_{leaf} after 1 h of rehydration (Table 1). In contrast to the **Light trans** treatment (final Ψ_{leaf} = -0.23 MPa for WT and -0.35 MPa for OE), ABA promoted a more rapid recovery of Ψ_{leaf} in OE leaves than WT leaves (-0.19 MPa for WT and -0.17 MPa for OE). A positive effect of ABA on Ψ_{leaf} recuperation has been already described by Lovisolo et al. (2008) and Chitarra et al. (2014) in grapevine. In the OE leaves, the

362 better recovery of Ψ_{leaf} appeared to be coupled to a low hydraulic resistance, in agreement with data
363 reported by Martre et al. (2002).

364 iii) In this study, we observed no effect of transpiration on changes in leaf Rh, contrary to that
365 reported for AQPs in root (Sakurai-Ishikawa et al. 2011; Laur and Hacke 2013). This might be due
366 to the low vapour-pressure deficit in the laboratory atmosphere or to the real absence of an effect of
367 this parameter in leaf.

368

369 *The effect of aquaporins on hydraulic capacitance (C)*

370 The C plays an important role in drought tolerance, as does Rh, by affecting the amount of water
371 destined to buffer the change in the transpiration stream elicited by the atmospheric conditions and
372 especially the ability to extend survival after stomatal closure (Bartlett et al. 2012; Gleason et al.
373 2014). Since sapwood has been recognised as the major source of stored water, several studies have
374 addressed the importance of C in this tissue in depth. Generally, C varies in sapwood between 40
375 and 900 kg m⁻³ MPa⁻¹ (Scholz et al. 2007; Čermák et al. 2007; McCulloch et al. 2014) and is
376 inversely related to the wood density. The contribution to the total daily transpiration reported in
377 literature varies between 5% and 45% (Goldstein et al. 1997; Phillips et al. 2003; Verbeeck et al.
378 2007). However, the importance of the leaf C was highlighted by Gleason et al. (2014), who
379 suggested that the majority of water lost during dehydration derives from leaves. By comparing the
380 C in grapevine leaves in this study (up to 250 mg H₂O gDW⁻¹ MPa⁻¹), classifies them within the
381 upper half of the ranking proposed by Blackman and Brodribb (2011), among species with a low C,
382 although the measured units are not the same (the conversion was performed by considering that 1
383 m² of leaf area corresponds to 31 g DW, from Fig. S2).

384 A novel aspect highlighted by this study is the effect of AQP on C, which, as far as we know, has
385 never been reported so far. The ability of a tissue to be a capacitor, and to buffer the xylem tension
386 and prevent embolism was debated as an important trait that might distinguish plants and their
387 susceptibility to water stress (Sperry et al. 2008; McCulloch et al. 2014).

388 In the pressurisation tests, differences in C were observed between WT and OE leaves only when
389 the pressure applied was low (+0.5 MPa, Fig. 3b), in contrast to what was observed for Rh (Fig. 3a).
390 The absence of effect when the applied pressure reached +1 MPa might be due to the high
391 dehydration imposed on the leaves. Indeed, C is a variable parameter that decreases together with
392 water status (Fig. 7b), potentially becoming comparable in the two lines when the leaves were
393 dehydrated to -1 MPa.

394 In the recovery trials (Fig. 5b), WT leaves showed the lowest C in dark conditions compared with
395 the other treatments; probably, the high Rh observed in WT hindered the water uptake and thus, the

396 recovery of C. In theory, a longer recovery time might lead to the complete recovery of C in WT
397 leaves, although the leaves were in dark conditions. However, in dark conditions, the final Ψ_{leaf}
398 values fully recovered in both lines; even though this occurred in WT without a complete recovery
399 in C. Thus, in this latter case, the amount of water inside the WT leaves was lower than that in OE
400 leaves, and probably, the amount of water outflow pressurizing once more the leaf could be lower
401 than that observed during the first pressurisation.

402 The aim of the last experiment was to check whether OE leaves dehydrate more rapidly in dark
403 conditions (where major differences between WT and OE leaves were observed). However, the
404 overexpression of *VvPIP2;4N* did not lead to a more rapid leaf dehydration, and the computed C
405 ($\Delta\text{RWC}/\Delta\Psi \cdot \text{dry weight}$) confirmed higher values in the OE line.

406

407 *Hypothesis and ecological significance*

408 Using transgenic plants and the novel method proposed in this study, a positive relationship between
409 C and AQPs in grapevine leaves was demonstrated. The mechanisms underlying this interaction are
410 not yet clear, however, an initial hypothetical mechanism might involve the reflection coefficient
411 (σ) of the plasma-membrane. This parameter is considered to be the ability of a channel (AQP in
412 our case) to be permeable to, or reflect a solute. It is determined by the arginine selectivity filter at
413 the end of the pore (Zeuthen et al. 2013) and by the solute size. In the membrane of transgenic
414 plants in this study, the higher concentration of AQPs can lead to a higher permeation of small
415 solutes (Gomes et al. 2009), resulting in a low reflection coefficient, and consequently allowing a
416 higher water flow (coupled to osmolyte flow) through the lipid bilayer. Although this consideration
417 depends on the contribution of *VvPIP2;4N* to the total water transport across the plasma membrane.
418 The theoretical framework for the implication of σ on C is provided as supplemental information.
419 Recently, Maurel et al. (2015) proposed a pivotal role for AQPs in buffering cell osmoregulation,
420 by reviewing and re-interpreting AQP function as osmo-sensor in guard cells during stomatal
421 movements or in growing pollen tubes. The mechanism was only speculated, but an AQP-mediated
422 increasing cell C, conferring to the cells higher buffer capacitance, could give light to this still un-
423 described phenomenon.

424 A second hypothesis, which does not exclude the first, is that AQPs might connect several cells, and
425 increase the volume of the reservoir. In grapevine, AQPs can increase the link between the symplast
426 and apoplast in the areoles, the enclosed areas between the interconnected veins, thereby improving
427 the leaf capacitance. In figure 8 a schematic representation of the leaf hydraulic pathway is
428 represented with its simplification (right side), showing the increase of the total capacitor due to the
429 sum of different capacitors.

Previously, studies have attributed a role to AQPs in iso/anisohydric behaviour, due to changes in leaf conductance (Sade et al. 2009; Vandeleur et al. 2009; Chaumont and Tyremann 2014). In other studies (Ogasa et al. 2013; McCulloh et al. 2014), plants are categorised according to their C, for their investment in structural features to maintain the transpiration stream (anisohydry) or their sensitivities to embolisms (Tombesi et al. 2014). Perrone et al. (2012) considered the ‘Brachetto’ WT as an anisohydric cultivar, and the transgenic lines could be interpreted as being even more anisohydric. In this study, AQPs conferred a greater C, and hence, a greater degree of anisohydry, highlighting the possible link between C, Rh, aquaporins and iso/anisohydric responses to water stress. Clearly, the implication of Rh and C on whole leaf hydraulics deserves further attention.

Author Contributions

H.C., C.L. and M.V. conceived and planned the study. M.V. performed the experiment and analysed the data. M.V. wrote the first draft of the manuscript. G.G. and I.P. produced plant materials, carried out the molecular analysis and reviewed the manuscript. A.P. helped in the physical dissertations. H.C. theorized the hypothesis presented and reviewed the manuscript. C.L. reviewed the manuscript and obtained funds to support the project.

Acknowledgements

M.V. wishes to thank the staff of INRA of Clermont Ferrand for their great assistance, and Philippe Label for providing ABA; Francesca Cardinale and Silvia Cavalletto for reviewing the manuscript and for ‘physics’ explanations, respectively, and Sonia Livigni for discussions on post-translational regulation and for her understanding. M.V. was partly granted during his PhD by SIGEVI project (PSR, misura 124, azione 1) founding by Piedmont region and UE.

References

Angeles G, Bond B, Boyer JS, Brodribb T, Brooks JR, Burns MJ, Cavender-Bares J, Clearwater M, Cochard H, Comstock J, Davis SD, Domec J-C, Donovan L, Ewers F, Gartner B, Hacke U, Hinckley T, Holbrook NM, Jones HG, Kavanagh K, Law B, Lopez-Portillo J, Lovisolo C, Martin T, Martinez-Vilalta J, Mayr S, Meinzer FC, Melcher P, Mencuccini M, Mulkey S, Nardini A, Neufeld HS, Passioura J, Pockman WT, Pratt RB, Rambal S, Richter H, Sack L, Salleo S, Schubert A, Schulte P, Sparks JP, Sperry J, Teskey R, Tyree M (2004) The cohesion-tension theory. *New Phytol* 163: 451–452

464 Bartlett MK, Scoffoni C, Sack L (2012) The determinants of leaf turgor loss point and prediction of
 465 drought tolerance of species and biomes: a global meta-analysis. *Ecol Lett* 15: 393–405
 466 Blackman CJ, Brodribb TJ (2011) Two measures of leaf capacitance: insights into the water
 467 transport pathway and hydraulic conductance in leaves. *Funct Plant Biol* 38: 118–126
 468 Čermák J, Kučera J, Bauerle WL, Phillips N, Hinckley TM (2007) Tree water storage and its
 469 diurnal dynamics related to sap flow and changes in stem volume in old-growth Douglas-fir
 470 trees. *Tree Physiol* 27: 181–198
 471 Chaumont F and Tyerman SD (2014) Aquaporins: highly regulated channels controlling plant water
 472 relations. *Plant Physiol* 164: 1600–1618
 473 Chitarra W, Balestrini R, Vitali M, Pagliarani C, Perrone I, Schubert A, Lovisolo C (2014) Gene
 474 expression in vessel-associated cells upon xylem embolism repair in *Vitis vinifera* L. petioles.
 475 *Planta* 239: 887–899
 476 Cochard H, Venisse JS, Barigah TS, Brunel N, Herbette S, Guilliot A, Tyree MT, Sakr S (2007)
 477 Putative role of aquaporins in variable hydraulic conductance of leaves in response to light. *Plant*
 478 *Physiol*, 143: 122–133
 479 Comstock JP (2002) Hydraulic and chemical signalling in the control of stomatal conductance and
 480 transpiration. *J Exp Bot* 53: 195–200
 481 Ding X, Iwasaki I, Kitagawa Y (2004) Overexpression of a lily PIP1 gene in tobacco increased the
 482 osmotic water permeability of leaf cells. *Plant Cell Environ* 27: 177–186
 483 Dixon HH (1914) Transpiration and the ascent of sap in plants. Macmillan, London.
 484 Fan W, Li J, Jia J, Wang F, Cao C, Hu J, Mu Z (2015) Pyrabactin regulates root hydraulic
 485 properties in maize seedlings by affecting PIP aquaporins in a phosphorylation-dependent
 486 manner. *Plant Physiol Bioch* 94: 28–34
 487 Flexas J, Scoffoni C, Gago J, Sack L (2013) Leaf mesophyll conductance and leaf hydraulic
 488 conductance: an introduction to their measurement and coordination. *J Exp Bot* 64: 3965–3981
 489 Gleason SM, Blackman CJ, Cook AM, Laws CA, Westoby M (2014) Whole-plant capacitance,
 490 embolism resistance and slow transpiration rates all contribute to longer desiccation times in
 491 woody angiosperms from arid and wet habitats. *Tree Physiol* 34: 275–284
 492 Goldstein G, Andrade JL, Meinzer FC, Holbrook NM, Cavelier J, Jackson P, Silvera K (1997) Stem
 493 water storage and diurnal patterns of water use in tropical forest canopy trees. *Plant Cell Environ*
 494 21: 397–406
 495 Gomes D, Agasse A, Thiébaud P, Delrot S, Gerós H, Chaumont F (2009) Aquaporins are
 496 multifunctional water and solute transporters highly divergent in living organisms. *Biochimica*
 497 *Biophysica Acta* 1788: 1213–1228

498 Guyot G, Scoffoni C, Sack L (2012) Combined impacts of irradiance and dehydration on leaf
 499 hydraulic conductance: insights into vulnerability and stomatal control. *Plant Cell Environ* 35:
 500 857–871

501 Hachez C, Heinen RB, Draye X, Chaumont F (2008) The expression pattern of plasma membrane
 502 aquaporins in maize leaf highlights their role in hydraulic regulation. *Plant Mol Biol* 68: 337–
 503 353

504 Honert TH van den (1948) Water transport in plants as a catenary process. *Discuss. Paraday Soc*, 3,
 505 146.

506 Hose E, Steudle E, Hartung W (2000) Absciscic acid and hydraulic conductivity of maize roots: a
 507 study using cell- and root-pressure probes. *Planta* 211: 874–882

508 Jang JY, Kim DG, Kim YO, Kim JS, Kang H (2004) An expression analysis of a gene family
 509 encoding plasma membrane aquaporins in response to abiotic stresses in *Arabidopsis thaliana*.
 510 *Plant Mol Biol* 54: 713–725

511 Kaldenhoff R, Ribas-Carbo M, Sans JF, Lovisolo C, Heckwolf M, Uehlein N (2008) Aquaporins
 512 and plant water balance. *Plant Cell Environ* 31: 658–666

513 Koide RT, Robichaux RH, Morse SR, Smith CM (2000) Plant water status, hydraulic resistance and
 514 capacitance. In *Plant Physiological Ecology: Field Methods and Instrumentation* (eds R.W.
 515 Pearcy, J.R. Ehleringer, H.A. Mooney & P.W. Rundel), pp. 161–183. Kluwer, Dordrecht, The
 516 Netherlands.

517 Lange OL, Nobel PS, Osmond CB, Ziegler H (1982) Physiological plant ecology II; water relations
 518 and carbon assimilation. *Encyclopedia of Plant Physiology*, vol. 12B. Springer, Berlin
 519 Heidelberg New York.

520 Lee SH, Chung GC, Jang JY, Ahn SJ, Zwiazek JJ (2012) Overexpression of PIP2;5 aquaporin
 521 alleviates effects of low root temperature on cell hydraulic conductivity and growth in
 522 *Arabidopsis*. *Plant Physiol* 159: 477–488

523 Lopez D, Venisse JS, Fumanal B, Chaumont F, Guillot E, Daniels MJ, Cochard H, Julien JL,
 524 Gousset-Dupont A (2013) Aquaporins and leaf hydraulics: Poplar sheds new light. *Plant Cell*
 525 *Physiol* 54: 1963–1975.

526 Laur J, Hacke UG (2013) Transpirational demand affects aquaporin expression in poplar roots. *J*
 527 *Exp Bot* 64: 2283–2293

528 Lovisolo C, Perrone I, Hartung W, Schubert A (2008) An absciscic acid-related reduced transpiration
 529 promotes gradual embolism repair when grapevines are rehydrated after drought. *New Phytol*
 530 180: 642–651

531 Martinez-Ballesta MdC, Carvajal M (2014) New challenges in plant aquaporin biotechnology. *Plant*
 532 *Sci* 217–218: 71–77
 533 Martre P, Morillon R, Barrieu F, North GB, Nobel PS, Chrispeels MJ (2002) Plasma membrane
 534 aquaporins play a significant role during recovery from water deficit. *Plant Physiol* 130: 2101–
 535 2110
 536 Maurel C, Boursiac Y, Luu D-T, Santoni V, Shahzad Z, Verdoucq L (2015) Aquaporins in Plants.
 537 *Physiol Reviews* 95 (4) 1321-1358
 538 McCulloh AK, Daniel MJ, Frederick CM, David RW (2014) The dynamic pipeline: hydraulic
 539 capacitance and xylem hydraulic safety in four tall conifer species. *Plant Cell Environ* 37: 1171–
 540 1183
 541 Millar AJ, Short SR, Chua NH, Kay SA (1992) A novel circadian phenotype based on firefly
 542 luciferase expression in transgenic plants. *Plant Cell* 4: 1075–1087
 543 Nardini A, Salleo S, Andri S. (2005) Circadian regulation of leaf hydraulic conductance in
 544 sunflower (*Helianthus annuus* cv. Margot). *Plant Cell Environ* 28: 750–759
 545 Nardini A, Raimondo F, Lo Gullo MA, Salleo S (2010) Leaf miners help us understand leaf
 546 hydraulic design. *Plant Cell Environ* 33: 1091–1100
 547 Ogasa M, Miki N, Murakami Y, Yoshikawa K (2013) Recovery performance in xylem hydraulic
 548 conductivity is correlated with cavitation resistance for temperate deciduous tree species. *Tree*
 549 *Physiol* 33: 335–344
 550 Pantin F, Monnet F, Jannaud D, Costa JM, Renaud J, Muller B, Simonneau T, Genty B (2013) The
 551 dual effect of abscisic acid on stomata. *New Phytol* 197: 65–72
 552 Parent B, Hachez C, Redondo E, Simonneau T, Chaumont F, Tardieu F (2009) Drought and
 553 abscisic acid effects on aquaporin content translate into changes in hydraulic conductivity and
 554 leaf growth rate: a trans-scale approach. *Plant Physiol* 149: 2000–2012
 555 Péret B, Li G, Zhao J, Band LR, Voß U, Postaire O, Luu DT, Da Ines O, Casimiro I, Lucas M,
 556 Darren MW, Lazzerini L, Nacry P, King JR, Jensen OE, Schaffer AR, Maurel C, Bennet MJ
 557 (2012) Auxin regulates aquaporin function to facilitate lateral root emergence. *Nature Cell Biol*
 558 14: 991–998
 559 Perrone I, Gambino G, Chitarra W, Vitali M, Pagliarani C, Riccomagno N, Balestrini R, Kaldenhoff
 560 R, Uehlein N, Gribaudo I, Schubert A, Lovisolo L (2012) The grapevine root-specific aquaporin
 561 VvPIP2;4N controls root hydraulic conductance and leaf gas exchange upon irrigation but not
 562 under water stress. *Plant Physiol* 160: 965–977

563 Phillips NG, Ryan MG, Bond BJ, McDowell NG, Hinckley TM, Čermák J (2003) Reliance of
564 stored water increases with tree size in three species in the Pacific Northwest. *Tree Physiol* 23:
565 237–245

566 Postaire O, Tournaire-Roux C, Grondin A, Boursiac Y, Morillon R, Schäffner AR, Maurel C (2010)
567 A PIP1 aquaporin contributes to hydrostatic pressure-induced water transport in both the root
568 and rosette of *Arabidopsis*. *Plant Physiology* 152: 1418–1430

569 Prado K, Maurel C (2013) Regulation of leaf hydraulics: from molecular to whole plant levels.
570 *Front Plant Sci* 4: 255

571 Prado K, Boursiac Y, Tournaire-Roux C, Monneuse JM, Postaire O, Da Ines O, Schäffner AR, Hem
572 S, Santoni V, Maurel C (2013) Regulation of *Arabidopsis* leaf hydraulics involves light-
573 dependent phosphorylation of aquaporins in veins. *Plant Cell* 25: 1029–1039

574 Robichaux HR (1984) Variation in the tissue water relation of two sympatric Hawaiian *Dubautia*
575 species and their natural hybrid. *Oecologia* 65: 75–81

576 Sack L, Cowan PD, Jaikumar N, Holbrook NM (2003) The "hydrology" of leaves: co-ordination of
577 structure and function in temperate woody species. *Plant Cell Environ* 26: 1343–1356.

578 Sack L. & Holbrook N.M. (2006) Leaf hydraulics. *Annu Rev Plant Biol*, 57, 361–381.

579 Sade N, Vinocur BJ, Diber A, Shatil A, Ronen G, Nissan H, Wallach R, Karchi H, Moshelion M
580 (2009) Improving plant stress tolerance and yield production: is the tonoplast aquaporin
581 SLTIP2;2 a key to isohydric to anisohydric conversion? *New Phytol* 181: 651–661

582 Sade N, Shatil A, Attia Z, Maurel C, Boursiac Y, Kelly G, Granot D, Yaaran A, Lerner S,
583 Moshelion M (2014) The role of plasma membrane aquaporins in regulating the bundle sheath-
584 mesophyll continuum and leaf hydraulics. *Plant Physiol* 166: 1609–1620

585 Sakurai-Ishikawa J, Murai-Hatano M, Hayashi H, Ahamed A, Fukushi K, Matsumoto T, Kitagawa
586 Y (2011) Transpiration from shoots triggers diurnal changes in root aquaporin expression. *Plant*
587 *Cell Environ* 34: 1150–1163

588 Scoffoni C, Vuong C, Diep S, Cochard H, Sack L (2014) Leaf shrinkage with dehydration:
589 coordination with hydraulic vulnerability and drought tolerance. *Plant Physiol* 164: 1772–1788

590 Siefritz F, Otto B, Bienert GP, van der Krol A, Kaldenhoff R. (2004) The plasma membrane
591 aquaporin NtAQP1 is a key component of the leaf unfolding mechanism in tobacco. *Plant J* 37:
592 147–155

593 Siefritz F, Tyree MT, Lovisolo C, Schubert A, Kaldenhoff R (2002) PIP1 Plasma Membrane
594 Aquaporins in Tobacco: From Cellular Effects to Function in Plants. *Plant Cell* 14 (4): 869-876

595 Shatil-Cohen A, Attia Z, Moshelion M (2011) Bundle-sheath cell regulation of xylem-mesophyll
 596 water transport via aquaporins under drought stress: a target of xylem-borne ABA? *Plant J* 67:
 597 72–80

598 Sellin A, Ounapuu E, Karusion A (2010) Experimental evidence supporting the concept of light-
 599 mediated modulation of stem hydraulic conductance. *Tree Physiol* 30: 1528–1535

600 Sperry JS, Adler FR, Campbell GS, Comstock JP (1998) Limitation of plant water use by
 601 rhizosphere and xylem conductance: results from a model. *Plant Cell Environ* 21: 347–359

602 Sperry JS, Meinzer FC, McCulloh AK (2008) Safety and efficiency conflicts in hydraulic
 603 architecture: scaling from tissues to trees. *Plant Cell Environ* 31: 632–645

604 Thompson AJ, Andrews J, Mulholland BJ, McKee JM, Hilton HW, Horridge JS, Farquhar GD,
 605 Smeeton RC, Smillie IR, Black CR, Taylor IB (2007) Overproduction of abscisic acid in tomato
 606 increases transpiration efficiency and root hydraulic conductivity and influences leaf expansion.
 607 *Plant Physiol* 143: 1905–1917

608 Tombesi S, Nardini A, Farinelli D, Palliotti A (2014) Relationships between stomatal behavior,
 609 xylem vulnerability to cavitation and leaf water relations in two cultivars of *Vitis vinifera*.
 610 *Physiol Plant* 152: 453–464

611 Tyree MT, Hammel HT (1972) The measurement of the turgor pressure and the water relations of
 612 plants by the pressure bomb technique. *J Exp Bot* 23: 267–282

613 Tyree MT, Ewers WF (1991) The hydraulic architecture of tree and other woody plants. *New*
 614 *Phytol* 119: 345–360

615 Vandeleur RK, Mayo G, Shelden MC, Gilliam M, Kaiser BN, Tyerman SD (2009) The role of
 616 plasma membrane intrinsic protein aquaporins in water transport through roots: diurnal and
 617 drought stress responses reveal different strategies between isohydric and anisohydric cultivars
 618 of grapevine. *Plant Physiol* 149: 445–460

619 Vandeleur RK, Sullivan W, Athman A, Jordans C, Gilliam M, Kaiser BN, Tyerman SD (2014)
 620 Rapid shoot-to-root signalling regulates root hydraulic conductance via aquaporins. *Plant Cell*
 621 *Environ* 37: 520–538

622 Verbeeck H, Steppe K, Nadezhdina N, Op de Beeck M, Deckmyn G, Meiresonne L, Lemeur R,
 623 Cermák J, Ceulemans R, Janssens IA (2007) Stored water use and transpiration in Scots pine: a
 624 modeling analysis with ANAFORE. *Tree Physiol* 27: 1671–1685

625 Voicu MC, Cooke JE, Zwiazek JJ (2009) Aquaporin gene expression and apoplastic water flow in
 626 bur oak (*Quercus macrocarpa*) leaves in relation to the light response of leaf hydraulic
 627 conductance. *J Exp Bot* 60 4063–4075

628 Xu, Y, Johnson CH (2001) A clock-and light-regulated gene that links the circadian oscillator to
629 LHCB gene expression. *The Plant Cell* 13(6): 1411-1426
630 Zeuthen T, Alsterfjord M, Beitz E, MacAulay N (2103) Osmotic water transport in aquaporins:
631 evidence for a stochastic mechanism. *J Physiol* 591.20: 5017–5029
632

633 **Captions to figures and tables**

634 **Table 1:** leaf water potential (Ψ_{leaf} , MPa) $\pm$ SE before and after the pressurisation and rehydration
635 treatments. ANOVA followed by Tukey’s post-hoc test was applied to assess significant differences
636 among lines and treatments in the final Ψ_{leaf} obtained ($* = P < 0.05$). The experiments concerned
637 two phases: in the first phase, the leaves were pressurised in the pressure chamber and water
638 flowing out from the leaf through the petiole was measured; in the second phase, the pressure was
639 released and water inflow into the leaf was measured by the weight-loss of water in the cylinder.

	Treatment	Line	native Ψ_{leaf}	Pressure applied	Final Ψ_{leaf}	ANOVA	n=
Dehydration phase	+0.5	WT	-0.01 $\pm$ 0	+0.5	-0.5 $\pm$ 0		6
		OE	-0.01 $\pm$ 0	+0.5	-0.5 $\pm$ 0		
	+1	WT	-0.01 $\pm$ 0	+0.99	-1 $\pm$ 0		16
		OE	-0.01 $\pm$ 0	+0.99	-1 $\pm$ 0		
Rehydration phase	Dark	WT	-1 $\pm$ 0	none	-0.08 $\pm$ 0.01	d	4
		OE	-1 $\pm$ 0	none	-0.08 $\pm$ 0.01	d	
	Light VPD$\approx$0	WT	-1 $\pm$ 0	none	-0.20 $\pm$ 0.02	b	4
		OE	-1 $\pm$ 0	none	-0.14 $\pm$ 0.01	c	
	Light trans	WT	-1 $\pm$ 0	none	-0.23 $\pm$ 0.01	b	4
		OE	-1 $\pm$ 0	none	-0.35 $\pm$ 0.08	a	
	Light trans ABA	WT	-1 $\pm$ 0	none	-0.19 $\pm$ 0.09	b	4
		OE	-1 $\pm$ 0	none	-0.17 $\pm$ 0.04	bc	

640

641

Figure 1: the experimental setup used in this study. Nitrogen gas, regulated through valves and monitored by a manometer, was used to compress the leaf in the pressure chamber. Water flowing out from the petiole increased the weight of the water-filled cylinder (left). Following rehydration (right), the chamber was removed, allowing the rehydration of the previously pressurised leaf. The rehydration was conducted under different stimuli (light, transpiration, ABA).

Figure 2: time-course of the cumulative water flow out of leaves in WT leaves (filled symbols) and OE leaves (empty symbols) in the +0.5 experiment. Data were normalised according to the dry weight (circles, left y-axis) or leaf area (triangles, right y-axis). Symbols represent the means $\pm$ SE. (n = 6).

654 **Figure 3:** a) hydraulic resistance (Rh), and b) capacitance (C) in WT leaves (black columns) and
 655 OE leaves (grey columns) obtained by pressurising the leaves as described in dehydration phase fo
 656 the experimental design: rehydrated leaves pressurised to +0.5 MPa (+0.5) or +1.0 MPa (+1).
 657 Columns represent the means ($n = 6$ for +0.5 and $n = 16$ for +1.0) $\pm$ SE Means were obtained by
 658 averaging the measurements performed at different times of day (8:00–10.00; 11:00–13.00; 14:00–
 659 16.00). Asterisks mark significant differences between means (* = $P < 0.05$ ** = $P < 0.01$).

660

661

663 **Figure 4:** frame a; time-course of the water flow into the petioles of WT leaves (filled symbols) and
 664 OE leaves (empty symbols) in the four recovery treatments: dark (circles), light low transpirative
 665 conditions ($VPD \approx 0$; triangles), light transpirative conditions (square) and light transpirative
 666 conditions in the presence of ABA (rhombus) ($n=4$). Frame b highlights the first 600 s of the time-
 667 course (the mean values of of Rh and C are displayed in Fig. 5).
 668

672 **Figure 5:** a) hydraulic resistance (Rh), and b) capacitance (C) in WT (black columns) and OE (grey
 673 columns) obtained from the rehydration of leaves after dehydration to $\Psi_{\text{leaf}} = -1$ MPa (+1), as
 674 described in the rehydration phase of the experimental design. Recovery treatments were performed
 675 in dark, in light low transpirative, light transpirative or light transpirative conditions after adding
 676 ABA (final concentration 100 μmol) to the cylinder where the cut petioles were submerged.
 677 Columns represent the means ($n = 4$) $\pm$ SE. Different letters mark significant differences ($P < 0.05$)
 678 between means according to ANOVA after data normalisation. In frame b, $P < 0.001$.

682 **Figure 6:** expression of endogenous and transgenic *PIP*-type AQP genes in WT and OE lines under
683 dark and light transpirative conditions in rehydrated leaves. Relative expression levels of *VvPIP1;1*,
684 *VvPIP2;1*, *VvPIP2;2*, *VvPIP2;3*, endogenous *VvPIP2;4N*, and transgenic *VvPIP2;4N* were
685 determined by qRT-PCR in leaves. The PCR data were normalised with those for UBI transcripts.
686 Data are expressed as the mean $\pm$ SE; different letters denote significant differences at $P \leq 0.05$.

687

688

689 **Figure 7:** relationship between Ψ_{leaf} and RWC (a) and Ψ_{leaf} and C (b). Data were obtained from leaves
 690 allowed to dehydrate in darkness in the laboratory atmosphere. Filled circles represent the WT leaves; open
 691 circles represent OE leaves. Solid and dotted lines correspond to the regression of WT and OE, respectively.

692
 693
 694
 695 **Figure 8:** schematic representation of the leaf hydraulic pathway. The leaf is divided in several
 696 compartments represented by resistors (vascular system, R_v ; Bundle sheath cells, BSCs R_b ; apoplast, R_a ;
 697 symplast R_{s1} R_{s2} ; and stomata R_{st}) and capacitors (symplast, C_{s1} C_{s2}). Dashed lines indicate additional
 698 parts that can be added to the system. When transpiration is stopped, the system could be simplified by
 699 summing the capacitors and the reciprocal resistors (right side).

702 **Figure S1:** time-course of the cumulative water flow out of leaves in WT leaves (filled symbols) and OE
 703 leaves (empty symbols) in the +0.5 experiment. Data (corresponding to the ones in Fig. 2) were plotted
 704 according to the time of the day when the experiment was performed: morning (8:00–10:00, circles), noon
 705 (11:00–13:00, triangles) and afternoon (14:00–16:00, squares). Symbols represent the means $\pm$ SE (n = 2).

710 **Figure S2:** relationship between dry weight and leaf area (circles) and fresh weight and leaf area
711 (triangles) in WT (filled symbols) and OE leaves (empty symbols). Regression lines are shown for
712 WT (solid trend line) and OE (dotted trend line).

716 **Figure S3:** a) hydraulic resistance (Rh), and b) capacitance (C) in WT leaves (black columns) and
 717 OE leaves (grey columns) obtained by pressurising the leaves to +0.5 MPa (+0.5). Columns
 718 represent the means ($n = 2$) $\pm$ SE. Means were obtained by averaging the measurements performed
 719 at different times of day (morning 8:00–10:00; noon 11:00–13:00; afternoon 14:00–16:00 hours).
 720 Averages were obtained from the same datasets of Fig. 2 and 3 (only +0.5 treatment) and S1.

723 **Reflection coefficient implication on hydraulic capacitance**

724 We provide here the theoretical framework demonstrating the relation between the bulk leaf reflection
725 coefficient s and the bulk leaf capacitance C . The demonstration is based on a reanalyze of the Pressure-
726 Volume curve theory (Tyree and Hamel 1972).

727 Whole leaf water potential Y is usually considered as the algebraic sum of the pressure turgor potential P and
728 the osmotic potential P :

729

730 $Y = P + P$ (s1)

731

732 However, this equation is correct only when the reflection coefficient is equal to one. If s less than unity
733 then:

734

735 $Y = P + sP$ (s2)

736

737 When leaf dehydrate, its relative water content (RWC) decreased and the total relative loss of water R is
738 equal to :

739

740 $R = 1 - RWC$ (s3)

741

742 Assuming a constant apoplastic water content fraction (af), we can compute the relative water content loss
743 R_s of the symplasmic compartment as:

744

745 $R_s = R / (1 - af)$ (s4)

746

747 R_s is equal to 0 when the leaf is fully turgid and equals to 1 when the symplasmic compartment is empty.

748 Leaf capacitance C is defined as:

749

750 $C = dR_s / dY$ (s5)

751

752 or, as a proxy, as:

753

754 $C = (R_s(Y_1) - R_s(Y_2)) / (Y_2 - Y_1)$ (s6)

755

756 We will focus here on our +0.5 experiment, where $Y_1 = 0$ and $Y_1 = -0.5\text{MPa}$

757

758 Defining P_0 as the osmotic potential at full leaf turgor and e as the bulk leaf modulus of elasticity, we can

759 express P and P as a function of R_s as:

760

761 $P = -s(P_0 + eR_s); P > 0$ (s7)

762 $P = sP_0 / (1 - R_s)$ (s8)

763

764 Equations s1, s7 and s8 are used to construct a Höfler diagram (figure s1):

766

767

768 Figure S4: Höfler diagram showing the changes in whole leaf water potential (Y), pressure potential (P) and

769 osmotic potential (Π) as a function of relative symplasmic water content. The parameters used to construct

770 the diagram are shown in the insert. These parameters were obtained on *Vitis* leaves similar to those used in
771 this study.

772 Combining s1, s7 and s8 we have:

773

774
$$Y = -s(P_0 + eR_s) + sP_0 / (1 - R_s) \quad \text{for } P > 0 \quad (s9)$$

775
$$Y = sP_0 / (1 - R_s) \quad \text{for } P = 0 \quad (s10)$$

776

777 By solving equations s9 and s10 we can express R_s as a function of Y as:

778

779
$$R_s = \frac{\sigma(\varepsilon - \Pi_0) - \Psi - \sqrt{(\Psi + \sigma(\Pi_0 - \varepsilon))^2 + 4\sigma\varepsilon\Psi}}{2\sigma\varepsilon} \quad \text{for } P > 0 \quad (s11)$$

780

781
$$R_s = 1 - sP_0/Y \quad \text{for } P = 0 \quad (s12)$$

782

783 Exact solutions of C can then be derived from s11 and s12 using s5 or s6.

784 A proxy of C can also be obtained if we assume that for low R_s values equation s8 can be approximated by:

785

786
$$P \approx sP_0 \quad (s13)$$

787

788 then, by combining s2, s7 and s13 we have:

789

790
$$Y \approx -seR_s \quad (s14)$$

791

792 then it comes:

793

794 $C \approx 1/se$ (s15)

795

796 The relative change of whole leaf capacitance C_{rel} when the reflection coefficient decreases from 1 to s is:

797

798 $C_{rel} \approx 1/s$ (s16)

799

800 The relations between C_{rel} derived from s11 and s16 and s are shown in figure s2. The approximation is
801 robust but valid only when R_s is low and s is high (>0.5).

802 Therefore, dividing s by two will double approximately the whole leaf capacitance and this effect is largely
803 independent of e , P_0 and Y .

804

805

806

807 Figure S5. Effect of the reflection coefficient s on the relative change in leaf capacitance. The exact relation
808 is shown in green and the approximation given in equation s16 is shown in red.

809