

This is the author's final version of the contribution published as:

Mori, Emiliano; Mazzoglio, Peter John; Rima, Patrizia Carla; Aloise, Gaetano; Bertolino, Sandro. Bark-stripping damage by *Callosciurus finlaysonii* introduced into Italy. *MAMMALIA*. 80 (5) pp: 507-514.  
DOI: 10.1515/mammalia-2015-0107

The publisher's version is available at:

<https://www.degruyter.com/view/j/mamm.2016.80.issue-5/mammalia-2015-0107/mammalia-2015-0107.pdf>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

**Mammalia, 2016, 80: 507-514**

**Bark-stripping damage by *Callosciurus finlaysonii* introduced into Italy**

Emiliano Mori, Peter John Mazzoglio, Patrizia Carla Rima, Gaetano Aloise, Sandro Bertolino\*

**\*Corresponding author: Sandro Bertolino**, Department of Agriculture, Forest and Food Sciences, University of Turin, Largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy

**Running title:** Bark-stripping damage by *Callosciurus finlaysonii*.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

**Abstract:** The Finlayson’s squirrel *Callosciurus finlaysonii* was introduced into Italy during the 1980s and has established two viable populations. The diet of this species includes a high proportion of tree barks, suggesting an intensive debarking behaviour. We reported a severe bark-stripping impact in both colonized areas, and we tested whether a preference for some tree species existed. Results of this work show the presence of a wide spectrum of damaged species, without any strong preference, mainly with large wounds. Old deciduous plants and conifers, which presented a hard bark, were usually avoided.

**Keywords:** alien species; debarking; Finlayson’s squirrel; tree species selection; Italy.

**Introduction**

Bark-stripping behaviour is commonly observed among herbivorous species, such as ungulates (Motta 1996; Månsson and Jarnemo 2013), lagomorphs (Allman 1946; Chapman et al. 1982) and rodents (Gill 1992; Sharma and Prasad 1992; Baxter and Hansson 2001). Concerning rodents, bark portions are often removed when other more preferred food resources are not available, e.g. during snow cover or during the warm season, as well as to reach the phloem sap, rich in sugary components (Kenward and Parish 1986; Baxter and Hansson 2001). In the latter case, debarking is markedly exerted during the spring, when the sap flow increases in deciduous plants (Kenward and Parish 1986). The reasons for bark-stripping behaviours are not always clear and a number of explanations have been proposed (e.g. Kenward 1983).

Bark removal can be exerted along the entire stem circumference or in a spiral pattern around the trunk/branches, with different consequences for the plant. Decortications affecting only a part of the circumference induce local diebacks; by contrast, when annular decortications interrupt the sap flow, the dieback of the distal portions and the death of the whole plant may occur. As a response to

27 the damage, plants react with healing phenomena and with the formation of scar tissue around the  
28 lesions (Mullick 1977; Biggs et al. 1984). Trees irreparably damaged are easily blown down by  
29 wind or attacked by pathogens (e.g. cryptogams), which may easily invade the plant through bark  
30 wounds (Tubeuf 1897; Purohit et al. 2001); in case of completely healed wounds, reductions in the  
31 commercial quality of wood also occur because of the presence of defects. Bark-stripping of  
32 selective tree species also alters the composition of forests, as well as hinder the establishment of  
33 new woodlands (Gill et al. 1995; Kerr and Niles 1998).

34 Among arboreal rodents, many squirrel species strip barks, e.g. Eurasian red squirrel *Sciurus*  
35 *vulgaris* (Pulliainen and Salonen 1963), fox squirrel *S. niger* (Allen 1943), Eastern grey squirrel *S.*  
36 *carolinensis* (Kenward and Parish 1986; Mountford 2006), American red squirrel *Tamiasciurus*  
37 *hudsonicus* (Sullivan and Vyse 1987) and red-bellied tree squirrel *Callosciurus erythraeus* (Zhu et  
38 al. 1990).

39 In Europe, the native red squirrel only occasionally causes a significant damage to forest  
40 vegetation or to arboriculture systems (Moller 1983; Gurnell 1987). By contrast, although in Italy  
41 and in its native range the damage is limited (Kenward 1989; Signorile and Evans 2007; Bertolino  
42 2008), the introduced Eastern grey squirrel is responsible for severe damage to the timber industry  
43 and forests in Great Britain and Ireland (Williams et al. 2010; Mayle and Broome 2013).  
44 Considering the differences in bark-stripping behaviour among areas of introduction, if the species  
45 will be left to expand in Europe, predicting the severity of its impact to forestry is challenging  
46 (Bertolino et al. 2008, 2014; Di Febbraro et al. 2013). The Eastern grey squirrel is not the only  
47 squirrel species introduced into Europe; at least three other species have established viable  
48 populations in one or more countries (red-bellied tree squirrel, Finlayson's squirrel *C. finlaysonii*  
49 and Siberian chipmunk *Tamias sibiricus*, Bertolino 2009; Bertolino and Lurz 2013). Therefore, an  
50 evaluation of the possible damage by introduced tree squirrels is particularly needed to better draw  
51 a complete assessment of their impacts to biodiversity and human activities.

52 The Finlayson's squirrel is a species naturally distributed in Indochina, from Myanmar to  
53 Vietnam (Thorington et al. 2012). Introduced populations are currently recorded in Singapore,  
54 Japan and Italy (Bertolino et al. 1999; Oshida et al. 2007; Bertolino and Lurz 2013). In Italy, two  
55 populations are present in the Northern and Southern parts of the peninsula; a study on the diet of  
56 this species in one of those areas already highlighted a bark-stripping behaviour in winter (Bertolino  
57 et al. 2004). Aims of this study were to evaluate (i) the impact of bark stripping by the Finlayson's  
58 squirrels in the areas of presence in Italy, and (ii) the preference for some plant species.

59

## 60 **Materials and Methods**

61

### 62 *Study areas*

63

64 Field work was carried out at both areas of occurrence of the Finlayson's squirrel in Italy (Figure 1).  
65 In the north of the country, the species is localized in the city of Acqui Terme (Province of  
66 Alessandria, Northern Italy) and its suburbs. In the south, the Finlayson's squirrel expanded its  
67 range along the coastline, both northwards and southwards of Maratea (Province of Potenza,  
68 Southern Italy), the city of first introduction (Aloise and Bertolino 2005). The study site in Northern  
69 Italy is an urban park (2 ha), located in the city center of Acqui Terme; here the survey was  
70 conducted in the years 1998-1999. The most represented tree species were *Celtis australis*, *Platanus*  
71 *sp.*, *Tilia cordata*, *Pinus spp.* and *Cedrus spp.* (Table 1). In Southern Italy, nine wooded areas were  
72 surveyed along the coast in the year 2004 (Supplementary Figure 1): 1 site in the point of the first  
73 introduction, 3 sites within 5.25 km northwards and 5 sites within 6.50 km southwards  
74 (Supplementary Figure 2). At that time, the species had spread 9 km northwards and 7 km  
75 southwards. These areas include a narrow belt of coastal woods, mainly composed of *Pinus*  
76 *halepensis*, *Quercus ilex* and *Ceratonia siliqua* (Table. 2). Valleys connect these coastal woodlands  
77 with the deciduous woodlands (*Quercus cerris*, *Q. virgiliana*, *Castanea sativa*) of the hinterland.

78

## 79 *Data collection*

80

81 All the plants of the study sites were identified at the species or genus level, counted and checked  
82 for the presence of debarking damage on the trunk and branches by Finlayson's squirrels. We  
83 differentiated the damage to the trunk or to the branches, because a different impact on the survival  
84 of the plant may occur. The extension of the damage was assessed on a sliding scale: no damage,  
85 less than 50 cm<sup>2</sup> of debarked surface, 50-500 cm<sup>2</sup>, > 500 cm<sup>2</sup>. A chi-square test was then applied to  
86 assess differences in the percentage of damage extension within and between the two study areas. In  
87 Northern Italy, the extent of the damage was also evaluated after one and nine months from an  
88 intervention of phytosanitary cutting on 42 littleleaf lindens *Tilia cordata*, 9 European hackberries  
89 *Celtis australis*, 1 Northern white cedar *Thuja occidentalis* and 1 Norway spruce *Picea excelsa*, to  
90 assess whether they had any effect on the debarking activity. During these interventions, some trees  
91 were removed and all branches were cut in the others, thus we expected a reduction on the  
92 debarking activity by *C. finlaysonii* on these species, at least before new branches had grown.

93 In Southern Italy, we also assessed whether the damage decreased in the various plots with an  
94 increasing distance from the centre of first introduction of the species (Maratea).

95 The Ivlev's Electivity Index (E, Ivlev 1961) was computed to assess the selection of Finlayson's  
96 squirrel for tree species, through the statistical software R 3.1.1, package *gplots* (Chiatante 2014).  
97 The values of this index range from -1 to +1, with values between -1 and 0 indicating avoidance and  
98 values from 0 to +1 indicating preference for a certain resource. We followed the suggestion by  
99 Lazzaro (1987) and Lückstädt and Reito (2002), who showed that a true selection or avoidance can  
100 be claimed only at values > 0.3 or < -0.3 respectively. Species represented by one individual were  
101 excluded from the analysis.

102

## 103 **Results**

104

105 ***Northern Italy***

106

107 A total of 308 trees were monitored, and bark stripping occurred in 242 (78.6%) trees. Finlayson's  
108 squirrel completely avoided *Pinus strobus*, *Celtis australis*, *Tilia cordata*, *Cercis siliquastrum*, *Ilex*  
109 *aquifolium*, *Acer saccharinum*, and *Platanus* sp. (all, E = -1, Table 1) at the level of the trunk;  
110 whereas *Picea excelsa*, *C. siliquastrum*, *A. saccharinum* and *Platanus* sp. were avoided at the  
111 branch level. All the other species were selected for bark-stripping, with E values generally > 0.8  
112 (Table 1).

113 Overall, 5.05% of all damaged trees appeared affected on a small area, 71.21% (mainly on  
114 *Celtis australis* and *Tilia cordata*) fell into the intermediate category, and the remaining 23.74%  
115 showed extensive decortications; the intermediate extension was significantly higher than the other  
116 categories (all species, trunks:  $\chi^2_{(2)} = 5.57$ , P < 0.05; branches:  $\chi^2_{(2)} = 266$ , P < 0.001) (Table 1,  
117 Figure 4).

118 The phytosanitary cutting affected the behaviour of the Finlayson's squirrels (Figure 2). For the  
119 first month after the intervention (T<sub>1</sub>), squirrels intensified debarking on evergreen species, while  
120 those on the deciduous plants decreased; nine months after the management intervention (T<sub>2</sub>), the  
121 damage decreased on evergreen plants and increased again on broadleaves ( $\chi^2_{(2)} = 90.28$ , P <  
122 0.001).

123

124 ***Southern Italy***

125

126 A total of 209 trees belonging to six species were surveyed and checked for debarked areas (Table  
127 2). Bark stripping by Finlayson's squirrel occurred on 107 (51.2%) trees, mainly *Ceratonia siliqua*  
128 trunk (E = 0.63) and branches (E = 0.77) and *Olea europaea* (E = 0.35 trunk, E = 0.85 branches).  
129 *Quercus ilex* was avoided both at the trunk (E = -0.86) and at the branch level (E = -0.51). *Quercus*

130 *virgiliana* ( $E = 0.52$ ) and *Pinus halepensis* ( $E = 0.39$ ) were only selected at the branch level; all the  
131 other species were used proportionally to their availability. Overall, about 43% of the damage  
132 occurred on a surface of over 500 cm<sup>2</sup> and mainly on *Ceratonia siliqua*, while the damage on the  
133 other species was always of a smaller extent (Table 2) (all species, trunks:  $\chi^2_{(2)} = 6.89$ ,  $P < 0.05$ ;  
134 branches:  $\chi^2_{(2)} = 6.77$ ,  $P < 0.05$ , Figure 4).

135 The percentages of trees with damage on trunks, branches or both, were not correlated with the  
136 distance from the site of introduction (all,  $P > 0.05$ , Figure 3, see also the Supplemental Figure 2).

137 Overall, the damage was higher in Northern than in Southern Italy at the branches ( $\chi^2_{(2)} = 69.13$ ,  
138  $P < 0.001$ ) but not at the trunk level ( $\chi^2_{(2)} = 2.06$ ,  $P < 0.36$ ; Figure 4).

139

## 140 **Discussion and Conclusions**

141

142 The Finlayson's squirrel has been introduced into two areas of Italy during the 1980s (Bertolino et  
143 al. 1999; Aloise and Bertolino 2005). Twenty years later, the damage produced to trees through  
144 bark-stripping was severe at both sites. In particular, all the species with few exceptions were  
145 debarked in a urban park in Acqui Terme (Northern Italy), with a marked preference for deciduous  
146 plants. The damage in natural woodlands of Southern Italy involved a smaller number of plants and  
147 was preferentially exerted on *Olea europaea* and *Ceratonia siliqua*, as well as on branches of  
148 *Quercus virgiliana* and *Pinus halepensis*. In several of the cases observed, bark stripping was likely  
149 to jeopardize the survival of part of the plant or even of the whole tree.

150 The Finlayson's squirrel is a frugivorous species, although its diet may vary according to the  
151 season and to the environmental availability of food resources (Bertolino et al. 2004). Bark-  
152 stripping occurs mainly during late autumn and winter, when the availability of seeds, fruits and  
153 blossoms decreases (Bertolino et al. 2004). In Northern Italy, where the population is still restricted  
154 to a small area, decortications occur also during the warm seasons, when other food sources may be  
155 less available. Mills (1938) suggested that sap would not be a survival food, but it would be

156 consumed because of its tasty sugary flavour. Our results confirmed the presence of a large  
157 spectrum of plant species damaged by the Finlayson's squirrel in Italy, which does not seem to  
158 show any strong preference. The few avoided species presented hard barks or, as in the case of *Acer*  
159 *saccharinum* in Northern Italy, a species usually debarked by squirrels (Brenneman 1954), they  
160 were isolated plants which may be reached by squirrels only by moving on the ground. Finlayson's  
161 squirrel is mostly arboreal and does not like to move on the ground (Lekagul and McNeely 1988;  
162 Thorington et al. 2012), a behaviour even discouraged by the presence of dogs in the park. As for  
163 plane trees *Platanus* sp., their bark peels off in large plates also after a slight contact with the teeth  
164 of the squirrel: so, clear young bark is brought to the surface, but it is not directly attacked by  
165 squirrels.

166 Most of the wounds caused by this tree squirrel affected tree portions larger than 50 cm<sup>2</sup> and  
167 often larger than 500 cm<sup>2</sup>, resulting in serious damage. The attacked plants may only heal moderate  
168 wounds and the bark removal determines a lower resistance of the trees to mechanical stresses and  
169 to parasite attacks. Behavioural and ecological reasons inducing Finlayson's squirrel to consume  
170 bark are not completely understood yet. Some species, such as *Celtis australis* and *Tilia cordata*,  
171 were debarked mainly during the autumn, when the sap flow is minimal, and for a small extent;  
172 wounds were mainly located on the branches where the bark may be more tender with respect to the  
173 trunk. Furthermore, only a few scraps were left on the ground, suggesting a direct interest towards  
174 the bark, thus not to the sap, as a food resource. In autumn and winter, consistently, fruit  
175 consumption was lower than in the rest of the year (Bertolino et al. 2004). By contrast, other species  
176 have been frequently used during spring when the sap flow is greatest; in these cases, the portions  
177 of removed bark were conspicuous and many scraps were found on the ground, in analogy with the  
178 behaviour of grey and red-bellied squirrels (Kuo et al. 1982; Kenward and Parish 1986).

179 Silvicultural interventions on lindens and hackberries carried out in Northern Italy influenced  
180 the debarking activities: after one month, a reduction of damage to the hardwood and a heavier  
181 damage on evergreen species has been recorded. Fruits of hackberries and lindens are an important

182 part of the diet of the Finlayson's squirrel (Bertolino et al. 2004); therefore, the cutting of 50 trees  
183 belonging to these species may have led the squirrels to feed on bark as an alternative food source.

184 Overall, despite a more localized distribution range in Italy, the debarking damage by  
185 Finlayson's squirrels seems to be higher with respect to that by grey squirrels (Signorile and Evans  
186 2007). Considering the vegetation characteristics of the areas of occurrence of the Finlayson's  
187 squirrel, the density of population and the likelihood of expansion (Bertolino et al. 2004; Aloise and  
188 Bertolino 2005), it is likely that this type of damage will increase in the next future, with a higher  
189 impact in Southern Italy, where the range expansion is recorded both along the coastline and  
190 towards the hinterland (Aloise and Bertolino 2005; Aloise and Bertolino 2008; Aloise et al. 2011).  
191 Furthermore, the amount of damage seems not to decrease with the distance from the site of first  
192 introduction, thus depending on local ecological factors, such as food resources and different  
193 attractiveness/palatability of tree species.

194 As far as we know, this work represents the first report of bark-stripping damage produced by  
195 the Finlayson's squirrel (cf. Bertolino et al. 2015); however, this is probably due to the absence of  
196 specific studies on the species. In fact, the congener red-bellied tree squirrel is known to produce  
197 severe damage both in the native range and in many areas of introduction (Zhu et al. 1990;  
198 Bertolino and Lurz 2013)

199 Once an alien species has spread over large areas, eradication and numerical or spatial control  
200 are generally hard programmes to be carried out (Genovesi and Shine 2004; Bertolino et al. 2015).  
201 Currently, the eradication of the Finlayson's squirrel from Italy seems to be still possible, and  
202 should be undertaken before the species spreads further, resulting in extensive damage. It must be  
203 remarked that both populations are close to one of the most productive hazelnut-growing areas  
204 (Langhe and Roero in Northwestern Italy, Salerno and Avellino provinces in Southern Italy).  
205 Moreover, in Northern Italy, the hazelnut-growing area is already menaced by the grey squirrel  
206 (Bertolino 2014). The introduction of this tree squirrel in Italy is due to intentional releases or

207 escapes of animals kept in captivity by private citizens; being the complete avoidance of such  
208 escapes almost impossible, restrictive measures in the trade of exotic species should be adopted.

209

## 210 **References**

211

212 Allen, D.L. 1943. Michigan fox squirrel management. Michigan Department of Conservation,  
213 Game Division Publ. 100:1-404.

214 Allman, D. 1946. Observations on damage by hares at Clonegal Forest. Irish For. 3: 92.

215 Aloise, G. and S. Bertolino. 2005. Free ranging population of the Finlayson's squirrel *Callosciurus*  
216 *finlaysonii* (Horsfield, 1824) (Rodentia, Sciuridae) in South Italy. Hystrix, Ital. J. Mammal.  
217 16:70-74.

218 Aloise, G. and S. Bertolino. 2008. Espansione della popolazione di *Callosciurus finlaysonii*  
219 (Horsfield, 1824) (Rodentia, Sciuridae) della costa tirrenica meridionale. Atti VI Congresso  
220 Italiano di Teriologia, Cles (Trento), 16-18<sup>th</sup> April 2008. Hystrix, Ital. J. Mammal. Supp.  
221 (2008):65.

222 Aloise, G., L. Lombardi and E. Fulco. 2011. Populations expansion of an exotic species in Southern  
223 Italy: the Finlayson's Squirrel *Callosciurus finlaysonii*. II International Congress Problematic  
224 Wildlife: Conservation and management. Genazzano (Rome, Italy), February 3-5 2011.  
225 Abstracts: 104-105.

226 Baxter, R. and L. Hansson. 2001. Bark consumption by small rodents in the northern and southern  
227 hemispheres. Mammal Rev. 31:47-59.

228 Bertolino, S. 2008. Introduction of the American grey squirrel (*Sciurus carolinensis*) in Europe: a  
229 case study in biological invasion. Current Sci. 95:903–906.

230 Bertolino, S. 2009. Animal trade and non-indigenous species introduction: the world-wide spread of  
231 squirrels. Diversity Distrib. 15:701–708.

232 Bertolino, S. and P.W. Lurz. 2013. *Callosciurus* squirrels: worldwide introductions, ecological  
233 impacts and recommendations to prevent the establishment of new invasive populations.  
234 Mammal Rev. 43:22-33.

235 Bertolino, S., I. Currado and P.J. Mazzoglio. 1999. Finlayson's (Variable) squirrel *Callosciurus*  
236 *finlaysonii* in Italy. Mammalia 63:522-525.

237 Bertolino, S., P.J. Mazzoglio, M. Vaiana and I. Currado. 2004. Activity budget and foraging  
238 behavior of introduced *Callosciurus finlaysonii* (Rodentia, Sciuridae) in Italy. J. Mammal. 85:58-  
239 63.

240 Bertolino, S., P. Colangelo, E. Mori and D. Capizzi. 2015. Good for management, not for  
241 conservation: an overview of research, conservation and management of Italian small mammals.  
242 Hystrix, Ital. J. Mammal. in press.

243 Bertolino, S., P.W.W. Lurz, R. Sanderson and S. Rushton. 2008. Predicting the spread of the  
244 American grey squirrel (*Sciurus carolinensis*) in Europe: a call for coordinated European  
245 approach. Biol. Cons. 141: 2564-2575.

246 Bertolino, S., N. Cordero di Montezemolo, D.G. Preatoni, L.A. Wauters and A. Martinoli. 2014. A  
247 grey future for Europe: *Sciurus carolinensis* is replacing native red squirrels in Italy. Biol. Inv.  
248 16:53-62.

249 Biggs, A.R., W. Merrill and D.D. Davis. 1984. Discussion: response of bark tissues to injury and  
250 infection. Can. J. Forest Res. 14: 351-356.

251 Brenneman, W.S. 1954. Tree damage by squirrels: silviculturally significant? J. For. 52:604.

252 Chapman J.A., J.G. Hockman and W.R. Edwards. 1982. The cottontails. In: (J.A. Chapman, G.A.  
253 Feldhamer, eds.) Wild mammals of North America: biology, managements and economics. The  
254 John Hopkins University Press, Baltimore, USA. pp. 83-123.

255 Chiatante, G. 2014. Habitat selection of Dartford warbler *Sylvia undata* on Elba island (Tuscan  
256 Archipelago, Italy). Bird Study 61:438-443.

257 Di Febbraro, M., P.W. Lurz, P. Genovesi , L. Maiorano, M. Girardello and S. Bertolino. 2013. The  
258 use of climatic niches in screening procedures for introduced species to evaluate risk of spread: a  
259 case with the American eastern grey squirrel. PloS one 8:e66559.

260 Genovesi, P. and C. Shine. 2004. European Strategy on Invasive Alien Species. Nature and  
261 Environment, n. 137. Council of Europe publishing (ed.), Strasbourg, France.

262 Gill, R.M.A. 1992. A review of damage by mammals in north temperate forests. 2. Small mammals.  
263 Forestry 65:281-308.

264 Gill, R.M.A., J. Gurnell and R.C. Trout. 1995. Do woodland mammals threaten the development of  
265 new woods? In: (R. Ferris-Kaan, eds.) The Ecology of woodland Creation. Wiley, Chichester  
266 (UK). pp. 201–224.

267 Gurnell, J. 1987. The natural history of squirrels. London, UK: Helm Editions.

268 Ivlev, V.S. 1961. Experimental ecology of the feeding of fishes. Yale University Press, New Haven,  
269 Connecticut, USA.

270 Kenward, R.E. 1983. The causes of damage by red and grey squirrels. Mammal Rev. 13:159-166.

271 Kenward, R.E. 1989. Bark-stripping by grey squirrels in Britain and North America: why does the  
272 damage differ. In: (R.E. Kenward, eds.) Mammals as Pests. Chapman and Hall. pp. 44-154.

273 Kenward, R.E. and T. Parish. 1986. Bark stripping by grey squirrels (*Sciurus carolinensis*). J. Zool.  
274 210:473-481.

275 Kerr, G. and J. Niles. 1998. Growth and provenance of Norway maple (*Acer platanoides*) in  
276 lowland Britain. Forestry 71:219-224.

277 Kuo PC, Kao C, Liu CF, Hwang FD. 1982. Correlation of the damage by Formosan red-bellied  
278 squirrel with chemical composition of the wood: part III. Sugar content of bark. Memoirs of the  
279 College of Agriculture National Taiwan University. 22:25-36.

280 Lazzaro, X. 1987. A review of planktivorous fishes: their evolution, feeding behaviours,  
281 selectivities, and impacts. Hydrobiol. 146:97-167.

282 Lekagul, B. and J.A. McNeely. 1988. Mammals of Thailand. Bangkok, Thailand: Darnsutha Press.

283 Lückstädt, C. and T. Reiti. 2002. Investigations on the feeding behavior of juvenile milkfish  
284 (*Chanos chanos* Forsskål) in brackishwater lagoons on South Tarawa, Kiribati. Verhandlungen  
285 der Gesellschaft für Ichthyologie Band 3:37-43.

286 Månsson J. and A. Jarnemo. 2013. Bark-stripping on Norway spruce by red deer in Sweden: level  
287 of damage and relation to tree characteristics. Scand. J. Forest. Res. 28:117-125.

288 Mayle, B.A. and A.C. Broome. 2013. Changes in the impact and control of an invasive alien: the  
289 grey squirrel (*Sciurus carolinensis*) in Great Britain, as determined from regional surveys. Pest  
290 Manag. Sci. 69:323-333.

291 Moller, H. 1983. Foods and foraging behaviour of red (*Sciurus vulgaris*) and grey (*Sciurus*  
292 *carolinensis*) squirrels. Mammal Rev. 13:83-98.

293 Motta, R. 1996. Impact of wild ungulates on forest regeneration and tree composition of mountain  
294 forests in the Western Italian Alps. Forest. Ecol. Manag. 88:127-134.

295 Mountford, E.P. 2006. Long-term patterns and impacts of grey squirrel debarking in Lady Park  
296 Wood young-growth stands (UK). Forest. Ecol. Manag. 232:100-113.

297 Mills, E.M. 1938. Free injury by squirrels. Bull. Mass. Agr. Exp. Stn. 353:79-80.

298 Mullick, D.B. 1977. The non-specific nature of defense in bark and wood during wounding, insect  
299 and pathogen attack. In: (D.B. Mullick, eds.) The structure, biosynthesis, and degradation of  
300 wood. Springer USA. pp. 395-441.

301 Oshida, T., H. Torii, L.K. Lin, J.K. Lee, Y.J. Chen, H. Endo and M. Sasaki. 2007. A preliminary  
302 study on origin of *Callosciurus* squirrels introduced into Japan. Mammal Study. 32:72-82.

303 Pulliainen, C. and K. Salonen. 1963. On ring barking of pine by squirrel (*Sciurus vulgaris*) in  
304 Finland. Annales Academiae Scientiarum Fennicae (Series A). 72:1-29.

305 Purohit, A., R.K. Maikhuri, K.S. Rao and S. Nautiyal. 2001. Impact of bark removal on survival of  
306 *Taxus baccata* L. (Himalayan yew) in Nanda Devi Biosphere Reserve, Garhwal Himalaya, India.  
307 Curr. Sci. Bangalore 81:586-590.

308 Sharma, D. and S.N. Prasad. 1992. Tree debarking and habitat use by porcupine (*Hystrix indica*  
309 Kerr) in Sariska National Park in Western India. *Mammalia* 56:351-362.

310 Signorile, A.L. and J. Evans. 2007. Damage caused by the American grey squirrel (*Sciurus*  
311 *carolinensis*) to agricultural crops, poplar plantations and semi-natural woodland in Piedmont,  
312 Italy. *Forestry* 80:89-98.

313 Sullivan, T.P. and A. Vyse. 1987. Impact of red squirrel feeding damage on spaced stands of  
314 lodgepole pine in the Cariboo Region of British Columbia. *Can. J. Forest. Res.* 17:666-674.

315 Thorington R.W. jr., Koprowski J.L., Steele M.A., Whatton J.F. (2012). Finlayson's squirrel  
316 *Callosciurus finlaysonii* (Horsfield, 1823). In: *Squirrels of the world*. The Johns Hopkins  
317 University Press, Baltimore, Maryland, USA: p. 139-142.

318 Tubeuf, K. 1897. Diseases of plants induced by cryptogamic parasites: introduction to the study of  
319 pathogenic fungi, slime-fungi, bacteria, and algae. Longmans, Green and Company Press,  
320 London.

321 Williams, F., R. Eschen, A. Harris, D. Djeddour, C. Pratt, R.S. Shaw, S. Varia, J. Lamontagne-  
322 Godwin, S.E. Thomas and S.T. Murphy. 2010. The economic cost of invasive non-native species  
323 on Great Britain. CABI report.

324 Zhu, Y., W. Zhang and X. Zhu. 1990. Bark-stripping damage to forest trees by red-bellied squirrel  
325 (*Callosciurus erythraeus*) in Zhejiang Province. *Acta Theriol. Sin.* 10:276-281.

326 **Tables**

327

328 **Table 1** Incidence of damage caused by bark stripping of the Finlayson's squirrel in Northern Italy and results of Ivlev's Electivity Index analysis

329 (E) for plant species selection on trunk and branches.

330

Species	Trees		Damage on trunks			Surface of exposed wood (%)			Damage on branches			Surface of exposed wood (%)		
	N	N (%)	N	N (%)	E	<50 cm <sup>2</sup>	51-500 cm <sup>2</sup>	>500 cm <sup>2</sup>	N	N (%)	E	<50 cm <sup>2</sup>	51-500 cm <sup>2</sup>	>500 cm <sup>2</sup>
<i>Abies alba</i>	5	1.6	5	100	0.97	20.0	80.0	0.0	5	100	0.97	0.0	100	0.0
<i>Picea excelsa</i>	12	3.9	1	8.3	0.36	0.0	8.3	0.0	0	0.0	-1.00	0.0	0.0	0.0
<i>Cedrus atlantica</i>	3	0.9	1	33.3	0.94	0.0	0.0	33.3	3	100	0.98	0.0	66.7	33.3
<i>Cedrus deodara</i>	16	5.2	3	18.8	0.57	0.0	0.0	18.8	13	81.3	0.88	0.0	25.0	56.3
<i>Pinus strobus</i>	5	1.6	0	0.0	-1.00	0.0	0.0	0.0	5	100	0.98	0.0	80.0	20.0
<i>Pinus nigra</i>	14	4.6	14	100	0.91	0.0	78.6	21.4	14	100	0.91	42.9	57.1	0.0
<i>Pinus laricio</i>	19	6.2	13	68.4	0.83	0.0	26.3	42.1	19	100	0.88	31.6	36.8	31.6
<i>Libocedrus decurrens</i>	4	1.3	2	50.0	0.94	0.0	0.0	50.0	4	100	0.97	25.0	0.0	75.0
<i>Thuja occidentalis</i>	4	1.3	4	100	0.97	0.0	0.0	100	4	100	0.97	0.0	100	0.0
<i>Cephalotaxus cuspidata</i>	14	4.6	10	71.4	0.88	0.0	21.4	50.0	14	100	0.91	0.0	42.9	57.1
<i>Celtis australis</i>	91	29.6	0	0.0	-1.00	0.0	0.0	0.0	91	100	0.54	0.0	100	0.0

<i>Prunus</i> sp.	11	2.3	11	100	0.96	100	0.0	0.0	11	100	0.96	0.0	100	0.0
<i>Tilia cordata</i>	56	18.2	0	0.0	-1.00	0.0	0.0	0.0	56	100	0.69	0.0	100	0.0
<i>Quercus ilex</i>	2	0.7	0	0.0	-1.00	0.0	0.0	0.0	2	100	0.99	0.0	100	0.0
<i>Cercis siliquastrum</i>	4	1.3	0	0.0	-1.00	0.0	0.0	0.0	0	0.0	-1.00	0.0	0.0	0.0
<i>Ilex aquifolium</i>	2	0.7	0	0.0	-1.00	0.0	0.0	0.0	0	0.0	-1.00	0.0	0.0	0.0
<i>Ginkgo biloba</i>	2	0.7	1	50.0	0.97	50.0	0.0	0.0	1	50.0	0.97	50.0	0.0	0.0
<i>Acer saccharinum</i>	2	0.7	0	0.0	-1.00	0.0	0.0	0.0	0	0.0	-1.00	0.0	0.0	0.0
<i>Platanus</i> sp.	42	13.6	0	0.0	-1.00	0.0	0.0	0.0	0	0.0	-1.00	0.0	0.0	0.0
<b>TOTAL</b>	<b>308</b>		<b>65</b>	<b>21.1</b>		<b>4.2</b>	<b>7.8</b>	<b>9.1</b>	<b>242</b>	<b>78.6</b>		<b>4.6</b>	<b>64.9</b>	<b>9.1</b>

331

332

333

334

335

336

337

338

339 **Table 2** Results of Ivlev's Electivity index analysis (E) for plant species selection on trunk and branches, and incidence of damage caused by bark  
 340 stripping of the Finlayson's squirrel in the sites in the Southern area of introduction.

341

Species	Damage on trunks			Surface of exposed wood (%)			Damage on branches			Surface of exposed wood (%)		
	N	N(%)	E	<50 cm <sup>2</sup>	51-500 cm <sup>2</sup>	>500 cm <sup>2</sup>	N	N(%)	E	<50 cm <sup>2</sup>	51-500 cm <sup>2</sup>	>500 cm <sup>2</sup>
<i>Pinus halepensis</i>	27	12.9	0.18	5	18.5	0.18	8	29.6	0.39	11.1	7.4	11.1
<i>Pinus sp.</i>	12	5.7	0.18	1	8.3	0.18	1	8.3	0.18	8.3	0.0	0.0
<i>Quercus ilex</i>	91	43.5	-0.86	3	3.3	-0.86	13	14.3	-0.51	3.3	6.6	4.4
<i>Quercus virgiliana</i>	44	21.1	-0.40	4	9.1	-0.40	29	65.9	0.52	9.1	29.6	27.3
<i>Ceratonia siliqua</i>	25	12.0	0.63	13	52.0	0.63	23	92.0	0.77	8.0	16.0	68.0
<i>Olea europaea</i>	10	4.8	0.35	1	10.0	0.35	6	60.0	0.85	40.0	20.0	0.0
<b>TOTAL</b>	<b>209</b>			<b>27</b>	<b>12.9</b>		<b>80</b>	<b>38.3</b>		<b>8.1</b>	<b>12.9</b>	<b>17.2</b>

342

343

344

345 **Figure legends**

346

347 **Figure 1** Distribution of the Finlayson's squirrel in Italy.

348

349 **Figure 2** Trend of bark-stripping damage by Finlayson's squirrel to different plant categories,  
350 before (T<sub>0</sub>, black bars), one month after (T<sub>1</sub>, white bars) and nine months after the phytosanitary  
351 intervention at Acqui Terme (T<sub>2</sub>, grey bars).

352

353 **Figure 3** Percentages of bark-stripping damage in nine wood patches in Southern Italy; black bars  
354 refer to branches, grey ones to the trunk. Location 0 refers to the site of first introduction of the  
355 Finlayson's squirrel; sites 1, 4, 6 are northwards, the others are southwards.

356

357 **Figure 4** Percentage of bark-stripping at trunk and branch levels. Black bars refer to the Northern  
358 Italy study site, grey ones to Southern Italy.


359

360




361

362




363

364




365

366



Surface of debarked trees on trunks



Surface of debarked trees on branches

367

368