

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Effect of bacterial canker caused by *Pseudomonas syringae* pv. *actinidiae* on postharvest quality and rots of kiwifruit 'Hayward'**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1581072> since 2017-05-12T13:42:47Z

*Published version:*

DOI:10.1016/j.postharvbio.2015.11.010

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in POSTHARVEST BIOLOGY AND TECHNOLOGY, 113, 2016, 10.1016/j.postharvbio.2015.11.010.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.postharvbio.2015.11.010

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S092552141530171X>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1581072>

1 **Effect of bacterial canker caused by *Pseudomonas syringae* pv. *actinidiae* on postharvest quality**  
2 **and rots of kiwifruit ‘Hayward’**

3

4 Simona Prencipe<sup>1,2</sup>, Luca Nari<sup>3</sup>, Graziano Vittone<sup>3</sup>, Maria Lodovica Gullino<sup>1,2</sup>, Davide Spadaro<sup>1,2\*</sup>

5

6 <sup>1</sup> DISAFA – Dipartimento di Scienze Agrarie, Forestali ed Alimentari, Università degli Studi di  
7 Torino, Largo P. Braccini 2, 10095 Grugliasco (TO), Italy.

8 <sup>2</sup> Centro di competenza per l’Innovazione in campo agro-ambientale (AGROINNOVA), Università  
9 degli Studi di Torino, Largo P. Braccini 2, 10095 Grugliasco (TO), Italy.

10 <sup>3</sup> Agrion - Fondazione per la ricerca l’innovazione e lo sviluppo tecnologico dell’agricoltura  
11 piemontese, Via Falicetto 24, 12030 Manta (Cn), Italy.

12

13

14

15

16

17

18

19 **\* Corresponding author**

20 Davide Spadaro

21 E-mail: [davide.spadaro@unito.it](mailto:davide.spadaro@unito.it)

22 Telephone: +39-011-6708942

23 Fax: +39-011-6709307

24 **Abstract**

25 Kiwifruits produced from plants infected by *Pseudomonas syringae* pv *actinidiae* (*Psa*), the agent of  
26 bacterial canker, do not show visible disease symptoms and can be commercialized. Firmness, total  
27 soluble solids (TSS), and titratable acidity (TA) of *Actinidia deliciosa* ‘Hayward’ fruit, infected or  
28 not by *Psa*, were quantified, for two years, at harvest, at 90 days storage, and during shelf life.  
29 Analysis of dry matter (DM) and calcium at harvest, and postharvest rots, caused by *Botrytis cinerea*,  
30 during shelf life were also assessed. DM was higher and calcium content was lower in the kiwifruit  
31 from orchards infected by bacterial canker. During storage, firmness decreased and TSS increased  
32 rapidly in fruit from infected orchards. These parameters, together with the lower TA, favoured the  
33 higher susceptibility to grey mould rots. Besides the incidence of bacterial canker, climate conditions  
34 throughout the cropping seasons greatly affected the postharvest quality of fruit. Also the storage  
35 conditions modified the fruit quality: controlled atmosphere (CA) and 1-MCP treatment delayed fruit  
36 softening compared to normal atmosphere. TSS on fruit from healthy orchards was not influenced by  
37 the storage conditions. Fruit from *Psa* diseased orchards showed significantly lower TSS in CA  
38 storage and after 1-MCP treatment, compared with normal atmosphere storage. In this work, we  
39 showed that the presence of bacterial canker affects postharvest quality, shelf life, and susceptibility  
40 to postharvest rots of fruit.

41

42 *Keywords:*

43 *Actinidia deliciosa*, bacterial canker, *Botrytis cinerea*, controlled atmosphere, 1-methylcyclopropene,  
44 *Pseudomonas syringae*.

## 1. Introduction

The global production of kiwifruit was over 1.4 million tons in 2012, and Italy, with 384,000 tons and 24,800 hectares (FAOSTAT, 2014), produced more kiwifruit than any other country, apart from China. ‘Hayward’ is the most important kiwifruit cultivar, because of size, taste, longer storage, and high antioxidant capacity and vitamin C (Testolin and Ferguson, 2009).

*Pseudomonas syringae* pv. *actinidiae* (*Psa*), the agent of bacterial canker on kiwifruit, was first isolated and described in Japan in 1984 (Takikawa et al., 1989). In Italy, Scortichini (1994) first reported the presence of *Psa* on ‘Hayward’ kiwifruit in 1994, but the first epidemic outbreak of bacterial canker, caused by the highly virulent biovar 3 of *Psa*, occurred in 2008 (Balestra et al., 2009; Vanneste et al., 2013). During 2010-2014, around 1,922 hectares of kiwifruit were uprooted in the different regions of Italy due to bacterial canker (personal communication). The disease quickly reached pandemic proportions by spreading to France, Spain, Portugal, Switzerland, New Zealand, Chile, Turkey, South Korea, and Japan (EPPO, 2012). Kiwifruits from diseased plants do not show disease symptoms and can be commercialized. The presence of *Psa* could affect postharvest quality, storage shelf life and susceptibility to postharvest rots of fruit, which depend on the physiological state of the fruits, similarly as described for *Xylella fastidiosa* on grapevine (Choi et al., 2013) and citrus (Ribeiro et al., 2003).

In *Actinidia deliciosa* ‘Hayward’, total soluble sugar (TSS; percentage) is a physiological marker from fruit development to ripening, and a value of 6.2% has been established as optimal for harvest (Burdon et al., 2013). Also dry matter (DM) is an important parameter used as maturity marker predictive of fruit flavour (Patterson and Currie, 2011). DM and TSS are closely related parameters on ripe kiwifruit. Other chemical parameters, commonly used as markers of fruit quality, are firmness and sugar:acid ratio. The physiological parameters of kiwifruits are influenced by complex interactions between pedoclimatic conditions, agricultural practices, harvesting maturity, ethylene production, and accumulation of calcium (Montanaro et al., 2012; Burdon et al. 2014).

71 Postharvest quality and rots of kiwifruit could also be related to concentration of minerals, such as  
72 calcium, magnesium, and nitrogen (Spadaro et al., 2010). Calcium is an important macronutrient in  
73 plants, involved in development and softening of fruit and it influences their quality. It is an essential  
74 nutrient due to its function in stabilization of membrane systems, increase of cell turgor, resistance to  
75 cell wall degradation by enzymes, and reduction of fruit transpiration (Ghani et al., 2011). The  
76 calcium content in fruits is closely related to the microclimatic conditions around the fruits, which  
77 influence their transpiration (Xiloyannis et al., 2010).

78 Kiwifruit can be stored over 6 months at  $0\pm 1^{\circ}\text{C}$  (Feng et al., 2006). During storage, fruit rots can  
79 cause serious economic losses (Spadaro et al., 2011), and *Botrytis cinerea*, causal agent of grey  
80 mould, is the most important postharvest pathogen (Pyke et al., 1994).

81 Fruit quality losses are mainly due to the fruit metabolic activity during postharvest, which is  
82 influenced by the storage conditions. Cold storage, controlled atmosphere (CA), and use of  
83 postharvest treatments based on ethylene inhibitors, such as 1-methylcyclopropene (1-MCP), could  
84 be used to increase the fruit storage life and to extend the ability to supply the market all year round,  
85 while maintaining the main fruit characteristics required by consumers (Watkins, 2006).

86 In order to evaluate the effects of *Psa* on postharvest quality and rots of kiwifruit, the quality  
87 parameters and the incidence of grey mould were assessed on kiwifruits harvested from healthy and  
88 *Psa* infected orchards, and stored under normal or controlled atmosphere. Orchards were considered  
89 healthy throughout the manuscript when bacterial canker symptoms were not visible or negligible.  
90 The physiochemical parameters considered were firmness, TSS, titratable acidity, DM, and calcium  
91 content. In addition, we verified the effects of 1-MCP treatment on postharvest quality and storage  
92 rots of kiwifruit from healthy and *Psa* infected orchards.

93

## 94 **2. Materials and methods**

95

### 96 *2.1. Fruit and bacterial canker incidence*

97 Twelve orchards of *A. deliciosa* ‘Hayward’, six infected by *Psa* and six healthy, were selected in  
98 north western Italy (Piedmont region) for harvesting kiwifruit for two years (2013 and 2014).  
99 The number of vines, the age, and the agronomic practices carried out are reported for each orchard  
100 of *Actinidia deliciosa* “Hayward” (supplementary Table 1). Three repetitions of 100 plants were  
101 observed in each orchard to calculate the incidence (%) of *Psa*-infected plants. Plants were considered  
102 symptomatic when spots with a chlorotic halo were observed on at least two leaves (Table 1). To  
103 verify the presence of *Psa*, from symptomatic leaves collected from all orchards, isolation was carried  
104 out by using Pseudomonas Selective Agar with CFC supplement (Merck, Germany). Single colonies  
105 were grown in Luria Bertani broth (Merck) on a rotary shaker (150 rpm) at 27°C for two days. DNA  
106 was extracted by using Instagene™ Matrix (Biorad, United States) according to the manufacturer’s  
107 protocol. Primers PsaF1/PsaR2 and PsaF3/PsaR4 were used for PCR amplification, following Rees-  
108 George et al. (2010). The obtained amplicons were run on a gel and compared with positive controls.  
109 The genomic DNA was also used to amplify the 16S ribosomal gene with the universal primers 27f  
110 and 1492r (Weisburg *et al.*; 1991) and the PCR products were sequenced and compared with those  
111 deposited in GenBank using the BLAST programme (Zhang et al., 2000). We also determined the  
112 virulence and the haplotype through *cts* sequencing, according to Vanneste et al. (2010).  
113 Five fruit were harvested from 100 vines per orchard. Fruits were stored in controlled conditions or  
114 in commercial packinghouses.

115

### 116 2.3. *Trials in controlled conditions*

117 Kiwifruits from every orchard (500 fruit, around 45 kg) were divided in three boxes (15 kg each).  
118 Fruits were stored – within 60 h from harvest – in experimental cold chambers for 90 days at 1.0°C  
119 ±1 and RH 98%. After 90 days, fruits were transferred at 10°C for 20 days shelf life, to favour the  
120 pathogen development. Firmness, total soluble sugar (TSS), and titratable acidity (TA) were  
121 measured at harvest, end of storage, and after shelf life. Grey mould incidence was assessed after

122 shelf life. The trial was carried out for two years (2013 and 2014). Statistical analysis was performed  
123 using Student's t-test at 95%, 99%, and 99.9% confidence level.

124

#### 125 *2.4. Trials in packinghouses*

126 During the first year, the fruit from one healthy orchard (Healthy 6 in Table 1) and one *Psa* diseased  
127 orchard (Diseased 6 in Table 1) were divided in three replications of 15 kg boxes. The fruit boxes  
128 were stored – one day after harvest – in two commercial packinghouses located in Piedmont, northern  
129 Italy (Sanifrutta, Castigliole Saluzzo (CN) and Rosatello, Lagnasco (CN)), under three conditions:  
130 normal atmosphere ( $1.0 \pm 1$  °C; RH 98%), controlled atmosphere alone ( $-0.8 \pm 0.8$  °C; RH 98%; O<sub>2</sub>  
131  $2.5 \pm 0.5$  kPa, CO<sub>2</sub>  $3.5 \pm 0.5$  kPa), and controlled atmosphere after treatment with SmartFresh™ Powder  
132 (a.i.: 3.3%, 1-MCP; Rohm & Hass). Kiwifruit were treated in 37 L plastic barrels with  $0.5 \mu\text{L L}^{-1}$   
133 SmartFresh™ according to manufacturer's instruction at room temperature for 24 hours before  
134 closing the storage room. Fruits were stored in cold chambers for 90 days, and later they were kept  
135 in simulated shelf life conditions at 10°C for 20 days. Firmness and total soluble sugar (TSS) were  
136 measured at harvest, end of storage, and shelf life. Statistical analysis was carried out using Student's  
137 t-test at 95%, 99% and 99.9% confidence level, depending on the experiment.

138

#### 139 *2.5. Quality parameters and chemical analysis*

140 Firmness, expressed in N, was measured using penetrometer FRUIT TEST™ FT 327 (EFFEGI,  
141 Alfonsine, Italy) with a 8 mm diameter plunger tip on two sides of fruits. TSS (%) was measured  
142 using the digital refractometer NR-151 (DBR95, Singapore), by squeezing one drop of juice from  
143 each end of kiwifruit. TA was obtained by titration with NaOH 0.1 M (pH 8.0) using 6 mL clear juice  
144 diluted with distilled water up to 30 mL final volume. Acidity was calculated as percent of citric acid.  
145 Each parameter was measured in four fruit per box ( $n = 12$ ) per orchard. The analyses were performed  
146 twice. At harvest, percentage of calcium and dry matter (Table 1) were determined by Laboratorio  
147 Agrochimico Regionale of Piedmont Region (Ceva, Italy).

148

## 149 2.6. *Phytopathological analysis*

150 The incidence of *Botrytis cinerea* was assessed on the fruits of the trial in controlled conditions. Fruit  
151 showing stem-end rots were used to isolate *B. cinerea* on Potato dextrose agar (PDA, Merck,  
152 Germany) with 0.025 g L<sup>-1</sup> streptomycin (Applichem, Germany). Isolations were made from 10 fruits  
153 per repetition and 3 repetitions per orchard. Morphological identification of the pathogen species,  
154 from monoconidial cultures, was confirmed with PCR amplification of ITS region using primer  
155 ITS1/ITS4 (White et al., 1990) following standard protocols. The sequences obtained were compared  
156 with those already deposited in GenBank.

157

## 158 3. Results

159

### 160 3.1. *Incidence of Psa and chemical analyses at harvest*

161

162 The average incidence (%) of bacterial canker on kiwifruits plants, in 2013, was 1.3% for healthy  
163 orchards and 86.5% for diseased ones, while in 2014 it was 0.5% and 58.0% for healthy and diseased  
164 orchards, respectively (Table 1). No differences between healthy and diseased orchards were found  
165 with respect to agronomic practices, including vine spacing, irrigation, training system, pesticide  
166 treatments, shelter, rootstock, number of pollinators, and girdling (Supplementary Table 1). Also the  
167 average age and height were not significantly different between healthy and diseased orchards. The  
168 fertilizer regime on healthy orchards was 80 to 90 nitrogen unit per hectare, while it was reduced in  
169 *Psa*-diseased orchards to 50-60 nitrogen unit per hectare. The lower nitrogen input did not affect the  
170 final nitrogen content of fruit, which was 0.78% and 0.71% in healthy and diseased fruit in 2013, and  
171 0.83% and 0.79%, respectively, in 2014 (data not shown). Plants showing bacterial canker symptoms  
172 were confirmed for the presence of *Psa*. Bacterial colonies isolated from symptomatic tissues were  
173 used for amplification and sequencing of the 16S ribosomal gene. All the sequences showed 100%

174 identity with other sequences of *Psa* and two of them were deposited in GenBank (accession numbers  
175 KP794939 and KP794940). The isolates confirmed their virulence on plantlets of kiwifruit ‘Hayward’  
176 and they showed to belong to the haplotype I of gene *cts* (data not shown).

177 At harvest, fruit from diseased orchards showed on average higher DM and lower calcium content,  
178 in both years, compared with those from healthy orchards (Table 1). In 2013, DM was 13.8% for  
179 healthy fruits and 16.3% for diseased ones ( $P \leq 0.001$ ), while in 2014 it was 14.7% and 15.6%  
180 ( $P \leq 0.05$ ), respectively. Calcium content (%) in 2013 was 0.32% for healthy fruits and 0.24% for  
181 diseased ones ( $P \leq 0.05$ ), while in 2014 it was 0.39% and 0.34%, respectively.

182 Differences in climate between the two years of study were relevant (Supplementary Figures 1 and  
183 2). The cropping season 2013 was characterized by lower minimum temperatures in winter, ranging  
184 from  $-3^{\circ}\text{C}$  to  $1^{\circ}\text{C}$ , and higher relative humidity, with precipitation concentrated in spring, with a total  
185 of 343 mm. In autumn and summer, precipitation and temperatures were similar to the average of the  
186 last years. In contrast, the cropping season 2014 showed higher minimum temperatures in winter,  
187 ranging from  $-1^{\circ}\text{C}$  to  $3^{\circ}\text{C}$ , without frosts, and warmer minimum and maximum pre-harvest  
188 temperatures in autumn. The harvesting period was characterized by higher precipitation, with 404  
189 mm, and a maximum of 290 mm in November 2014.

190

### 191 3.2. Fruit quality in controlled conditions

192 The quality parameters measured for the trials performed in 2013 and 2014 in controlled conditions  
193 are reported in Table 2 and Table 3, respectively. Firmness and TA values decreased from harvest to  
194 shelf life, while TSS increased following fruit ripening. Fruits from diseased orchards showed lower  
195 firmness and TA, and higher TSS, compared with healthy fruits. Significant differences in TSS  
196 ( $P \leq 0.001$ ) were found at the end of storage in 2013 (Table 2) between the fruits from healthy (11.9%)  
197 and *Psa* infected orchards (13.8%). Also TA was significantly different ( $P \leq 0.001$ ) between fruit from  
198 healthy (0.84%) and diseased (0.48%) orchards. During simulated shelf life, TSS on healthy and  
199 diseased kiwifruit increased, respectively, to 11.6% and 14.2% ( $P \leq 0.001$ ).

200 During the second year of experimentation (Table 3), significant differences ( $P\leq 0.001$ ) were found  
201 at the end of storage with TSS values of 12.2% for healthy and 13.5% for diseased orchards, and in  
202 shelf life ( $P\leq 0.05$ ) with, respectively, 13.4% and 14.2%. For TA, significant differences occurred  
203 only at harvest ( $P\leq 0.001$ ) with values of 2.68% and 1.46% respectively, on fruits from healthy and  
204 diseased orchards. Firmness was significantly different at the end of storage with values of 6.94 N  
205 and 5.36 N for fruits from healthy and diseased orchards ( $P\leq 0.01$ ), and in shelf life with values,  
206 respectively, of 2.87 N and 1.76 N ( $P\leq 0.05$ ).

207

### 208 *3.3. Fruit quality in semi-commercial conditions*

209 The results of the trials performed in commercial packinghouses are shown in Table 4. In normal  
210 atmosphere storage (NA), no differences were found for firmness at the end of storage and in shelf  
211 life, with values of 22.5 N and 12.5 N for healthy fruits and 18.9 N and 12.1 N for fruits from diseased  
212 orchards. In contrast, significant differences ( $P\leq 0.001$ ) were observed for TSS at the end of storage  
213 and in shelf life, with values of 11.5% and 12.0% for healthy fruits and 13.5% and 14.4% for diseased  
214 fruits, respectively.

215 The results under controlled atmosphere (CA) showed, the same trend, with a decrease of firmness  
216 and increase of TSS. Firmness values were significantly different after simulated shelf life ( $P\leq 0.05$ ),  
217 with 5.49 N for healthy fruits and 4.61 N for diseased ones. Significant TSS differences were observed  
218 both at the end of storage and after shelf life, with values of 11.6% and 12.8% for healthy fruits, and  
219 of 12.4% and 13.7% for fruits from diseased orchards, respectively ( $P\leq 0.001$ ).

220 The fruits stored under controlled atmospheres after treatment with 1-MCP showed significant  
221 differences both for firmness ( $P\leq 0.05$ ) and TSS ( $P\leq 0.001$ ) at the end of storage and after simulated  
222 shelf life. Firmness of healthy fruits, with values of 69.1 N at the end of storage and 18.9 N after shelf  
223 life, were significantly higher compared with fruits from diseased orchards, with values of 65.2 N and  
224 16.7 N, respectively. TSS was 11.7% for healthy fruits and 12.4% for diseased fruits at the end of  
225 storage, while, after shelf life it was 12.6% and 13.8%, respectively.

226

### 227 3.4. Postharvest rots

228 Phytopathological survey showed higher incidence of postharvest rots in fruit from diseased orchards  
229 with 23% rotten fruit in the first year and 45% in the second one, compared with 2% and 16% in fruit  
230 from healthy orchards ( $P \leq 0.001$ ). Some sequences obtained by PCR amplification of the internal  
231 transcribed spacer (ITS1, 5.8S gene and ITS2) of the fungal pathogen isolated from diseased fruit  
232 (Accession numbers KP794937 and KP794938) showed 100% identity to sequences deposited in  
233 GenBank as *Botryotinia fuckeliana*, the teleomorph of *B.cinerea*, which is the commonly accepted  
234 name of the pathogen.

235

## 236 4. Discussion

237

238 Kiwifruit quality and diseases could be influenced by climatic conditions during the cropping season.  
239 As a consequence of a colder winter and rainy spring, 2013 was characterized by a higher incidence  
240 of *Psa*, compared to 2014, with milder winter and drier spring. According to Serizawa and Ichikawa  
241 (1993a, b), favourable climatic conditions for *Psa* are cool temperatures, persistent rain, and high  
242 humidity, which characterized the 2013 season.

243 Changes in physiochemical parameters were observed during kiwifruit storage. Kiwifruit TA  
244 decreased gradually during storage (Park et al., 2006), because of organic acid consumption for  
245 respiration (Moing et al., 2001), while TSS increased over storage, due to starch conversion into sugar  
246 (Wills and Rigney, 2007).

247 Plants respond to biotic stresses, i.e a plant pathogen, with modifications in the physio-chemical  
248 parameters of sink organs, such as fruit or tuber, as reported for potato infected by *Erwinia carotovora*  
249 (Nourian et al., 2002). Also kiwifruit showed an influence in the chemical composition and the quality  
250 of fruits due to the presence of *Psa*. The chemical analyses performed revealed substantial differences  
251 between kiwifruit from orchards infected or not by bacterial canker. Higher DM and lower calcium

252 concentration were observed in fruit from diseased plants. DM modification could be related to lower  
253 water uptake from roots linked to the presence of *Psa* in the vascular system of kiwifruit (Spinelli et  
254 al., 2011; Donati et al., 2014). The correlation between the presence of a plant pathogen in the vascular  
255 system and its influence in plant transpiration is reported for several pathosystems (Ribeiro et al.,  
256 2003; Choi et al., 2013). DM is also related to climatic conditions, orchard location, fertilization, and  
257 harvest time (Spadaro et al., 2010).

258 Lower calcium content contributes to favourable conditions to infections and physiochemical  
259 disorder and, consequently, higher incidence of postharvest diseases (Biggs, 1999). Gerasopoulos et  
260 al. (1996) showed positive correlation between calcium and firmness in many fruits. Lower calcium  
261 content in kiwifruit from diseased plants could indirectly compromise the cell wall integrity of fruits,  
262 increasing the susceptibility to postharvest pathogens (Quiles et al., 2004). Calcium could also  
263 contribute to lower susceptibility to pathogens, by reducing the germination of fungal species, as  
264 reported by Droby et al. (1997) for *Penicillium digitatum*. For these reasons, probably, a higher  
265 presence of *B. cinerea* rots was observed in fruit from diseased orchards, both in 2013 and 2014,  
266 similarly to Barboni et al. (2010).

267 In the fruit from *Psa* diseased orchards, TA was lower and TSS was higher compared to the fruit from  
268 healthy orchards. The bacterial pathogen presence could affect the fruit, which has higher respiration  
269 rate, higher sugar solubilisation, and, consequently, higher TSS (Shahkoomahally and Ramezani,  
270 2015).

271 During the second year, TA in fruits, particularly in the healthy ones, was significantly higher  
272 compared to 2013, due to higher temperatures. Organic acid metabolism occurring from harvest to  
273 storage is influenced by temperatures. Higher temperatures induce higher fruit respiration rate and  
274 enzymatic activity (Walton and De Jong, 1990), that reduce TA (Marsh et al., 2004).

275 Moreover, higher grey mould incidence and lower firmness in the second year is related to the  
276 abundant precipitation of November 2014. As reported by Lysiak (2013), high precipitation at harvest

277 favoured postharvest diseases on fruit, because the rain contributed to reduce the firmness and the  
278 wet environment could favour the pathogen spore penetration.

279 Significant differences in fruit quality were also related to storage conditions. CA delays fruit  
280 softening (Manolopoulou and Papadopoulou, 1998), which is related to the combined effect of lower  
281 respiration and transpiration (Hertog et al., 2004). Kiwifruit treated with 1-MCP had, as expected,  
282 even higher firmness compared with CA-stored fruit, as reported by Watkins (2006). TSS was not  
283 significantly different on healthy fruit stored under NA, CA or after 1-MCP treatment. Fruit from *Psa*  
284 diseased orchards showed significantly lower TSS in CA storage and after 1-MCP treatment,  
285 compared with normal atmosphere storage. Throughout the experiment in different storage  
286 conditions, firmness was lower and TSS was higher in fruit from *Psa* diseased orchards compared  
287 with fruit from healthy orchards.

288 In conclusion, kiwifruit produced from plants infected by bacterial canker did not show visible disease  
289 symptoms. For the first time, we showed that the presence of *Psa* greatly affects the postharvest  
290 quality, the shelf life, and the susceptibility to postharvest rots of fruit. Dry matter was higher and  
291 calcium content was lower in the kiwifruit from diseased orchards. During storage faster firmness  
292 decrease and TSS increase were registered. These parameters and the lower TA favoured the  
293 susceptibility to postharvest rots, and particularly to *B. cinerea*. Besides the incidence of bacterial  
294 canker, also climate conditions throughout the cropping season greatly affected the postharvest  
295 quality of fruit. The choice of the storage conditions, including controlled atmosphere and 1-MCP  
296 treatment, could help to maintain the shelf life of kiwifruit, with higher firmness and lower TSS.  
297 Future work will focus on the effect of bacterial canker on some physiological parameters of kiwifruit,  
298 such as respiration and transpiration rates.

299

### 300 **Acknowledgments**

301 The Authors wish to thank the financial support of Piedmont Region Union in the framework of the  
302 project “PRO.ACT.IN.” (PSR FEASR 2007/2013, European Fund for Rural Development, Measure

303 124, Action 1). The Authors gratefully acknowledge the orchards and packinghouses owners, and  
304 Michele Girauda who contributed to the experimental trials. The Authors acknowledge also Prof.  
305 Angelo Garibaldi for his accurate revision of the manuscript.

306

307 **References**

- 308 Balestra, M., Mazzaglia, A., Quattrucci, A., Renzi, M., Rossetti, A., 2009. Occurrence of  
309 *Pseudomonas syringae* pv. *actinidiae* in Jin Tao kiwi plants in Italy. *Phytopathol. Mediterr.* 48, 299-  
310 301.
- 311 Barboni, T., Cannac, M., Chiaramonti, N., 2010. Effect of cold storage and ozone treatment on  
312 physicochemical parameters, soluble sugars and organic acids in *Actinidia deliciosa*. *Food Chem.*  
313 121, 946-951.
- 314 Biggs, A.R., 1999. Effects of calcium salts on apple bitter rot caused by two *Colletotrichum*  
315 spp. *Plant Dis.* 83, 1001-1005.
- 316 Burdon, J., Lallu, N., Pidikala, P., Barnett, A. 2013. Soluble solids accumulation and  
317 postharvest performance of ‘Hayward’ kiwifruit. *Postharvest Biol. Technol.* 80, 1-8.
- 318 Burdon, J., Pidakala, P., Martin, P., McAtee, P.A., Boldingh, H.L., Hall, A., Schaffera, R.J.,  
319 2014. Postharvest performance of the yellow-fleshed “Hort16A” kiwifruit in relation to fruit  
320 maturation. *Postharvest Biol. Technol.* 92, 98-106.
- 321 Choi, H.-K., Iandolino, A., da Silva, F.G., Cook, D.R., 2013. Water Deficit Modulates the  
322 response of *Vitis vinifera* to the Pierce’s Disease pathogen *Xylella fastidiosa*. *Mol. Plant Microbe*  
323 *Interact.* 26, 643–657.
- 324 Donati, I., Buriani, G., Cellini, A., Mauri, S., Costa, G., Spinelli, F., 2014. New insights on the  
325 bacterial canker of kiwifruit (*Pseudomonas syringae* pv. *actinidiae*). *J. Berry Res.* 4, 53–67.
- 326 Droby, S., Wisniewski, M.E., Cohen, L., Weiss, B., Touitou, D., Eilam, Y., Chalutz, E., 1997.  
327 Influence of CaCl<sub>2</sub> on *Penicillium digitatum*, grapefruit peel tissue, and biocontrol activity of *Pichia*  
328 *guilliermondii*. *Phytopathology* 87,310-315.
- 329 EPPO, 2012. *Pseudomonas syringae* pv. *actinidiae*. Bacterial canker of kiwifruit. Available at:  
330 [http://www.eppo.int/QUARANTINE/Alert\\_List/bacteria/P\\_syringae\\_pv\\_actinidiae.htm](http://www.eppo.int/QUARANTINE/Alert_List/bacteria/P_syringae_pv_actinidiae.htm) Accessed  
331 on August 28, 2015.

332 FAOSTAT, 2014. Available at: <http://faostat3.fao.org/faostat-gateway/go/to/browse/Q/QC/E>  
333 Accessed on August 28, 2015.

334 Feng, J., Maguire, K.M., MacKay, B.R., 2006. Discriminating batches of “Hayward” kiwifruit  
335 for storage potential. *Postharvest Biol. Technol.* 41, 128-134.

336 Gerasopoulos, D., Chouliaras, V., Lionakis, S., 1996. Effects of preharvest calcium chloride  
337 sprays on maturity and storability of Hayward kiwifruit. *Postharvest Biol. Technol.* 7, 65-72.

338 Ghani, M.A., Awang, Y., Sijam, K., 2011. Disease occurrence and fruit quality of pre-harvest  
339 calcium treated red flesh dragon fruit (*Hylocereus polyrhizus*). *Afr. J. Biotechnol.* 10, 1550-1558.

340 Hertog, M.L.A.T.M., Nicholson, S.E., Jeffery, P.B., 2004. The effect of modified atmospheres  
341 on the rate of firmness change of ‘Hayward’ kiwifruit. *Postharvest Biol. Technol.* 31, 251-261.

342 Lysiak, G., 2013. The influence of harvest maturity and basic macroelement content in fruit on  
343 the incidence of diseases and disorders after storage of the ‘Ligol’ apple cultivar. *Folia Hort.* 25, 31-  
344 39.

345 Manolopoulou, H., Papadopoulou, P., 1998. A study of respiratory and physico-chemical  
346 changes of four kiwi fruit cultivars during cool storage. *Food Chem.* 63, 529-534.

347 Marsh, K., Attanayake, S., Walker, S., Gunson, A., Boldingh, H., MacRae, E., 2004. Acidity  
348 and taste in kiwifruit. *Postharvest Biol. Technol.* 32, 159-168.

349 Moing, A., Renaud, C., Gaudillere, M., Raymond, P., Roudeillac, P., Denoyes-Rothan, B.,  
350 2001. Biochemical changes during fruit development of four strawberry cultivars. *J. Am. Soc. Hort.*  
351 *Sci.* 126, 394-403.

352 Montanaro, G., Dichio, B., Xiloyannis, C., Lang, A., 2012. Fruit transpiration in kiwifruit:  
353 environmental drivers and predictive model. *AoB Plants*. doi: 10.1093/aobpla/pls036

354 Nourian, F., Kushalappa, A.C., Ramaswamy, H.S., 2002. Physical, physiological and chemical  
355 changes in potato as influenced by *Erwinia carotovora* infection. J. Food Process. Pres. 26, 339-359.

356 Park, Y.S., Jung, S.T., Gorinstein, S., 2006. Ethylene treatment of 'Hayward' kiwifruits  
357 (*Actinidia deliciosa*) during ripening and its influence on ethylene biosynthesis and antioxidant  
358 activity. Sci. Hortic. 108, 22-28.

359 Patterson, K.J., Currie, M.B., 2011. Optimizing kiwifruit vine performance for high  
360 productivity and superior fruit taste. Acta Hortic. 913, 257-268.

361 Pyke, N.B., Manktelow, D.G., Elmer, P.A.G., Tate, K.G., 1994. Postharvest dipping of  
362 kiwifruit in iprodione to control stem-end rot caused by *Botrytis cinerea*. New Zeal. J. Crop Hort. 22,  
363 81-86.

364 Quiles, A., Hernando, I., Perez-Munuera, I., Llorca, E., Larrea, V., Lluch, M.A. 2004. The  
365 effect of calcium and cellular permeabilization on the structure of the parenchyma of the osmotic  
366 dehydrated 'Granny Smith' apple. J. Sci. Food Agr. 84, 1765-1770.

367 Rees-George, J., Vanneste, J.L., Cornish, D.A., Pushparajah, I.P.S., Yu, J., Templeton, M.D.,  
368 Everett, K.R., 2010. Detection of *Pseudomonas syringae* pv *actinidiae* using polymerase chain  
369 reaction (PCR) primers based on the 16S-23S rDNA intertranscribed spacer region and comparison  
370 with PCR primers based on other gene regions. Plant Pathol. 59, 453-464.

371 Ribeiro, R.V., Machado, E.C., Oliveira, R.F., 2003. Early photosynthetic responses of sweet  
372 orange plants infected with *Xylella fastidiosa*. Physiol. Mol. Plant Pathol. 62, 167-173.

373 Scortichini, M. 1994. Occurrence of *Pseudomonas syringae* pv. *actinidiae* on kiwifruit in Italy.  
374 Plant Pathol. 43, 1035-1038.

375 Serizawa, S., Ichikawa, T., 1993a. Epidemiology of bacterial canker of kiwifruit 4. Optimum  
376 temperature for disease development of new canes. Ann. Phytopathol. Soc. Jpn. 59, 694-701.

377 Serizawa, S., Ichikawa, T., 1993b. Epidemiology of bacterial canker of kiwifruit: 2. The most  
378 suitable times and environments for infection on new canes. Ann. Phytopathol. Soc. Jpn. 59, 460-  
379 468.

380 Shahkoomahally, S., Ramezani, A., 2015. Changes in physico-chemical properties related to  
381 quality of kiwifruit (*Actinidia Deliciosa* cv. Hayward) during cold storage. Int. J. Fruit Sci. 15, 187-  
382 197.

383 Spadaro, D., Galliano, A., Pellegrino, C., Gilardi, G., Garibaldi, A., Gullino, M.L., 2010. Dry  
384 matter and mineral composition, together with commercial storage practices, influence the  
385 development of skin pitting caused by *Cadophora luteo-olivacea* on kiwifruit 'Hayward'. J. Plant  
386 Pathol. 92, 339-346.

387 Spadaro, D., Pellegrino, C., Garibaldi, A., Gullino, M.L., 2011. Development of SCAR primers  
388 for the detection of *Cadophora luteo-olivacea* on kiwifruit and pome fruit and of *C. malorum* on  
389 pome fruit. Phytopathol. Mediterr. 50, 430-441.

390 Spinelli, F., Donati, I., Vanneste, J.L., Costa, M., Costa, G., 2011. Real time monitoring of the  
391 interactions between *Pseudomonas syringae* pv. *actinidiae* and *Actinidia* species. Acta Hort. 913,  
392 461-465.

393 Takikawa, Y., Serizawa, S., Ichikawa, T., 1989. *Pseudomonas syringae* pv. *actinidiae* pv. nov.:  
394 the causal bacterium of canker of kiwifruit in Japan. Ann. Phytopathol. Soc. Jpn. 55, 437-444.

395 Testolin, R., Ferguson, A.R., 2009. Kiwifruit production and marketing in Italy. New Zeal. J.  
396 Crop Hort. 37, 1-32.

397 Vanneste, J.L., You, J., Cornish, D.A., 2010. Molecular characterization of *Pseudomonas*  
398 *syringae* pv. *actinidiae* isolated from the recent outbreak of bacterial canker in Italy. New Zeal. Plant  
399 Prot. 63, 7-14.

400 Vanneste, J.L., You, J., Cornish, D.A., 2013. Identification, virulence, and distribution of two  
401 biovars of *Pseudomonas syringae* pv. *actinidiae* in New Zealand. Plant Dis. 97, 708-719.

402 Walton, E.F., De Jong, T.M., 1990. Growth and compositional changes in kiwifruit berries from  
403 three Californian locations. Ann. Bot. 66, 285-298.

404 Watkins, C.B., 2006. The use of 1-methylcyclopropene (1-MCP) on fruits and vegetables.  
405 Biotech. Adv. 24, 389-409.

406 Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA  
407 amplification for phylogenetic study. *J. Bacteriol.* 173, 697–703.

408 White, T.J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal  
409 ribosomal RNA genes for phylogenetics. In: *PCR Protocols: a guide to methods and applications.*  
410 (Innis M.A., Gelfand D.H., Sninsky J.J., White T.J. eds). Academic Press, New York, USA, 315-322.

411 Wills, R., Rigney, C., 2007. Effect of calcium on activity of mitochondria and pectic enzymes  
412 isolated from tomato fruits. *J. Food Biochem.* 3, 103-110.

413 Xiloyannis, C., Montanaro, G., Dichio, B., 2010. Sustainable orchard management, fruit quality  
414 and carbon footprint. *Acta Hort.* 913, 269-274.

415 Zhang, Z., Schwartz, S., Wagner, L., Miller, W., 2000. A greedy algorithm for aligning DNA  
416 sequences. *J. Comput. Biol.* 7, 203-14.

417 **Tables**

418

419 **Table 1** – Geographical location and incidence of plants infected by bacterial canker (%) of twelve *Actinidia deliciosa* “Hayward” orchards considered  
 420 in this study (2013 and 2014). Average dry matter (DM; %) and calcium content (Ca<sup>2+</sup>; %) are reported for the kiwifruits harvested from these  
 421 orchards.

Orchard	Geographical location	Geographical coordinates	2013			2014		
			DM (%)	Ca <sup>2+</sup> (%)	Infected plants(%)	DM (%)	Ca <sup>2+</sup> (%)	Infected plants(%)
Healthy 1	Campiglione Fenile (TO)	44° 48' 10" N, 7° 19' 29" E	13.53	0.35	0	13.98	0.31	0
Healthy 2	Campiglione Fenile (TO)	44° 48' 10" N, 7° 19' 29" E	13.23	0.28	4	14.09	0.35	2
Healthy 3	Campiglione Fenile (TO)	44° 48' 10" N, 7° 19' 29" E	13.87	0.29	1	15.36	0.37	0
Healthy 4	Campiglione Fenile (TO)	44° 48' 10" N, 7° 19' 29" E	13.58	0.32	2	14.5	0.47	1
Healthy 5	Campiglione Fenile (TO)	44° 48' 10" N, 7° 19' 29" E	14.13	0.34	0	15.01	0.5	0
Healthy 6	Envie (CN)	44° 40' 59" N 7° 22' 19" E	14.21	0.32	1	15.45	0.32	0
		<b>Mean</b>	13.76	0.32	1.3	14.73	0.39	0.5
Diseased 1	Bagnolo Piemonte (CN)	44° 45' 42" N 7° 18' 54" E	15.17	0.32	66	16.62	0.39	26
Diseased 2	Envie (CN)	44° 40' 59" N 7° 22' 19" E	17.12	0.22	86	15.72	0.35	49
Diseased 3	Envie (CN)	44° 40' 59" N 7° 22' 19" E	17.23	0.26	90	15.08	0.3	58
Diseased 4	Revello (CN)	44° 39' 20" N 7° 23' 34" E	16.09	0.22	94	16.18	0.32	63
Diseased 5	Revello (CN)	44° 39' 20" N 7° 23' 34" E	16.48	0.26	95	15.64	0.37	65
Diseased 6	Envie (CN)	44° 40' 59" N 7° 22' 19" E	15.3	0.16	88	14.58	0.34	55
		<b>Mean</b>	16.23	0.24	86	15.64	0.34	58
		<b>P value</b>	***	*	***	*	ns	***

422

423 \*, \*\*, \*\*\* Differences between given mean values with Student's t-test are significant for  $P$  value  $\leq 0.05$ ,  $\leq 0.01$ ,  $\leq 0.001$ . Not significant (ns,

424  $P \geq 0.05$ ). **Table 2** – Firmness, total soluble sugar, and titratable acidity of kiwifruit ‘Hayward’ harvested from six healthy and six *Psa* infected orchards

425 at harvest, end of storage, and shelf life. Fruit were harvested on 23 October 2013. The trial was performed under experimental conditions during  
 426 2013. Each value of firmness, total soluble sugar and titratable acidity is the mean of n=12 replicates.

Orchard	Firmness [N]					Total soluble sugar [%]			Titratable Acidity [%]			Grey mould (%)
	Harvest	End of storage		Shelf life		Harvest	End of storage	Shelf life	Harvest	End of storage	Shelf life	
Healthy 1	95.7 ±6.37	8.04 ±2.45	5.29 ±0.39	5.80 ±0.25	10.2 ±0.91	11.4 ±0.73	0.94 ±0.03	0.95 ±0.21	0.96 ±0.13	0		
Healthy 2	71.2 ±6.96	6.86 ±2.25	5.29 ±0.69	5.70 ±0.32	12.0 ±0.41	10.9 ±0.55	0.95 ±0.14	1.05 ±0.37	0.79 ±0.19	3		
Healthy 3	80.0 ±8.23	8.53 ±3.14	5.29 ±0.69	4.80 ±0.45	12.3 ±0.73	12.0 ±0.57	0.81 ±0.02	0.98 ±0.24	0.75 ±0.09	3		
Healthy 4	73.3 ±7.64	9.41 ±5.00	5.19 ±0.39	7.30 ±0.33	12.2 ±0.51	11.4 ±0.58	0.81 ±0.04	0.93 ±0.24	0.81 ±0.06	1		
Healthy 5	70.9 ±7.06	11.0 ±4.41	5.29 ±0.39	6.80 ±0.28	12.4 ±0.41	11.8 ±0.56	0.82 ±0.08	0.78 ±0.14	0.79 ±0.09	4		
Healthy 6	81.8 ±7.84	10.3 ±2.55	7.64 ±0.98	6.06 ±0.36	12.1 ±0.85	12.0 ±0.79	0.83 ±0.07	0.74 ±0.15	0.93 ±0.08	2		
<b>Healthy Average</b>	78.8 ±7.35	9.02 ±3.30	5.67 ±0.59	6.08 ±0.88	<b>11.9</b> ±0.84	<b>11.6</b> ±0.45	0.86 ±0.07	0.90 0.12	<b>0.84</b> ±0.09	<b>2</b>		
Diseased 1	80.7 ±6.37	14.3 ±2.16	5.29 ±0.69	6.60 ±0.41	13.0 ±0.58	12.9 ±0.76	1.17 ±0.06	0.87 ±0.16	0.53 ±0.01	29		
Diseased 2	86.3 ±6.86	6.66 ±2.55	5.59 ±0.59	6.40 ±0.35	14.4 ±0.43	15.3 ±0.58	0.88 ±0.12	1.01 ±0.22	0.48 ±0.14	26		
Diseased 3	75.7 ±6.17	8.13 ±1.67	5.19 ±0.49	6.20 ±0.44	14.9 ±0.67	15.9 ±0.82	0.86 ±0.03	0.76 ±0.14	0.48 ±0.04	15		
Diseased 4	73.2 ±7.45	9.21 ±1.57	6.08 ±0.59	6.20 ±0.4	13.5 ±0.81	14.3 ±0.61	1.01 ±0.04	0.75 ±0.19	0.51 ±0.03	15		
Diseased 5	72.4 ±7.94	11.4 ±5.19	5.78 ±0.69	7.60 ±0.32	13.7 ±0.64	14.1 ±1.12	0.82 ±0.06	0.92 ±0.21	0.50 ±0.07	28		
Diseased 6	77.9 ±7.06	10.4 ±3.82	7.55 ±1.37	6.60 ±0.26	13.3 ±0.83	12.7 ±0.76	1.07 ±0.17	0.84 ±0.13	0.35 ±0.06	22		
<b>Diseased Average</b>	77.7 ±6.97	10.0 ±2.83	5.91 ±0.74	6.60 ±0.52	<b>13.8</b> ±0.70	<b>14.2</b> ±1.28	0.97 ±0.14	0.86 ±0.10	<b>0.48</b> ±0.06	<b>23</b>		
<b>P value</b>	ns	ns	ns	ns	***	***	ns	Ns	***	***		

427

428 \*\*\*Differences between given mean values with Student's t-test are significant for  $P$  value  $\leq 0.001$ . Not significant (ns,  $P \geq 0.05$ ).

429 **Table 3** – Firmness, total soluble sugar, and titratable acidity of kiwifruits ‘Hayward’ harvested from six healthy and six *Psa* infected orchards at  
 430 harvest, end of storage, and shelf life. Fruit were harvested on 16 October 2014. The trial was performed under experimental conditions during 2014.  
 431 Each value of firmness, total soluble sugar and titratable acidity is the mean of n= 12 replicates.

Orchard	Firmness [N]			Total soluble sugar [%]			Titratable Acidity [%]			Grey mould (%)
	Harvest	End of storage	Shelf life	Harvest	End of storage	Shelf life	Harvest	End of storage	Shelf life	
Healthy 1	54.1 ±5.59	7.64 ±2.74	3.53 ±2.06	7.59 ±0.40	12.2 ±0.63	13.2 ±0.50	2.92 ±0.12	1.39 ±0.02	1.66 ±0.29	17
Healthy 2	59.8 ±4.21	7.64 ±2.94	3.72 ±2.35	7.48 ±0.53	11.6 ±0.71	12.1 ±0.60	2.94 ±0.30	1.32 ±0.02	1.30 ±0.03	12
Healthy 3	59.6 ±4.90	5.19 ±0.98	2.94 ±2.06	7.62 ±0.44	12.2 ±0.58	13.1 ±0.52	2.91 ±0.12	1.44 ±0.03	1.34 ±0.03	17
Healthy 4	54.4 ±6.86	7.84 ±2.35	2.55 ±1.76	7.88 ±0.65	12.0 ±0.47	13.6 ±0.53	2.65 ±0.36	1.41 ±0.03	1.48 ±0.02	10
Healthy 5	54.8 ±5.00	7.94 ±3.23	2.35 ±1.27	8.53 ±0.37	12.8 ±0.73	13.9 ±0.49	2.52 ±0.14	1.50 ±0.01	1.43 ±0.01	19
Healthy 6	59.7 ±5.78	5.39 ±2.16	2.16 ±0.88	8.17 ±0.43	12.1 ±0.52	14.3 ±0.47	2.14 ±0.07	1.42 ±0.08	1.39 ±0.02	23
<b>Healthy Average</b>	57.1 ±5.39	<b>6.94</b> ±2.40	<b>2.87</b> ±1.73	7.88 ±0.40	<b>12.2</b> ±0.39	<b>13.4</b> ±0.77	<b>2.68</b> ±0.32	1.41 ±0.06	1.43 ±0.13	<b>16</b>
Diseased 1	45.8 ±6.47	4.90 ±2.25	1.27 ±0.39	8.21 ±0.57	13.7 ±1.02	13.9 ±0.55	1.72 ±0.29	1.38 ±0.04	1.39 ±0.03	48
Diseased 2	59.7 ±4.31	6.37 ±2.35	1.96 ±0.98	8.03 ±0.31	13.8 ±0.25	13.8 ±0.46	1.52 ±0.26	1.45 ±0.04	1.48 ±0.06	34
Diseased 3	54.0 ±2.84	6.27 ±2.45	1.57 ±0.69	7.83 ±0.20	13.3 ±0.26	14.3 ±0.70	1.53 ±0.03	1.40 ±0.01	1.44 ±0.04	56
Diseased 4	52.2 ±5.88	5.19 ±2.35	2.55 ±1.27	7.92 ±0.53	13.4 ±1.24	14.1 ±0.47	1.39 ±0.12	1.45 ±0.03	1.44 ±0.01	53
Diseased 5	54.8 ±4.21	5.68 ±2.55	-	8.01 ±0.42	14.0 ±1.38	15.3 ±1.20	1.42 ±0.16	1.55 ±0.06	1.42 ±0.05	38
Diseased 6	60.4 ±6.86	3.72 ±2.65	1.47 ±0.49	7.67 ±0.45	13.3 ±1.17	14.0 ±0.70	1.20 ±0.12	1.34 ±0.04	1.23 ±0.04	38
<b>Diseased Average</b>	54.5 ±5.10	<b>5.36</b> ±2.43	<b>1.76</b> 0.76	7.95 ±0.19	<b>13.5</b> ±0.29	<b>14.2</b> ±0.54	<b>1.46</b> ±0.17	1.43 ±0.07	1.40 ±0.09	<b>45</b>
<b>P value</b>	ns	**	*	ns	***	*	***	ns	ns	***

432 \*, \*\*, \*\*\* Differences between given mean values with Student's t-test are significant for  $P$  value  $\leq 0.05$ ,  $\leq 0.01$ ,  $\leq 0.001$ . Not significant (ns,  $P \geq 0.05$ ).

433 **Table 4** - Firmness and total soluble sugar (TSS) at three time points (harvest, end of storage and shelf life) of kiwifruits ‘Hayward’ harvested from  
 434 healthy and *Psa* infected orchards and stored under normal atmosphere (1.0°C ±1 and RH 98%), controlled atmosphere alone (-0.8±0.8°C; RH 98%;  
 435 O<sub>2</sub> 2.5±0.5%, CO<sub>2</sub> 3.5±0.5%), and controlled atmosphere after treatment with SmartFresh™ Powder (a.i.: 3.3%, 1-MCP; Rohm & Hass). Each value  
 436 of firmness, total soluble sugar and titratable acidity is the mean of n=12 replicates.

437

Firmness [N]

	Normal atmosphere			Controlled atmosphere		Controlled atmosphere + 1-MCP	
	Harvest	End of storage	Shelf life	End of storage	Shelf life	End of storage	Shelf life
Healthy	78.8 ±9.51	22.5 ±8.62	12.5 ±5.10	56.4 ±6.37	17.0 ±5.49	69.1 ±4.51	18.9 ±3.14
Diseased	77.7 ±5.19	18.9 ±6.27	12.1 ±2.35	57.1 ±6.17	14.7 ±4.61	65.2 ±6.76	16.7 ±2.06
<b>P value*</b>	ns	ns	ns	ns	*	*	*

Total soluble sugar (TSS) [%]

	Normal atmosphere			Controlled atmosphere		Controlled atmosphere + 1-MCP	
	Harvest	End of storage	Shelf life	End of storage	Shelf life	End of storage	Shelf life
Healthy	6.08 ±0.88	11.5 ±0.79	12.0 ±0.62	11.6 ±0.48	12.8 ±0.73	11.7 ±0.67	12.6 ±0.49
Diseased	6.60 ±0.52	13.5 ±0.78	14.4 ±0.65	12.4 ±0.87	13.7 ±0.68	12.4 ±0.79	13.8 ±0.86
<b>P value*</b>	ns	***	***	***	***	***	***

438

439 \*, \*\*, \*\*\* Differences between given mean values with Student's t-test are significant for  $P$  value  $\leq 0.05$ ,  $\leq 0.01$ ,  $\leq 0.001$ . ns. Not significant ( $P \geq 0.05$ ).

440

441 **Supplementary material**

442

443 **Supplementary Table 1.** Number of vines, age and agronomic practices of the twelve *Actinidia deliciosa* “Hayward” orchards considered in this  
444 study.

445

Orchard	Vines (n°)	Vine spacing (vines/ha)	Age	Irrigation	Fertilisers	Training system	Pesticide treatments	Shelter	Average height (m)	Rootstock	Pollinators (n°)	Girdled trunk
Healthy 1	880	500	20				Copper	no	1.6	no	1:4	no
Healthy 2	780	460	14		Normal		Copper	no	1.5	no	1:4	no
Healthy 3	460	480	17	localized irrigation	fertilizer regime: 80-90 N unit/ha	pergola (V-trellis)	Copper	no	1.6	no	1:3	no
Healthy 4	860	520	15				Copper	no	1.6	no	1:4	no
Healthy 5	430	500	17				Copper	no	1.6	no	1:3	no
Healthy 6	590	460	21				Copper	no	1.5	no	1:4	no
Diseased 1	740	480	15				Copper	no	1.6	no	1:4	no
Diseased 2	910	470	19				Reduced		Copper	no	1.5	no
Diseased 3	560	510	17	localized irrigation	fertilizer regime: 50-60 N unit/ha	pergola (V-trellis)	Copper	no	1.6	no	1:4	no
Diseased 4	580	490	21				Copper	no	1.6	no	1:3	no
Diseased 5	630	500	20				Copper	no	1.5	no	1:4	no
Diseased 6	960	475	18				Copper	no	1.6	no	1:4	no

446

447 **Supplementary Figure captions**

448

449 **Supplementary Fig. 1** – Minimum, mean and maximum temperatures [°C] and precipitation [mm]  
450 of Piedmont region, northern Italy during 2013.

451

452 **Supplementary Fig. 2** - Minimum, mean and maximum temperatures [°C] and precipitation [mm]  
453 of Piedmont region, northern Italy during 2014.

454