AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Climate change will increase the potential conflict between skiing and high-elevation bird species in the Alps

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1591706> since 2017-03-24T10:41:28Z

Published version:

DOI:10.1111/jbi.12796

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Brambilla, Mattia; Pedrini, Paolo; Rolando, Antonio; Chamberlain, Dan E..
Climate change will increase the potential conflict between skiing and
high-elevation bird species in the Alps. JOURNAL OF BIOGEOGRAPHY.
None pp: 1-11.
DOI: 10.1111/jbi.12796

The publisher's version is available at:

<http://doi.wiley.com/10.1111/jbi.12796>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1591706>

1 **Original article**

2

3

4 **Climate change will increase the potential conflict between skiing and high-**
5 **elevation bird species in the Alps**

6

7 Mattia Brambilla^{1,2*}, Paolo Pedrini¹, Antonio Rolando³, Dan E. Chamberlain³

8

9 ¹ Museo delle Scienze, Sezione Zoologia dei Vertebrati, Corso della Scienza e del Lavoro 3, I-38123

10 Trento, Italy

11 ² Fondazione Lombardia per l'Ambiente, Settore Biodiversità e Aree protette, Largo 10 luglio 1976

12 1, I-20822 Seveso, MB, Italy

13 ³ Università di Torino, Dipartimento di Scienze della Vita e Biologia dei Sistemi, Via Accademia

14 Albertina 13, I-10123 Torino, Italy

15

16 *corresponding author; e-mail: brambilla.mattia@gmail.com

17

18 **Running header:** *Climate change, skiing and birds in the Alps*

19

20 **word count:** 6961

21 **ABSTRACT**

22 **Aim** To assess the extent of the possible future conflict between skiing and biodiversity driven by
23 climate change, human adaptation and species' distribution shifts.

24 **Location** Italian Alps.

25 **Methods** We assessed the extent of the possible future conflict between skiing and biodiversity by
26 predicting locations likely to be suitable for both skiing and for high-elevation birds in the Italian
27 Alps by modelling ski-piste and species presence in relation to climate, topography and habitat.
28 Potential conflict was assessed by comparing the overlap of areas projected as suitable for skiing
29 and those suitable for five high-elevation bird species under different scenarios of climate change
30 for the year 2050.

31 **Results** Areas suitable for both ski-pistes and birds were projected to contract towards upper
32 elevations, which for birds resulted in an average decrease of 58% - 67% of suitable area. The
33 degree of overlap between species and skiing was projected to increase, especially for the most
34 valuable sites, i.e. those hosting the most species, or the most threatened species.

35 **Main conclusions** Given the alarming range contractions forecast for high elevation species, and
36 the potential impact of ski-pistes on those species, it is essential to safeguard high mountain
37 grasslands against negative effects of ski development. An effective conservation strategy at a
38 landscape scale needs to consider prevention of ski-piste construction in sites of high conservation
39 value. The approach developed here provides a means by which such a strategy could be
40 formulated, and which could be potentially applied elsewhere to investigate the effect of human
41 adaptation on biodiversity.

42

43 **Keywords:** alpine grassland; bird conservation; global warming; human adaptation; mountain; ski-
44 piste

45 INTRODUCTION

46

47 Climate change induced by anthropogenic emissions of greenhouse gases is among the most severe
 48 threats to ecosystems and species at a global level (IPCC, 2013). Indirect impacts arising from
 49 human adaptation to new climates (i.e. taking appropriate action to prevent or minimize the adverse
 50 effects of climate change) pose imminent and important threats to biodiversity (Bradley *et al.*, 2012;
 51 Chapman *et al.*, 2014), and may impair species' ability to cope with climate change (Watson, 2014).
 52 However, the impacts driven by human adaptation are rarely considered in the conservation
 53 literature (Watson & Segan, 2013; Chapman *et al.*, 2014), although notable exceptions exist (e.g.
 54 Bradley *et al.*, 2012; Wetzel *et al.*, 2012). Humans continue to respond to climate change, and such
 55 a response is to some degree predictable (Watson, 2014). Therefore, researchers need to focus not
 56 only on the direct effect of changing climate, but also on the climate-related variation in human-
 57 mediated threats (Turner *et al.*, 2010; Watson, 2013). The latter can have quite immediate and
 58 overwhelming impacts, being of equal or greater intensity for species and ecosystems than the
 59 direct impacts of climate change (Turner *et al.*, 2010; Watson & Segan, 2013; Chapman *et al.*,
 60 2014).

61 High-mountain regions harbour a relatively high biodiversity (Dirnböck *et al.*, 2011) and a
 62 high percentage of both endemic (Essl *et al.*, 2009) and vulnerable species (Viterbi *et al.*, 2013).
 63 Many species are restricted to the upper elevations of their former range because of human
 64 pressures (Martin, 2001) and thus careful management of mountains is crucial for conservation
 65 (Rolando *et al.*, 2013). Mountain regions are particularly threatened by climate change (Brunetti *et*
 66 *al.*, 2009) and show a higher rate of warming compared to the global average (Böhm *et al.*, 2001).
 67 Species and habitats are already undergoing elevational range shifts due to climate warming, which
 68 is predicted to have important impacts on biodiversity at high elevation (Sekercioglu *et al.*, 2008;
 69 Dirnböck *et al.*, 2011; Chamberlain *et al.*, 2013). However, habitat changes caused by human action
 70 may have more severe consequences than climate change (Jetz *et al.*, 2007), or may interact with it,

71 showing synergistic effects (Mantyka-Pringle *et al.*, 2012).

72 Recreational activities, and in particular winter sports, represent one of the main threats to
 73 wild species in many mountain regions (e.g. Buckley *et al.*, 2000; Arlettaz *et al.*, 2013). There is
 74 increasing evidence of negative effects of skiing activities from a range of taxa (Rixen & Rolando,
 75 2013) via habitat destruction/alteration/fragmentation, soil degradation, and ski-related urban
 76 development (Rolando *et al.*, 2013). Ski-pistes for downhill skiing in the Alps are associated with
 77 both lower bird species richness and bird abundance (Rolando *et al.*, 2007; Caprio *et al.*, 2011;
 78 Rolando *et al.*, 2013). The negative effect of ski-pistes is mostly tied to the removal of vegetation
 79 and soil (Fig. S1 in Appendix S1); this is not invariably performed on all ski-pistes, but has become
 80 the standard practice for modern ski-piste construction (Negro *et al.*, 2013). Altered soil structure,
 81 harsh climate and plant species traits together prevent the re-establishment of vegetation, and grass
 82 cover for wild species remains extremely low for long periods even with modern restoration
 83 techniques such as hydro-seeding (Negro *et al.*, 2013).

84 Skiing activities are likely to be affected by climate change, and in particular by temperature
 85 rise and variation in precipitation regimes (Behringer *et al.*, 2000; Uhlmann *et al.*, 2009), and as
 86 such they have the potential to cause indirect impacts on biodiversity deriving from human
 87 adaptation. Due to decreasing snowfall and/or less reliable snow cover at lower altitudes, the area
 88 suitable for skiing is likely to show a range contraction and an upwards altitudinal shift (Elasser &
 89 Messerli, 2001; Disch *et al.*, 2007; Scott *et al.*, 2008), as already evident in some alpine areas
 90 (Pozzi, 2009; Marty, 2013). Given that a similar pattern of range contraction is likely to occur in the
 91 distribution of many mountain species (e.g. Sekercioglu *et al.*, 2008; Chamberlain *et al.*, 2013), this
 92 could lead to an increase in the potential conflict between winter sports and wildlife. However, the
 93 potential consequences for mountain wildlife of changes in ski developments as a response to
 94 climate change have not been fully assessed.

95 We aim to describe the increase in the potential impact of ski-pistes on alpine biodiversity
 96 by modelling the potential future distributions of both species and ski-pistes in relation to projected

97 climate changes in the European Alps. Our goal is to highlight where conflicts between downhill
98 skiing and nature conservation are most likely to occur over the next decades, by constructing
99 spatially explicit models of conflict zones between ski-pistes for downhill skiing and wildlife
100 (Braunisch *et al.*, 2011), using high-elevation birds as an example group. We first model the current
101 and future distribution of some high-altitude bird species potentially vulnerable to ski-piste
102 development in relation to climate, habitat and topography. Then, we model the current and future
103 distribution of ski-pistes in relation to climate and topography; finally, we adopt an approach to
104 reduce possible future conflicts by evaluating where they may arise. Among the areas that should be
105 considered as high-priority for the conservation of high-elevation species in a warmer climate, we
106 identify those which are likely to be suitable for ski-pistes in the future, and hence those where ski-
107 developments should be restricted or regulated.

108 MATERIALS AND METHODS

109

110 Study area, fieldwork and model species

111 The study area comprised a large area of the southern Alps in northern Italy included within the
 112 borders defined by the Alpine Convention (Fig. 1). As model species, we selected passerine birds
 113 that in the Alps are mostly or exclusively tied to grassland and other open habitats at high elevation,
 114 and that could be potentially affected by climate change and by the occurrence of ski-pistes (Tables
 115 S1 and S2 in Appendix S1): water pipit *Anthus spinoletta*, alpine accentor *Prunella collaris*,
 116 northern wheatear *Oenanthe oenanthe*, black redstart *Phoenicurus ochruros*, and snowfinch
 117 *Montifringilla nivalis*. Bird data were collated from different studies that were carried out between
 118 2000 and 2015 in all the main sectors of the study region (see Appendix S1). The number of
 119 occurrences for each species was: water pipit 658; alpine accentor 235; northern wheatear 443;
 120 black redstart 1428; snowfinch 74.

121

122 Modelling species distributions

123 Species distributions were modelled using MaxEnt (release 3.3.3k; Phillips *et al.*, 2006), which can
 124 deal appropriately with the climate variables potentially relevant for species distribution (Braunisch
 125 *et al.*, 2013) and is routinely adopted to analyse species distribution using data collected with
 126 different survey methodologies, as here (Appendix S1) or collected by means of unknown field
 127 methods (e.g. Engler *et al.*, 2014). All bird data were collected at a spatial resolution ≤ 100 -m.

128 To make distribution models as general and robust as possible, we applied the approach
 129 proposed by Radosavljevic & Anderson (2014), adopting a masked geographically structured
 130 evaluation of models. We divided our study area into four longitudinal belts (see Fig. S8 in
 131 Appendix S2), and used records and background points (10 000) from eastern and central-western
 132 portions to build models (training data), and records from western and central-eastern portions to
 133 evaluate models (test data).

Environmental variables included land cover, bioclimatic and topographical variables (Table S3). Ten land cover types were selected from the CORINE Land Cover (CLC2006; EEA, 2007) database in order to describe the habitat composition of each cell, assuming constant land cover over the time period (up to 2050). Two bioclimatic variables, annual temperature and precipitation, were calculated for each grid cell from the values downloaded from WorldClim v.1.4 (Hijmans *et al.*, 2005; <http://www.worldclim.org>; resolution 30 arc-seconds, corresponding to less than 1-km at this latitude). Slope and solar radiation (calculated taking 21st June as the reference day), were extracted from a Digital Terrain Model (resolution 20 m) of the study area. Before distribution modelling, we checked for multicollinearity among environmental predictors in order to minimize the risk of overfitting species' responses to climate. Variance Inflation Factors (VIFs) were evaluated, and highly collinear variables ($VIF > 5$) were omitted, following Zuur *et al.* (2009). This procedure included an evaluation of seasonal, in addition to annual, bioclimatic variables, but the former showed high levels of collinearity, therefore only annual values were used in distribution modelling (see below).

All environmental variables were calculated for 40 m x 40 m cells; for each cell, values represented i) the sum of cover (total area) per each type of land cover, ii) the average slope, and iii) the mean solar radiation, all referring to a radius of c. 100 m from each cell. For bioclimatic variables, we assigned to each 40 m x 40 m cell the value of the 30 arc-seconds cell with which it overlapped. Given the raster structure, this resulted in variables measured over an area of slightly more than 2 ha, a grain that matches fairly well with the supposed territory sizes, and hence the main areas of activity, of the species considered (Cramp, 1998; Gustin *et al.*, 2010). To reduce the risk of overfitting, we fitted models considering only linear and quadratic terms (dropping interaction, threshold and hinge functions), irrespective of sample size.

A spatial test of the representativeness of the climatic niche defined by the data

Given that we focussed on a portion of the range of our study species, we could potentially

overestimate the effect of climate change on those species (Barbet-Massin *et al.*, 2010). To avoid this risk, using our data (all locations per each species) and only climatic factors, we modelled species distribution across a large part of Europe (using tile no. 16 of the wordclim database as a study area, from southern Scandinavia to north Africa) and compared it with independent estimates of species distributions at larger scales (e.g. Cramp, 1998). Moreover, we compared the distribution predicted for Italy with the distribution data derived from the reporting to the EU under the Birds Directive (Nardelli *et al.*, 2015; data downloaded from <http://cdr.eionet.europa.eu/it/eu/art12/envuzmuow/>).

We compared models obtained using only either annual temperature, or temperature and rainfall, and selected the model better describing the European and Italian distribution for each species (Appendix S1) using visual assessment (in accord with the general approach of Zuur *et al.* 2009). We obtained reliable climate models for all species except northern wheatear, which was excluded from subsequent analyses.

Future distribution models

To simulate future conditions under climate warming, we chose two scenarios with different values of representative concentration pathways (RCPs; van Vuuren *et al.*, 2011), with respectively low and high rates of climate change. We selected the RCP values of +4.5 and +8.5 W/m² (Brambilla *et al.*, 2015), corresponding to an average increase of +1.4 and 2.0°C respectively in global temperature (IPCC 2013), and of +3.3 and 4°C in our study area, consistent with previously recorded rates of warming in the Alps that are approximately double the global average (Brunetti *et al.*, 2009). We downloaded relative bioclimatic variables for 2050, according to the Hadley Global Environment Model 2 (HadGEM2-ES), at a resolution of 30" from www.worldclim.org. Our choice of using 2050 as a future reference was due to the need to measure impacts on a rather short-term timescale, given that human decisions are rarely established on the basis of long-term predictions and assessments, and that the life-cycle of species is usually much shorter than the timescale often

186 considered in studies on the effect of climate change (Chapman *et al.*, 2014).

187 We selected logistic model output to allow for a binary reclassification (into suitable and
188 unsuitable sites) of the continuous estimate of habitat suitability from MaxEnt using recommended
189 thresholds for presence-only models (e.g. Liu *et al.*, 2013; Appendix 2), selected by mapping the
190 predicted occurrences on the basis of the individual thresholds, and comparing them with the
191 current species range (Nardelli *et al.*, 2015). To assess model validity, we considered the respective
192 performance over the test data set. We expected good models to show a substantial stability of the
193 area under the curve (AUC) of the receiver-operator plot between training and test data sets (see
194 above and Appendix S2). In general, very small differences in AUC values suggested model
195 stability, with the partial exception of black redstart (Table S4 in Appendix S2).

196

197 **A model for ski-piste distribution**

198 We used all available information on the location of ski-pistes to identify their current
199 distribution. Then, we manually placed in a GIS environment (by looking at detailed and updated
200 aerial photographs) points over ski-pistes known to be currently in use, each separated by a
201 minimum of 500m and a maximum of 1000m from the next nearest point; this procedure resulted in
202 610 points placed along ski-pistes throughout the entire study area. As we knew both occurrence
203 and true absence sites, we adopted a presence-absence modelling approach. We used multivariate
204 adaptive regression splines (MARS, Milborrow, 2011) in R (R Development Core Team, 2013) to
205 develop a model for ski-piste occurrence (Appendix 2). We compared the 610 points representing
206 ski-pistes with 610 randomly placed points representing areas free from ski-pistes. The latter were
207 placed at 5-10 kilometers from ski-piste points, to simultaneously avoid overlap with ski-areas and
208 ensure that absence points were in mountain areas potentially suitable for ski-pistes, and were
209 outside protected areas to avoid biases due to regulatory instead of morphological/climatic factors.
210 We tested the potential effect of mean slope, mean radiation, annual temperature, annual rainfall,
211 and the relative interactions. Minimum temperature and precipitation of the coldest quarter were

also tested, but they were removed because of high VIF values. Variables were calculated using the same environmental layers and resolution adopted for species distribution modelling. The *evimp* command was used to evaluate relative variable importance (Milborrow, 2011; Jedlikowski *et al.*, 2014) and to confirm model validity (Appendix S2). Minimum and maximum suitable areas actually used for ski-pistes varied between 2.7 to almost 30 000 ha (average 2 231 ha $\pm$ 3 877 SD). Then, we projected the model over the whole study area both for current and future climatic conditions.

Evaluating the risk of potential overlap between birds and ski-pistes

We evaluated the overlap between the area potentially suitable for ski-pistes and variables representing the distribution of target species considering: i) the area potentially suitable for each target species, ii) the area suitable for different numbers of target species (from one to four), iii) the areas suitable for all the species except for snowfinch (the species displaying the most extreme variation), iv) the conservation priority areas for high-elevation passerine birds. For the latter, we identified those areas which would harbour the largest number of species, or the most affected species (snowfinch; see Results).

Ski-piste expansion is more likely to occur in the proximity of already existing ski-resorts, because of accessibility and other practical reasons. Therefore, we re-ran the above analyses for each species, limiting areas potentially suitable for ski-pistes to within 5-km of existing ski-pistes (i.e. considering the 610 points mapped to define the current occurrence of ski-pistes; see above). We selected the 5 km buffer as this figure matched the average linear extent of existing ski-resorts (mean 4.75 km $\pm$ 3.00 SD, N = 48). In summary, four different scenarios were considered, low and high rates of warming where any climatically and topographically suitable area could be considered for future ski-piste development, and low and high rates of warming where only climatically and topographically suitable areas within close proximity of an existing ski-piste could be considered for future ski development.

238 RESULTS

239

240 Average temperature had consistently high contributions to predicting distributions across species
 241 (Appendix S2). Consequently, all species were projected to undergo a more or less marked
 242 reduction in potentially suitable areas in the future, ranging from 24% (black redstart, RCP +4.5) to
 243 97% (snowfinch, RCP +8.5) of the current range, with an average decrease across species of 58% -
 244 67% according to the two scenarios, with the scenario of higher rates of warming having the highest
 245 impact (Table 1). As expected for species tied to grassland and rocky habitats, a negative effect of
 246 forest habitats and/or a positive effect of natural grassland and other open habitats was found in all
 247 species (Table S4).

248 Ski-piste occurrence was associated with annual temperatures below 6-7° (with a peak in
 249 occurrence probability at 3.1°), slopes lower than 27-28°, and was affected by the interactions
 250 between annual temperature and both slope and mean radiation (Appendix S2): ski-piste occurrence
 251 probability was particularly low in areas less favourable to snow accumulation, i.e. the steepest
 252 gradients and relatively warmer areas, as well as in areas with high solar radiation. The area
 253 potentially suitable for ski-pistes was also projected to decrease from 529 000 ha to 254 000 ha
 254 (RCP +4.5) or 196 000 ha (RCP +8.5), thus being more than halved in 2050 compared to current
 255 conditions. The potential location of ski-pistes and the distribution of the target species were
 256 projected to show a contraction towards upper elevations (Appendix S2).

257 In addition to the overall contractions in species range, for some species the models also
 258 predicted a decrease in the proportion of species distributions unsuitable for ski-piste developments,
 259 i.e. the overlap between areas potentially suitable for alpine birds and areas potentially suitable for
 260 ski-pistes will increase. This was the case for water pipit, alpine accentor and snowfinch (under the
 261 RCP +4.5 scenario), whereas the potential overlap for black redstart was predicted to decrease
 262 (Table 1). The areas potentially suitable for all the target species were projected to undergo a large
 263 decline (91%-97% according to scenario), and the potential conflict with ski-pistes for these areas

264 to increase, from 66% to 68%-70%. Similar results were found for the areas potentially suitable for
265 three out of four species, with a less dramatic overall decrease (44%-60%) coupled with a greater
266 increase in potential conflict with ski-pistes, from 44% to 61%-64% (Table 2). The pattern of
267 variation in areas suitable for all species combined except snowfinch was similar, although of lower
268 magnitude: a decrease in potentially suitable areas of 56%-70% and an increase in potential overlap
269 with ski-pistes from 60% to 63%-65% were predicted.

270 The procedure for priority area definition identified c. 118000 hectares, dispersed throughout
271 the study area. Of those priority sites, 50% are currently also potentially suitable for ski-pistes; in
272 2050, the proportion of these areas also suitable for ski-pistes will increase to 63%-65% (RCP +4.5
273 and RCP +8.5 respectively).

274 Repeating the analyses for areas within 5 km of existing ski-pistes, the pattern of variation in
275 overlap between areas suitable for a given species (or for a given number of species) and the areas
276 suitable for ski-pistes mirrored the general pattern, although overlap was obviously lower (Tables 3
277 and 4). The potential overlap between priority areas and areas suitable for pistes was projected to
278 increase from 16% to 20%-21%. This projected increase in overlap was found despite a 51%-63%
279 decrease in the areas suitable for pistes (from 172 448 ha to 83 907-63 656 ha) within the adjacent
280 5-km area, i.e. fewer areas adjacent to ski-pistes will be potentially suitable for ski-piste
281 development, but there will still be an increase in the overlap with areas potentially suitable for bird
282 species.

283 **DISCUSSION**

284

285 Future climate change is likely to increase the potential conflict between high elevation bird species
 286 and skiing activities. Model outcomes suggested a shrinkage towards higher elevations and a
 287 contraction in range for both high-elevation bird distributions and locations suitable for ski-pistes in
 288 the Italian Alps by 2050. Moreover, the overlap between areas potentially suitable for high-
 289 elevation birds and those for skiing is projected to increase to the extent that most of the area above
 290 the treeline could be potentially subject to human-wildlife conflict: 61-70% of the area predicted as
 291 potentially suitable in the future for three or four species and two thirds of conservation priority
 292 areas will also be potentially suitable for ski-pistes. Limiting potentially suitable areas for ski-pistes
 293 to within 5 km of existing ones still suggested an increase in the overlap between areas potentially
 294 suitable for three bird species (not black redstart) and areas potentially suitable for ski-pistes.
 295 Obviously, the absolute overlap over the whole potential range of each species is much weaker, but
 296 importantly, this confirmed that an increase in the potential conflict due to climate change between
 297 ski-pistes and high-elevation bird species should be expected in any case.

298 Black redstart makes wide use of urban habitats in addition to open mountain habitats,
 299 unlike the other species considered, and is projected to disappear from middle-elevation sites, but to
 300 retain a reasonable extent of potentially suitable areas both at high elevations and in urban habitats.
 301 However, the other three species are projected to lose a large part of their potential distribution.
 302 Some of these species are already declining or contracting their range in Italy (Gustin *et al.*, 2010;
 303 Rete Rurale Nazionale & LIPU, 2014; Nardelli *et al.*, 2015), and as a result of such shifts, the status
 304 of mountain grassland birds is particularly concerning (Chamberlain *et al.*, 2013; Rete Rurale
 305 Nazionale & LIPU, 2014). The pattern of range contraction was projected to be quite similar across
 306 species (Appendix S2), and it is likely that a similar distributional change may also be relevant to
 307 other high-elevation species of open habitats in the future, including birds (e.g. rock ptarmigan
 308 *Lagopus muta helvetica*) and other taxa for which high altitude grasslands hold a high diversity

309 and/or species of conservation interest (e.g. flowers, butterflies, carabids, dung beetles - Nagy *et al.*,
 310 2003; Tocco *et al.*, 2013). To that extent, our results can be considered to represent threats not just
 311 to alpine birds, but to biodiversity of high altitude open habitats in general.

312 A similar pattern of range contraction is predicted for ski-pistes. The ski industry will be
 313 affected by climate change (Behringer *et al.*, 2000; Disch *et al.*, 2007; Uhlmann *et al.*, 2009), and
 314 our results suggest a potentially marked contraction of the areas suitable for ski-pistes. Notably,
 315 given that climatic variables are representative of the period 1950-2000 (Hijmans *et al.*, 2005) and
 316 that the ski industry is particularly sensitive to climate variations, the potential distribution
 317 modelled on the basis of the 'current' conditions gave a good representation of the distribution of
 318 areas suitable for skiing in the second half of the past century. Some of the ski-pistes located in the
 319 'marginal' parts of our study area (e.g. in the pre-Alps in Lombardy), in sites predicted by the model
 320 to be on the edge of the suitable area and to be less suitable for ski-pistes than those located in the
 321 'core' of the Alpine region, have already been decommissioned in recent years, because of the
 322 prolonged lack of adequate snow cover, in particular at lower elevations (Pozzi, 2009; Marty, 2013).
 323 Future projections suggest that other sites at medium and low elevations will probably be
 324 decommissioned (Pozzi, 2009; Marty, 2013), in keeping with our own results. Even if the use of
 325 artificial snow (a non-sustainable adaptation; Disch *et al.*, 2007) may to some extent buffer against
 326 decreased snow cover, it is likely that ski-pistes will contract towards higher elevations to track the
 327 availability of suitable temperatures and snow cover.

328

329 **Model assumptions**

330 We used scenarios derived from a single global climate model to evaluate potential conflicts
 331 with ski-pistes. Future species distribution models based on scenarios derived from an alternative
 332 global model (MIROC-ESM-CHEM; Watanabe et al. 2011) led to consistent, though less severe,
 333 predictions (Appendix 2). Although this general agreement and the fact that the predictions about
 334 temperature increase in the Alps are rather consistent across different models (Giorgi & Lionello,

2008; in our study area, the correlation between average temperature values according to the scenario RCP +8.5 of the two climate models was 0.98), we cannot exclude some possible minor differences in future predictions according to scenarios taken from other global models. However, we used different scenarios (RCP +4.5 and +8.5 W/m²) and different assumptions (restricted or unrestricted development of ski-pistes) to provide a range of scenarios, in keeping with our generally conservative approach (see below). It is unlikely that the possible variation from using different circulation models would be greater than that associated with different emission scenarios/RCP values used – whilst there would be some variation, we expect that the general patterns would very likely be the same.

In our approach, we assumed constant land cover over the period considered. Significant land cover changes due to climate change at high elevation can require a fairly long time to become discernible due to lagged responses (e.g. Cannone *et al.*, 2007), although there is clearly much geographic variation (e.g. Harsch *et al.*, 2009; Carlson *et al.*, 2014). Nevertheless, we feel our approach is justified in that it gives a conservative estimate of species distribution decrease, as the most likely scenario of climate-induced habitat change is loss of open grassland habitat as treelines continue to shift upslope. Without management intervention, this may cause potentially severe reductions in the area of alpine grasslands, and increased fragmentation of remaining patches, with subsequent negative impacts on grassland species (Chamberlain *et al.*, 2013). At the same time, ski-pistes are not likely to be constrained by habitat changes, as they can be constructed on any habitat that is topographically suitable. A model that considered both altitudinal shifts in the treeline in conjunction with expansion of ski-pistes to higher altitudes would therefore almost certainly result in more severe model outcomes compared to our conservative approach. It is all the more striking then that even under the relatively conservative scenarios adopted here, there are nonetheless some major declines and increased conflicts predicted.

We considered only ski-pistes, but their impact extends beyond the piste itself, as the development of a ski-area also includes the development of infrastructure (e.g. roads, hotels,

361 restaurants; Rolando *et al.*, 2013). Off-piste skiing may also be expected to respond to climate
 362 changes in the same way, which also has negative consequences for biodiversity (Arlettaz *et al.*,
 363 2007). The fact that wider impacts of development of infrastructure and of off-piste skiing were not
 364 explicitly included in the approach are further factors which makes the estimates of negative
 365 impacts fairly conservative. Our model outcomes can therefore be considered to be relatively
 366 optimistic given the underlying assumptions.

367 Regarding the distribution models, the use of a rather coarse land-cover map, the lack of
 368 explicit information about model transferability, and the relatively small sample size for snowfinch
 369 should lead to some caution when interpreting the model outcomes. First, despite the low resolution
 370 of the Corine Land Cover map, the species-habitat relationships we modelled were all coherent with
 371 the basic species' ecology, and the reliable output we obtained (e.g. in terms of high AUC values)
 372 further confirmed that at the scale considered, the use of such data coupled with climatic data
 373 provides valuable insights into species distributions. Second, although we did not explicitly
 374 evaluate model transferability across different sub-regions, the general consistency of the
 375 discriminatory ability of models across the geographically independent training and testing data
 376 (Table S4) suggested an overall validity of the species-environment relationships over the study
 377 area. Furthermore, consistency in the species trends along altitudinal gradients in different Alpine
 378 regions (unpubl. data) suggests that model transferability is likely to be high. Third, sample size was
 379 relatively low for the snowfinch ($N = 74$) compared to the other species. However, MaxEnt is less
 380 sensitive to sample size than other methods (Wisz *et al.*, 2008), and there are several examples in
 381 the literature of robust MaxEnt models with much lower sample sizes (e.g. Guisan *et al.*, 2007;
 382 Wisz *et al.*, 2008; Brambilla, 2015), and indeed, snowfinch models evaluated by visual assessment
 383 and by AUC had good predictive ability. Nevertheless, future insights based on a larger sample may
 384 potentially increase model accuracy and hence would be desirable for this scarce high altitude
 385 specialist.

386 Future specific work is required to further understand the ultimate mechanisms driving

species occurrence. In common with the general climate envelope modelling approach, we implicitly assume that climate, either directly or indirectly (e.g. via resources) is a key determinant of species distributions, although such approaches have limitations on estimating the true niche of a species (Schurr *et al.*, 2012). We can conclude little on the precise mechanisms that may influence species distributions. There is a need for further studies, especially in mountain environments which are relatively poorly studied, to understand mechanisms underpinning apparently climate-limited species distributions and hence to identify potential compensatory or mitigation measures (Chamberlain *et al.* 2012).

395

396 **Applications**

Given the alarming range contractions forecast for high elevation species in general, and the increasing potential impact of ski-pistes on those species, it is essential to develop conservation strategies to safeguard high alpine habitats against negative effects of ski development. There is a need to promote better management of existing ski-pistes to minimise their negative impacts, through grassland restoration and minimization of deleterious management practices (Negro *et al.*, 2013; Rixen & Rolando, 2013). However, the priority should be to secure the persistence of climatically and structurally suitable sites for those threatened species, unaltered by development. We identified c. 118 000 hectares that are currently suitable for the most threatened species (snowfinch) and/or for all the other species belonging to the species assemblage (mountain grassland birds), and that will remain suitable in the future, whatever the scenario considered; those areas should be regarded as conservation priorities, where the development of ski infrastructures should be avoided. Unfortunately, the potential pressure on those areas will be high, as two thirds of them will also be suitable for future ski-piste development. The scenarios which considered that potentially suitable areas for ski-pistes will only be in the proximity of existing ski-pistes predicted a consequently lower overlap with areas potentially suitable for high-elevation bird species. However, many areas close to and including existing ski-pistes may not be climatically suitable for

413 adequate snow conditions in the future, hence there is likely to be further pressure to develop new
414 ski resorts far from existing ones.

415 The skiing industry provides economic benefits to local mountain communities which
416 otherwise could have limited economic capacity (Elsasser and Messerli 2001). Adaptation of skiing
417 activities in response to climate change via the construction of pistes in more climatically suitable
418 areas could be therefore desirable for the local economy. Given that there are some areas which are
419 predicted to be potentially suitable for ski-pistes, but which are not within the conservation priority
420 areas, a way to minimize the impact of new developments would be to perform a first selection of
421 areas for new ski-pistes among those sites. Clearly, other factors should then be taken into account
422 (e.g. occurrence of other species or habitats of conservation concern, accessibility, economic
423 feasibility, etc.), but our outputs could provide spatially explicit guidance in avoiding planned
424 development in the most important sites for the conservation of high-elevation biodiversity.

425 A clear conservation strategy is required to preserve suitable conditions in the priority areas
426 for the protection of alpine biodiversity (Fig. 2). The winter tourism industry is already adapting to
427 climate changes, via a range of different measures adopted to offset adverse economic impacts,
428 putting new pressures on mountain environments (Marty, 2013). Consequently, effective
429 conservation strategies, implemented at a landscape scale, need to consider prevention of ski-piste
430 construction in sites characterized by high conservation value. The approach developed here
431 provides a means by which such a strategy could be formulated.

432

433 **ACKNOWLEDGMENTS**

434

435 G. Assandri, E. Bassi, M. Belardi, A. Iemma provided additional field data. MB was partly
436 supported by the LIFE+ project "LIFE11 NAT/IT/044 GESTIRE". Most of data from Trento
437 province have been made available through the WebGIS developed within the LIFE+ project
438 "LIFE11/IT/187 T.E.N." and MB and PP were also partly supported by this project and by the
439 Accordo di Programma per la Ricerca PAT/MUSE 2014. DEC was funded by the People
440 Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme
441 FP7/2007-2013/.

442

443

REFERENCES

- Arlettaz, R., Patthey, P., Baltic, M., Leu, T., Schaub, M., Palme, R. & Jenni-Eiermann, S. (2007) Spreading free-riding snow sports represent a novel serious threat for wildlife. *Proceedings of the Royal Society B*, **274**, 1219–1224.
- Arlettaz, R., Patthey, P. & Braunisch, V. (2013) Impacts of Outdoor Winter Recreation on Alpine Wildlife and Mitigation Approaches: A Case Study of the Black Grouse. *The Impacts of Skiing and Related Winter Recreational Activities on Mountain Environments* (ed. by C. Rixen and A. Rolando), pp 137-154. Bentham.
- Barbet-Massin, M., Thuiller, W., & Jiguet, F. (2010). How much do we overestimate future local extinction rates when restricting the range of occurrence data in climate suitability models?. *Ecography*, **33**, 878-886.
- Behringer, J., Buerki, R. & Fuhrer, J. (2000) Participatory integrated assessment of adaptation to climate change in Alpine tourism and mountain agriculture. *Integrated Assessment*, **1**, 331-338.
- Böhm, R., Auer, I., Brunetti, M., Maugeri, M., Nanni, T. & Schöner, W. (2001) Regional temperature variability in the European Alps; 1769–1998 from homogenized instrumental time series. *International Journal of Climatology*, **21**, 1779–1801.
- Bradley, B.A., Estes, L.D., Hole, D.G., Holness, S., Oppenheimer, M., Turner, W.R., Beukes, H., Schulze, R.E., Tadross, M.A. & Wilcove, D.S. (2012) Predicting how adaptation to climate change could affect ecological conservation: secondary impacts of shifting agricultural suitability. *Diversity and Distributions*, **18**, 425-437.
- Brambilla, M. (2015) Landscape traits can contribute to range limit equilibrium: habitat constraints refine potential range of an edge population of Black-headed Bunting *Emberiza melanocephala*. *Bird Study*, **62**, 132-136.
- Brambilla, M., Bergero, V., Bassi, E. & Falco, R. (2015) Current and future effectiveness of Natura

- 470 2000 network in the central Alps for the conservation of mountain forest owl species in a
 471 warming climate. *European Journal of Wildlife Research*, **61**, 35-44.
- 472 Braunisch, V., Coppes, J., Arlettaz, R., Suchant, R., Schmid, H. & Bollmann, K. (2013) Selecting
 473 from correlated climate variables: a major source of uncertainty for predicting species
 474 distributions under climate change. *Ecography*, **36**, 971-983.
- 475 Braunisch, V., Patthey, P. & Arlettaz, R. (2011) Spatially explicit modeling of conflict zones
 476 between wildlife and snow sports: prioritizing areas for winter refuges. *Ecological*
 477 *Applications*, **21**, 955-967.
- 478 Brunetti, M., Lentini, G., Maugeri, M., Nanni, T., Auer, I., Böhm, R. & Schöener, W. (2009) Climate
 479 variability and change in the Greater Alpine Region over the last two centuries based on
 480 multi-variable analysis. *International Journal of Climatology*, **29**, 2197-2225.
- 481 Buckley, R.C., Pickering, C.M. & Warnken, J. (2000). Environmental Management for Alpine
 482 Tourism and Resorts in Australia. *Tourism and development in mountain regions* (ed. by P.
 483 Godde, M. Price and F.M. Zimmerman), pp 27-45. Cabi Publishing, Oxon.
- 484 Cannone, N., Sgorbati, S. & Guglielmin, M. (2007) Unexpected impacts of climate change on
 485 alpine vegetation. *Frontiers in Ecology and the Environment*, **5**, 360-364.
- 486 Caprio, E., Chamberlain, D.E., Isaia, M. & Rolando, A. (2011) Landscape changes caused by high
 487 altitude ski-pistes affect bird species richness and distribution in the Alps. *Biological*
 488 *Conservation*, **144**, 2958-2967.
- 489 Carlson, B.Z., Georges, D., Rabatel, A, Randin, C.F., Renaud, J., Delestrade, A., Zimmermann,
 490 N.E., Choler, P. & Thuiller, W. (2014) Accounting for tree line shift, glacier retreat and
 491 primary succession in mountain plant distribution models. *Diversity and Distributions*, **20**,
 492 1379-1391.
- 493 Chamberlain, D., Arlettaz, R., Caprio, E., Maggini, R., Pedrini, P., Rolando, A. & Zbinden, N.
 494 (2012) The altitudinal frontier in avian climate change research. *Ibis*, **154**, 205-209.
- 495 Chamberlain, D.E., Negro, M., Caprio, E. & Rolando, A. (2013) Assessing the sensitivity of alpine

- 496 birds to potential future changes in habitat and climate to inform management strategies.
 497 *Biological Conservation*, **167**, 127-135.
- 498 Chapman, S., Mustin, K., Renwick, A.R., Segan, D.B., Hole, D.G., Pearson, R.G. & Watson, J.E.
 499 (2014) Publishing trends on climate change vulnerability in the conservation literature reveal
 500 a predominant focus on direct impacts and long time-scales. *Diversity and Distributions*, **20**,
 501 1221-1228.
- 502 Cramp, S. (1998) *The Birds of the Western Palearctic, CD-Rom Edition*. Oxford University Press,
 503 Oxford.
- 504 Dirnböck, T., Essl, F. & Babitsch, W. (2011) Disproportional risk for habitat loss of high-altitude
 505 endemic species under climate change. *Global Change Biology*, **17**, 990–996.
- 506 Disch, D., Reppe, S. & Liebing, A. (2007) *Climate change in the Alps Facts - Impacts - Adaptation*.
 507 Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU).
- 508 EEA (European Environment Agency) (2007) *CLC2006 technical guidelines*. EEA Technical report
 509 No. 17.
- 510 Elsasser, H. & Messerli, P. (2001) The vulnerability of the snow industry in the Swiss Alps.
 511 *Mountain Research and Development*, **21**, 335-339.
- 512 Engler, J.O., Rödder, D., Stiels, D. & Förschler, M.I. (2014) Suitable, reachable but not colonised:
 513 seasonal niche duality in an endemic mountainous songbird. *Journal of Ornithology*, **155**,
 514 657-669.
- 515 Essl, F., Staudinger, M., Stöhr, O., Schratt-Ehrendorfer, L., Rabitsch, W. & Niklfeld, H. (2009)
 516 Distribution patterns, range size and niche breadth of Austrian endemic plants. *Biological*
 517 *Conservation*, **142**, 2547–2558.
- 518 Giorgi, F. & Lionello, P. (2008) Climate change projections for the Mediterranean region. *Global*
 519 *and Planetary Change*, **63**, 90-104.

- 520 Guisan, A., Zimmermann, N.E., Elith, J., Graham, C.H., Phillips, S., & Peterson, A.T. (2007) What
 521 matters for predicting the occurrences of trees: techniques, data, or species' characteristics? *Ecological Monographs*, **77**, 615-630.
- 522
- 523 Gustin, M., Brambilla, M. & Celada, C. (eds) (2010) *Valutazione dello Stato di Conservazione*
 524 *dell'avifauna italiana. Volume II. Passeriformes*. Ministero dell'Ambiente, della Tutela del
 525 Territorio e del Mare & LIPU/BirdLife Italia, Roma.
- 526 Harsch, M.A., Hulme, P.E., McGlone, M.S. & Duncan, R.P. (2009) Are treelines advancing? A
 527 global meta-analysis of treeline response to climate warming. *Ecology Letters*, **12**, 1040–
 528 1049.
- 529 Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high resolution
 530 interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**,
 531 1965–1978.
- 532 IPCC (2013) Climate change 2013: the physical science basis. *Working group I contribution to the*
 533 *fifth assessment report of the intergovernmental panel on climate change* (ed. by T.F. Stocker,
 534 D. Qin, G.-K. Plattner, M.M.B. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and
 535 P.M. Midgley), pp. 1–14. Cambridge University Press, Cambridge.
- 536 Jedlikowski, J., Brambilla, M. & Suska-Malawska, M. (2014) Nest site selection in Little Crake
 537 *Porzana parva* and Water Rail *Rallus aquaticus* in small midfield ponds. *Bird Study*, **61**,
 538 171-181.
- 539 Jetz, W., Wilcove, D.S. & Dobson, A.P. (2007) Projected impacts of climate and land-use change on
 540 the global diversity of birds. *PLoS Biology*, **6**, e157.
- 541 Liu, C., White, M. & Newell, G. (2013) Selecting thresholds for the prediction of species
 542 occurrence with presence-only data. *Journal of Biogeography*, **40**, 778 – 789.
- 543 Mantyka-Pringle, C.S., Martin, T.G. & Rhodes, J.R. (2012) Interactions between climate and habitat
 544 loss effects on biodiversity: a systematic review and meta-analysis. *Global Change Biology*,
 545 **18**, 1239–1252.

- 546 Martin, K. (2001) Wildlife in alpine and sub-alpine habitats. *Wildlife-Habitat Relationships in*
 547 *Oregon and Washington* (eds Johnson DH, O'Neil TA), pp 285-310. Oregon State University
 548 Press, Corvallis, OR.
- 549 Marty, C. (2013) Climate Change and Snow Cover in the European Alps. *The Impacts of Skiing and*
 550 *Related Winter Recreational Activities on Mountain Environments* (ed. by C. Rixen and A.
 551 Rolando), pp 30-44. Bentham.
- 552 Nagy, L., Grabherr, G., Körner, C. & Thompson, D.B.A. (eds.) (2003) *Alpine Biodiversity in*
 553 *Europe*. Ecological Studies 167 XXXI, Springer Verlag, Berlin.
- 554 Nardelli, R., Andreotti, A., Bianchi, E., Brambilla, M., Brecciaroli, B., Celada, C., Dupré, E.,
 555 Gustin, M., Longoni, V., Pirrello, S., Spina, F., Volponi, S. & Serra, L. (2014) Rapporto
 556 sull'applicazione della Direttiva 147/2009/CE in Italia: dimensione, distribuzione e trend delle
 557 popolazioni di uccelli (2008-2013). Roma, Italy: ISPRA, Serie Rapporti, 219/2015.
- 558 Negro, M., Rolando, R., Barni, E., Bocola, D., Filippa, G., Freppaz, M., Isaia, M., Siniscalco, C. &
 559 Palestini, C. (2013) Differential responses of ground dwelling arthropods to ski-piste
 560 restoration by hydroseeding. *Biodiversity Conservation*, **22**, 2607-2634.
- 561 Patthey, P., Wirthner, S., Signorell, N. & Arlettaz, R. (2008) Impact of outdoor winter sports on the
 562 abundance of a key indicator species of alpine ecosystems. *Journal of Applied Ecology*, **45**,
 563 1704-1711.
- 564 Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006) Maximum entropy modeling of species
 565 geographic distributions. *Ecological Modelling*, **190**, 231–259.
- 566 Pozzi, A. (2009) *The Effect of Climate Change on Alpine destinations in Lombardy*. IREALP.
 567 available at <http://www.climalptour.eu/>
- 568 Radosavljevic, A. & Anderson, R.P. (2014). Making better Maxent models of species distributions:
 569 complexity, overfitting and evaluation. *Journal of Biogeography*, **41**, 629-643.
- 570 Rete Rurale Nazionale & LIPU (2014) *Uccelli comuni in Italia. Aggiornamento degli andamenti di*
 571 *popolazione e del Farmland Bird Index per la Rete Rurale Nazionale dal 2000 al 2013*.

- 572 www.reterurale.it/farmlandbirdindex
- 573 Rixen, C. & Rolando, A. (eds) 2013 *The Impacts of Skiing and Related Winter Recreational*
 574 *Activities on Mountain Environments*. Bentham, DOI: 10.2174/97816080548861130101.
- 575 Rolando, A., Caprio, E. & Negro, M. (2013) The Effect of Ski-Pistes on Birds and Mammals. In:
 576 *The Impacts of Skiing and Related Winter Recreational Activities on Mountain Environments*
 577 (ed. by C. Rixen and A. Rolando), pp 101-122. Bentham.
- 578 Rolando, A., Caprio, E., Rinaldi, E. & Ellena, I. (2007) The impact of high-altitude ski-runs on
 579 alpine grassland bird communities. *Journal of Applied Ecology*, **44**, 210-219.
- 580 Rolando, A., Laiolo, P. & Carisio, L. (2003) Urbanization and the flexibility of the foraging ecology
 581 of the Alpine Chough *Pyrrhocorax graculus* in winter. *Revue d'écologie*, **58**, 337-352.
- 582 Schurr, F.M., Pagel, J., Cabral, J.M., Groenevald, J., Bykova, O., O'Hara, R.B., Hartig, F., Kissling,
 583 W.D., Linder, H.P., Midgley, G.F., Schröder, B., Singer, A. & Zimmermann, N.E. (2012) How to
 584 understand species' niches and range dynamics: a demographic research agenda for
 585 biogeography. *Journal of Biogeography*, **29**, 2146-2162.
- 586 Scott, D., Dawson, J. & Jones, B. (2008) Climate change vulnerability of the US Northeast winter
 587 recreation – tourism sector. *Mitigation, Adaptation Strategies to Global Change*, **13**, 577–596.
- 588 Sekercioglu, C.H., Schneider, S.H., Fay, J.P. & Loarie, S.R. (2008) Climate change, elevational
 589 range shifts and bird extinctions. *Conservation Biology*, **22**, 140-150.
- 590 Tocco, C., Negro, M., Rolando, A. & Palestini, C., (2013). Does natural reforestation represent a
 591 potential threat to dung beetle diversity in the Alps? *Journal of Insect Conservation*, **17**, 207-217.
- 592 Turner, W.R., Bradley, B.A., Estes, L.D., Hole, D.G., Oppenheimer, M. & Wilcove, D.S. (2010)
 593 Climate change: helping nature survive the human response. *Conservation Letters*, **3**, 304-
 594 312.
- 595 Uhlmann, B., Goyette, S. & Beniston, M. (2009) Sensitivity analysis of snow patterns in Swiss ski
 596 resorts to shifts in temperature, precipitation and humidity under conditions of climate
 597 change. *International Journal of Climatology*, **29**, 1048-1055.

- 598 Van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C.,
 599 Kram, T., Krey, V., Lamarque, J.F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S.J.
 600 & Rose, S. K. (2011) The representative concentration pathways: an overview. *Climate*
 601 *Change*, **109**, 5–31.
- 602 Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T.,
 603 Kawase, H., Abe, M., Yokohata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S. &
 604 Kawamiya, M. (2011). MIROC-ESM 2010: model description and basic results of CMIP 5-20
 605 c3m experiments. *Geoscientific Model Development*, **4**, 845-872.
- 606 Viterbi, R., Cerrato, C., Bassano, B., Bionda, R., Hardenberg, A., Provenzale, A. & Bogliani, G.
 607 (2013) Patterns of biodiversity in the northwestern Italian Alps: a multi-taxa approach.
 608 *Community Ecology*, **14**, 18-30.
- 609 Watson, J.E. (2014) Human Responses to Climate Change will Seriously Impact Biodiversity
 610 Conservation: It's Time We Start Planning for Them. *Conservation Letters*, **7**, 1-2.
- 611 Watson, J.E. & Segan, D.B. (2013) Accommodating the human response for realistic adaptation
 612 planning: response to Gillson et al. *Trends in Ecology and Evolution*, **28**, 573-574.
- 613 Wetzal, F.T., Kissling, W.D., Beissmann, H. & Penn, D.J. (2012) Future climate change driven sea-
 614 level rise: secondary consequences from human displacement for island biodiversity. *Global*
 615 *Change Biology*, **18**, 2707-2719.
- 616 Wisz, M.S., Hijmans, R.J., Li, J., Peterson, A.T., Graham, C.H., & Guisan, A. (2008) Effects of
 617 sample size on the performance of species distribution models. *Diversity and Distributions*,
 618 **14**, 763-773.
- 619
- 620

621 **Supporting Information**

622 Additional Supporting Information may be found in the online version of this article:

623 S.1. Dataset of bird occurrence and selection of target species

624 S.2. Modelling distribution of ski-piste and bird species

625

626 **BIOSKETCH**

627 Our research focus is on animal ecology and conservation in high altitude habitats, with particular
628 emphasis on the effects of environmental change and of direct and indirect human impacts on alpine
629 faunal biodiversity.

630 Author contributions: M.B. and D.E.C. collected part of the field data; P.P. and A.R. managed
631 fieldwork in Trentino and Piemonte, respectively; M.B. took a lead on the analyses; M.B. and
632 D.E.C. wrote a first draft of the paper; all authors contributed to the final version of the paper.

633

634

635

636

637 **Table 1** Predicted decrease in species range and overlap with cells potentially suitable for ski-pistes
 638 in current and future scenarios according to the climates predicted under RCPs of +4.5 and +8.5
 639 (general overlap and overlap considering sites potentially suitable for ski-pistes only within 5 km
 640 from existing ones).

641

	suitable area (ha*1000)			Distribution change		overlap with areas suitable for ski-pistes			overlap with areas suitable for ski-pistes (< 5 km)		
	current	+4.5	+8.5	+4.5	+8.5	current	+4.5	+8.5	current	+4.5	+8.5
water pipit	617	264	184	-57%	-70%	53%	61%	64%	17%	19%	20%
alpine accentor	615	261	193	-57%	-69%	43%	48%	49%	14%	15%	15%
black redstart	902	685	622	-24%	-31%	47%	31%	26%	16%	10%	9%
snowfinch	318	27	10	-91%	-97%	42%	47%	42%	14%	15%	15%

642 **Table 2** Suitable area for different levels of species richness, and the relative overlap with areas
 643 potentially suitable for skiing under current and future conditions (general overlap and overlap
 644 considering sites potentially suitable for ski-pistes only within 5 km from existing ones).

645

no. of species	area (ha * 1000)			overlap with ski			overlap with ski (< 5 km)		
	current	+4.5	+8.5	current	+4.5	+8.5	current	+4.5	+8.5
1	431	476	473	24%	13%	11%	9%	5%	4%
2	161	87	68	40%	51%	52%	7%	28%	16%
3	308	171	124	44%	61%	64%	14%	19%	20%
4	193	18	6	66%	70%	68%	23%	23%	25%

646

647

648

649 **Figure 1** Location of the study area in Italy, defined as the provinces included in the Alpine
650 Convention in Piedmont, Lombardy and Trentino. Extent 40,569 km², elevation range 30-4,600 m
651 asl.

652

653

654 **Figure 2** Spatial relationship between conservation priority sites (sites suitable for snowfinch and/or
655 all the other high-elevation species in 2050) and sites suitable for ski-pistes in 2050 (upper: RCP
656 +4.5; lower: RCP: +8.5), on the southern Alps, and in two sample areas (for RCP +4.5), Val
657 d'Ossola (left) and Valtellina (right).