AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Rapid on-site identification of the biocontrol agent of the Asian chestnut gall wasp

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1601009> since 2016-10-12T11:13:52Z

Published version:

DOI:10.1080/09583157.2016.1195335

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Colombari, F.; Villari, C.; Simonato, M.; Cascone, P.; Ferracini, C.; Alma, A.; Guerrieri and A Battisti, E. Guerrieri and A. Battisti. Rapid on-site identification of the biocontrol agent of the Asian chestnut gall wasp. BIOCONTROL SCIENCE AND TECHNOLOGY. 26 (9) pp: 1285-1297.
DOI: 10.1080/09583157.2016.1195335

The publisher's version is available at:

<http://www.tandfonline.com/doi/full/10.1080/09583157.2016.1195335>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

Rapid on-site identification of the biocontrol agent of the Asian chestnut gall wasp

Colombari F.¹, Villari C.^{1,2}, Simonato M.¹, Cascone P.³, Ferracini C.⁴, Alma A.⁴, Guerrieri E.³, Battisti A.¹

¹ Department of Agronomy Food Natural resources Animals and Environment (DAFNAE), University of Padova, Agripolis, 35020 Legnaro (PD), Italy; ² Department of Plant Pathology, The Ohio State University, Columbus, OH 43210, USA; ³ Institute for Sustainable Plant Protection, National Research Council of Italy, 80055 Portici (NA), Italy; ⁴ Department of Agricultural, Forest and Food Sciences (DISAFA), University of Torino, 10095 Grugliasco (TO), Italy.

CONTACT Fernanda Colombari ✉ fernanda.colombari@unipd.it

Rapid on-site identification of the biocontrol agent of the Asian chestnut gall wasp

Abstract In classical biocontrol programs a rapid and correct identification of the introduced antagonist is a key issue during both the release and establishment monitoring phases. It is often difficult to distinguish morphologically cryptic species or immature stages, and, in this case, an accurate diagnosis can be provided by molecular diagnostic methods. Among the conventional and real-time PCR based methods, loop-mediated isothermal amplification (LAMP) is a particularly suitable technique as it allows a rapid amplification of target DNA directly in the field. During the program implemented in Italy against the Asian Chestnut Gall Wasp (ACGW) *Dryocosmus kuriphilus*, we developed a real-time LAMP assay, combined with a simple DNA extraction, for rapid in-field identification of larvae, pupae and adults of the biocontrol agent, the parasitoid *Torymus sinensis*. Validation of the assay comprised adults as well as preimaginal stages of parasitoids obtained from ACGW galls collected from different localities and results confirmed the effectiveness of the LAMP assay to rapidly and specifically identify the target parasitoid in the field. This assay will be a valuable tool for quick on-site checking of the parasitism rate.

Key words: *Torymus sinensis*, *Dryocosmus kuriphilus*, LAMP (Loop Mediated Isothermal Amplification), ACGW, biocontrol, parasitism rate,

Introduction

The measure of the success of classical biocontrol programs is largely determined by whether and to what extent the target pest's density changes after the establishment of introduced natural enemies (Guit, Wratten, & Barbosa, 2000; Hoddle, 2004; Mahr, Whitaker, & Ridgway, 2008). Identification, preliminary safety testing and release of control agents are undoubtedly matters of great concern (Van Driesche & Hoddle, 2000). Nonetheless, once the agent is approved for release, an important issue is quantifying post-release the effectiveness of parasitoids and predators in reducing pest abundance (Stiling & Cornelissen, 2005; Furlong & Zalucki, 2010). To objectively assess the effectiveness of biocontrol in terms of costs and benefits relative to conventional control, standardized measures of success are needed in post-release

surveys repeated over time (Sweetman, 1935; Van Driesche & Hoddle, 2000). The evaluation of parasitism rate can be performed following ‘rearing’ or ‘dissection’ methods applied to field-collected insects or plant material if host species feed in concealed or semi-concealed situations (i.e. leaf miners, gallmakers, borers). The former method can take a long time and could be delayed by many factors (i.e. diapause, weather conditions), whereas the latter is relatively quick and can guide decisions on whether more or different agents should be released (Day, 1994; Guit et al., 2000).

Whichever method is adopted, a correct identification of the parasitoids obtained is the critical step in the evaluation phase of the biocontrol program implemented, as it is a decisive factor when assessing the efficacy and host specificity of control agents and their possible interactions with the pest/native natural enemy complex (Delucchi, Rosen, & Schlinger, 1976; Agustí et al., 2005; Garipey, Kuhlmann, Gillott, & Erlandson, 2008). Strong support in systematics and taxonomy is essential to correctly identify parasitoids at the species level (Van Driesche & Hoddle, 2000). However, using morphological features to distinguish closely related taxa or members of cryptic species complexes is often difficult, if not impossible, especially for immature stages (Dawah & Rothfritz, 1996; Agustí et al., 2005; Mathé-Hubert, Gatti, Poirié, & Malausa et al., 2013). Several molecular diagnostic methods that have been developed and largely implemented over the last 20 years can assist, complement and even replace morphologically based approaches (Agustí et al., 2005; Garipey et al., 2008; Jenkins, Chapman, Micallef, & Reynolds, 2012).

In particular, the loop-mediated isothermal amplification (LAMP) is based on specific amplification of target DNA without the need for thermal cycling steps, thus allowing reactions to be performed in a portable heating block (Notomi et al., 2000; Tomlinson, Barker, & Boonham, 2007). The possibility of using LAMP directly in the

field, with short reaction time and no need for a long training period for the staff involved in the survey, gives considerable advantages compared to other conventional or real-time PCR based methods (Jenkins et al., 2012; Tomlinson, Dickinson, & Boonham, 2010; Tomlinson et al., 2013;). As the DNA polymerase in LAMP reactions is not influenced by the co-presence of inhibitors or non-target DNA (Kogovšek et al., 2015; Lenarčič, Morisset, Mehle, & Ravnikar, 2013; Notomi et al., 2000) crude DNA extracts can be used, thus increasing the usefulness and portability of the method in the field (Danks & Boonham, 2007).

The Asian Chestnut Gall Wasp (ACGW) *Dryocosmus kuriphilus* Yasumatsu (Hymenoptera: Cynipidae), native to China is an invasive species and included in the quarantine list of European Union (EC, 2000). After being accidentally introduced into Japan, Korea, North America, and Nepal (in the forties, fifties, seventies, and nineties, respectively), it arrived in north-western Italy in 2002 and then spread rapidly throughout Italy and several European countries (Slovenia: 2005; France: 2005; Hungary and Switzerland: 2009; Spain, Croatia and the Netherlands: 2010; Czech Republic: 2012; Austria and Germany: 2013; Portugal and Turkey: 2014; United Kingdom: 2015; Belgium: 2016) (EPPO, 2015; EPPO, 2016a, 2016b). ACGW represents a very serious threat to chestnut stands in all the invaded countries as high numbers of galls on leaves and shoots reduces tree vigour and severely affects nut and timber productivity (Kato & Hijii, 1997; Maltoni, Mariotti, & Tani, 2012; Battisti, Benvegnù, Colombari, & Haack, 2014). Management of ACGW infestations, after the unsuccessful attempts to reduce pest densities by other measures, currently relies only on classical biological control methods (Moriya, Inoue, & Mabuchi, 1989). The release of the parasitoid *Torymus sinensis* Kamijo (Hymenoptera: Torymidae) from the native region of the gall wasp has been shown to reduce damage below a tolerable threshold

level in Japan (less than 30%; Gyoutoku & Uemura, 1985; Moriya, Shiga, & Adachi, 2003) as well as in the USA, where a decline of pest population density has been also reported (Cooper & Rieske, 2007). In Italy, adults of *T. sinensis* were released in the first introduction site of ACGW starting in 2005, and later in all the other invaded areas (Quacchia, Moriya, Bosio, Scapin, & Alma, 2008; MiPAAF, 2010).

We developed a real-time LAMP assay combined with a crude DNA extraction for the identification directly in the field of larvae, pupae and adults of *T. sinensis*. The method was developed in order to quickly identify the preimaginal stages of the parasitoid and to evaluate the parasitism rate, so as to aid prompt management decisions regarding possible further releases of the parasitoid. Moreover, it may help the quality assessment of the parasitoid rearing program.

Materials and methods

The real-time LAMP assay was first developed using pure DNA extracted from voucher specimens by a salting out protocol. Then, the assay was validated using crude DNA extracted with a simple procedure from adults emerged from the galls (rearing method) and from preimaginal stages obtained by dissecting collected galls from the field.

Insect material

For the LAMP protocol development, 13 *T. sinensis*, 29 other chalcid adult parasitoids belonging to the same genus, family, and superfamily (21, 1, and 7 species, respectively; Table 1) as well as two *D. kuriphilus* were used as voucher specimens for molecular characterization. All the specimens were stored in 70% alcohol after their emergence from chestnut or oak galls (i.e. *T. geranii* Walker emerged from galls of *Biorhiza pallida* Olivier) collected in pure or mixed chestnut stands in Veneto (Crespano

del Grappa - TV), Piemonte (Avigliana - TO), and Campania (Serino - AV, Sicignano degli Alburni - SA) regions.

Validation of the LAMP protocol was first tested on 30 adults of *T. sinensis* and 15 adults of other native chalcid parasitoids belonging to Torymidae (*Megastigmus dorsalis* Fabricius and *Glyphomerus stigma* Fabricius), Eurytomidae (*Eurytoma pistaciae* Rondani and *E. brunniventris* Ratzeburg) and Eupelmidae (*Eupelmus urozonus* Dalman, *E. annulatus* Nees, *E. rostratus* Ruschka). Three adults of *D. kuriphilus* were also tested. All these specimens emerged from chestnut galls collected during the summer and winter at the rearing centre of the Veneto Region (Crespano del Grappa); *G. stigma* emerged from galls of *Diplolepis rosae* Linnaeus. In addition, a sample of 10 to 13 adults of *T. sinensis*, obtained from each of 8 Italian and 1 French population were further tested (N=101), including a few individuals from Korea (N=6) and Japan (N=3). All the specimens were first identified using various diagnostic morphological characters (Kamijo, 1982; de Vere Graham & Gijswijt, 1998).

The LAMP assay was then tested on larvae and pupae of the populations for which they were available (Table 2). At least 1,000 galls were collected during late winter at 7 Italian release sites in 4 regions (Veneto, Valle d'Aosta, Piemonte, Liguria) and one site in France. After collection, a subsample of 300 galls for each site was stored at 4° C to arrest the development of gall inhabitants at larval or pupal stages. Galls randomly chosen from each subsample were then dissected. Dissections were carried out until between 12 and 16 immature individual parasitoids were obtained and the number dissected recorded.

DNA extraction

Two types of DNA extractions were performed. For the LAMP assay development, a pure DNA extraction was carried out on 42 adult parasitoids (Table 1) using a salting

out protocol (Patwary, Kenchington, Bird, & Zouros, 1994). Conversely, for the LAMP assay validation, a crude DNA extraction was performed following a simple protocol intended for applications under field settings on: i) the adult specimens reported in the previous section i.e. 30 adults of *T. sinensis* and 18 adults of other chalcid parasitoids; 110 adults of *T. sinensis* from 11 different populations; ii) the immatures dissected from galls (Table 2). Whole insect bodies were individually placed into 1.5-ml Eppendorf tubes containing 200 µl of double-distilled water and ground with a plastic sterile pestle to obtain a crude homogenate. Tubes were then shaken by hand without vortexing for few seconds before taking the volume to be tested. Approximate DNA concentrations were determined at 260 nm using the Nano-drop 2000 spectrophotometer (Nano-drop Technologies, Wilmington, DE, USA).

Design of LAMP primers

Two sets of primers satisfying LAMP requirement criteria were designed (Notomi et al., 2000). A first set of primers, specific for *T. sinensis*, was developed on the sequences of internal transcribed spacer 2 (ITS2) in the nuclear ribosomal region. Primers were designed on the *T. sinensis* sequences retrieved from GenBank (accession numbers AB200273, AB200274, and AB200275), and aligned with homologous sequences of other species of Torymidae and Pteromalidae (i.e., *T. geranii*, GenBank accession number AB200280; *T. flavipes*, GenBank accession numbers HM574233 and HM574237; *Mesopolobus xanthocerus* (Thomson), GenBank accession number HM573972). In preliminary analyses (results not shown), some falsepositives with *T. affinis* were obtained when loop primers only were included in the reactions, although the use of these primers is usually recommended to improve both the specificity and the speed of the reaction (Nagamine, Hase, & Notomi, 2002). These primers were excluded from the primer set.

To avoid falsenegative results (Tomlinson et al., 2010), and to check quality of both DNA and reagents, a set of primers, hereafter referred to as external amplification control (EAC) primer set (D'Agostino, 2013), was designed on the 28S region of other chalcid wasps using sequences retrieved from Munro et al. (2011). Preliminary tests showed a wider inclusiveness of this primer set. Nonetheless, this was not a matter of concern because the purpose of the design was to ensure reliability of the assay by excluding the presence of inhibitors or suboptimal reaction conditions (i.e. not sufficient DNA template).

All the sequences alignments were edited using the ClustalW algorithm in the software MEGA, version 6, (Tamura, Stecher, Peterson, Filipski, & Kumar, 2013). Primers were designed using LAMP Primer Explorer software (version 4; Fujitsu System Solutions Ltd., Tokyo, Japan) and synthesized by Invitrogen.

LAMP development and diagnostic performance

The DNA extracted by the salting out method from the specimens listed in Table 1 was used for the evaluation of the sensitivity and specificity of both the primer sets. The reaction mixture for the *T. sinensis* LAMP test contained 15 µl of Isothermal MasterMix 1× (OptiGene), 0.2 µM of each external primer, 2 µM of each internal primer, and 1 µl of template DNA. The same conditions were used for the EAC assay, with a further addition of two loop primers, 1 µM each. Total volume for both reactions was 25 µl. LAMP reactions were performed on a Genie II device (OptiGene).

Optimal conditions for LAMP reaction were determined by performing runs in a range of temperatures for different time periods. The identity of amplification products was evaluated in terms of annealing temperature determined through the measure of fluorescence during a slow annealing step (0.05°C/s) starting at 98°and ending at 84°C.

The sensitivity of both the assays was tested through 10-fold serial dilutions of template DNA in double-distilled water, starting from a higher concentration of 10 ng μl^{-1} to a lower concentration of 100 fg μl^{-1} . Each run contained double-distilled water as negative control (Tomlinson et al., 2013). The specificity of both the *T. sinensis* and the EAC primer sets was evaluated using the specimens reported in Table 1. Both sensitivity and specificity were tested on pure DNA extracted by the salting out method.

Validation of LAMP with crude DNA

Evaluation of the specificity of the LAMP assay on crude DNA was first performed on the 48 specimens (30 adults of *T. sinensis* and 18 adults of other chalcid parasitoids) emerged at the rearing centre. Reactions were repeated three times for each specimen to confirm the diagnosis. For those specimens that were not amplified by the *T. sinensis*-specific set of primers, a EAC primer set test was performed in order to ensure quality of both DNA and reagents. The assay was then tested on a further 110 adults of *T. sinensis* belonging to 11 different populations and 80 larvae and 25 pupae obtained by the dissection method from 8 populations (Table 2).

To confirm the results obtained by LAMP assays, 20 specimens (10 immature and 10 adults), randomly selected among the above mentioned 263 samples, were identified by sequencing of the DNA barcode region (Hebert, Cywinska, & Ball, 2003). For each specimen, 100 μl of the crude DNA extract was used for a further DNA extraction using Qiagen DNeasy Plant mini kit (Valencia, CA, USA). The DNA obtained was then eluted in 50 μl of AE buffer and directly used to amplify and sequence the barcode region of the *cox1* gene (Hebert et al., 2003), using the universal primers and the amplification condition as reported in Folmer, Black, Hoeh, Lutz, & Vrijenhoek (1994). PCR products were checked through electrophoresis on 1.0% agarose gels stained with SYBR® (Invitrogen) and then purified using **exonuclease and**

215 antarctic phosphatase (GE Healthcare). PCR products were then sequenced at BMR
216 Genomics Service (Padova, Italy).

217 *Statistical analyses*

218 Throughout the text, temperature values are expressed in degrees Celsius and time
219 values in minutes and seconds. All average values are reported as mean $\pm$ standard
220 deviation (SD), unless otherwise specified.

221 One-way analysis of variance (ANOVA), followed by a Tukey's HSD (Honest
222 Significant Differences) test, was used to compare the mean annealing temperatures of
223 the 110 *T. sinensis* adults from the eleven different populations. An alpha level of 0.05
224 was considered statistically significant. Statistical analyses were performed using
225 STATISTICA, version 8 (Statsoft Inc., Tulsa OK, USA).

226 **Results**

227 *LAMP primers design*

228 The designed *T. sinensis* primer set contained two external primers (TS_F3 and TS_B3)
229 and two inner primers (TS_FIP and TS_BIP). The EAC primer set was composed of two
230 external (CH_F3 and CH_B3) and two internal primers (CH_FIP and CH_BIP), with
231 the addition of two loop primers (CH_Fl and CH_BL) in order to accelerate the
232 amplification reaction. Primer sequences for both the primer sets are reported in Table
233 3.

234 *LAMP development and diagnostic performance*

235 Optimal temperature and running time for both the LAMP reactions were an isothermal
236 condition of 67°C maintained for 19 and 23 min. for the *T. sinensis* and the EAC LAMP
237 assays, respectively. The specificity and the sensitivity of both primer sets and the

diagnostic performance of the LAMP assays were then tested on the DNA extracted by the salting out method.

Specific primers worked exclusively on *T. sinensis* DNA and did not amplify any of the non-target sequences. All the 13 *T. sinensis* DNA samples were correctly identified while EAC primers amplified all the 44 DNA samples analyzed. Amplification was not observed in the negative control reactions. The serial dilutions of DNA tested with both the assays, showed an analytical sensitivity of 10 pg μl^{-1} . The annealing temperatures of the amplification products were $88.82 \pm 0.09^\circ\text{C}$ and $87.51 \pm 0.18^\circ\text{C}$ for *T. sinensis* and EAC LAMP assays, respectively.

Validation of LAMP with crude DNA

Validation of the method on 48 adult parasitoids showed that 30 *T. sinensis* individuals were positive to the *T. sinensis* assay, whereas all tested insects were positive to the EAC assay. Average positive reaction times of crude DNA were $16:34 \pm 00:42$ for *T. sinensis* assay, and $14:14 \pm 1:43$ for EAC assay. Annealing temperatures were $88.98 \pm 0.06^\circ\text{C}$ and $87.40 \pm 0.10^\circ\text{C}$ for *T. sinensis* and EAC assays, respectively. There was a slight variation in the starting average DNA concentrations of crude extracts ($1.17 \pm 0.17 \text{ ng } \mu\text{l}^{-1}$), reaction times ($16:28 \pm 1:47$), and mean annealing temperatures ($88.84^\circ\text{C} \pm 0.14^\circ\text{C}$) among the 11 populations of *T. sinensis*. Interestingly, the mean annealing temperature of *T. sinensis* adults differed significantly among populations ($F_{(10, 99)} = 2.04$, $p < 0.05$), with the Korean specimens showing the highest average value, although the Tukey's test was not significant for any pair of means.

Forpreimaginal stages (larvae and pupae) from 8 available populations (Table 2), average DNA concentrations of crude extracts, reaction times, and mean annealing temperatures were: $2.72 \pm 1.27 \text{ ng } \mu\text{l}^{-1}$, $16:47 \pm 1:40$, $88.84^\circ\text{C} \pm 0.17^\circ\text{C}$ for larvae; $2.05 \pm 0.07 \text{ ng } \mu\text{l}^{-1}$, $15:21 \pm 1:52$, $88.76^\circ\text{C} \pm 0.16^\circ\text{C}$ for pupae. At some sites, it was recorded

a low number of larval samples that generated a positive amplification out of the total number of samples tested. DNA barcoding of the 20 samples confirmed the accuracy of the LAMP identification. Positive LAMP results always corresponded to *T. sinensis*, whereas negative results corresponded to species belonging to the genera *Eupelmus* (Hymenoptera: Eupelmidae) and *Mesopolobus* (Hymenoptera: Pteromalidae).

Discussion

The LAMP assay enabled real-time detection of *T. sinensis*, the biocontrol agent of the ACGW, through a simple procedure designed to give rapid on-site results for samples without the need for time-consuming analyses or rearing in the laboratory. The use of a quick technique, together with the simple equipment required and the possibility to discriminate insect material obtained from various life stages, make the method valuable for making decisions when evaluating effectiveness of biocontrol programs directly in the field.

The whole procedure developed in this study, from sample preparation to the evaluation of amplification products, was completed in less than 30 minutes. The specific LAMP primer set showed high specificity to *T. sinensis*, with no positive reactions when other species were tested. However, we cannot exclude that the primer set designed specifically for *T. sinensis* works for the closely related *T. beneficus*, which cannot be discriminated on the basis of morphological characters (Yara, 2004).

Unfortunately, it has not been possible to retrieve and test any specimen of *T. beneficus* to better validate the *T. sinensis* primer set. For this purpose, it would be interesting to look at the possibility to obtain an annealing temperature specific for the DNA region of the target species, as a way to unambiguously identify the samples and, concurrently, to detect possible variants (Ririe, Rasmussen, & Wittwer, 1997). In our case, the Korean population showed the highest mean annealing temperature, although this result needs

to be supported by a higher number of samples. If confirmed, this result may reveal the occurrence of two different strains of *T. sinensis* in Korea and Japan, as the European populations came from Japan, given their introduction history (Murakami, Ohkubo, Moriya, Gyoutoku, Kim, & Kim, 1995; Yara, 2004; Quacchia et al., 2008).

The diagnostic sensitivity level of the assay was high enough to correctly identify the parasitoid in any of the life stages considered, without the need for complex DNA extraction and quantification. Successful outcomes were easily obtained by simply grinding the insects in double distilled water and by directly adding these crude homogenates to pre-prepared strips containing isothermal field stable reagents. Results were positive when specimens both stored in alcohol and freshly collected were used, demonstrating that even DNA of relatively low purity gives positive amplification products (Huang, Hsu, Haymer, Lin, & Wu, 2009). We found that the negative results were successfully amplified by the EAC primer set, minimizing the risk of false negative results.

The samples tested were not sufficient to give reliable estimates of parasitism. However, the proportion of positive larval samples roughly reflects the parasitoid success at the different sites. A higher occurrence of *T. sinensis* was indeed recorded where the parasitoid was released first. Alternatively, the high proportion of positive pupal samples can be explained by the period of gall collection (i.e. late winter). In this period only *T. sinensis* occupies the galls as the other parasitoids generally use other hosts for overwintering (Luo, Huang, & Liao, 1987; Shiga, 2009). A proper identification of *T. sinensis* using a reliable on-site molecular technique is of great importance as the release of the parasitoid is, at present, the only viable long-term management option against *D. kuriphilus* in the invaded areas (Yara, 2006; Gibbs et al., 2011). Although the biocontrol of ACGW is successful in those areas where *T. sinensis*

has been introduced (Gyoutoku & Uemura, 1985; Moriya et al., 2003; Cooper & Rieske, 2007; Ferracini et al., 2015a), many recently invaded regions have now to deal with this problem (EPPO, 2014; EPPO, 2015). Consequently, a correct identification of both preimaginal and adult stages of the parasitoid is fundamental during all the phases of the biological control program (Danks, 1988; Gordh & Beardsley, 1999). This is particularly true if we consider that other parasitoid species, such as local oak gall wasp parasitoids, are rapidly recruited as enemies by ACGW once it arrives in a new area and that *T. sinensis* may cause unwanted non-target effects (Aebi et al., 2006; Quacchia et al., 2012; Matošević & Melika, 2013; Panzavolta et al., 2013; Palmeri et al., 2014; Ferracini et al., 2015b).

The LAMP assay we developed is suitable for diagnostic as well as research use (Lenarčič et al., 2013) as it can be used both for a rapid quality check of the released stock in new areas and a quick in-field monitoring of the parasitism rate of the biological control agent. Moreover, this method bypasses the need for specialist knowledge or a long period of training for the staff involved (Jenkins et al., 2012) avoiding, in particular, the long procedure in the morphological identification of preimaginal stages.

Acknowledgements

This work relies on the contribution of several persons. G. Narduzzo (Veneto Region's Forest Service), P. Paolucci and P. Dall'Ara (DAFNAE Padova), C. Salvadori (Fondazione Mach Trento), the staff of the rearing area Centro Polifunzionale Onè (Crespano del Grappa), Veneto Region's Phytosanitary and Forest Services provided precious support during field activities. E. Ferrari, E. Gonella, M. A. Saladini (DISAFA Torino) and L. Iodice (IPP Napoli) provided kind help for collection and shipment of insect specimens. Myron Zalucki carefully revised and edited the text. Two anonymous reviewers significantly improved the quality of the manuscript.

Funding

The research was funded by a grant of the Ministero delle Politiche Agricole Alimentari e Forestali of the Italian government and by the Regione del Veneto Servizi Fitosanitari.

References

- Aebi, A., Schönrogge, K., Melika, G., Alma, A., Bosio, G., Quacchia, A., ... Stone, G. (2006). Parasitoid Recruitment to the Globally Invasive Chestnut Gall Wasp *Dryocosmus kuriphilus*. In K. Ozaki, J. Yukawa, T. Ohgushi & P. Price (Eds.), *Galling arthropods and their associates: Ecology and Evolution* (pp. 103-121). Tokio: Springer Japan.
- Agustí, N., Bourguet, D., Spataro, T., Delos, M., Eychenne, N., Folcher, L., & Arditi, R. (2005). Detection, identification and geographical distribution of European corn borer larval parasitoids using molecular markers. *Molecular Ecology*, 14, 3267-3274.
- Battisti, A., Benvegnù, I., Colombari, F., & Haack, R. A. (2014). Invasion by the chestnut gall wasp in Italy causes significant yield loss in *Castanea sativa* nut production. *Agricultural and Forest Entomology*, 16, 75-79.
- Cooper, W. R., & Rieske, L. K. (2007). Review of the historic and current status of the Asian chestnut gall wasp in North America. *Journal of the American Chestnut Foundation*, 21(2), 28-34.
- EC (2000) EC Council Directive 2000/29/EC of 8 May 2000 on protective measures against the introduction into the community of organisms harmful to plants or plant products and against their spread within the community. *Official Journal of the European Communities*, 50, 1-159.
- D'Agostino, M. (2013). Quality control in the analytical laboratory: analysing food-and waterborne viruses. In N. Cook (Ed.), *Viruses in food and water: risks, surveillance and control* (pp. 126-138). New Delhi: Woodhead Publishing.

364 Danks, C., & Boonham, N. (2007). Purification method and kits. Patent
365 WO/2007/104962.

366 Danks, H. V. (1988). Systematics in Support of Entomology. *Annual Review of*
367 *Entomology*, 33, 271-294.

368 Dawah, H. A., & Rothfritz, H. (1996). Generic-level identification of final instar larvae
369 of Eurytomidae and their parasitoids associated with grasses (Poaceae) in N.W.
370 Europe (Hymenoptera: Braconidae, Eulophidae, Eupelmidae, Eurytomidae,
371 Ichneumonidae, Pteromalidae). *Journal of Natural History*, 30, 1517-1526.

372 Day, W. H. (1994). Estimating mortality caused by parasites and diseases of insects:
373 comparisons of the dissection and rearing methods. *Environmental Entomology*,
374 23, 543-550.

375 Delucchi, V., Rosen, D., & Schlinger, E. I. (1976). Relationship of systematics to
376 biological control. In C. B. H. S. Messenger (Ed.), *Theory and Practice of*
377 *Biological Control* (pp. 81-91). New York: Academic Press.

378 de Vere Graham, M. W. R., & Gijswijt, M. J. (1998). Revision of the European species
379 of *Torymus* Dalman (s. lat) (Hymenoptera: Torymidae). Leiden: Nationaal
380 Natuurhistorisch Museum.

381 EPPO (European and Mediterranean Plant Protection Organization) (2015). First report
382 of *Dryocosmus kuriphilus* in the United Kingdom. EPPO Reporting Service
383 (Report No. 6). Retrieved January 27, 2016, from [https://gd.eppo.int/
384 reporting/article-4773](https://gd.eppo.int/reporting/article-4773)

385 EPPO (European and Mediterranean Plant Protection Organization) (2016a). PQR -
386 EPPO database on quarantine pests. Retrieved January 27, 2016, from the EPPO
387 Databases.

388 EPPO (European and Mediterranean Plant Protection Organization) (2016b). First

389 report of *Dryocosmus kuriphilus* in Belgium. EPPO Reporting Service (Report
390 No. 2). Retrieved March 24, 2016,

391 Ferracini, C., Gonella, E., Ferrari, E., Saladini, M. A., Picciau, L., Tota, F., Pontini, M.,
392 & Alma, A. (2015a). Novel insight in the life cycle of *Torymus sinensis*,
393 biocontrol agent of the chestnut gall wasp. *BioControl*, 60, 169-177.

394 Ferracini, C., Ferrari, E., Saladini, M. A., Pontini, M., Corradetti, M., & Alma, A.
395 (2015b). Non-target host risk assessment for the parasitoid *Torymus sinensis*.
396 *BioControl*, 60, 583–594.

397 Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for
398 amplification of mitochondrial cytochrome c oxidase subunit I from diverse
399 metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294-
400 299.

401 Furlong, M. J., & Zalucki, M. P. (2010). Exploiting predators for pest management: the
402 need for sound ecological assessment. *Entomologia Experimentalis et Applicata*,
403 135, 225-236.

404 Garipey, T., Kuhlmann, U., Gillott, C., & Erlandson, M. (2008). A large-scale
405 comparison of conventional and molecular methods for the evaluation of host–
406 parasitoid associations in non-target risk-assessment studies. *Journal of Applied*
407 *Ecology*, 45, 708-715.

408 Gibbs, M., Schönrogge, K., Alma, A., Melika, G., Quacchia, A., Stone, G., & Aebi, A.
409 (2011). *Torymus sinensis*: a viable management option for the biological control
410 of *Dryocosmus kuriphilus* in Europe? *BioControl*, 56, 527-538.

411 Gordh, G., Beardsley, J. W. (1999). Taxonomy and Biological Control In: T. S.
412 Bellows, T. W. Fisher, L. E. Caltagirone, D. L. Dahlsten, G. Gordh, & C. B.

413 Huffaker (Eds.), *Handbook of Biological Control* (pp. 45-55). San Diego:
414 Academic Press.

415 Guit, G. M., Wratten, S. D., & Barbosa, P. (2000). Success in conservation biological
416 control of arthropods. In: G. Gurr & S. Wratten (Eds.), *Biological control:
417 Measures of success* (pp. 105-132). Dordrecht: Springer Netherlands.

418 Gyoutoku, Y., & Uemura, M. (1985). Ecology and biological control of the chestnut
419 gall wasp, *Dryocosmus kuriphilus* Yasumatsu (Hymenoptera: Cynipidae). 1.
420 Damage and parasitization in Kumamoto Prefecture. *Proceedings of the
421 Association for Plant Protection of Kyushu*, 31, 213–215.

422 Hebert, P. D., Cywinska, A., & Ball, S. L. (2003). Biological identifications through
423 DNA barcodes. *Proceedings of the Royal Society of London B: Biological
424 Sciences*, 270, 313-321.

425 Hoddle, M. S. (2004). Restoring balance: using exotic species to control invasive exotic
426 species. *Conservation Biology*, 18, 38-49.

427 Huang, G. C., Hsu, J. C., Haymer, D., Lin, G. C., & Wu, W. J. (2009). Rapid
428 identification of the Mediterranean fruit fly (Diptera: Tephritidae) by loop-
429 mediated isothermal amplification. *Journal of Economic Entomology*, 102,
430 1239-1246.

431 Jenkins, C., Chapman, T. A., Micallef, J. L., & Reynolds, O. L. (2012). Molecular
432 techniques for the detection and differentiation of host and parasitoid species
433 and the implications for fruit fly management. *Insects*, 3, 763-788.

434 Kato, K., & Hijii, N. (1997). Effects of gall formation by *Dryocosmus kuriphilus*
435 Yasumatsu (Hym., Cynipidae) on the growth of chestnut trees. *Journal of
436 Applied Entomology*, 121, 9-15.

437 Kamijo, K. (1982). Two new species of *Torymus* (Hymenoptera, Torymidae) reared
438 from *Dryocosmus kuriphilus* (Hymenoptera, Cynipidae) in China and Korea.
439 *Kontyû*, 50, 505–510.

440 Kogovšek, P., Hodgetts, J., Hall, J., Prezelj, N., Nikolić, P., Mehle, N., ... Boonham, N.
441 (2015). LAMP assay and rapid sample preparation method for on-site detection
442 of flavescence dorée phytoplasma in grapevine. *Plant Pathology*, 64, 286-296.

443 Lenarčič, R., Morisset, D., Mehle, N., & Ravnikar, M. (2013). Fast real-time detection
444 of potato spindle tuber viroid by RT-LAMP. *Plant Pathology*, 62, 1147-1156.

445 Luo, Y. Q., Huang, J. F., & Liao, D. X. (1987). Studies on the distribution and biology
446 of *Torymus sinensis* Kamijo. *Journal of Beijing Forestry University*, 9, 45-57.

447 Mahr, D. L., Whitaker, P., & Ridgway, N. (2008). *Biological control of insects and*
448 *mites: An introduction to beneficial natural enemies and their use in pest*
449 *management*, 116 pp. Madison, WI: Cooperative Extension Publishing,
450 University of Wisconsin, USA.

451 Maltoni, A., Mariotti, B., & Tani, A. (2012). Case study of a new method for the
452 classification and analysis of *Dryocosmus kuriphilus* Yasumatsu damage to
453 young chestnut sprouts. *iForest - Biogeosciences and Forestry*, 5, 50-59.

454 Mathé-Hubert, H., Gatti, J.-L., Poirié, M., & Malausa, T. (2013). A PCR-based method
455 for estimating parasitism rates in the olive fly parasitoids *Psytalia concolor* and
456 *P. lounsburyi* (Hymenoptera: Braconidae). *Biological Control*, 67, 44-50.

457 Matošević, D., & Melika, G. (2013). Recruitment of native parasitoids to a new invasive
458 host: First results of *Dryocosmus kuriphilus* parasitoid assemblage in Croatia.
459 *Bulletin of Insectology*, 66, 231-238.

460 MiPAAF - Ministero delle Politiche Agricole Alimentari e Forestali (2010). Piano del
461 settore castanicolo 2010/2013 - 2. Riferimenti tecnici di attuazione della lotta

biologica al *Dryocosmus kuriphilus* del castagno con *Torymus sinensis*.

Retrieved January 27, 2016, from <https://www.politicheagricole.it/flex/cm/>

[pages/ServeBLOB.php/L/IT/IDPagina/3277](https://www.politicheagricole.it/flex/cm/pages/ServeBLOB.php/L/IT/IDPagina/3277)

Moriya, S., Inoue, K., & Mabuchi, M. (1989). The use of *Torymus sinensis* to control chestnut gall wasp, *Dryocosmus kuriphilus*, in Japan. *FFTC Technical Bulletin*, 118, 1-12.

Moriya, S., Shiga, M., & Adachi, I. (2003). Classical biological control of the chestnut gall wasp in Japan. In: R. G. Van Driesche (Ed.), *Proceedings of the 1st international symposium on biological control of arthropods* (pp. 407-415). Washington: USDA Forest Service.

Munro, J. B., Heraty, J. M., Burks, R. A., Hawks, D., Mottern, J., Cruaud, A., ... Jansta, P. (2011). A molecular phylogeny of the Chalcidoidea (Hymenoptera). *PLoS ONE*, 6, e27023.

Murakami, Y., Ohkubo, N., Moriya, S., Gyoutoku, Y., Kim, H. C., & Kim, K. J. (1995). Parasitoids of *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae) in South Korea with particular reference to ecologically different types of *Torymus* (*Syntomaspis*) *sinensis* (Hymenoptera: Torymidae). *Applied Entomology and Zoology*, 30, 277-284.

Nagamine, K., Hase, T., & Notomi, T. (2002). Accelerated reaction by loop-mediated isothermal amplification using loop primers. *Molecular and cellular probes*, 16, 223-229.

Notomi, T., Okayama, H., Masubuchi, H., Yonekawa, T., Watanabe, K., Amino, N., & Hase, T. (2000). Loop-mediated isothermal amplification of DNA. *Nucleic Acids Research*, 28, e63.

486 Palmeri, V., Cascone, P., Campolo, O., Grande, S. B., Laudani, F., Malacrinò, A., &
 487 Guerrieri, E. (2014). Hymenoptera wasps associated to the Asian Gall Wasp
 488 (AGW) of chestnut *Dryocosmus kuriphilus* in Calabria (Italy). *Phytoparasitica*,
 489 42, 699-702.

490 Panzavolta, T., Bernardo, U., Bracalini, M., Cascone, P., Croci, F., Gebiola, M., ...
 491 Guerrieri, E. (2013). Native parasitoids associated with *Dryocosmus kuriphilus*
 492 in Tuscany, Italy. *Bulletin of Insectology*, 66, 195-201.

493 Patwary, M. U., Kenchington, E. L., Bird, C. J., & Zouros, E. (1994). The use of
 494 random amplified polymorphic DNA markers in genetic studies of the sea
 495 scallop *Placopecten magellanicus* (Gmelin, 1791). *Journal of Shellfish*
 496 *Research*, 13, 547-553.

497 Quacchia, A., Ferracini, C., Nicholls, J. A., Piazza, E., Saladini, M. A., Tota, F., ...
 498 Alma, A. (2012). Chalcid parasitoid community associated with the invading
 499 pest *Dryocosmus kuriphilus* in north-western Italy. *Insect Conservation and*
 500 *Diversity*, 6, 114-123.

501 Quacchia, A., Moriya, S., Bosio, G., Scapin, I., & Alma, A. (2008). Rearing, release and
 502 settlement prospect in Italy of *Torymus sinensis*, the biological control agent of
 503 the chestnut gall wasp *Dryocosmus kuriphilus*. *BioControl*, 53, 829-839.

504 Ririe, K. M., Rasmussen, R. P., & Wittwer, C. T. (1997). Product differentiation by
 505 analysis of DNA melting curves during the polymerase chain reaction.
 506 *Analytical Biochemistry*, 245, 154-60.

507 Shiga, M. (2009). Life history of an introduced parasitoid, *Torymus sinensis*, and
 508 dynamics of the host-parasitoid system. In: S. Moriya (Ed.), *Proceedings of the*
 509 *Japan-Italy Joint International Symposium 'A Global Serious Pest of Chestnut*

510 *Trees: Yesterday, Today and Tomorrow* (pp 21-22). Ibaraki: Asahi Printing
 511 Corporation, Japan.

512 Stiling, P., & Cornelissen, T. (2005). What makes a successful biocontrol agent? A
 513 meta-analysis of biological control agent performance. *Biological Control*, 34,
 514 236-246.

515 Sweetman, H. L. (1935). Successful examples of biological control of pest insects and
 516 plants. *Bulletin of Entomological Research*, 26, 373-377.

517 Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA6:
 518 molecular evolutionary genetics analysis version 6.0. *Molecular Biology and*
 519 *Evolution*, 30, 2725-2729.

520 Tomlinson, J. A., Barker, I., & Boonham, N. (2007). Faster, simpler, more-specific
 521 methods for improved molecular detection of *Phytophthora ramorum* in the
 522 field. *Applied and Environmental Microbiology*, 73, 4040-4047.

523 Tomlinson, J. A., Dickinson, M. J., & Boonham, N. (2010). Rapid detection of
 524 *Phytophthora ramorum* and *P. kernoviae* by two-minute DNA extraction
 525 followed by isothermal amplification and amplicon detection by generic lateral
 526 flow device. *Phytopathology*, 100, 143-149.

527 Tomlinson, J. A., Ostoj-Starzewska, S., Adams, I. P., Miano, D. W., Abidrabo, P.,
 528 Kinyua, Z., ... Boonham, N. (2013). Loop-mediated isothermal amplification
 529 for rapid detection of the causal agents of cassava brown streak disease. *Journal*
 530 *of Virological Methods*, 191(2), 148-154.

531 Van Driesche, R. G., & Hoddle, M. S. (2000) Classical arthropod biological control:
 532 Measuring success, step by step. In: G. Gurr & S. Wratten (Eds.), *Biological*
 533 *control: Measures of success* (pp. 39-75). Dordrecht: Springer Netherlands. .

534 Yara, K. (2004). Relationship between the introduced and indigenous parasitoids

535 *Torymus sinensis* and *T. beneficus* (Hymenoptera: Torymidae) as inferred from
536 mt-DNA (COI) sequences. *Applied Entomology and Zoology*, 39, 427-433.
537 Yara, K. (2006). Identification of *Torymus sinensis* and *T. beneficus* (Hymenoptera:
538 Torymidae), introduced and indigenous parasitoids of the chestnut gall wasp
539 *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae), using the ribosomal ITS2
540 region. *Biological Control*, 36, 15-21.

Table 1. Insect material tested for the development of the LAMP assays with pure DNA extracted by the salting out protocol.

Superfamily	Family	Species	Number of tested samples
Chalcidoidea	Torymidae	<i>Torymus sinensis</i>	13
		<i>affinis</i> *	4
		<i>auratus</i>	2
		<i>cyaneus</i> *	1
		<i>erucarum</i>	2
		<i>favardi</i> *	2
		<i>flavipes</i>	2
		<i>formosus</i> *	1
		<i>geranii</i>	2
		<i>notatus</i>	4
		<i>scutellaris</i>	1
		<i>Megastigmus dorsalis</i>	1
	Eupelmidae	<i>Eupelmus urozonus</i>	1
		<i>annulatus</i>	1
	Eurytomidae	<i>Eurytoma pistaciae</i>	1
		<i>brunniventris</i>	1
	Ichneumonidae	<i>Orthopelma mediator</i>	1
	Pteromalidae	<i>Mesopolobus tibialis</i>	1
	Ormyridae	<i>Ormyrus nitidulus</i>	1

* Indicates congeneric species of *T. sinensis* emerged from cynipid galls on oak.

546 Table 2. Individuals of *T. sinensis* from different populations tested for the validation of
547 the LAMP method with crude DNA. For each sample the number of samples that
548 generated a positive amplification out of the total number of samples tested is reported.

Country	Region	Site	Larvae	Pupae
France		St. Dalmas de Tende	8/10	2/2
Italy	Valle d'Aosta	Forte di Bard	4/7	5/5
	Piedmont	Avigliana	2/8	3/4
	Liguria	Millesimo	7/11	1/1
	Veneto	Cavaso del Tomba	2/14	-
		San Mauro di Saline	1/12	-
		Pianezze	2/15	-
		Seren del Grappa	2/3	11/13
Total			80	25

549

550 Table 3. Primers used for the loop-mediated isothermal amplification (LAMP) assays.

LAMP assay target	Primer	Sequence (5' – 3')
<i>Torymus sinensis</i>	TS_F3	CGCAAGATGGATGAGAGAGAG
	TS_B3	GCAAACAGAGAGCTCCGG
	TS_FIP	TCAAAACACTCACGAGGCGCGTCGC- TCGAAACAATGGCG
	TS_BIP	TACGCACACGCACACGCTACTCGAC- GCAAACAACACG
Chalcid wasps	CH_F3	GGTGAACATATGCCTGGTCAG
	CH_B3	TTCGCTTTACCAGATGAGACTC
	CH_FIP	CCGACGATCGATTGACGTCAGAC- GAAGTCAGGGGAAACC
	CH_BIP	ACTGGGTATAGGGGCGAAAGACTAA- TCAAGCGAGTGCCAGCTATC
	CH_FL	CGCTACGGACCTCCATCAG
	CH_BL	GAACCATCTAGTAGCTGGTTCC

551