AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Osteology, fossil record and palaeodiversity of the European lizards

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1627296> since 2017-05-19T14:48:55Z

Published version:

DOI:10.1163/15685381-00003085

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Villa, Andrea; Tschopp, Emanuel; Georgalis, Georgios L.; Delfino, Massimo.
Osteology, fossil record and palaeodiversity of the European lizards.
AMPHIBIA-REPTILIA. 38 pp: 79-88.
DOI: 10.1163/15685381-00003085

The publisher's version is available at:

<http://booksandjournals.brillonline.com/content/journals/10.1163/15685381-00003085>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

Osteology, fossil record and palaeodiversity of the European lizards

Andrea Villa^{1,*}, Emanuel Tschopp^{1,2,3}, Georgios L. Georgalis^{4,1}, Massimo Delfino^{1,5}

¹ Dipartimento di Scienze della Terra, Università degli Studi di Torino, via Valperga Caluso 35, 10125 Torino, Italy

² GeoBioTec, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Caparica, Portugal

³ Museu da Lourinhã, Rua João Luís de Moura, 2530-157 Lourinhã, Portugal

⁴ Department of Geosciences, University of Fribourg/Freiburg, Chemin du Musée 6, 1700 Fribourg, Switzerland

⁵ Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Edifici ICTA-ICP, Carrer de les Columnes s/n, Campus de la UAB, 08193 Cerdanyola del Vallès, Barcelona, Spain

* Corresponding author: a.villa@unito.it

Type of manuscript: Article

Total words: 4703 in the manuscript, 225 in the abstract

Abstract. The capability of palaeontologists to identify fossil remains of a particular group of vertebrates strongly depends on the knowledge they have of its comparative osteology and on the actual presence of diagnostic differences among the considered taxa. This could have a relevant influence on the study of palaeodiversity, since a low recognisability causes a loss of data when trying to reconstruct the history of taxa that lived on Earth in the past. Currently, more than 6000 extant species of lizards and worm lizards are known, and new ones continue to be discovered, mainly based on molecular data. But are we able to recognise this high diversity using osteology? As far as European taxa are concerned, the osteological recognisability of non-snake squamates is very low: only 31% of the extant European taxa can be identified based on their skeletal morphology. This is balanced partially by the fact that most recognisable taxa have been actually recognised in the fossil record, suggesting that the lost data are mainly due to the scarce knowledge of the comparative osteology of these reptiles and less influenced by other biases, such as taphonomic or collection biases. In this context, specimen-level phylogenetic analysis has proved to be a useful tool to identify diagnostic combinations of osteological features, at least for lacertid species, as evidenced by a case study focused on the genus *Lacerta*.

Keywords: Identification, Squamates, Reptiles, Palaeontology, Biodiversity

Introduction

Non-snake squamates (i.e., lizards and worm lizards) exist since at least the Middle Jurassic (Rage, 2013), and are represented by more than 6000 extant species worldwide (Uetz and Hošek, 2016) and a number of extinct taxa (for a summary of extinct species see the now dated review of Estes, 1983, but also Böhme and Ilg, 2003). In order to identify the current diversity of these reptiles, herpetologists can rely on various different tools, in particular external morphology, behaviour, molecular and distributional data. Lately, the rate of new lizard descriptions was rising considerably

(Pincheira-Donoso et al., 2013). But what about the palaeodiversity? How precisely can we recognise the diversity of taxa that lived on Earth in the past? Except for a few remarkable examples (e.g., specimens in amber that preserve also the integument like *Succinilacerta succinea* (Boulenger, 1917) or those recently described by Daza et al., 2016), vertebrate palaeontologists are mostly restricted to osteological features to identify fossil remains and study palaeodiversity. This makes the task of palaeoherpetologists much more complex and error-prone than that of neoherpetologists, given that it strongly depends on the degree of knowledge of the comparative osteology of the considered group and on the actual presence of diagnostic differences among the taxa. We here try to quantify the current osteological recognisability of the extant European non-snake squamates, in order to evaluate to what degree it affects our ability to comprehend the past diversity of these reptiles in the continent. The degree of recognisability is also evaluated at different hierarchical levels and within different clades, in order to understand where the information is scarcer and new studies are needed. Last but not least, we try to assess the utility of phylogenetic analyses to identify diagnostic osteological features for lizards, and lacertids in particular.

Materials and methods

We included 75 species of lizards and 3 worm lizards in our analysis. The included species correspond to the European amphibians and reptiles reported by Sillero et al. (2014), plus the lizards and worm lizards living on the Greek islands along the coast of Asiatic Turkey as reported by Speybroeck et al. (2016). Recently introduced taxa have been excluded (as e.g., *Chamaeleo africanus*). The definition of Europe largely follows Arnold and Ovenden (2002), but with the exclusion of Madeira and the Canary islands in the west, because they do not belong to the continent in geological terms.

The evaluation of the recognisability of the different taxa is based on existing bibliography, as well as on personal observations (see supplementary text S1 for a list of the studied specimens). Details about the osteology of European lizards and worm lizards have been mentioned in more than 90 papers,

either dealing directly with comparative osteology (e.g., Evans, 2008) or simply reporting features in the remarks of the identification of fossil material of extant taxa (e.g., Blain, 2009). For a complete list of these works, see supplementary text S2. In order to avoid circular reasoning (as outlined by Bell et al., 2010), geographical and/or chronological criteria are not considered useful for the identification of a taxon, since the range of a species may strongly vary during time.

The recognisable taxa are not always represented by species, but often by different kinds of higher taxonomic ranks. Because of this, the recognisability is here analysed in two ways: 1) in terms of Operational Taxonomic Units (OTUs), regardless whether they represent family, genus or species level; and 2) at the level of species, genus, and family separately. The first analysis tells us how many taxonomic bins can be recognised when having skeletons of all the 78 extant European lizards, in order to depict the recognisable diversity based only on osteology. This diversity is then compared with the actual one recognised by neoherpetologists using other tools, in which OTUs are represented by the 78 extant species of European lizards and worm lizards. The second analysis quantifies how many species, genera, and families of lizards can actually be diagnosed with osteological characters, and, assuming that diagnostic characters are present, in which taxon we need more work to be done to increase this recognisability. Finally, given the widespread individual variation in lizard skeletons, and thus the low probability to find unambiguous, osteological autapomorphies for the species, we implemented a case study to evaluate the utility of specimen-level phylogenetic analysis for the identification of diagnostic combinations of osteological features of the European species of the genus *Lacerta* (as defined by Arnold et al., 2007). We compiled a novel phylogenetic matrix based on osteology only, including 159 character statements taken from earlier works (see references in the supplementary text S2) and personal observations of ET. These character statements were scored in 37 OTUs consisting of single specimens (see supplementary text S1), which were identified before skeletonization, based on external morphology by the collectors or curators of the respective collections. The matrix was compiled in the software Mesquite v. 3.04 (Maddison and Maddison, 2015), and includes nine specimens of *Lacerta agilis*, nine specimens of *Lacerta bilineata*, five specimens of *Lacerta schreiberi*, four specimens of *Lacerta trilineata*, and four specimens of *Lacerta viridis*, which have all been scored based on first hand observations. The outgroup consists of specimens of the European lacertids *Timon lepidus* (1), *Podarcis muralis* (2), *Archaeolacerta bedriagae* (1), *Algyroides nigropunctatus* (1), and *Psammodromus algirus* (1). The matrix was analysed with TNT v.1.1 (Goloboff et al., 2008). We used an extended implied weighting approach (Goloboff, 2014) in order to reduce the influence of highly variable characters and

missing data. The tree search was performed using the New Technology Search, enabling all algorithms, and stabilizing the consensus tree five times with a factor of 75. In order to find all most parsimonious trees (MPTs), a second iteration of tree bisection and reconnection was performed.

Results

Recognisability

The recognisability of European lizards and worm lizards appears to be rather low (Tab. 1): the recognisable OTUs represent only 31% (24 out of 78 taxa) of the current European lizard diversity. A large number of lacertids and scincids, as well as all species of the genus *Anguis*, can only be identified at genus or even at family level (species complex in the case of *Anguis*). It has to be noted that agamids and chamaeleonids are poorly identifiable too, but since a single autochthonous species of each family is currently present in the continent, this has no evident effect on the count of the recognisability of extant taxa. Finally, the most recognisable group is Gekkota: all four species, belonging to four different genera, are clearly distinguishable from one another.

The second analysis shows that recognisability highly depends on the hierarchical levels. Whereas all families of lizards are recognisable based on osteological features, only 45% of the genera and 17% of the extant species are known to have diagnostic features in their skeletons (Fig. 1). Also the degree of recognisability varies greatly within the different families: nearly half (43%) of the scincid and lacertid species can only be identified as belonging to Scincidae and Lacertidae, and the two European species of agamids and chamaeleonids cannot be distinguished from other, non-European members of their family based on osteology. On the other hand, osteological characters of the anguids and blanids allow to recognise all European taxa at least at the genus level (Fig. 2).

As for the European fossil record, a large number of the recognisable OTUs have been identified (20 out of 24, 83%; 26% of the 78 extant species, meaning a loss of data of 74%; Tab. 1). Among the 24 OTUs, in fact, only those corresponding to *Euleptes europaea*, *Mediodactylus kotschy*, *Lacerta schreiberi* and worm lizards of the *Blanus strauchi* complex have never been recovered as fossils.

Phylogenetic analysis

The first run of the case study of *Lacerta* produced a single MPT of a length of 39.25405, and recovered the specimen of *T. lepidus* among the specimens of *Lacerta*. Given that all the recent phylogenetic analyses including these two genera find them as sister taxa (e.g., Carranza et al., 2004; Arnold et al., 2007; Kapli et al., 2011; Mendes et al., 2016), our result indicates that either we did not find enough osteological characters to unite *Lacerta* to the exclusion of *Timon*, or that the osteological variability is too high to obtain a meaningful tree topology based on osteological characters only. In order to test this, we performed a second analysis, including a constraint forcing the software to find all the specimens of *Lacerta* as a monophyletic group. This second run resulted in nine MPTs with a length of 39.80892. With the exclusion of a single specimen of *L. viridis*, all the other specimens form monophyletic clades with their respective species members in the strict consensus tree (Fig. 3).

Discussion

The impact of recognisability on the study of fossil European non-snake squamates

The knowledge we have of past ecosystems is incomplete, because of the different biases that cause loss of information in the palaeontological record (among others, Dunhill et al., 2012). Among them, taphonomic bias has a major effect, in particular when it comes to small animals (Soligo and Andrews, 2005; Cooper et al., 2006). When an excavation is not directly focused on the recovery of small vertebrate remains, a collection bias may also occur, caused by the difficulty of discovering and collecting

small bones without proper methods and tools. This latter issue is particularly significant when old works are considered, whereas an increased attention towards these fossils became established in XX century. Finally, also a historical bias probably influenced the fossil record of poorly studied vertebrates, such as lizards: studies on their osteology were, in fact, rare until about the 1970s, with a significant increase from the 1980s onwards (Fig. 4). This increase in the number of lizard osteology papers still continues: nearly 25% of all the descriptions used for the current study were published between 2010 and 2016. Obviously, this lack of osteological data until recently hampered the identification of fossil taxa, even if the increasing number of description is highly promising for the future. Also, current taxonomic trends based on molecular analyses tend to raise subgenera and subspecies to genus and species level, respectively (e.g., Arribas and Carranza, 2004; Arnold et al., 2007), and it remains to be seen if this is also reflected in their osteology.

The mentioned biases result in a significant loss of data, especially in small, fossil vertebrate taxa whose comparative osteology is poorly known, such as amphibians or reptiles. Delfino (2004) estimated the loss to be of 58% when comparing the Italian palaeoherpetological fossil record with the extant Italian herpetofauna. Within lizards, only the 36% of the diversity of extant Italian species has been recognised in the fossil record (Delfino, 2004), which amounts to a loss of data of 64%.

Results are even worse when fossils of all European lizards and worm lizards are considered, with an increase in the lost information of 10% (74% versus the above-mentioned 64% in Italy). The same holds true for the percentage of recognisability of extant lizards, which is 44% for Italian species (Delfino, 2004) and only 31% for European ones. The trend of decreasing recognisability when going from higher to lower taxonomic levels (Fig. 1) clearly shows that detailed osteological studies are strongly needed at genus and species level. As far as lacertids are concerned, this issue

is exacerbated by the fact that they are the most speciose family of lizards in Europe (56 species out of 78, 72%) and that, among them, the sole genus *Podarcis* includes 21 species, which can currently not be distinguished based on osteology. Moreover, the recognisability of *Podarcis* itself can be also questioned, because of the poor knowledge of the osteology of other small-sized European lacertids that can be potentially confused with it. Nonetheless, as will be shown below, detailed osteological study combined with novel methodological approaches like specimen-level phylogenetic analysis appears to be well suited to identify diagnostic combinations of osteological traits for lacertid species. This promising result indicates that at least for lacertids, if not for all lizards, the main issue decreasing recognisability may not be the lack of osteological traits in their skeletons, but the lack of our current osteological knowledge. As for anguids, this may hold true for the genus *Anguis* too, whose species have been mostly erected based on molecular data (Gvoždík et al., 2010, 2013), and for which no comparative osteological study has yet been published. Finally, despite not distinctly affecting the recognisability of extant European lizards, the fact that both European agamids and chamaeleonids can only be recognised at the family level will likely cause a loss in our knowledge of the past diversity of these two groups because of the absence of characters useful for distinguishing them from extant taxa that are currently extralimital, or from other species that have become extinct in the meantime.

The degree of lost data about the palaeodiversity of non-snake squamates in Europe, at least when extant taxa are considered, is therefore very high, and most probably even higher considering that our estimation is made under the implied premise of an equal probability of fossilization for all taxa. However, as already stated by Delfino (2004), this is probably not true: some taxa (such as *Iberolacerta* spp.) live in environments that complicate fossilization, such as high-altitude forests or areas with acid soils, and nearly no sedimentation. Therefore, these species have lower

probabilities of being preserved in the fossil record. Moreover, some osteological features of certain species (e.g., the scarce robustness of the bones of small gekkotans and scincids) could prevent their preservation. It is also true that fossils are often represented by isolated skeletal elements or incomplete specimens, missing the necessary bones or bone features to identify them. Therefore, even if a species is theoretically recognisable based on its osteology, taphonomic bias often prevents identification of the fossils. Nonetheless, both in the Italian and the European fossil record, most recognisable OTUs have been recovered as fossils (Tab. 1). This further corroborates that the loss of diversity is not significantly due to an absence of remains (i.e., taphonomic and collection biases may be less influent), but mainly due to the lack of detailed knowledge of potentially diagnostic, osteological features.

Given that, it appears evident that an improvement of our knowledge of the comparative osteology of lizards and worm lizards (but also of other reptiles and amphibians) is crucial in order to increase our understanding of the past diversity of these animals (as already pointed out by Bell and Mead, 2014). In particular, agamids, chamaeleonids, lacertids, scincids and *Anguis* need further investigations. The paucity of scholars interested in the comparative osteology of lizards in past times has surely prevented us to have a better capability to recognise fossil representatives of these animals, also because few researchers could not cover the whole extent of lizard diversity. New studies dealing with the osteology of extant species will surely benefit palaeontologists working with Pleistocene and Holocene fossils, which appear to be mostly attributable to modern taxa, but they will have an impact on the study of older fossils as well, because they will allow researchers to better recognise extinct forms and their relationships. Moreover, a better knowledge of the comparative osteology of extant species could also be useful for neoherpetologists to better understand the relationships between them, adding to other information such as external morphology and molecular

data, as shown by our phylogenetic analysis. Finally, only apomorphy-based identifications of fossil and sub-fossil material will allow to overcome the circular reasoning in palaeobiogeographical reconstructions, which led to the flawed recognition of herpetofaunal stability throughout the Pleistocene and Holocene of North American taxa (Bell et al. 2010). We should therefore be cautious with referrals of European Quaternary remains to extant taxa.

The case of Lacerta

Although the high variability within *Lacerta* species was confirmed by our phylogenetic analysis, the second run of the phylogenetic analysis with the constrained searches shows that specimen-level phylogeny can be a useful tool to identify diagnostic combinations of osteological features for lizard species. Given that the vast majority of specimens formed monophyletic clades with the other specimens of the same species, the set of character states uniting these monophyletic clades can be interpreted to be a diagnostic combination of osteological features of the respective species. It is important to stress that most single features of these diagnostic combinations can be variable within a clade, or shared with specimens of other species, so that they can rarely be considered unique autapomorphies of a particular species. However, the combination of all these osteological features is unique, and can be added to differential, morphological diagnoses of the species. The single specimen of *L. viridis*, which was found within the clade formed by the specimens of *L. bilineata*, is very incomplete, and does not preserve cranial material. It is possible that the erroneous placement of this specimen is due to that lack of cranial material, which is generally considered to be more diagnostic at species level than postcranial bones. Also, *L. bilineata* has only relatively recently been separated as distinct species from *L. viridis* (Rykena, 1991; Amann et al., 1997; Marzahn et al., 2016), and no morphological characters have been reported to date to distinguish the two species. Our results therefore appear to corroborate the general

assumption that cranial material is more readily identifiable to the species level than postcranial bones. Finally, it is important to note that only a constrained search managed to find the specimens of a single species in monophyletic clades. This result indicates that in the case of *Lacerta*, osteological characters alone do not seem to be enough to recover a reliable tree topology at the species level. Although the referral of the specimens to a single species was successful in nearly all cases (actually all the specimens that included cranial elements), the position of the *Lacerta* species as found in our strict consensus tree might thus not represent true phylogenetic relationships. More methodological work will be needed to address this discrepancy, but this would be out of the scope of the present study. In any case, the current analysis highlights the utility of specimen-level phylogeny to identify diagnostic combinations of osteological features for lacertid species. It also shows that lizard skeletons bear taxonomically significant features, and that detailed osteological studies are promising for the recognition of diagnostic traits and finally for identifying lizard remains in the fossil record.

Acknowledgements

Numerous persons have contributed to make this work possible, either by providing literature or by making available prepared specimens or CT-scans. Among others, we can mention: Salvador Bailon, Corentin Bochaton, Virginie Bouetel, Annemarie Ohler and Laure Pierre (Muséum national d'Histoire naturelle, Paris); Federico Bernardini (Museo Storico della Fisica e Centro di Studi e Ricerche “Enrico Fermi”, Roma); Rebecca Biton (Hebrew University of Jerusalem); Hugues-Alexandre Blain (Institut Català de Paleoecologia Humana i Evolució Social, Tarragona); Àngel Hernández Luján, Arnau Bolet and Josep Fortuny (Institut Català de Paleontologia Miquel Crusafont, Cerdanyola del Vallès); Christy A. Hipsley (Museum Victoria); Martin Kirchner and Johannes Müller (Museum für Naturkunde, Berlin); Marta Calvo

Revuelta and Borja Sanchiz (Museo Nacional de Ciencias Naturales, Madrid); Roberto Sindaco (Museo Civico di Storia Naturale, Carmagnola); Claudio Tuniz ('Abdus Salam' International Centre for Theoretical Physics, Trieste); Heinz Grillitsch and Georg Gassner (Naturhistorisches Museum Wien, Austria); Sónia Gabriel and Simon Davis (Laboratório Arqueociências Lisboa); Christian Klug and Winand Brinkmann (Paläontologisches Institut und Museum, Universität Zürich, Switzerland). Access to the collections has been made possible also thanks to Synthesys grants: Muséum national d'Histoire naturelle, Paris (FR-TAF-5007) to AV, Naturhistorisches Museum Wien (AT-TAF-5725) to ET, Museo Nacional de Ciencias Naturales, Madrid (ES-TAF-1975) to MD. Savvas Zotos and the organizers of the 9th Symposium on the Lacertids of the Mediterranean Basin and the 1st Symposium on Mediterranean Lizards (20-23/06/2016, Lemesos, Cyprus) are thanked for having promoted the inclusion of palaeontology in the topics of the meeting. The editor, Sylvain Ursenbacher (University of Basel), and two reviewers, Jean-Claude Rage (Muséum National d'Histoire Naturelle, Paris) and Andrej Čerňanský (Comenius University, Bratislava), improved this paper with their comments. This work was supported by Fondi di Ateneo dell'Università di Torino (2014-2015) and Generalitat de Catalunya (2014 SGR 416 GRC) to MD. ET has received funding from the European Union's Seventh Framework programme for research and innovation under the Marie Skłodowska-Curie grant agreement No 609402 - 2020 researchers: Train to Move (T2M).

References

- Amann, T., Rykena, S., Joger, U., Nettmann, H.K., Veith, M. (1997): Zur artlichen trennung von *Lacerta bilineata* Daudin, 1802 und *L. viridis* (Laurenti, 1768). Salamandra **33**: 255-268.
- Arribas, O., Carranza, S. (2004): Morphological and genetic evidence of the full species status of *Iberolacerta cyreni martinezricai* (Arribas, 1996). Zootaxa **634**: 1-24.

Arnold, E.N., Arribas, O., Carranza, S. (2007): Systematics of the palaearctic and oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* **1430**: 3-86.

Arnold, N., Ovenden, D. (2002): A field guide to the reptiles and amphibians of Britain and Europe. London, HarperCollins Publisher.

Bell, C.J., Gauthier, J.A., Bever, G.S. (2010): Covert biases, circularity, and apomorphies: a critical look at the North American Quaternary Herpetofaunal Stability Hypothesis. *Quat. Int.* **217**: 30-36.

Bell, C.J., Mead, J.I. (2014): Not enough skeletons in the closet: collections-based anatomical research in an age of conservation conscience. *Anat. Rec.* **297**: 344-348.

Blain, H.-A. (2009): Contribution de la paléoherpétofaune (Amphibia & Squamata) à la connaissance de l'évolution du climat et du paysage du Pliocène supérieur au Pléistocène moyen d'Espagne. *Treballs del Museu de Geologia de Barcelona* **16**: 39-170.

Böhme, M., Ilg, A. (2003): fosFARbase, www.wahre-staerke.com/ (2015)

Boulenger, G.A. (1917): A revision of the lizards of the genus *Nucras*, Gray. *Ann. S. Afr. Mus.* **13**: 195-215.

Carranza, S., Arnold, E.N., Amat, F. (2004): DNA phylogeny of *Lacerta* (*Iberolacerta*) and other lacertine lizards (Reptilia: Lacertidae): did competition cause long-term mountain restriction? *Syst. Biodivers.* **2**: 57-77.

Cooper, R.A., Maxwell, P.A., Crampton, J.S., Beu, A.G., Jones, C.M., Marshall, B.A. (2006): Completeness of the fossil record: Estimating losses due to small body size. *Geology* **34**: 241-244.

Daza, J.D., Stanley, E.L., Wagner, P., Bauer, A.M., Grimaldi, D.A. (2016): Mid-Cretaceous amber fossils illuminate the past diversity of tropical lizards. *Sci. Adv.* **2**: e1501080.

Delfino, M. (2004): Herpetological paleodiversity: general considerations about the Italian fossil record of modern species. *Ital. J. Zool.* **71**: 13-16.

Dunhill, A.M., Benton, M.J., Twitchett, R.J., Newell, A.J. (2012): Completeness of the fossil record and the validity of sampling proxies at outcrop level. *Palaeontology* **55**: 1155-1175.

Estes, R. (1983): *Handbuch der Paläoherpetologie 10A. Sauria terrestria, Amphisbaenia.* München, Friedrich Pfeil.

Evans, S.E. (2008): The skull of lizards and Tuatara. In: *Biology of the Reptilia*, pp. 1-347. Gans, C., Gaunt, A., Eds., Ithaca, New York.

Goloboff, P.A. (2014): Extended implied weighting. *Cladistics* **30**: 260-272.

Goloboff, P.A., Farris, J.S., Nixon, K.C. (2008): TNT, a free program for phylogenetic analysis. *Cladistics* **24**: 774-786.

Gvoždík, V., Benkovský, N., Crottini, A., Bellati, A., Moravec, J., Romano, A., Sacchi, R., Jandzik, D. (2013): An ancient lineage of slow worms, genus *Anguis* (Squamata: Anguidae), survived in the Italian Peninsula. *Mol. Phylogenet. Evol.* **69**: 1077-1092.

Gvoždík V., Jandzik, D., Lymberakis, P., Jablonski, D., Moravec, J. (2010): Slow worm, *Anguis fragilis* (Reptilia: Anguidae) as a species complex: genetic structure reveals deep divergences. *Mol. Phylogenet. Evol.* **55**: 460-472.

Kapli, P., Poulakakis, N., Lymberakis, P., Mylonas, M. (2011): A re-analysis of the molecular phylogeny of Lacertidae with currently available data. *Basic and Applied Herpetology* **25**: 97-104.

Maddison, W.P., Maddison, D.R. (2015). Mesquite: a modular system for evolutionary analysis. Version 3.04. <http://mesquiteproject.org>

Marzahn, E., Mayer, W., Joger, U., Ilgaz, Ç., Jablonski, D., Kindler, C., Kumlucaş, Y., Nistri, A., Schneeweiss, N., Vamberger, M., Žagar, A., Fritz, U. (2016):

Phylogeography of the *Lacerta viridis* complex: mitochondrial and nuclear markers provide taxonomic insights. *J. Zool. Syst. Evol. Res.* **54**: 85-105.

Mendes, J., Harris, D.J., Carranza, S., Salvi, D. (2016). Evaluating the phylogenetic signal limit from mitogenomes, slow evolving nuclear genes, and the concatenation approach. New insights into the Lacertini radiation using fast evolving nuclear genes and species trees. *Mol. Phylogenet. Evol.* **100**: 254-267.

Pincheira-Donoso, D., Bauer, A.M., Meiri, S., Uetz, P. (2013): Global taxonomic diversity of living reptiles. *PLoS ONE* **8**: e59741.

Rage, J.-C. (2013): Mesozoic and Cenozoic squamates of Europe. *Palaeobio. Palaeoenv.* **93**: 517-534.

Rykena, S. (1991): Kreuzungsexperimente zur prüfung der artgrenzen im genus *Lacerta* sensu stricto. *Mitteilungen aus dem Museum für Naturkunde in Berlin. Zoologisches Museum Und Institut Für Spezielle Zoologie (Berlin)* **67**: 55-68.

Sillero, N., Campos, J., Bonardi, A., Corti, C., Creemers, R., Crochet, P.-A., Crnobrnja Isailović, J., Denoël, M., Ficetola, G.F., Gonçalves, J., Kuzmin, S., Lymberakis, P., de Pous, P., Rodríguez, A., Sindaco, R., Speybroeck, J., Toxopeus, B., Vieites, D.R., Vences, M. (2014): Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia* **35**: 1-31.

Soligo, C., Andrews, P. (2005): Taphonomic bias, taxonomic bias and historical non-equivalence of faunal structure in early hominin localities. *J. Hum. Evol.* **49**: 206-229.

Speybroeck, J., Beukema, W., Bok, B., Van der Voort, J. (2016). Field guide to the amphibians and reptiles of Britain and Europe. London, Bloomsbury Publishing.

Uetz, P., Hošek J. (eds.): The Reptile Database, <http://www.reptile-database.org>, accessed June 29, 2016

388 **Table 1)** Recognisability of European lizards and worm lizards and their presence in the
 389 fossil record.

Extant species	Recognisable OTU	Fossil record
<i>Laudakia stellio</i>	Agamidae indet.	Yes
<i>Chamaeleo chamaeleon</i>	Chamaeleonidae indet.	Yes
<i>Euleptes europaea</i>	<i>E. europaea</i>	No
<i>Hemidactylus turcicus</i>	<i>H. turcicus</i>	Yes
<i>Mediodactylus kotschy</i>	<i>M. kotschy</i>	No
<i>Tarentola mauritanica</i>	<i>T. mauritanica</i>	Yes
<i>Acanthodactylus erythrurus</i>	<i>A. erythrurus</i>	Yes
<i>Algyroides fitzingeri</i>	Lacertidae indet.	Yes
<i>Algyroides marchi</i>		
<i>Algyroides moreoticus</i>		
<i>Algyroides nigropunctatus</i>		
<i>Anatololacerta anatolica</i>		
<i>Anatololacerta pelasgiana</i>		
<i>Archaeolacerta bedriagae</i>		
<i>Dalmatolacerta oxycephala</i>		

Darevskia armeniaca

Darevskia lindholmi

Darevskia praticola

Darevskia saxicola

Dinarolacerta

montenegrina

Dinarolacerta

mosorensis

Hellenolacerta graeca

Iberolacerta aranica

Iberolacerta aurelio

Iberolacerta bonnali

Iberolacerta cyreni

Iberolacerta galani

Iberolacerta horvathi

Iberolacerta

martinezricai

Iberolacerta monticola

Zootoca vivipara

<i>Eremias arguta</i>	Eremiadini indet.	Yes
-----------------------	-------------------	-----

Ophisops elegans

<i>Lacerta agilis</i>	<i>L. agilis</i>	Yes
-----------------------	------------------	-----

<i>Lacerta bilineata</i>	<i>L. bilineata</i>	Yes
--------------------------	---------------------	-----

<i>Lacerta viridis</i>	<i>L. viridis</i>	Yes
------------------------	-------------------	-----

<i>Lacerta schreiberi</i>	<i>L. schreiberi</i>	No
---------------------------	----------------------	----

<i>Lacerta trilineata</i>	<i>L. trilineata</i>	Yes
<i>Podarcis bocagei</i>	<i>Podarcis</i> sp. (?)	Yes
<i>Podarcis carbonelli</i>		
<i>Podarcis cretensis</i>		
<i>Podarcis erhardii</i>		
<i>Podarcis filfolensis</i>		
<i>Podarcis gaigeae</i>		
<i>Podarcis hispanicus</i>		
<i>Podarcis levendis</i>		
<i>Podarcis lilfordi</i>		
<i>Podarcis liolepis</i>		
<i>Podarcis melisellensis</i>		
<i>Podarcis milensis</i>		
<i>Podarcis muralis</i>		
<i>Podarcis</i>		
<i>peloponnesiacus</i>		
<i>Podarcis pityusensis</i>		
<i>Podarcis raffonei</i>		
<i>Podarcis siculus</i>		
<i>Podarcis tauricus</i>		
<i>Podarcis tiliguerta</i>		
<i>Podarcis vaucheri</i>		
<i>Podarcis waglerianus</i>		
<i>Psammodromus algirus</i>	<i>Psammodromus</i> sp.	Yes

<i>Psammodromus</i>		
<i>hispanicus</i>		
<i>Timon lepidus</i>	<i>T. lepidus</i>	Yes
<i>Ablepharus kitaibelii</i>	Scincidae indet.	Yes
<i>Ophiomorus</i>		
<i>punctatissimus</i>		
<i>Trachylepis aurata</i>		
<i>Chalcides bedriagai</i>	<i>Chalcides</i> sp.	Yes
<i>Chalcides chalcides</i>		
<i>Chalcides striatus</i>		
<i>Chalcides ocellatus</i>	<i>C. ocellatus</i>	Yes
<i>Anguis cephallonica</i>	<i>Anguis</i> gr. <i>A.</i> <i>fragilis</i>	Yes
<i>Anguis colchica</i>		
<i>Anguis fragilis</i>		
<i>Anguis graeca</i>		
<i>Anguis veronensis</i>		
<i>Pseudopus apodus</i>	<i>P. apodus</i>	Yes
<i>Blanus cinereus</i>	<i>Blanus</i> sp. (Western Group)	Yes
<i>Blanus mariae</i>		
<i>Blanus strauchi</i>	<i>B. strauchi</i> complex	No
78	24 (31%)	20 (26%)

Figure 1) Recognisability of extant European lizards at family, genus and species levels. Numbers inside the columns represent the exact percentage of recognisable taxa and their number compared to the total number of family, genera and species respectively (between parentheses).

Figure 2) Hierarchical level of the recognisability in the different families of European lizards and worm lizards: percentage of extant species recognisable at family level is depicted in black, percentage at genus level in dark grey, percentage at species level in white, percentage at subfamily or species complex level in light grey. Numbers inside the columns represent the exact percentage of extant species recognisable at a specific level and their number compared to the total number of species included in the family (between parentheses). Abbreviations: Ag, Agamidae; Ch, Chamaeleonidae; Sp, Sphaerodactylidae; Ge, Gekkonidae; Ph, Phyllodactylidae; La, Lacertidae; Sc, Scincidae; An, Anguidae; Bl, Blanidae.

Figure 3) Strict consensus tree of 9 MPTs resulting from a constrained search forcing all specimens of *Lacerta* into a monophyletic clade (as confirmed by phylogenetic studies based on molecular data). All but one specimen (*L. viridis* MNCN 16504, marked in white) were recovered in monophyletic clades together with the other members of their species (highlighted by the grey rectangles). The character states uniting these clades at their base (indicated by dark grey dots) can be interpreted as autapomorphic combination of osteological traits of the respective species, and used to identify fossil material. The questionable position of MNCN 16504 is probably due to the lack of cranial material (the specimen only preserves a partial vertebral column).

419 **Figure 4)** Number of articles dealing with European lizard osteology per 10 years from
420 the 1850s to today.

421

Osteology, fossil record and palaeodiversity of the European lizards

Andrea Villa¹, Emanuel Tschopp^{1, 2, 3}, Georgios L. Georgalis^{4, 1}, Massimo Delfino^{1, 5}

¹ Dipartimento di Scienze della Terra, Università degli Studi di Torino, via Valperga Caluso 35, 10125 Torino, Italy

² GeoBioTec, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Caparica, Portugal

³ Museu da Lourinhã, Rua João Luís de Moura, 2530-157 Lourinhã, Portugal

⁴ Department of Geosciences, University of Fribourg/Freiburg, Chemin du Musée 6, 1700 Fribourg, Switzerland

⁵ Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Edifici ICTA-ICP, Carrer de les Columnes s/n, Campus de la UAB, 08193 Cerdanyola del Vallès, Barcelona, Spain

Supplementary material

Supplementary text S1

List of studied specimens. The specimens marked with an asterisk were included in the phylogenetic analysis. Abbreviations: CIPA - Laboratório Arqueociências Lisboa, Portugal; HUJ.OST - Osteological collection of the Hebrew University of Jerusalem, Israel; MCCI - Museo Civico di storia naturale di Carmagnola, Italy; MDHC - Massimo Delfino Herpetological Collection, Department of Earth Sciences, University of Torino, Italy; MNCN - Museo Nacional de Ciencias Naturales, Madrid, Spain; MNHN - Muséum national d'Histoire naturelle, Paris, France; NHMW - Naturhistorisches Museum Wien, Austria; PIMUZ - Paläontologisches Institut und Museum der Universität Zürich, Switzerland; UAM.R - Universidad Autónoma de Madrid (Reptiles), Spain; ZMS - Zoologische Staatssammlung München, Germany.

Laudakia stellio: MDHC 245; HUJ.OST-Z-5, 423, 424.

Chamaeleo chamaeleon: MNHN 241, 1942-103, 2002-24, 1887-875; HUJ.OST-Z-380, 425.

Euleptes europaea: MDHC 384, 388, 389.

Hemidactylus turcicus: MDHC 26, 238

Mediodactylus kotschy: MDHC 201, 285.

Tarentola mauritanica: MDHC 97, 98, 119, 194, 302.

Acanthodactylus erythrurus: UAM.R.AC VII; EBD 1266 (specimen from the collection of Salvador Bailon).

Algyroides fitzingeri: MDHC 351.

Algyroides moreoticus: MDHC 174.

Algyroides nigropunctatus: MDHC 171, 242, 243; NHMW 797*.

Archaeolacerta bedriagae: MDHC 167*; unnumbered specimen from the collection of Salvador Bailon.

Eremias arguta: MNHN 1944-168.

Iberolacerta bonnali: UAM.R.Lm28A.

Iberolacerta cyreni: UAM.R.Lm4.

Iberolacerta monticola: UAM.R.Lm77, Lm92.

Lacerta agilis: CIPA 1550*; MDHC 176*, 177*, 178*, 230*, 231*; MNCN 15979*; NHMW 802*; PIMUZ A/III 0902*.

Lacerta bilineata: MDHC 15*, 48*, 73*, 77*, 84*, 381*; MNCN 16505*; PIMUZ A/III 1276*; UAM.R.Q21*.

Lacerta schreiberi: CIPA 778*, 1256*, 1511*, 1517*; UAM.R.S-6*.

Lacerta trilineata: MDHC 240*, 241*, 295*, 356*.

Lacerta viridis: MNCN 16504*; NHMW 778*, 906*, 32879-3*.

Ophisops elegans: MDHC 281, 282; unnumbered specimen from the collection of Salvador Bailon.

Podarcis bocagei: UAM.R.PB48.

Podarcis filfolensis: MDHC 385.

Podarcis hispanicus: UAM.R.H30; two unnumbered specimens from the collection of Salvador Bailon.

Podarcis lilfordi: two unnumbered specimens from the collection of Salvador Bailon.

Podarcis melisellensis: MDHC 217, 218.

Podarcis muralis: MDHC 6, 65, 66, 72, 81, 89, 90, 222, 267, 311*, 312*, 313, 395, 413; MNHN 1992.192.

Podarcis siculus: MDHC 25, 91, 125, 229.

Podarcis tauricus: MDHC 244.

Podarcis tiliguerta: MDHC 153, 154.

Podarcis waglerianus: MDHC 390; MNHN 1992.189, 1992.190.

Psammodromus algirus: MNHN 1992.41; NHMW 788*; UAM.R.Ps9.

Timon lepidus: MDHC 216*; MNHN 1988.6629, 1991.4010, 1991.4242; unnumbered specimen stored in the Institut Català de Paleontologia Miquel Crusafont.

Zootoca vivipara: MDHC 179; UAM.R.Lv24.

Ablepharus kitaibelii: MDHC 239.

Chalcides bedriagai: unnumbered specimen from the collection of Salvador Bailon.

Chalcides chalcides: MDHC 94, 329, 398, 408.

Chalcides ocellatus: MDHC 193, 250; MNHN 1992.193; specimen number 28 from the collection of Salvador Bailon.

Chalcides striatus: MDHC 404; MNCN 16508; unnumbered specimen from the collection of Salvador Bailon.

Trachylepis aurata: MDHC 280; MNHN 1887-863.

Anguis gr. A. fragilis: MDHC 45, 49, 67, 102, 213, 221, 236, 237, 310, 367, 402.

Pseudopus apodus: MDHC 214, 215; MNHN 1918.95, 1992.199; PIMUZ A/III0975.

Blanus cinereus: MDHC 156; ZSM 175-1993-1, 175-1993-2, 227-1975, 548-2003, 652-0-1, 652-0-2, 653-0-1, 653-0-2.

Blanus mariae: ZSM 27-1988-1, 27-1988-2.

Blanus strauchi: MCCI R-1635, 1668; MDHC 93, 286, 287, 288.

Supplementary text S2

List of the published works dealing with the osteology of extant European lizards and worm lizards. The references marked with an asterisk provided useful information for the creation of the phylogenetic matrix.

Anderson, S.C., Leviton, A.E. (1966): A review of the genus *Ophiomorus* (Sauria: Scincidae), with descriptions of three new forms. *Proceedings of the California Academy of Sciences, Fourth series* **XXXIII**: 499-534.

*Arnold, E.N. (1973): Relationships of the Palaearctic lizards assigned to the genera *Lacerta*, *Algyroides* and *Psammodromus* (Reptilia: Lacertidae). *Bulletin of the British Museum of Natural History, Zoology* **25**: 291-366.

Arnold, E.N. (1983): Osteology, genitalia and relationship of *Acanthodactylus* (Reptilia: Lacertidae). *Bulletin of the British Museum of Natural History, Zoology* **44**: 291-339.

Arnold, E.N. (1984): Evolutionary aspects of tail shedding in lizards and their relatives. *J. Nat. Hist.* **18**: 127-169.

*Arnold, E.N. (1989): Towards a phylogeny and biogeography of the Lacertidae: relationships within an Old-World family of lizards derived from morphology. *Bulletin of the British Museum of Natural History, Zoology* **55**: 209-257.

*Arnold, E.N. (1997): Interrelationships and evolution of the east Asian grass lizards, *Takydromus* (Squamata: Lacertidae). *Zool. J. Linn. Soc.-Lond.* **119**: 267-296.

*Arnold, E.N. (1998): Structural niche, limb morphology and locomotion in lacertid lizards (Squamata, Lacertidae); a preliminary survey. *Bulletin of the Natural History Museum Zoology Series* **64**: 63-90.

*Arnold, E.N., Arribas, O., Carranza, S. (2007): Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* **1430**: 1-86.

*Arribas, O.J. (1998): Osteology of the Pyrenean Mountain Lizard and comparison with other species of the collective genus *Archaeolacerta* MERTENS, 1921 s.l. from Europe and Asia Minor (Squamata: Sauria: Lacertidae). *Herpetozoa* **11**: 47-70.

Arribas, O.J., Odierna, G. (2004): Karyological and osteological data supporting the specific status of *Iberolacerta (cyreni) martinezricai*. *Amphibia-Reptilia* **25**: 359-367.

Baig, K.J., Wagner, P., Ananjeva, N.B., Böhme, W. (2012): A morphology-based taxonomic revision of *Laudakia* Gray, 1845 (Squamata: Agamidae). *Vertebrate Zoology* **62**: 213-260.

Bailon, S. (1991): Amphibiens et reptiles du Pliocène et du Quaternaire de France et d'Espagne: mise en place et evolution des faunes. PhD thesis, Université Paris VII.

*Barahona, F.F. (1996): Osteología craneal de lacértidos de la Península Ibérica e Islas Canarias: análisis sistemático filogenético. Unpublished PhD thesis, Universidad Autonoma de Madrid.

*Barahona, F., Barbadillo, L.J. (1997): Identification of some Iberian lacertids using skull characters. *R. Esp. Herp.* **11**: 47-62.

Barahona, F., Barbadillo, L.J. (1998): Inter- and intraspecific variation in the post-natal skull of some lacertid lizards. *J. Zool.* **245**: 393-405.

- Barbadillo, L.J. (1989): Los Reptilia (Sauria y Amphisbaenia) de los yacimientos Plio-pleistocénicos de la cuenca Guadix-Baza (sudeste español). Trabajos sobre el Neogeno-Cuaternario **11**: 151-165.
- Barbadillo, L.J., Barahona, F. (1994): The number of cervical vertebrae in lacertid lizards: some unusual cases. Herpetol. J. **4**: 166.
- *Barbadillo, L.J., Sanz, J.L. (1983): Análisis osteométrico de las regiones sacra y presacra de la columna vertebral en los Lagartos Ibéricos *Lacerta viridis* Laurenti, *Lacerta lepida* Daudin y *Lacerta schreiberi* Bedriaga. Amphibia-Reptilia **4**: 215-239.
- Bauer, A.M., Good, D.A., Branch, W.R. (1997): The taxonomy of the southern african leaf-toed geckos (Squamata: Gekkonidae), with a review of old world “*Phyllodactylus*” and the description of five new genera. Proceedings of the California Academy of Science **49**: 447-497.
- Blain, H.-A. (2009): Contribution de la paléoherpétofaune (Amphibia & Squamata) à la connaissance de l'évolution du climat et du paysage du Pliocène supérieur au Pléistocène moyen d'Espagne. Treballs del Museu de Geologia de Barcelona **16**: 39-170.
- Blain, H.-A. (2012-2014): Anfíbios y escamosos de Cueva Victoria. - Amphibians and squamate reptiles from Cueva Victoria. Mastia **11-12-13**: 175-197.
- Blain, H.-A., Bailon, S., Agustí, J. (2007): Anurans and squamate reptiles from the latest early Pleistocene of Almenara-Casablanca-3 (Castellón, East of Spain). Systematic, climatic and environmental considerations. Geodiversitas **29**: 269-295.
- Bochaton, C., Buffrénil, V. de, Lemoine, M., Bailon, S., Ineich, I. (2015): Body Location and Tail Regeneration Effects on Osteoderms Morphology. Are They Useful Tools for Systematic, Paleontology, and Skeletochronology in Diploglossine Lizards (Squamata, Anguidae)? J. Morphol. **276**: 1333-1344.
- Bochaton, C., Boistel, R., Casagrande, F., Grouard, S., Bailon, S. (2016): A fossil *Diploglossus* (Squamata, Anguidae) lizard from Basse-Terre and Grande-Terre Islands (Guadeloupe, French West Indies). Scientific Reports **6**: 28475.
- *Böhme, M. (2010): Ectothermic vertebrates (Actinopterygii, Allocaudata, Urodela, Anura, Crocodylia, Squamata) from the Miocene of Sandelzhausen (Germany, Bavaria) and their implications for environment reconstruction and palaeoclimate. Paläont. Z. **84**: 3-41.
- Bolet, A., Delfino, M., Fortuny, J., Almécija, S., Robles, J.M., Alba, D.M. (2014): An Amphisbaenian Skull from the European Miocene and the Evolution of Mediterranean Worm Lizards. PLoS ONE **9**: e98082.
- Bons, J., Geniez, P. (1995): Contribution to the systematic of the lizard *Acanthodactylus erythrurus* (Sauria, Lacertidae) in Morocco. Herpetol. J. **5**: 271-280.
- Boulenger, G.A. (1885): Catalogue of the lizards in the british museum (Natural History). London, order of the Trustees.
- *Boulenger, G.A. (1916): On the lizards allied to *Lacerta muralis*, with an account of *Lacerta agilis* and *L. parva*. The Transactions of the Zoological Society of London **21**: 1-104.
- *Boulenger, G.A. (1920): Monograph of the Lacertidae., London, British Museum Natural History.
- Bruner, E., Costantini, D., Fanfani, A., Dell'Omo, G. (2005): Morphological variation and sexual dimorphism of the cephalic scales in *Lacerta bilineata*. Acta Zool.-Stockholm **86**: 245-254.

Buffrénil, V. de, Bardet, N., Pereda-Suberbiola, X., Bouya, B. (2008): Specialization of bone structure in *Pachyvaranus crassispondylus* Arambourg, 1952, an aquatic squamate from the Late Cretaceous of the southern Tethyan margin. *Lethaia* **41**: 59-69.

Buffrénil, V. de, Dauphin, Y., Rage, J.-C., Sire, J.-Y. (2011): An enamel-like tissue, osteodermine, on the osteoderms of a fossil anguid (Glyptosaurinae) lizard. *CR Palevol* **10**: 427-437.

Buffrénil, V. de, Sire, J.-Y., Rage, J.-C. (2010): The Histological Structure of Glyptosaurine Osteoderms (Squamata: Anguidae), and the Problem of Osteoderm Development in Squamates. *J. Morphol.* **271**: 729-737.

*Calori, L. (1858): Sullo scheletro della *Lacerta viridis* Linn., sulla riproduzione della coda nelle lucertole, e sulle ossa cutanee del teschio dei sauri. *Memorie Della Accademia Delle Scienze dell'Istituto Di Bologna* **9**: 345-382.

*Camp, C.L. (1923): Classification of the lizards. *B. Am. Mus. Nat. Hist.* **48**: 289-480.

Caputo, V. (2004): The cranial osteology and dentition in the scincid lizards of the genus *Chalcides* (Reptilia, Scincidae). *Ital. J. Zool.* **71**: 35-45.

Caputo, V., Lanza, B., Palmieri, R. (1995): Body elongation and limb reduction in the genus *Chalcides* Laurenti 1768 (Squamata Scincidae): a comparative study. *Trop. Zool.* **8**: 95-152.

Čerňanský, A. (2010a): A revision of chamaeleonids from the Lower Miocene of the Czech Republic with description of a new species of *Chamaeleo* (Squamata, Chamaeleonidae). *Geobios-Lyon* **43**: 605-613.

Čerňanský, A. (2010b): Earliest world record of green lizards (Lacertilia, Lacertidae) from the Lower Miocene of Central Europe. *Biologia* **65**: 737-741.

Čerňanský, A. (2011): A revision of the chameleon species *Chamaeleo pfeili* Schleich (Squamata; Chamaeleonidae) with description of a new material of chamaeleonids from the Miocene deposits of southern Germany. *B. Geosci.* **86**: 275-282.

Čerňanský, A., Bauer, A.M. (2010): *Euleptes gallica* Müller (Squamata: Gekkota: Sphaerodactylidae) from the lower Miocene of North-West Bohemia, Czech Republic. *Folia Zool.* **59**: 323-328.

Čerňanský, A., Klembara, J., Smith, K.T. (2016): Fossil lizard from central Europe resolves the origin of large body size and herbivory in giant Canary Island lacertids. *Zool. J. Linn. Soc.-Lond.* **176**: 861-877.

Čerňanský, A., Rage, J.-C., Klembara, J. (2015): The Early Miocene squamates of Amöneburg (Germany): the first stages of modern squamates in Europe. *J. Syst. Palaeontol.* **13**: 97-128.

Čerňanský, A., Smith, K.T., Klembara, J. (2014a): Variation in the position of the jugal medial ridge among lizards (Reptilia: Squamata): its functional and taxonomic significance. *Anat. Rec.* **297**: 2262-2272.

Čerňanský, A., Boistel, R., Fernandez, V., Tafforeau, P., Le Noir, N., Herrel, A. (2014b): The atlas-axis complex in chamaeleonids (Squamata: Chamaeleonidae), with description of a new anatomical structure of the skull. *Anat. Rec.* **297**: 369-396.

Conrad, J.L. (2004): Skull, mandible, and hyoid of *Shinisaurus crocodilurus* Ahl (Squamata, Anguimorpha). *Zool. J. Linn. Soc.-Lond.* **141**: 399-434.

Conrad, J.L. (2008): Phylogeny and systematics of Squamata (Reptilia) based on morphology. *B. Am. Mus. Nat. Hist.* **310**: 1-182.

- Cope, E.D. (1892): The osteology of the Lacertilia. P. Am. Philos. Soc. **XXX**: 184-221.
- Cooper, J.S., Poole, D.F.G., Lawson, R. (1970): The dentition of agamid lizards with special reference to tooth replacement. J. Zool. **162**: 85-98.
- Daza, J.D., Bauer, A.M., Snively, E.D. (2014): On the Fossil Record of the Gekkota. Anat. Rec. **297**: 433-462.
- Delfino, M., Kotsakis, T., Arca, M., Tuveri, C., Pitruzzella, G., Rook, L. (2008): Agamid lizards from the Plio-Pleistocene of Sardinia (Italy) and an overview of the European fossil record of the family. Geodiversitas **30**: 641-656.
- *De Stefano, G. (1903): I sauri del Quercy appartenenti alla collezione Rossignol. Atti Soc. ital. sci. nat., Mus. civ. stor. nat. Milano **42**: 382-418.
- Edmund, A.G. (1969): Dentition. In: Gans C. (Ed.), Biology of the Reptilia, Vol. I, Morphology A. Academic Press, London and New York: 117-200.
- El-Toubi, M.R. (1947): Sacral Ribs of Lacertilia. Nature **159**: 342.
- Estes, R. (1970): Die fauna der miozänen Spaltenfüllung von Neudorf an der March (CSSR). Reptilia (Lacertilia). Sitzungsberichte der Akademie der Wissenschaften mathematisch-naturwissenschaftliche Klasse **178**: 77-82.
- Estes, R., Williams, E.E. (1984): Ontogenetic variation in the molariform teeth of lizards. J. Vertebr. Paleont. **4**: 96-107.
- Etheridge, R. (1967): Lizard caudal vertebrae. Copeia **4**: 699-721.
- Evans, S.E. (2008): The skull of lizards and Tuatara. In: Gans C., Gaunt A. (Eds.), Biology of the Reptilia. Ithaca, New York.
- Gans, C., Montero, R. (2008): An atlas of amphisbaenian skull anatomy. In: Gans C., Gaunt A.S., Adler K. (Eds.), Biology of the Reptilia. Volume 21. Morphology I. The Skull and Appendicular Locomotor Apparatus of Lepidosauria. Society for the Study of Amphibians and Reptiles, Ithaca, New York. Pp. 621-738.
- Gasc, J.P. (1966): Les reptiles. In: Atlas de préhistoire, 3, Faunes et flores préhistoriques de l'Europe occidentale. R. Lavocat Eds., 470-474, Paris.
- Gaupp, E. (1900): Das Chondrocranium von *Lacerta agilis*, ein beitrag zum Verstandnis des Amniotenschadels. Anatomisebe Hefte. I. Abteilung.
- *Gauthier, J.A., Kearney, M., Maisano, J.A., Rieppel, O., Behlke, A.D.B. (2012): Assembling the squamate tree of life: perspectives from the phenotype and the fossil record. Bull. Peabody Mus. Nat. Hist. **53**: 3-308.
- Gleed-Owen, C. (1997): The prehistory of the slow-worm in Britain. Krag News special edition, **October 1997**: 11-15.
- Greer, A.E., Arnold, C., Arnold, E.N. (2000): The systematic significance of the number of presacral vertebrae in the scincid lizard genus *Mabuya*. Amphibia-Reptilia **11**: 121-126.
- Greer, A.E., Wilson, G.D.F. (2001): Comments on the scincid lizard genus *Ophiomorus*, with a cladistic analysis of the species. Hamadryad **26**: 261-271.
- Herrel, A., Aerts, P., Fret, J., De Vree, F. (1999): Morphology of the Feeding System in Agamid Lizards: Ecological Correlates. Anat. Rec. **254**: 496-507.

- Hoffstetter, R. (1944): Sur les Scincidae fossiles. 1. Formes européennes et nord-américaines. Bulletin du Muséum National d'Histoire Naturelle **XVI**: 547-553.
- Hoffstetter, R. (1962): Observations sur les ostéodermes et la classification des anguidés actuels et fossiles (Reptiles, Sauriens). Bulletin du Muséum National d'Histoire Naturelle - 2^o série **34**: 149-157.
- Hoffstetter, R., Gasc, J.-P. (1969): Vertebrae and ribs of modern reptiles. pp. 201-310. In: Gans C., Bellairs A.d'A., Parsons T.S. (Eds.), Biology of the Reptilia. Volume 1. Morphology A. Academic Press, London and New York.
- Holman, J.A. (1998): Pleistocene amphibians and reptiles in Britain and Europe. Oxford monographs on geology and geophysics, **38**.
- Jollie, M.T. (1960): The head skeleton of the lizard. Acta Zool.-Stockholm **41**: 1-64.
- Kazemi, S.M., Farhadi Qomi, M., Kami, H.G., Anderson, S.C. (2011): A new species of *Ophiomorus* (Squamata: Scincidae) from Maranjab Desert, Isfahan Province, Iran, with a revised key to the genus. Amphibian and Reptile Conservation **5**: 23-33.
- Klembara, J. (2012): A new species of *Pseudopus* (Squamata, Anguinae) from the early Miocene of Northwest Bohemia (Czech Republic). J. Vertebr. Paleontol. **32**: 854-866.
- Klembara, J., Hain, M., Dobiašová, K. (2014): Comparative Anatomy of the Lower Jaw and Dentition of *Pseudopus apodus* and the Interrelationships of Species of Subfamily Anguinae (Anguimorpha, Anguinae). Anat. Rec. **297**: 516-544.
- Klembara, J., Rummel, M. (2016): New material of *Ophisaurus*, *Anguis* and *Pseudopus* (Squamata, Anguinae) from the Miocene of the Czech Republic and Germany and systematic revision and palaeobiogeography of the Cenozoic Anguinae. Geol. Mag.: 1-25.
- Kosma, R. (2004): The dentitions of recent and fossil scincomorphan lizards (Lacertilia, Squamata) – Systematics, Functional Morphology, Paleocology. Unpublished PhD Thesis, Universität Hannover.
- Kotsakis, T. (1981): Le lucertole (Lacertidae, Squamata) del Pliocene, Pleistocene e Olocene delle Baleari. Bolletí de la Societat d'Història Natural de les Balears **25**: 135-150.
- Lantz, L.A., Cyrén, O. (1919): On *Lacerta praticola*. The Annals and Magazine of Natural History **3**: 28-31.
- López-García, J.M., Blain, H.-A., Cuenca-Bescós, G., Alonso, C., Alonso, S., Vaquero, M. (2011): Small vertebrates (Amphibia, Squamata, Mammalia) from the late Pleistocene-Holocene of the Valdavara-1 cave (Galicia, northwestern Spain). Geobios-Lyon **44**: 253-169.
- Ljubisavljević, K., Urošević, A., Aleksić, I., Ivanović, A. (2010): Sexual dimorphism of skull shape in a lacertid lizard species (*Podarcis* spp., *Dalmatolacerta* sp., *Dinarolacerta* sp.) revealed by geometric morphometrics. Zoology **113**: 168-174.
- Malashichev, Y.B. (2001): Sacrum and pelvic girdle development in Lacertidae. Russian Journal of Herpetology **8**: 1-16.
- Mateo, J.A., López-Jurado, L.F. (1997): Dental Ontogeny in *Lacerta lepida* (Sauria, Lacertidae) and Its Relationship to Diet. Copeia **1997**: 461-465.
- Metzger, K. (2002): Cranial Kinesis in Lepidosaur: Skulls in Motion. pp. 15-46. In: Aerts P., D'Août K., Herrel A., Van Damme R. (Eds.), Topics in Functional and Ecological Vertebrate Morphology. Shaker Publishing.

Mezzasalma, M., Maio, N., Guarino, F.M. (2014): To Move or Not to Move: Cranial Joints in European Gekkotans and Lacertids, an Osteological and Histological Perspective. *Anat. Rec.* **297**: 463-472.

Moody, S., Roček, Z. (1980): *Chamaeleo caroliquarti* (Chamaeleonidae, Sauria): a new species from the Lower Miocene of Central Europe. *Vestník Ústředního Ústavu Geologického* **55**: 85-92.

*Müller, J. (1996) Eine neue art der echten eidechsen (Reptilia: Lacertilia: Lacertidae) aus dem Unteren Miozän von Poncenat, Frankreich. *Mainzer Geowissenschaftliche Mitteilungen* **25**: 79-88.

Müller, J. (2001): A new fossil species of *Euleptes* from the early Miocene of Montaigny, France (Reptilia, Gekkonidae). *Amphibia-Reptilia* **22**: 341-348.

Parker, K.W. (1879): The Croonian Lecture: On the Structure and Development of the Skull in the Lacertilia. Part I. On the Skull of the Common Lizards (*Lacerta agilis*, *L. viridis*, and *Zootoca vivipara*). *Phil. Trans. R. Soc. Lond.* **170**: 595-640.

*Rauscher, K.L. (1992): Die echsen (Lacertilia, Reptilia) aus dem Plio-Pleistozän von Bad Deutsch-Altenburg, Niederösterreich. *Beiträge zur paläontologie von Österreich* **17**: 81-177.

Roček, Z. (1980a): The dentition of the European glass lizard *Ophisaurus apodus* (Pallas, 1775) (Reptilia, Sauria: Anguidae), with notes on the pattern of tooth replacement. *Amphibia-Reptilia* **1**: 19-27.

Roček, Z. (1980b): Intraspecific and ontogenetic variation of the dentition in the green lizard *Lacerta viridis* (Reptilia, Squamata). *Věstník Československé společnosti zoologické* **44**: 272-277.

*Siebenrock, F. (1894). Das Skelet der *Lacerta simonyi* Steind. und der Lacertidenfamilie überhaupt. *Sitzungsberichte Der Kaiserlichen Akademie Der Wissenschaften* **103**: 205-292.

Siebenrock, F. (1895): Das skelet der Agamidae. *Sitzungsberichten der kaiserl. Akademie der Wissenschaften in Wien. Mathem.-Naturw. Classe* **CIV**: 1-108.

Stokely, P.S. (1947): Limblessness and Correlated Changes in the Girdles of a Comparative Morphological Series of Lizards. *Am. Midl. Nat.* **38**: 725-754.

Sumida, S.S., Murphy, R.W. (1987): Form and function of the tooth crown structure in gekkonid lizards (Reptilia, Squamata, Gekkonidae). *Can. J. Zoolog.* **65**: 2886-2892.

Van Damme, R., Vanhooydonck, B. (2002): Speed versus manoeuvrability: association between vertebral number and habitat structure in lacertid lizards. *J. Zool.* **258**: 327-334.

Wellborn, V. (1933): Comparative osteological examinations of gekkonids, eublepharids and uroplatids. Translated by Russel A.P., Bauer A.M., Deufel A. *Herpetological Translations*, 1. Breck Bartholomew, Bibliomania!

Yaryhin, O., Klembara, J. (2015): Different embryonic origin of the basiptyergoid process in two species of *Lacerta* (Squamata: Lacertidae). *Biologia* **70**: 530-534.

Young, L.R., Caputo, V., Giovannotti, M., Kohlsdorf, T., Vargas, A.O., May, G.E., Wagner, G.P. (2009): Evolution of digit identity in the three-toed Italian skink *Chalcides chalcides*: a new case of digit identity frame shift. *Evol. Dev.* **11**: 647-658.