AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Rabbits killing hares: An invasive mammal modifies native predator-prey dynamics

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1631365> since 2017-05-23T11:02:15Z

Published version:

DOI:10.1111/acv.12343

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Cerri Jacopo, Ferreti Marco, Bertolino Sandro. Rabbits killing hares: An invasive mammal modifies native predator-prey dynamics. *ANIMAL CONSERVATION*. None pp: 1-9.
DOI: 10.1111/acv.12343

The publisher's version is available at:

<http://onlinelibrary.wiley.com/doi/10.1111/acv.12343/fullpdf>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

Rabbits killing hares: an invasive mammal modifies native predator-prey dynamics.

Authors:

Cerri Jacopo¹ Istituto di Management, Scuola Superiore Sant'Anna 33, 56127, Pisa (Italy), e-mail: j.cerri@sssup.it;

Ferretti Marco² Regione Toscana, Corso Gramsci 110, 51100, Pistoia (Italy), email: marco.ferretti@regione.toscana.it;

Bertolino Sandro³ Department of Life Sciences and Systems Biology, Via Accademia Albertina 13, 10123 Torino, Italy. Email: sandro.bertolino@unito.it

Corresponding author: Cerri Jacopo*, Istituto di Management, Scuola Superiore Sant'Anna 33, 56127, Pisa (Italy), e-mail: j.cerri@sssup.it, mobile: (+39)3395692346;

Short title: Introduced cottontails modifies predator-prey dynamics;

Abstract

Invasive species management requires practical evidence of the impacts of introduced species over ecosystem structure and functioning. Theoretical ecology and empirical data support the potential of introduced mammals to drive native species to extinction, indeed the majority of practical evidence comes from insular environments, where conditions may differ from the mainland.

We analyzed the effects of an introduced lagomorph, the Eastern cottontail (*Sylvilagus floridanus*) on two native mammals, the European hare (*Lepus europaeus*) and the Red fox (*Vulpes vulpes*). We used relative abundances collected over 8 years at 30 protected areas in Italy. A Generalized Linear Mixed Model was fit to test various hypotheses about the relationships between cottontails, foxes and climatic conditions over the abundance of native hares.

Our model showed that the relationship between foxes and hares became more and more negative, as cottontail abundance increased. As no direct competition between introduced cottontails and native hares emerged, we believe that indirect dynamics like apparent competition exists between the two lagomorphs. Climatic conditions, expressed through the North Atlantic Oscillation, did not affect the relationship between cottontail and hare abundances. As the impact of parasites on mammal populations is generally climate-dependent, we believe that cottontails do not play a direct role in the cycle of parasites affecting hares.

Our results provide a clue that an invasive mammal, the Eastern cottontail, is modifying the predator-prey relationship between two native species in a non-insular environment. The existence of such dynamics should lead wildlife managers to account for the effect of introduced species in their decision-making, directing control activities on cottontails and not on native foxes.

Keywords: apparent competition, conservation, invasion ecology, invasive species, *Lepus europaeus*, *Sylvilagus floridanus*, *Vulpes vulpes*, wildlife management.

Introduction

Introduced species are considered one of the main drivers of biodiversity loss (Kumschick *et al.*, 2015) and their spread and impacts are likely to increase in an increasingly connected world (Hulme, 2009). The introduction of species changes the composition and functionality of ecosystems, and can drive native species to extinction through different mechanisms, such as predation, competition or diseases transmission (Bellard, Genovesi & Jeschke, 2016).

Interspecific competition operates primarily on individuals, with a reduction of their fitness; these effects will be translated at the population level, negatively influencing demographic parameters and possibly determining the decline or even the local extinction of one of the two species (Gurnell *et al.*, 2004). The competitive process could be the results of interference or exploitation competition. Interference competition involve direct interactions between species, while exploitation competition imply indirect negative interactions arising from the use of a common resource (Schoener, 1983; Lang & Benbow, 2013). Traditionally, ecologists demonstrated the occurrence of competition with removal experiments that, however, are difficult to perform (Schoener, 1983). Alternatively, data on demographic patterns (Forsyth & Hickling, 1998) and on species distributions (Bertolino *et al.*, 2014) can be assessed for evidence of competition by examining whether the presence of one species negatively influences the population demography and the distribution of the other. However, competitive interactions may be more complex. Species that do not interact directly and do not exploit the same resources can still influence each other if they share common natural enemies, such as predators, parasites, or pathogens (Zhang, Fan & Kuang, 2006). For example, the so-called 'apparent competition' occur as an indirect effect when species that do not directly compete for resources affect each other by being prey for the same predator (Courchamp, Langlais & Sugihara, 2000; Gibson, 2006; Noonburg & Byers, 2005; Lang & Benbow, 2013). Competition, either direct or indirect, between introduced and native mammals has been found to occur on various contexts. Direct competitive interactions has been observed between introduced and native squirrels in Europe (Gurnell *et al.*, 2004), as well as between marsupial and eutherian carnivores in Australia (Glen & Dickman, 2008). Apparent competition is another major consequence of introduced mammals, especially on islands when both a prey and a predator are introduced causing hyperpredation processes (Courchamp *et al.*, 2000; Zhang *et al.*, 2006). Other forms of indirect competition between native and invasive species include those mediated by diseases (Brummer *et al.*, 2010; Fournier-Chambrillon *et al.*, 2004; Strauss, White & Boots, 2012), whose transmission can be regulated by climatic conditions (Beard & O'Neill, 2005; Kiesecker, Blaustein & Belden, 2001; Skerratt *et al.*, 2007). As the various competitive dynamics arising with the introduction of invasive mammals can produce long-standing effects on ecosystems, assessing their structure and magnitude is crucial to design effective policies aimed at reducing the impact of invasive species meanwhile minimizing collateral damage to ecosystems (Bergstrom *et al.*, 2009).

This work aims at modeling the ecological impact of an invasive mammal in Northern Italy. Notably, we would like to test the occurrence of competitive processes with a native species, as well as the occurrence of apparent competition. The Eastern cottontail (*Sylvilagus floridanus*), has been introduced in North-Western Italy during the 1960s and it is now widespread in Northern and Central Italy (Bertolino, Ingegno & Girardello, 2011; Bertolino *et al.*, 2011a). Its interactions with the native European hare (*Lepus europaeus*) are complex, because the two species select different macro- and micro-habitats (Bertolino, Cordero di Montezemolo & Perrone, 2011b, 2013; Vidus-Rosin *et al.*, 2011), but cottontails carry several viruses and parasites, which can potentially affect hares (Bertolino *et al.*, 2010; Lavazza *et al.*, 2015; Tizzani *et al.*, 2011, 2014; Zanet *et al.*, 2013). Finally, the red fox (*Vulpes vulpes*), a major predator for hares in Europe (e.g. Lindström *et al.*, 1994; Reynolds & Tapper, 1995), includes cottontails in its diet when they are available (Balestrieri, Remonti & Prigioni, 2006).

To the best of our knowledge, no study has examined the population dynamics of hares, cottontails and foxes in areas where all these species coexist. Nevertheless, we deem such an approach necessary to fully address the issue of potential impacts of introduced cottontails.

Particularly, we tested three hypothesis, based on previous knowledge about the potential impact of invasive cottontails on native hares in Italy.

In the first hypothesis (H_1), we theorized that invasive cottontails and native hares are direct competitors for environmental resources. Previous works do not provide evidence for this, however they analyzed a limited geographical scale (Bertolino *et al.*, 2011b, 2013) and had a cross-sectional design (Vidus-Rosin *et al.*, 2011) that we deemed poor in revealing complex biotic interactions, compared to large-scale data and time series (Wisz *et al.*, 2013). As a consequence of H_1 we expected that a negative, clear, correlation occurred between cottontail and hare abundances.

In the second hypothesis (H_2), we predicted that cottontails play a role in the transmission of parasitic diseases to native hares. Parasites and some infective diseases (e.g pseudotuberculosis) have a higher impact on hare populations in cold and wet years (Chroust, 1984; Wibbelt & Frölich, 2005). Therefore, we expected that the abundance of cottontails had a negative correlation with the abundance of hares and that such correlation was stronger in wet and cold years, when body conditions of hare worsen and there is a higher number of parasites at the infective stage in the environment (Stromberg 1997).

In the third hypothesis (H_3) we hypothesized that cottontails negatively affect hares through indirect competition, modifying their relationships with foxes. In this case, we expected that the magnitude of the correlation between hare and fox abundances changed at different levels of cottontail abundance.

Materials and methods

Study areas and data collection

The study area was located in the lowlands of the province of Torino (Piedmont, Italy), where the wildlife office every year monitors small game at 30 protected areas through spotlight counts (Fig.1). From December to January, gamekeepers carry out 3 spotlight census from an off-road vehicle on fixed transects, recording hares, foxes and cottontails. We used the 2008-2015 data, expressing the relative abundance of each species as the number of individuals per kilometer of transect (Kilometric Index of Abundance, KIA), and averaging abundances for every year. Despite a lively debate about the effectiveness of relative indexes to model ecological dynamics (Anderson, 2001; Engeman, 2003), we deemed the KIA a reliable measure of population abundance, because we conducted longitudinal data analysis and because of its wide adoption in studies about the population ecology of lagomorphs and small carnivores (Barrio, Acevedo & Tortosa, 2010; Gortazar *et al.*, 2007; Preatoni *et al.*, 2012; Sobrino *et al.*, 2009).

Climatic data

Climatic conditions were expressed as winter (January-March) and summer (July-September) averages of the North Atlantic Oscillation (NAO; Lamb & Pepler, 1987) The NAO accounts for multiple climatic factors influencing many ecological dynamics in Europe and affecting the ecology of endothermic organisms, like mammals (Gordo, Barriocanal & Robson, 2011; Ottersen *et al.*, 2001; Schmidt, Asferg & Forchhammer, 2004; Straile & Stensteth, 2007).

Because our study area was close to the Mediterranean, where the effects of the NAO have been less studied than in Central Europe, before using the NAO as a covariate in our model we decided to test its associations with seasonal temperatures and rainfalls. We downloaded monthly values of the NAO from <http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml> and data about temperatures, rainfalls and moisture for the city of Torino from <http://archivio-meteo.distile.it/>. Associations between winter/summer NAO and local climate were always linear and significant Pearson's correlations are shown in Table 1.

Statistical analysis

As a first step to investigate a possible direct competition between the two lagomorphs, we evaluated the Pearson's correlation coefficients between KIAs of hares and cottontails in the 2008-2015 period for each area.

Then, we modeled the effect of cottontails, foxes and weather covariates on hare abundance with a Generalized Linear Mixed Model with a Gamma distribution of the error, a random intercept and a Log-link. The random structure of the model was selected on the basis of the lowest value of the Akaike's Informative Criterion (AIC; Zuur *et al.*, 2009). The choice of a Gamma distribution of the error was made by observing the distribution of the response variable, which was non-normal (Shapiro test: $W = 0.77$, $p < 0.01$) and positively skewed (2.46). Since no distance-dependent pattern emerged in trends correlations, we decided to discard spatial-explicit modeling. Weather covariates included winter NAO, as a measure of winter and summer conditions, and summer NAO, due to its correlation with spring weather. Winter weather was included in the model because harsh conditions can lead to higher energy needs of hares, reducing their body conditions (Hackländer, Arnold & Ruf, 2002a), making them more susceptible to diseases (Smith, Vaughan Jennings & Harris 2005) and reducing the milk production during the breeding season (Hackländer, Tataruch & Ruf, 2002b). Winter NAO index also accounted for summer conditions from July to September, when hare densities are at their maximum and density-dependent factors like coccides (Chroust, 1984) can operate. Spring weather was included in the model as it affects the survival of leverets by acting on thermoregulation (Hackländer *et al.*, 2002a). We also added an interaction between winter and summer NAO indexes, to account for inter-seasonal effects of weather conditions. Two other interaction terms were included to the model. The first one accounted for an interaction between cottontail abundance and winter NAO, representing our second hypothesis about the role of cottontails in disease transmission and their interactions with winter and summer conditions. The second interaction term accounted for an interaction between cottontail and fox abundances and expressed our third hypothesis of indirect competition.

Collinearity of predictors was checked by graphical exploration of correlations and with the Variance Inflation Factors (VIF). Predictors were not multicollinear and all of them were included in the initial model. As they were on different scales, predictors were standardized. Furthermore, graphical exploration of their relationships with the response variable enabled us to exclude higher-order interactions. The set of predictors was selected with the information-criteria approach, starting from the beyond-optimal model (Zuur *et al.*, 2009). Once we identified the best model, residuals were plotted for evaluating the fixed term and to eventually detect patterns. We also graphically explored the normality of random effects. We estimated Nakagawa's R^2 , to measure the proportion of data variability explained by the model (Nakagawa & Schielzeth, 2013). Statistics were computed with R (R Development Core Team 2015).

Results

We could not find any clear pattern in the temporal trend of local abundances of hares and cottontails (see Fig. S1 in Supplementary Material). A significant correlation was found only in six out of 30 sampled areas: five correlations were positive with $r = 0.74-0.97$ and only one negative with $r = 0.72$ (all $p < 0.05$, see Table S1 in Supplementary Material).

Residuals of GLMM did not show any clear pattern (see Fig. S2) and normality of the random effects was respected (see Fig. S3). Despite the fixed-term of the model accounted for a small proportion of the total variability of the data (marginal $R^2 = 0.15$), the whole model with the random term explained a good proportion of the variability in the data (conditional $R^2 = 0.73$).

Cottontail abundance was positively and significantly correlated to hare abundance ($\beta = 0.19 \pm 0.04$, $p < 0.01$) and the same applied for fox and hare abundances ($\beta = 0.75 \pm 0.21$, $p < 0.01$) (Table 2). However, the interaction term between cottontail and fox abundances was negative and significant ($\beta = -0.16 \pm 0.05$, $p < 0.01$), therefore the relationship between hares and foxes changed as cottontail abundance increased. The KIA of foxes was positively associated to the KIA of hares, in conditions of scarce cottontails ($\beta = 0.67 \pm 0.19$, for 0.55 cottontails/km), while the relationship between foxes and hares was increasingly negative at increasing cottontail abundances ($\beta = -0.04 \pm 0.15$, for 5 cottontails/km; $\beta = -0.82 \pm 0.33$, for 10 cottontails/km) (Fig.2).

From our model (Tab.2) winter values of the NAO index were negatively associated to hare abundance ($\beta = -0.43 \pm 0.13$, $p < 0.01$), but their interaction with cottontail abundance was not significant and had a low marginal effect ($\beta = 0.03 \pm 0.02$, $p = 0.09$). The interaction term between summer and winter values of the NAO was significantly and negatively associated to hare abundance ($\beta = -0.36 \pm 0.09$, $p < 0.01$).

Discussion

The spatial-temporal data analysis on the population dynamics of native hares and introduced cottontails does not support our first hypothesis about a direct interference between these two species. The local abundances of the two species were generally not correlated and, even when, the correlation was positive in five out of six cases. Also considering the areas altogether with a GLMM model, cottontails do not seem to be direct competitors of hares, as the correlation between the two species was positive. Instead, hares and cottontails seem to covary: this could mean that the two species are subjected to similar limiting factors, like summer rainfalls (Hackländer *et al.*, 2002 a,b; Jacobs & Dixon, 1981; Rödel & Dekker, 2012; Smith *et al.*, 2005).

On the other hand, our data supported the third hypothesis, with evidence of an impact of invasive cottontails on prey-predator dynamics of native hares and foxes. The relationship between foxes and hares, positive when cottontail are scarce, becomes strongly negative when cottontails increase their numbers. An initial positive correlation between foxes and hares in the presence of few or no cottontails, indicate that their population dynamics are more influenced by external factors, e.g. habitat quality (Bertolino *et al.*, 2011a), than by species interactions. Furthermore, we believe that two non-exclusive hypotheses can be advanced to explain the pattern we observed when cottontails increase.

In the first hypothesis, an increase in cottontail abundance would lead to a numerical response of foxes, magnifying their predatory impact on hares. This dynamic is known as “hyperpredation” and it is a particular case of apparent competition, a well-known impact of alien mammals that can produce profound changes in the relationship between native prey and predators (Norbury, 2001; Oliver, Luque-Larena & Lambin, 2009) and in the trophic interactions between native species (Roemer *et al.*, 2001). In hyperpredation, an introduced prey species, easy to kill and predation-resilient, triggers the extinction of a native prey species by rapidly increasing the population size of their predators (Courchamp *et al.*, 2000). Despite invasive species are often suboptimal preys for native predators, compared to native species, they can become important supplemental food resources, contributing to boost up predator populations (Pintor & Byers, 2015). In turn, such an increase in the abundance of native predators affects native preys, as increased predation pressure can have a density-independent and prolonged impact on their populations (Noonburg & Byers, 2005). Hyperpredation can be an overwhelming force for those native species who do not have very high growth rates or suitable behavioral anti-predator response (Courchamp *et al.*, 2000), driving them to collapse. In our case, we deem cottontails matching all the requirements for triggering hyperpredation of foxes on native hares. The reproductive rate of cottontails is high and can make their population resilient to prolonged and intensive predation by foxes (Balestrieri *et al.*, 2006) while native hares do not have high growth rates and may not be able to cope with high densities of foxes in intensive agricultural landscapes (Panek, 2009), like those in the study area.

In the second hypothesis, cottontails would trigger a case of “spillover” predation by foxes on native hares, around patches of permanent cover. Spillover predation occurs when predators emigrate from one source habitat where they find a primary prey, and they start foraging in sink habitats where they can find secondary preys (DeCesare *et al.*, 2009). Various field studies indicate that spillover predation is relatively common among generalist predators and that it can seriously affect prey populations occurring in proximity of permanent sources of food (Kirstan & Boarman, 2003; Oro & Martinez-Abraín, 2007; Sanz-Aguilar *et al.*, 2009). In our case, we believe that cottontails can become a main prey for foxes in the study area, because of their limited home range and the constant selection of permanent cover that can make them predictable preys (Bertolino *et al.*, 2013; Swihart, 1986). Therefore, foxes would alter their foraging behavior, choosing permanent patches of vegetation rich of profitable preys. As these habitats are also important resting sites for adult and young hares (Fernex, Nagel & Weber, 2011; Neumann *et al.*, 2011), an increased presence of foraging foxes is likely to result in a higher predation risk for hares.

These two hypotheses are not mutually exclusive and future studies, based on diet analysis and radiotracking, may properly quantify them. As indirect competition between mammal species often produces time-delayed impacts (Hansen *et al.*, 1999), time series analysis could also be another fruitful approach to disentangle interactions between cottontails, hares and foxes.

The lack of any interaction between cottontail abundance and the winter NAO on hare abundances partly rejects our second hypothesis about a significant role of cottontails as reservoirs for disease impacting hares. Cottontails are unlikely to play a clear role in the cycle of parasites like coccidies, reaching their infestation peak on hares in autumn and whose abundance is strongly affected by climatic conditions (Chroust, 1984). If cottontail had influenced coccidie loads of hares, the sign and magnitude of their relationship with hare abundance would have changed strongly with different climatic conditions. On the other hand, we cannot be completely sure about the lack of any role of cottontails in the transmission of other diseases, whose dynamics are less related to seasonal conditions. For example, cottontails can also be infected by EBHSV (Lavazza *et al.*, 2015) and the same applies to foxes eating infected lagomorphs (Chiari *et al.*, 2016). We recommend future studies exploring the potential role of cottontails in the epidemiology of EBHSV, as this virus is a serious threat for hare populations in Europe (Chiari *et al.*, 2014).

Our model also provides valuable insights about the impact of climatic conditions on hare abundance. Hares seem to be scarcer in years with a positive winter NAO, characterized by mild and dry winters but also by rainy summers. This is contrary to previous studies on the effect of winter weather (Schmidt *et al.*, 2004) but in line with evidence about the detrimental effect of summer rains (Hackländer *et al.*, 2002 a,b; Rödel & Dekker, 2012; Smith *et al.*, 2005). The interaction between winter and summer NAO was significant and part of the best set of predictors, suggesting that wet springs can magnify the negative effects of adverse weather in winter and/or summer, by decreasing the survival of leverets (Hackländer *et al.*, 2002a).

In this study, we provided evidence for an indirect effect of introduced cottontails on native hares through apparent competition. We conducted longitudinal data analysis on the abundance of the three species, while previous studies were cross-sectional and did not consider indirect interactions with foxes, failing in finding any impact (Bertolino *et al.*, 2011b; 2013; Vidus-Rosin *et al.*, 2011). We therefore highlight the need to use adequate data, with spatial replicate and temporal trends, to test hypotheses about the interactions between native and introduced species (Ricciardi *et al.*, 2013). When an invasive species becomes widespread it is important to evaluate possible impacts in the new ecosystems and considering its trophic interactions with new species, as these two tasks are crucial for policy-making and environmental management. In fact, obtaining adequate or upgraded information about the ecological impact of biological invaders, lays the foundation for their classification and their inclusion in black lists (Blackburn *et al.*, 2014; Hawkins *et al.*, 2015). Furthermore, knowing the interaction between native and invasive species is mandatory to design effective eradication campaigns or numerical control schemes, without unintended consequences (Zavaleta, Hobbs & Mooney, 2001).

As we found support for apparent competition between invasive cottontails and native hares, our research challenges some game management activities that are common in Northern Italy. Foxes are considered a main predator of hares, a possible negative effect of predation on hare populations was suggested, though other studies did not support this view (Lindström *et al.*, 1994; Reynolds & Tapper, 1995; Smtih *et al.*, 2005). Despite this disagreement between studies and the need of local evidences on the relationship between hares and foxes, irregular predator control is a common measure in Italy, as it is believed to increase game population densities, including hares. Previous research on invasive mammals has demonstrated that invasive prey species should be managed by keeping them at a low level of abundance through a constant culling effort, avoiding periodic shifts in their abundance, which exacerbate the impact of predators on native preys (Noorbury, 2001). This is exactly the opposite of the approach of wildlife managers to introduced cottontails, whose densities fluctuate throughout the year due to the hunting season or the poor implementation of local voluntary control schemes. The existence of trophic interactions between cottontails and native foxes should discourage wildlife managers from adopting irregular culling schemes for these two species. We believe that numerical control of foxes is unlikely to be effective to restore hare populations in the study area, because it is far from reaching the enforcement standards that are required for a serious impact on fox populations (Mahon, 2009) and because abundant cottontails can enable rapid recovery of fox populations. Similarly, we believe that carrying out irregular numerical control of cottontails, without a constant effort, may negatively affect native species without preventing a future recovery of cottontail populations. We suggest that a data-informed game management activity should be focused on a regular control of cottontail populations, and we encourage future human dimensions research aimed at designing the most effective control strategy (Santo *et al.*, 2015).

Acknowledgements

We would like to thanks Francesco Campra (Città Metropolitana di Torino), who provided us the population trends.

References

- Anderson, D. R. (2001). The need to get the basics right in wildlife field studies. *Wildlife Soc. B.*, **29**, 1294-1297.
- Balestrieri, A., Remonti, L. & Prigioni, C. (2006). Local feeding specialization of the red fox (*Vulpes vulpes*) in response to eastern cottontail (*Sylvilagus floridanus*) introduction (NW Italy). *Hystrix*, **16**, 113-126.
- Barrio, I. C., Acevedo, P., & Tortosa, F. S. (2010). Assessment of methods for estimating wild rabbit population abundance in agricultural landscapes. *Eur. J. Wildlife Res.*, **56**, 335-340.
- Bellard, C., Genovesi, P. & Jeschke, J.M. (2016). Global patterns in threats to vertebrates by biological invasions. *P. Roy. Soc. B-Biol. Sci.*, **283**, 201-245.
- Beard, K. H. & O'Neill, E. M. (2005). Infection of an invasive frog *Eleutherodactylus coqui* by the chytrid fungus *Batrachochytrium dendrobatidis* in Hawaii. *Biol. Conserv.*, **126**, 591-595.
- Bergstrom, D. M., Lucieer, A., Kiefer, K., Wasley, J., Belbin, L., Pedersen, T. K., & Chown, S. L. (2009). Indirect effects of invasive species removal devastate World Heritage Island. *J. Appl. Ecol.*, **46**, 73-81.
- Bertolino, S., Cordero di Montezemolo, N. & Perrone, A. (2011b). Daytime habitat selection by introduced eastern cottontail (*Sylvilagus floridanus*) and native European hare (*Lepus europaeus*) in northern Italy. *Zool. Sci.*, **28**, 414-419.
- Bertolino, S., Ingegno, B. & Girardello, M. (2011). Modelling the habitat requirements of invasive eastern cottontail (*Sylvilagus floridanus*) introduced to Italy. *Eur. J. Wildlife Res.*, **57**, 267-274.
- Bertolino, S., Hofmannová, L., Girardello, M. & Modry, D. (2010). Richness, origin and structure of an Eimeria community in a population of Eastern cottontail (*Sylvilagus floridanus*) introduced into Italy. *Parasitology*, **137**, 1179-1186.

- Bertolino, S., Perrone, A., Gola, L. & Viterbi, R. (2011a). Population density and habitat use of introduced eastern cottontail (*Sylvilagus floridanus*) in comparison with the native European hare (*Lepus europaeus*). *Zool. Stud.*, **50**, 315–326.
- Bertolino, S., Cordero di Montezemolo, N. & Perrone, A. (2013). Habitat use of coexisting introduced eastern cottontail and native European hare. *Mamm. Biol.*, **78**, 235–240.
- Bertolino, S., Cordero di Montezemolo, N., Preatoni, D.G., Wauters, L.A. & Martinoli, A. (2014). A grey future for Europe: *Sciurus carolinensis* is replacing native red squirrels in Italy. *Biol. Invasions*, **16**, 53–62.
- Blackburn, T. M., Essl, F., Evans, T., Hulme, P. E., Jeschke, J. M., Kühn, I., Kumschick, S., Mrugala, A., Nentwig, N., Pergl, J., Pyšek, P., Rabitsch, W., Ricciardi, A., Richardson, D.M., Sendek, A., Vilà, M., Wilson, J.R.U., Winter, M., Genovesi, P. & Bacher, S. (2014). A unified classification of alien species based on the magnitude of their environmental impacts. *PLoS Biol*, **12**, e1001850
- Bruemmer, C.M., Rushton, S.P., Gurnell, J., Lurz, P.W.W., Nettleton, P., Sainsbury, A. W., Duff, J.P., Gilray, J. & McInnes, C. J. (2010). Epidemiology of squirrelpox virus in grey squirrels in the UK. *Epidemiol. Infect.*, **138**, 941.
- Chiari, M., Ferrari, N., Giardiello, D., Avisani, D., Zanoni, M., Alborali, G.L., Lanfranchi, P., Guberti, V., Capucci, L. & Lavazza, A. (2014). Temporal dynamics of European brown hare syndrome infection in Northern Italian brown hares (*Lepus europaeus*). *Eur. J. Wildlife Res.*, **60**, 891–896.
- Chiari, M., Molinari, S., Cavadini, P., Bertasi, B., Zanoni, M., Capucci, L., & Lavazza, A. (2016). Red foxes (*Vulpes vulpes*) feeding brown hares (*Lepus europaeus*) infected by European brown hare syndrome virus (EBHSV) might be involved in the spread of the virus. *Eur. J. Wildlife Res.*, **62**, 761–765.
- Chroust, K. (1984). Dynamics of Coccidia Infection in Free Living and Cage-reared Hares. *Acta Vet. Brno*, **53**, 175–182.
- Courchamp, F., Langlais, M. & Sugihara, G. (2000). Rabbits killing birds: modelling the hyperpredation process. *J. Anim. Ecol.*, **69**, 154–164.
- DeCesare, N. J., Hebblewhite, M., Robinson, H. S., & Musiani, M. (2010). Endangered, apparently: the role of apparent competition in endangered species conservation. *Anim. Conserv.*, **13**, 353–62.
- Engeman, R. M. (2003). More on the need to get the basics right: population indices. *Wildlife. Soc. B.*, **31**, 286–288.
- Fernex, A., Nagel, P. & Weber, D. (2011). Sites with reduced predation risk to young hares within an agricultural landscape. *Mammalia*, **75**.
- Forsyth, D.M. & Hickling, G.J. (1998). Increasing Himalayan tahr and decreasing chamois densities in the eastern Southern Alps, New Zealand: evidence for interspecific competition. *Oecologia*, **113**, 377–382.
- Fournier-Chambrillon, C., Aasted, B., Perrot, A., Pontier, D., Sauvage, F., Artois, M., Cassiède, J.M, Chauby, X., Dal Molin, A., Simon, C. & Fournier, P. (2004). Antibodies to Aleutian mink disease parvovirus in free-ranging European mink (*Mustela lutreola*) and other small carnivores from southwestern France. *J. Wildlife Dis.*, **40**, 394–402.
- Gibson, L. (2006). The role of lethal control in managing the effects of apparent competition on endangered prey species. *Wildlife Soc. B.*, **34**, 1220–1224.
- Glen, A.S. & Dickman, C.R. (2008). Niche overlap between marsupial and eutherian carnivores: does competition threaten the endangered spotted-tailed quoll?. *J. Appl. Ecol.*, **45**, 700–707.
- Gordo, O., Barriocanal, C. & Robson, D. (2011). Ecological impacts of the North Atlantic Oscillation (NAO) in Mediterranean ecosystems. *Hydrological, Socioeconomic and Ecological Impacts of the North Atlantic Oscillation in the Mediterranean Region* (Eds S.M. Vicente-Serrano & R.M. Trigo), pp. 153–170. Springer Netherlands.

- Gortazar, C., Millán, J., Acevedo, P., Escudero, M. A., Marco, J. & De Luco, D. F. (2007). A large-scale survey of brown hare *Lepus europaeus* and Iberian hare *L. granatensis* populations at the limit of their ranges. *Wildlife Biol.*, **13**, 244-250.
- Gurnell, J., Wauters, L.A., Lurz, P.W.W. & Tosi, G. (2004). Alien species and interspecific competition: effects of introduced eastern grey squirrels on red squirrel population dynamics. *J. Anim. Ecol.*, **73**, 26–35.
- Hackländer, K., Arnold, W. & Ruf, T. (2002a). Postnatal development and thermoregulation in the precocial European hare (*Lepus europaeus*). *Comp. Physiol.*, **172**, 183-190.
- Hackländer, K., Tataruch, F. & Ruf, T. (2002b). The effect of dietary fat content on lactation energetics in the European hare (*Lepus europaeus*). *Physiol. Biochem. Zool.*, **75**, 19-28.
- Hansen, T. F., Stenseth, N. C., Henttonen, H., & Tost, J. (1999). Interspecific and intraspecific competition as causes of direct and delayed density dependence in a fluctuating vole population. *PNAS*, **96**, 986-991.
- Hawkins, C. L., Bacher, S., Essl, F., Hulme, P. E., Jeschke, J. M., Kühn, I., ... & Rabitsch, W. (2015). Framework and guidelines for implementing the proposed IUCN Environmental Impact Classification for Alien Taxa (EICAT). *Divers. Distrib.*, **21**, 1360-1363.
- Hulme, P.E. (2009). Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J. Appl. Ecol.*, **46**, 10-18.
- Kiesecker, J.M., Blaustein, A.R. & Belden, L.K. (2001). Complex causes of amphibian population declines. *Nature*, **410**, 681-684.
- Kristan, W. B., & Boarman, W. I. (2003). Spatial pattern of risk of common raven predation on desert tortoises. *Ecology*, **84**, 2432-2443.
- Kumschick, S., Gaertner, M., Vilà, M., Essl, F., Jeschke, J.M., Pyšek, P., Ricciardi, A., Bacher, S., Blackburn, T.M., Dick, J.T.A., Evans, T., Hulme, P.E., Kuhn, I., Mrugala, A., Pergl, J., Rabitsch, W., Richardson, D.M., Sendek, A. & Evans, T. (2015). Ecological impacts of alien species: quantification, scope, caveats, and recommendations. *BioScience*, **65**, 55-63.
- Lamb, P.J. & Pepler, R.A. (1987). North Atlantic Oscillation: concept and an application. *B. Am. Meteorol. Soc.*, **68**, 1218-1225.
- Lang, J.M. & Benbow, M.E. (2013). Species Interactions and Competition. *Nature Education Knowledge*, **4**, 8.
- Lavazza, A., Cavadini, P., Barbieri, I., Tizzani, P., Pinheiro, A., Abrantes, J., Esteves, P., Grilli, G., Gioia, M., Zanoni, M.G., Meneguz, P.G., Guittòn, J.S., Marchandeu, S., Chiari, M. & Capucci, L. (2015). Field and experimental data indicate that the eastern cottontail (*Sylvilagus floridanus*) is susceptible to infection with European brown hare syndrome (EBHS) virus and not with rabbit haemorrhagic disease (RHD) virus. *Vet. Res.*, **1**, 1-10.
- Lindström, E. R., Andren, H., Angelstam, P., Cederlund, G., Hörnfeldt, B., Jäderberg, L., Lemnel, P.A., Martinsson, B., Sköld, K. & Swenson, J. E. (1994). Disease reveals the predator: sarcoptic mange, red fox predation, and prey populations. *Ecology*, 1042-1049.
- Mahon, P. S. (2009). Targeted control of widespread exotic species for biodiversity conservation: the red fox (*Vulpes vulpes*) in New South Wales, Australia. *Ecol. Manage. Restor.*, **10**, S59-S69.
- Nakagawa, S. & Schielzeth, H. (2013). A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol. Evol.*, **4**, 133-142.
- Noonburg, E.G. & Byers, J.E. (2005). More harm than good: when invader vulnerability to predators enhances impact on native species. *Ecology*, **86**, 2555-2560.
- Norbury, G. (2001). Conserving dryland lizards by reducing predator-mediated apparent competition and direct competition with introduced rabbits. *J. Appl. Ecol.*, **38**, 1350-1361.
- Oliver, M., Luque-Larena, J.J. & Lambin, X. (2009). Do rabbits eat voles? Apparent competition, habitat heterogeneity and large-scale coexistence under mink predation. *Ecol. Lett.*, **12**, 1201-1209.
- Oro, D., & Martínez-Abraín, A. (2007). Deconstructing myths on large gulls and their impact on threatened sympatric waterbirds. *Anim. Conserv.*, **10**, 117-126.

- Ottersen, G., Planque, B., Belgrano, A., Post, E., Reid, P.C. & Stenseth, N.C. (2001). Ecological effects of the North Atlantic oscillation. *Oecologia*, **128**, 1-14.
- Panek, M. (2009). Factors affecting predation of red foxes *Vulpes vulpes* on brown hares *Lepus europaeus* during the breeding season in Poland. *Wildl. Biol.*, **15**, 345-349.
- Pintor, L. M., & Byers, J. E. (2015). Do native predators benefit from non-native prey?. *Ecol. Lett.*, **18**, 1174-1180.
- Preatoni, D. G., Tattoni, C., Bisi, F., Masseroni, E., D'Acunto, D., Lunardi, S., Grimod, I., Martinoli, A. & Tosi, G. (2012). Open source evaluation of kilometric indexes of abundance. *Ecol. Inform.*, **7**, 35-40.
- R Development Core Team 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available from <https://www.R-project.org/>
- Reynolds, J. C. & Tapper, S. C. (1995). Predation by foxes *Vulpes vulpes* on brown hares *Lepus europaeus* in central southern England, and its potential impact on annual population growth. *Wildlife Biol.*, **1**, 145-158.
- Ricciardi, A., Hoopes, M.F., Marchetti, M.P. & Lockwood, J.L. (2013). Progress toward understanding the ecological impacts of nonnative species. *Ecol. Monogr.*, **83**(3), 263-282.
- Roemer, G.W., Coonan, T.J., Garcelon, D.K., Bascompte, J. & Laughrin, L. (2001). Feral pigs facilitate hyperpredation by golden eagles and indirectly cause the decline of the island fox. *Anim. Conserv.*, **4**, 307-318.
- Rödel, H.G. & Dekker, J.J. (2012). Influence of weather factors on population dynamics of two lagomorph species based on hunting bag records. *Eur. J. Wildlife Res.*, **58**, 923-932.
- Santo, A. R., Sorice, M. G., Donlan, C. J., Franck, C. T., & Anderson, C. B. (2015). A human-centered approach to designing invasive species eradication programs on human-inhabited islands. *Global Environ. Chang.*, **35**, 289-298.
- Sanz-Aguilar, A., Martínez-Abraín, A., Tavecchia, G., Mínguez, E., & Oro, D. (2009). Evidence based culling of a facultative predator: efficacy and efficiency components. *Biol. Conserv.*, **142**, 424-431.
- Schmidt, N.M., Asferg, T. & Forchhammer, M.C. (2004). Long-term patterns in European brown hare population dynamics in Denmark: effects of agriculture, predation and climate. *BMC Ecol.*, **4**, 15.
- Schoener, T.W. (1983). Field experiments on interspecific competition. *Am. Nat.*, **122**, 240-285.
- Skerratt, L.F., Berger, L., Speare, R., Cashins, S., McDonald, K.R., Phillott, A.D., Hines, B.H. & Kenyon, N. (2007). Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth*, **4**, 125-134.
- Smith, R.K., Vaughan Jennings, N. & Harris, S. (2005). A quantitative analysis of the abundance and demography of European hares *Lepus europaeus* in relation to habitat type, intensity of agriculture and climate. *Mammal Rev.*, **35**, 1-24.
- Sobrinho, R., Acevedo, P., Escudero, M. A., Marco, J. & Gortázar, C. (2009). Carnivore population trends in Spanish agrosystems after the reduction in food availability due to rabbit decline by rabbit haemorrhagic disease and improved waste management. *Eur. J. Wildlife Res.*, **55**, 161-165.
- Straile, D. & Stenseth, N.C. (2007). The North Atlantic Oscillation and ecology: links between historical time-series, and lessons regarding future climate warming. *Clim. Res.*, **34**, 259-262.
- Strauss, A., White, A. & Boots, M. (2012). Invading with biological weapons: the importance of disease-mediated invasions. *Funct. Ecol.*, **26**, 1249-1261.
- Stromberg, B.E. (1997). Environmental factors influencing transmission. *Vet. Parasitol.*, **72**, 247-264.
- Swihart, R.K. (1986). Home range-body mass allometry in rabbits and hares (*Leporidae*). *Acta Theriol.*, **31**, 139-148.
- Tizzani, P., Menzano, A., Catalano, S., Rossi, L. & Meneguz, P.G. (2011). First report of *Obeliscoides cuniculi* in European brown hare (*Lepus europaeus*). *Parasitol. Res.*, **109**, 963-966.

- Tizzani, P., Catalano, S., Rossi, L., Duignan, P.J. & Meneguz, P. G. (2014). Invasive species and their parasites: eastern cottontail rabbit *Sylvilagus floridanus* and *Trichostrongylus affinis* (Graybill, 1924) from Northwestern Italy. *Parasitol. Res.*, **113**, 1301-1303.
- Vidus-Rosin, A., Meriggi, A., Cardarelli, E., Serrano-Perez, S., Mariani, M.C., Corradelli, A. & Barba, A. (2011). Habitat overlap between sympatric European hares (*Lepus europaeus*) and Eastern cottontails (*Sylvilagus floridanus*) in northern Italy. *Acta Theriol.*, **56**, 53-61.
- Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C.F., Forchhammer, M.C., Grytnes, J.A., Guisan, A., Heikkinen R.H., Høye T.T., Kuhn, I., Luoto, M., Maiorano, L., Nilsson, M.C., Normand, S., Ockinger, E., Schmidt, N.M., Termansen, M., Timmermann, A., Wardle, D.A., Aastrup, P. & Svenning, J.C. (2013). The role of biotic interactions in shaping distributions and realized assemblages of species: implications for species distribution modelling. *Biol. Rev.*, **88**, 15-30.
- Wittmer, H. U., Serrouya, R., Elbroch, L., & Marshall, A. J. (2013). Conservation strategies for species affected by apparent competition. *Conserv. Biol.*, **27**, 254-260.
- Zanet, S., Palese, V., Trisciuglio, A., Alonso, C. C. & Ferroglio, E. (2013). *Encephalitozoon cuniculi*, *Toxoplasma gondii* and *Neospora caninum* infection in invasive Eastern Cottontail Rabbits *Sylvilagus floridanus* in Northwestern Italy. *Vet. Parasitol.*, **197**, 682-684.
- Zavaleta, E. S., Hobbs, R. J., & Mooney, H. A. (2001). Viewing invasive species removal in a whole-ecosystem context. *Trends Ecol. Evol.*, **16**, 454-459.
- Zhang, J., Fan, M. & Kuang, Y. (2006). Rabbits killing birds revisited. *Math. Biosci.*, **203**, 100-123.
- Zuur, A., Ieno, E.N., Walker, N., Saveliev, A.A., Smith, G.M. (2009). Mixed Effects Models and Extensions in Ecology with R. *Springer*.

Table 1. Significant Pearson's correlations between NAO and local climate.

NAO index	Local climate	Pearson's correlation	
		Coefficient (r)	p-value
Winter (January-March)	Winter temperature (°C)	r = 0.79	p < 0.01
	Winter rainfalls (mm)	r = -0.52	p < 0.01
	Winter moisture (%)	r = -0.64	p < 0.01
	Summer rainfalls (mm)	r = 0.51	p < 0.01
Summer (July-September)	Summer temperature (°C)	r = -0.30	p < 0.01
	Summer rainfalls (mm)	r = 0.02	p < 0.01
	Spring temperature (°C)	r = -0.57	p < 0.01
	Spring rainfalls (mm)	r = 0.70	p < 0.01

Table 2. Generalized Linear Mixed Model output

Fixed effects				
	Coefficients	Standard error	t-value	p
Intercept	-0.80480	0.20251	-3.974	0.00007
Cottontail	0.18755	0.03623	5.177	0.0000002
Fox	0.75206	0.21159	3.554	0.000379
Summer NAO	-0.12221	0.07370	-1.658	0.097280
Winter NAO	-0.43381	0.12672	-3.423	0.000618
Cottontail x Fox	-0.15720	0.04586	-3.428	0.000602
Cottontail X Winter NAO	0.02711	0.01597	1.698	0.089553
Winter NAO X Summer NAO	-0.36104	0.09847	-3.667	0.000246
Random effects				
	Variance	Standard deviation		
Intercept (Site)	0.3331	0.5772		
Residual	0.3112	0.5579		
Model indexes				
AIC = 299.1	BIC =332.0	LogLik= -139.6	Deviance=279.1	Residual = 189

Fig. 1. Study area and the network of protected areas around the city of Torino where species were sampled.

Fig. 2. Effect of foxes on hare abundance, at different levels of cottontail abundance; hares data are log-transformed