AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Association Between Lifestyle and Systemic Arterial Hypertension in Young Adults: A National, Survey-Based, Cross-Sectional Study

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1637861> since 2017-05-24T12:02:07Z

Published version:

DOI:10.1007/s40292-016-0135-6

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**Association between lifestyle and systemic arterial hypertension in young adults:
a national, survey-based, cross-sectional study.**

Bruno RM ¹, Pucci G ², Rosticci M ³, Guarino L ⁴, Guglielmo C ⁴, Agabiti Rosei C ⁵, Monticone S ⁶, Giavarini A ⁷, Lonati C ⁷, Torlasco C ⁸, Fedecostante M ⁹, Manzi MV ¹⁰, Pezzutto F ¹¹, Artom N ¹², Battistoni A ¹³, Pignatelli G ¹³, Sanga V ¹⁴, Pengo MF ¹⁴, on behalf of the Young Investigators Group of the Italian Society of Arterial Hypertension.

¹ Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy

² Department of Medicine, University of Perugia, Italy

³ Department of Medical and Surgery Sciences, University of Bologna, Italy

⁴ Biomedical Department of Internal Medicine, University of Palermo, Italy

⁵ Department of Clinical and Experimental Sciences, University of Brescia, Italy

⁶ Department of Medical Sciences, University of Torino, Italy

⁷ Cardiovascular Department, IRCCS Ospedale Maggiore Policlinico, University of Milano, Italy

⁸ Department of Health Sciences, University of Milano-Bicocca, Italy

⁹ Italian National Research Centre on Aging "U.Sestilli", IRCCS-INRCA, Ancona, Italy

¹⁰ University of Napoli, Italy

¹¹ Department of Experimental and Clinical Medical Sciences, University of Udine

¹² Department of Internal Medicine, University of Genova, Italy

¹³ Department of Clinical and Molecular Medicine, Division of Cardiology, Faculty of Medicine and Psychology, Sant'Andrea Hospital, University of Rome Sapienza, Rome, Italy

¹⁴ Department of Medicine (DIMED), University of Padova, Italy

Word count: 4938 words

Key words: hypertension, young adults, dietary habits, lifestyle, blood pressure

Corresponding author:

Rosa Maria Bruno

Department of Clinical and Experimental Medicine, University of Pisa

Via Roma 67, 56123 Pisa, Italy

Email: rosam.bruno@gmail.com

Phone: +39 050 992914

Fax: +39 050 992409

Abstract

Introduction: The prevalence of systemic arterial hypertension in young adults is increasing worldwide in association with modifiable risk factors.

Aim: to assess the prevalence of high blood pressure (BP) in young adults participating to a screening campaign during the World Hypertension Day (17/05/2014), and to determine the possible association with lifestyle factors.

Methods: 493 individuals aged 18-35 years were selected in 13 Italian cities. All participants underwent BP measurement together with the administration of a questionnaire exploring: medical and drug history; traditional cardiovascular risk factors and diseases; dietary pattern; salt intake; sleep habits; mood disorders.

Results: High BP ($\geq 140/90$ mmHg) was found in 54 individuals, with a prevalence of 11% and awareness of 28%. Those with high BP values were more frequently men, reported a higher BMI and a greater use of corticosteroids and non-steroidal anti-inflammatory drugs, and had a lower anxiety score. Concerning dietary habits, they were more likely to eat cheese /cold cuts ≥ 3 times/week, to have their meals out ≥ 1 /day and to eat in fast foods ≥ 1 /week. In the multiple logistic regression analysis, male sex [OR 3.19, 95%CI(1.33-7.63)], BMI [OR 1.14 95%CI(1.04-1.25)], eating in fast foods [OR 3.10 95%CI(1.21-7.95)], and anxiety [OR 0.85 95%CI(0.75-0.97)], were independently associated with high BP.

Conclusions: High BP values were found in 11% young adults. Male sex, adiposity and alimentary habits were the main determinants of high BP values, indicating that young men are a suitable target for healthy lifestyle interventions.

INTRODUCTION

High blood pressure (BP) is the single biggest contributor to the global disease and mortality burden, leading to 9,4 millions of deaths each year [1]. Being tightly connected to the prevalence of modifiable risk factors, the majority of these deaths could be prevented by prompting a healthier lifestyle [2]. In contrast, the prevalence of systemic arterial hypertension continues to rise, especially in young adults [3]: in the United States it has reached 19% in 2008 in the 24-32 age group, according to the Add Health study [4]. More worryingly, even amongst the younger hypertension still remains under-diagnosed and undertreated [5-7]. In study conducted by general practitioners, a diagnosis of hypertension was made only in half individuals aged 18-24 years with high BP values [8]. Furthermore, young adults have consistently lower prevalence of hypertension control compared with middle-aged (40–59 years) and older (≥ 60 years) individuals [9]. Medical inertia towards hypertension in young adults might increase cumulative BP exposure, leading to early cardiovascular complication and myocardial dysfunction in middle age [10].

There is paucity of data regarding hypertension prevalence and determinants in young adults in Italy: large European surveys, with Italy among the countries involved, included individuals older than 35 years [11], patients in secondary prevention [12] or with established hypertension [13]. The Hypertension and Ambulatory Recording Venetia Study (Harvest) selected about 1,000 patients with established hypertension or borderline BP values, aged 18-45 years old, and gave a crucial insight about clinical characteristics of hypertension in young adults [14]; however hypertension prevalence and determinants in young adults in Italy is still unknown [6].

Diet, among the other environmental factors related to hypertension, is likely to play the predominant role in blood pressure homeostasis. Reduced salt intake, weight loss, and moderation of alcohol consumption are established dietary modifications that lower BP [15], but also increased potassium intake and consumption of dietary patterns based on the Dietary Approaches to Stop Hypertension (DASH) diet can prove beneficial [16]. However, a secular trend towards a decreased diet quality has been described over the past three decades, with the lowest scores observed in the young adults [17]. For example, a recent observational cross-sectional study showed an increased prevalence of hypertension among consumers of street food [18].

Other emerging, modifiable risk factors might also play a major role in the increased hypertension prevalence in young adults. For instance, sleep, and more specifically sleep loss, has been frequently associated with the development of hypertension, especially in individuals < 60 years old

[19]. A recent study found that, after controlling for confounders, insomnia combined with physiological hyper-arousal increased the odds of hypertension by 300% [20], though conflicting data exist [21]. Mood disorders have also been linked with cardiovascular diseases and hypertension may well be the intermediary link [22]. Although Bosworth *et al* [23] found an association between elevated cardiovascular reactivity during stress and the development of hypertension, other authors had not confirmed this hypothesis [24].

Thus, there is a strong need to understand the importance of these novel potential risk factors for hypertension in large-scale population cohort studies involving young adults; should their causative role be confirmed, specific measure will be required to halt the diffusion of arterial hypertension and cardiovascular diseases [6]. To the best of our knowledge, no studies are available on the prevalence and determinants of hypertension in young adults in Italy.

The aims of this study were: to investigate the prevalence of arterial hypertension in a nationwide sample of individuals aged between 18 and 35 years recruited in a multicentre study; to explore the association between hypertension and both traditional and emerging risk factors, with particular attention to lifestyle.

METHODS

Study population

The study was performed during the World Hypertension Day (WHD, 17/05/2014), a global initiative supported by the International Society of Hypertension (ISH) and World Hypertension League (WHL) and locally promoted by the Italian Society of Hypertension (Società Italiana dell'Iipertensione Arteriosa – SIIA). Citizens approaching to the stalls, set up on purpose in city squares, public hospitals and other places of interest, were provided with leaflets and given information about hypertension and cardiovascular risk factors and were offered a free consultation by an Hypertension specialist and blood pressure measurement. Individuals aged 18-35 years were asked to join in the survey. Inclusion criteria were: age between 18 and 35 years; signed informed consent. Exclusion criteria were: inability to complete the survey; missing BP values.

Measurements

The individuals aged 18-35 years conforming inclusion/exclusion criteria and accepting to participate in the survey were asked to fill in a short (about 10-min long) self-administered questionnaire including demographic and anthropometric data such as age, sex, self-reported body weight and height, medical history, current drug treatment, smoke habits, physical activity, alcohol intake and usual diet; furthermore, specific questions aimed at exploring depression and anxiety traits, sleep habits and the amount of time spent using computer or smartphones. In particular, specific items from validated questionnaires were embedded in order to explore the prevalence of: sleep-disordered breathing [25]; depression, anxiety and stress [26]; dietary sodium intake [27]. The English version of the questionnaire is available as on line supplement. Lastly, BP was measured three times by a Hypertension specialist in all centres, after at least 5 minutes of rest in the sitting position, according to the current guidelines [28], using an oscillometric validated device. Written informed consent was obtained from all recruited patients.

Statistical analysis

All questionnaires were anonymised and data, collected by each recruiting team, were sent to the coordinating centre based in Pisa for statistical analysis, performed by means of NCSS 8 software (NCSS; Kaysville, Utah; USA). Continuous, normally distributed, variables are presented as mean and standard deviation; continuous, not normally distributed, variables are presented as median and

25-75 percentile; categorical variables are presented by counts and percentages.

Systemic arterial hypertension was defined by the mean of three BP measurements ≥ 140 mmHg for the systolic and / or ≥ 90 mmHg for the diastolic BP or when the volunteer was habitually treated with BP lowering drugs [28]. Patients with systolic, diastolic or systo-diastolic hypertension were identified. The whole sample was then divided into two subgroups of normotensive and hypertensive patients. The two subgroups were compared using unpaired t-test for continuous, normally distributed variables, Wilcoxon Rank Sum Test for continuous, not normally distributed variables and Chi-square test for discrete variables. When feasible, items from the questionnaires (See supplemental material) were transformed in dichotomous variables. The following dummy variables were created:

- question #8: eat out ≥ 1 times/day; eat restaurant/canteen ≥ 1 times/week; eat fast food ≥ 1 times/week; eat bar/street food ≥ 1 times/week; (1=yes for all; 0=no)
- question #9: add salt (often / always =1; never / sometimes= 0)
- question #10: salt bread (4 or more rolls/slices=1; 3 or less = 0)
- question #11: cheese/cold cuts (3 or more times/week=1; 2 or less=0)
- question #12: feel thirsty (always-often=1; never/rarely=0)
- question #13: food insipid out (insipid=1; normal/salty=0)

In order to detect possible determinants of hypertension in the study sample, a multiple logistic regression model was built, considering hypertension as the dependent variable. The independent variable were chosen among those associated with $p < 0.05$ with hypertension in the univariate analysis. A gender-stratified analysis was also performed. A p value less than 0.05 was considered significant throughout the study.

Results

During the 2014 "World Hypertension Day", data were collected from 6356 individuals (age range 18-105 years) in 50 hypertension centers[29]. 15 Hypertension centers in 13 cities (Ancona, Bologna, Brescia, Genova, Milano, Napoli, Palermo, Padova, Pisa, Roma, Torino, Terni, Udine) participated in this study: 508 individuals aged 18-35 years were selected and were given the self-administered questionnaire described above.. Based on inclusion / exclusion criteria, 4 individuals were excluded as were aged <18 years and 11 individuals for missing BP values. The final analysis was therefore performed on 493 individuals. Hypertension was present in 54 individuals, for a prevalence of 11%: 4 were already on BP-lowering treatment, 30 had systolic hypertension, 10 diastolic hypertension and 10 systo-diastolic hypertension. Only 15 out of 54 individuals were aware of their hypertension status (Table 1). Forty-one out of 54 hypertensive individuals were men; furthermore, hypertensive individuals showed a greater BMI and similar occupational and educational status than normotensives.

According to the self-reported medical history, the prevalence of established cardiovascular disease or its equivalents (diabetes mellitus, kidney disease) was very low (<1%), but significantly greater in the hypertensive subgroup (Table 1), while there were no differences as far as non-cardiovascular diseases were concerned (data not shown). Drug history revealed a significantly greater use of non-steroidal anti-inflammatory drugs and/or corticosteroids in the hypertensive subgroup (7.7% vs 1.6%, $p=0.006$). High birth weight was reported by 4.7%, while low birth weight and/or preterm birth by 10.3% of the study population, with no significant difference between the hypertensive and normotensive subgroups.

Among lifestyle measures, no significant differences in alcohol consumption, smoking habit, and habitual physical activity were observed (Table 1). Interestingly, despite a similar reported time spent using computer (>5 hours/day: 24.5% vs 20.2%, $p=0.47$), a subjective perception of an excessive use tended to be more frequent in the hypertensive subgroup (30.8% vs 20.3%, $p=0.08$).

As far as mood and sleep disorders are concerned, we found a significantly lower prevalence of anxiety in the hypertensive population. Hypertensive individuals more often reported snoring or witnessed sleep apneas, whereas subjective sleep duration was similar to normotensive individuals (Table 2). By analyzing questionnaire items exploring salt intake, hypertensive individuals were more likely to eat cheese and/or cold cuts 3 or more times/week (Figure 1). Furthermore, they were more likely to have their meals out 1 or more times/day; in particular, the prevalence of eating in fast-foods one or more times/week was almost two-fold in hypertensive in comparison to normotensive individuals (Figure 2).

Gender differences in clinical and lifestyle characteristics were analyzed. In our study sample,

women were younger, more frequently students and with a higher level of education compared to men (Table 3). They also had lower BMI and BP values, but reported a similar prevalence of CV risk factors and CV disease. Overall, women practiced less physical activity than men, while alcohol and cigarette habits were equally prevalent. Drug history revealed a significantly greater use of non-steroidal anti-inflammatory drugs and/or corticosteroids in men (4.3% vs 0.7%, $p=0.01$), while oral contraceptive were used by 9.6% of women. Despite a similar reported time spent at the computer (>5 hours/day: 21.6% vs 20.0%, $p=0.66$), a subjective perception of an excessive use was more frequent among men than women (26.3% vs 17.6%, $p=0.02$).

As far as mood and sleep disorders are concerned, women reported higher stress scores compared to men, whereas snoring and sleep apneas were more frequent in men (Table 4). In terms of salt intake, men were more likely to eat cheese and/or cold cuts 3 or more times/week and to eat more bread. Furthermore, they were more likely to have their meals out 1 or more times/day; the frequency of eating in fast-foods, restaurants, bars or having street food was higher in men for all the above mentioned categories (Table 4).

A multiple logistic regression model was built in order to identify risk factors independently associated to higher BP values in the overall population. Male sex, BMI and eating in fast foods one or more times per week were associated with an increased probability of having high BP values, while anxiety appeared as a protective factor (Table 5).

We performed the same analysis in men and women separately, in order to identify gender-specific predictors of high BP. Compared to the normotensive, in hypertensive men the following lifestyle habits were more prevalent: eating in fast foods one or more times/week (28.9 vs 11.3%, $p=0.005$), adding salt to prepared food (57.5 vs 40.2%, $p=0.047$), consuming cheese/cold cuts (75.6 vs 53.7%, $p=0.01$) and using NSAID/corticosteroids (10.0 vs 2.9%, $p=0.046$); furthermore, BMI was higher in the hypertensive group [25.8 (22.5-28.9) vs 23.7 (22.1-25.9) kg/mq, $p=0.003$]. In a model including the aforementioned variables, BMI [OR 1.13 (95%CI 1.03-1.24)] and eating in fast foods one or more times/week [OR 3.62 (95%CI 1.34-9.79)] were confirmed as independent predictors, together with the consumption of cheese and/or cold cuts 3 or more times/week [OR 3.04 (95%CI 1.23-7.51)]. Conversely, we could not find any variable associated to high BP values in women.

Discussion

This cross-sectional survey was aimed at establishing the prevalence of high BP values in a sample of young adults recruited during the 2014 World Hypertension Day initiative in Italy and at exploring its relationship with lifestyle habits. The main result of this study is that the prevalence of high BP values among Italian individuals aged 18-35 years is a not negligible 11%, with a very low awareness and treatment percentage. The main predictors of the hypertensive status were gender, BMI and dietary habits such as eating in fast foods. Furthermore, we found remarkable gender differences, with women adhering to a globally healthier dietary lifestyle, with lower BMI and BP values. As a consequence, an association between incorrect dietary habits and high BP was found only in men.

This survey highlighted remarkable differences in hypertension prevalence, awareness and treatment between young adults and the overall population who had their BP measured in Italy during the WHD in the same years and recently reported by Tocci et al [30]. Indeed young adults, in spite of a lower hypertension prevalence (11% vs 55.4%), exhibited considerably lower rates of hypertension awareness (28.8% vs 67.4%) and treatment (7.7% vs 35.3%), in line with US data[3]. While hypertension prevalence has not substantially varied in Italy in the last decade, awareness and control has improved over time since 2004 WHD[30]. Though temporal trends in Italian young adults are not available, US data suggest a surge in prevalence [3, 4]. Thus, this survey suggests that an educational intervention aimed at improving knowledge and control of hypertension in young adults is strongly needed in Italy. In particular, screening campaigns in young adults and physician education to contrast medical inertia might be particularly useful [8].

The prevalence of systemic arterial hypertension among children and adolescents has increased in concert with the marked surge in obesity among the young [31], so that essential hypertension is the most prevalent form after the age of 6 years [32]. Accordingly, our study suggests that body weight and dietary habits are the main factors influencing BP in a cohort of young adults aged 18-35 years. Interestingly, the effect of dietary habits such as eating out often and eating cheese and cold cuts on BP is independent of BMI, suggesting that not only caloric intake, but also nutrient composition, especially sodium content, might play a role. Salt intake reduction is probably the single non-pharmacological intervention able to reduce BP, cardiovascular disease and mortality worldwide. The beneficial effects of salt intake reduction are constantly present regardless of age, gender and ethnicity [33]. Our results are in line with with a cross-sectional analysis collected in a cohort of 1248 adolescents aged 17 years old, showing that BMI and urinary sodium/potassium ratio were

significantly associated with BP values both in males and females [34]. Recently a retrospective survey demonstrated that among 300 young adults with incident hypertension, only 55% had documented lifestyle education within one year [35]. Conversely, our results clearly indicate that lifestyle education should be the most important intervention for BP reduction in young adults, and in particular in men. A poor quality-diet is constantly observed in young adults, as demonstrated by large cohort studies [17]. Aging is usually accompanied by positive changes in diet quality [17]: however, an unhealthy lifestyle in young age might not be innocent, since maintaining a healthy lifestyle throughout young adulthood is strongly associated with a low cardiovascular disease risk profile in middle age [36].

We did not find a correlation between alcohol consumption and BP values, unlike previous studies [34, 37]. The discrepancy with studies conducted in other countries might be due to the fact that, though alcohol is known to increase BP in a dose-dependent manner, wine consumption (traditionally more frequent in Italy) and other alcoholic beverages might have opposite effects on BP, due to their different polyphenol content [38]. Interestingly, in a study conducted in Italy, alcohol-related increase in BP was evident at the 24-hours blood pressure monitoring, but not at office BP [37].

The impact of emerging risk factors on BP appears to be negligible in the study population, although trends for a perceived excessive computer use and for reported snoring or sleep apnoeas were found in the hypertensive subgroup. As a matter of fact, larger population studies are required to reveal subtle effects. The inverse association found between anxiety score and BP values might be attributed to a population bias: anxious, otherwise healthy, people are more likely to volunteer for screening campaigns like the present one. Accordingly, in a retrospective analysis including 4362 patients from primary care, patients with anxiety and/or depression have a greater healthcare utilization, leading to a faster hypertension control [39].

It is worthwhile to comment upon gender differences in clinical and lifestyle profile in the present cohort. Women had globally healthier dietary habits, conceivably responsible for the lower BP and BMI values found in comparison with men. Hypertension in women was not associated to lifestyle characteristics, in contrast to men. A recent survey demonstrated that young women following a healthy lifestyle were virtually free of cardiovascular disease later on [40]. Taken together, these results suggest that gender differences in cardiovascular disease are attributable not only to biological/hormonal differences, but also to cultural and social factors influencing adherence to a healthy lifestyle. Effective strategies aimed at implementing lifestyle education especially in young men are warranted and might play a relevant role in hypertension prevention in the youth.

We must acknowledge some limitations. First, BP was measured on a single occasion, while to ascertain and confirm the diagnosis of arterial hypertension several measurements performed at least in two different occasions are necessary, often together with home blood pressure monitoring and, in selected cases, 24-hour ABPM [28]. Second, because of the nature of the World Hypertension Day, weight and height data were not measured directly, but referred, with the subsequent risk of weight underestimation and height overestimation; different settings (city squares, public hospitals and so on) might also have influenced BP values. Third, a selection bias, related to self-selection of participants who decided to have their BP measured in open checkpoints during the World Hypertension Day, cannot be excluded: thus, participants in this survey may not be representative of the general population. Finally, lack of blood and urine sample collection did not allowed a more accurate assessment of CV risk profile, sodium intake and secondary forms of hypertension.

Conclusions

In conclusion, this cross-sectional survey performed during the 2014 World Hypertension Day initiative, suggested a relevant prevalence of high BP values among Italian young adults, with a very low rate of awareness and treatment. The gap between actual and desired BP control in Italy appears to be wider in young adults than in the older population. Since adiposity and dietary habits are the main predictors of the hypertensive status, especially in men, public health campaigns for hypertension prevention should focus on this target to be effective, in order to reach the ambitious 70% goal for BP control, which was the target of SIIA actions for 2015[7], also in young adults.

Figure legends

Figure 1. Prevalence of salt intake - related habits to the overall study population

Figure 2. Prevalence of lifestyle habits related to eating out in the overall study population

Table 1. Clinical characteristics of the study population according to the hypertensive status

	Overall population (n=493)	Normotensive individuals (n=439)	Hypertensive individuals (n=54)	P value
Men (%)	43.8	39.9	75.9	<0.0001
Age (years)	25.7±4.4	25.6±4.4	26.5±4.6	0.14
BMI (kg/mq)	22.5 (20.3-25.0)	22.2 (20.2-24.5)	25.5 (22.4-28.6)	<0.0001
Highest level of education attained:				0.65
Degree or more	40.4	40.3	41.5	
High school diploma	54.5	54.9	50.9	
Less than high school	5.1	4.8	7.5	
Occupational status:				0.41
Unemployed	12.2	12.6	9.3	
Employed	34.8	33.8	42.6	
Student	53.0	53.7	48.1	
Hypertension awareness (%)	5.2	2.3	28.8	<0.0001
Use of BP-lowering drugs (%)	0.8	0.0	7.7	<0.0001
Systolic BP (mmHg)	119.3	116.3	144.3	<0.0001
Diastolic BP (mmHg)	72.6	75.1	83.0	<0.0001
FC (bpm)	75.9±11.5	75.0±11.0	83.8±12.2	<0.0001
Diabetes	0.8	0.5	3.8	0.01
Previous CV events	0.6	0.2	3.8	0.002
Chronic kidney disease	0.6	0.5	1.9	0.21
Hypercholesterolemia	5.4	4.9	9.3	0.18
Habitual physical activity (%):				
No	51.0	49.9	60.4	0.23
1-2 times/week	14.0	14.8	7.5	
>2 times/week	35.0	35.3	32.1	
Alcohol:>2 drinks/die (%)	27.2	27.5	24.5	0.65
Cigarette smoke (%)				
No	62.0	63.4	50.9	0.07
Ex	14.5	13.3	24.5	
Yes	23.5	23.3	24.5	

Table 2. Sleep and mood characteristics according to the hypertensive status

	Overall population (n=493)	Normotensive individuals (n=439)	Hypertensive individuals (n=54)	P value
Depression score	3 (1-5)	3 (1-5)	3 (0-4)	0.11
Anxiety score	3 (1-6)	3 (1-6)	2 (0-4)	0.02
Stress score	5 (3-8)	6 (3-9)	5 (3-6)	0.07
Short sleep duration (%)	35.2	34.2	43.4	0.18
Snoring /sleep apneas (%)	13.5	12.2	24.5	0.01
Sleep dissatisfaction (%)	26.5	27.2	20.4	0.28

Table 3. Clinical characteristics of the study population according to gender

	Overall population (n=493)	Women (n=277)	Men (n=216)	P value
Age (years)	25.7±4.4	24.9±4.2	26.7±4.5	<0.0001
BMI (kg/mq)	22.5 (20.3-25.0)	20.9 (19.6-23.0)	24.1 (22.3-26.2)	<0.0001
Highest level of education attained:				0.02
Degree or more	40.4	43.5	36.4	
High school diploma	54.5	53.6	55.6	
Less than high school	5.1	2.9	7.9	
Occupational status:				0.02
Unemployed	12.2	12.0	12.5	
Employed	34.8	29.7	41.2	
Student	53.0	58.3	46.3	
Hypertension awareness (%)	5.2	3.7	7.1	0.09
Use of BP-lowering drugs (%)	0.8	0.0	7.7	<0.0001
Systolic BP (mmHg)	119.3	113.2±11.8	127.2±14.5	<0.0001
Diastolic BP (mmHg)	72.6	70.4±8.0	75.5±8.9	<0.0001
FC (bpm)	75.9±11.5	76.6±10.5	74.9±12.7	0.11
Diabetes	0.8	1.1	0.5	0.44
Previous CV events	0.6	0.4	0.9	0.42
Chronic kidney disease	0.6	1.1	0.0	0.13
Hypercholesterolemia	5.4	5.9	4.7	0.56
Habitual physical activity (%):				
No	51.0	56.3	44.2	0.02
1-2 times/week	14.0	13.7	14.4	
>2 times/week	35.0	30.0	41.4	
Alcohol >2drinks/die (%)	27.2	25.2	29.7	0.27
Cigarette smoke (%)				
No	62.0	63.4	50.9	0.07
Ex	14.5	13.3	24.5	
Yes	23.5	23.3	24.5	

Table 4. Sleep and mood characteristics and dietary habits according to gender

	Overall population (n=493)	Women (n=277)	Men (n=216)	P value
Depression score	3 (1-5)	3 (1-6)	2 (1-5)	0.07
Anxiety score	3 (1-6)	3 (1-6)	3 (1-5)	0.03
Stress score	5 (3-8)	6 (3-10)	4 (2-7)	0.0004
Short sleep duration (%)	35.2	33.1	37.9	0.27
Snoring /sleep apneas (%)	13.5	7.6	21.0	<0.0001
Sleep dissatisfaction (%)	26.5	28.3	24.2	0.31
Eat out>1/day	41.6	31.0	55.4	<0.0001
Restaurant/canteen	43.6	39.0	49.8	0.02
Fast food	10.3	6.9	14.6	0.006
Bar street food	40.7	35.2	47.4	0.008
Add salt	40.2	37.7	43.5	0.20
Bread	22.4	15.4	31.2	<0.0001
Cheese / cold cuts	52.3	48.0	57.9	0.03
Thirsty	43.5	44.6	42.2	0.60

Table 5. Multiple logistic regression, considering hypertension as dependent variable, in the overall population.

	OR (95% CL)	Beta coefficient	R²	P value
Anxiety	0.85 (0.75-0.97)	-0.16	0.025	0.018
BMI	1.14 (1.04-1.25)	0.13	0.034	0.003
Cheese/cold cuts 3+times/week	1.99 (0.95-4.17)	0.69	0.013	0.069
Fast food 1+times/week	3.10 (1.21-7.95)	0.99	0.015	0.041
Eat out 1+/day	1.11 (0.51-2.44)	0.11	0.000	0.791
Snoring/apneas	1.35 (0.53-3.42)	0.30	0.001	0.529
Male sex	3.19 (1.33-7.63)	1.15	0.028	0.009
NSAIDs/corticosteroids	1.98 (0.46-8.49)	0.68	0.003	0.355

Bibliography

Bibliografia

1. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, Amann M, Anderson HR, Andrews KG, Aryee M, Atkinson C, Bacchus LJ, Bahalim AN, Balakrishnan K, Balmes J, Barker-Collo S, Baxter A, Bell ML, Blore JD, Blyth F, Bonner C, Borges G, Bourne R, Boussinesq M, Brauer M, Brooks P, Bruce NG, Brunekreef B, Bryan-Hancock C, Bucello C, Buchbinder R, Bull F, Burnett RT, Byers TE, Calabria B, Carapetis J, Carnahan E, Chafe Z, Charlson F, Chen H, Chen JS, Cheng AT, Child JC, Cohen A, Colson KE, Cowie BC, Darby S, Darling S, Davis A, Degenhardt L, Dentener F, Des Jarlais DC, Devries K, Dherani M, Ding EL, Dorsey ER, Driscoll T, Edmond K, Ali SE, Engell RE, Erwin PJ, Fahimi S, Falder G, Farzadfar F, Ferrari A, Finucane MM, Flaxman S, Fowkes FG, Freedman G, Freeman MK, Gakidou E, Ghosh S, Giovannucci E, Gmel G, Graham K, Grainger R, Grant B, Gunnell D, Gutierrez HR, Hall W, Hoek HW, Hogan A, Hosgood HD, 3rd, Hoy D, Hu H, Hubbell BJ, Hutchings SJ, Ibeanusi SE, Jacklyn GL, Jasrasaria R, Jonas JB, Kan H, Kanis JA, Kassebaum N, Kawakami N, Khang YH, Khatibzadeh S, Khoo JP, Kok C, Laden F, Lalloo R, Lan Q, Lathlean T, Leasher JL, Leigh J, Li Y, Lin JK, Lipshultz SE, London S, Lozano R, Lu Y, Mak J, Malekzadeh R, Mallinger L, Marcenes W, March L, Marks R, Martin R, McGale P, McGrath J, Mehta S, Mensah GA, Merriman TR, Micha R, Michaud C, Mishra V, Mohd Hanafiah K, Mokdad AA, Morawska L, Mozaffarian D, Murphy T, Naghavi M, Neal B, Nelson PK, Nolla JM, Norman R, Olives C, Omer SB, Orchard J, Osborne R, Ostro B, Page A, Pandey KD, Parry CD, Passmore E, Patra J, Pearce N, Pelizzari PM, Petzold M, Phillips MR, Pope D, Pope CA, 3rd, Powles J, Rao M, Razavi H, Rehfuess EA, Rehm JT, Ritz B, Rivara FP, Roberts T, Robinson C, Rodriguez-Portales JA, Romieu I, Room R, Rosenfeld LC, Roy A, Rushton L, Salomon JA, Sampson U, Sanchez-Riera L, Sanman E, Sapkota A, Seedat S, Shi P, Shield K, Shivakoti R, Singh GM, Sleet DA, Smith E, Smith KR, Stapelberg NJ, Steenland K, Stockl H, Stovner LJ, Straif K, Straney L, Thurston GD, Tran JH, Van Dingenen R, van Donkelaar A, Veerman JL, Vijayakumar L, Weintraub R, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams W, Wilson N, Woolf AD, Yip P, Zielinski JM, Lopez AD, Murray CJ, Ezzati M, AlMazroa MA, Memish ZA. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012 Dec 15;380(9859):2224-60.
2. Arena R, Guazzi M, Lianov L, Whitsel L, Berra K, Lavie CJ, Kaminsky L, Williams M, Hivert MF, Franklin NC, Myers J, Dengel D, Lloyd-Jones DM, Pinto FJ, Cosentino F, Halle M, Gielen S, Dendale P, Niebauer J, Pelliccia A, Giannuzzi P, Corra U, Piepoli MF, Guthrie G, Shurney D. Healthy Lifestyle Interventions to Combat Noncommunicable Disease-A Novel Nonhierarchical Connectivity Model for Key Stakeholders: A Policy Statement From the American Heart Association, European Society of Cardiology, European Association for Cardiovascular Prevention and Rehabilitation, and American College of Preventive Medicine. *Mayo Clinic proceedings*. 2015 Aug;90(8):1082-103.
3. Gooding HC, McGinty S, Richmond TK, Gillman MW, Field AE. Hypertension awareness and control among young adults in the national longitudinal study of adolescent health. *Journal of general internal medicine*. 2014 Aug;29(8):1098-104.
4. Nguyen QC, Tabor JW, Entzel PP, Lau Y, Suchindran C, Hussey JM, Halpern CT, Harris KM, Whitsel EA. Discordance in national estimates of hypertension among young adults. *Epidemiology*. 2011 Jul;22(4):532-41.
5. Hansen ML, Gunn PW, Kaelber DC. Underdiagnosis of hypertension in children and adolescents. *Jama*. 2007 Aug 22;298(8):874-9.
6. Battistoni A, Canichella F, Pignatelli G, Ferrucci A, Tocci G, Volpe M. Hypertension in Young People: Epidemiology, Diagnostic Assessment and Therapeutic Approach. *High blood pressure & cardiovascular prevention : the official journal of the Italian Society of Hypertension*.

2015 Jul 8.

7. Volpe M, Rosei EA, Ambrosioni E, Cottone S, Cuspidi C, Borghi C, De Luca N, Fallo F, Ferri C, Morganti A, Muiesan ML, Sarzani R, Sechi L, Virdis A, Tocci G, Trimarco B, Filippi A, Mancia G, Italian Society of H. 2012 consensus document of the Italian Society of Hypertension (SIIA): strategies to improve blood pressure control in Italy: from global cardiovascular risk stratification to combination therapy. *High blood pressure & cardiovascular prevention : the official journal of the Italian Society of Hypertension*. 2013 Mar;20(1):45-52.
8. Johnson HM, Thorpe CT, Bartels CM, Schumacher JR, Palta M, Pandhi N, Sheehy AM, Smith MA. Undiagnosed hypertension among young adults with regular primary care use. *Journal of hypertension*. 2014 Jan;32(1):65-74.
9. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Huffman MD, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Magid D, Marcus GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Nichol G, Paynter NP, Schreiner PJ, Sorlie PD, Stein J, Turan TN, Virani SS, Wong ND, Woo D, Turner MB, American Heart Association Statistics C, Stroke Statistics S. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation*. 2013 Jan 1;127(1):e6-e245.
10. Kishi S, Teixido-Tura G, Ning H, Venkatesh BA, Wu C, Almeida A, Choi EY, Gjesdal O, Jacobs DR, Jr., Schreiner PJ, Gidding SS, Liu K, Lima JA. Cumulative Blood Pressure in Early Adulthood and Cardiac Dysfunction in Middle Age: The CARDIA Study. *Journal of the American College of Cardiology*. 2015 Jun 30;65(25):2679-87.
11. Wolf-Maier K, Cooper RS, Kramer H, Banegas JR, Giampaoli S, Joffres MR, Poulter N, Primatesta P, Stegmayr B, Thamm M. Hypertension treatment and control in five European countries, Canada, and the United States. *Hypertension*. 2004 Jan;43(1):10-7.
12. Kotseva K, Wood D, De Bacquer D, De Backer G, Ryden L, Jennings C, Gyberg V, Amouyel P, Bruthans J, Castro Conde A, Cifkova R, Deckers JW, De Sutter J, Dilic M, Dolzhenko M, Erglis A, Fras Z, Gaita D, Gotcheva N, Goudevenos J, Heuschmann P, Laucevicius A, Lehto S, Lovic D, Milicic D, Moore D, Nicolaidis E, Oganov R, Pajak A, Pogosova N, Reiner Z, Stagmo M, Stork S, Tokgozoglu L, Vulic D, on behalf of the EI. EUROASPIRE IV: A European Society of Cardiology survey on the lifestyle, risk factor and therapeutic management of coronary patients from 24 European countries. *European journal of preventive cardiology*. 2015 Feb 16.
13. Bramlage P, Bohm M, Volpe M, Khan BV, Paar WD, Tebbe U, Thoenes M. A global perspective on blood pressure treatment and control in a referred cohort of hypertensive patients. *Journal of clinical hypertension*. 2010 Sep;12(9):666-77.
14. Palatini P, Graniero GR, Mormino P, Nicolosi L, Mos L, Visentin P, Pessina AC. Relation between physical training and ambulatory blood pressure in stage I hypertensive subjects. Results of the HARVEST Trial. *Hypertension and Ambulatory Recording Venetia Study*. *Circulation*. 1994 Dec;90(6):2870-6.
15. Appel LJ, Brands MW, Daniels SR, Karanja N, Elmer PJ, Sacks FM, American Heart A. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension*. 2006 Feb;47(2):296-308.
16. Salehi-Abargouei A, Maghsoudi Z, Shirani F, Azadbakht L. Effects of Dietary Approaches to Stop Hypertension (DASH)-style diet on fatal or nonfatal cardiovascular diseases--incidence: a systematic review and meta-analysis on observational prospective studies. *Nutrition*. 2013 Apr;29(4):611-8.
17. Sijtsma FP, Meyer KA, Steffen LM, Shikany JM, Van Horn L, Harnack L, Kromhout D, Jacobs DR, Jr. Longitudinal trends in diet and effects of sex, race, and education on dietary quality score change: the Coronary Artery Risk Development in Young Adults study. *The American journal of clinical nutrition*. 2012 Mar;95(3):580-6.
18. Buscemi S, Barile A, Maniaci V, Batsis JA, Mattina A, Verga S. Characterization of street food consumption in Palermo: possible effects on health. *Nutrition journal*. 2011;10:119.
19. Palagini L, Bruno RM, Gemignani A, Baglioni C, Ghiadoni L, Riemann D. Sleep loss and

hypertension: a systematic review. *Current pharmaceutical design*. 2013;19(13):2409-19.

20. Li Y, Vgontzas AN, Fernandez-Mendoza J, Bixler EO, Sun Y, Zhou J, Ren R, Li T, Tang X. Insomnia with physiological hyperarousal is associated with hypertension. *Hypertension*. 2015 Mar;65(3):644-50.
21. Haba-Rubio J, Marques-Vidal P, Andries D, Tobback N, Preisig M, Vollenweider P, Waeber G, Luca G, Tafti M, Heinzer R. Objective sleep structure and cardiovascular risk factors in the general population: the HypnoLaus Study. *Sleep*. 2015 Mar;38(3):391-400.
22. Van der Kooy K, van Hout H, Marwijk H, Marten H, Stehouwer C, Beekman A. Depression and the risk for cardiovascular diseases: systematic review and meta analysis. *International journal of geriatric psychiatry*. 2007 Jul;22(7):613-26.
23. Bosworth HB, Bartash RM, Olsen MK, Steffens DC. The association of psychosocial factors and depression with hypertension among older adults. *International journal of geriatric psychiatry*. 2003 Dec;18(12):1142-8.
24. Shinn EH, Poston WS, Kimball KT, St Jeor ST, Foreyt JP. Blood pressure and symptoms of depression and anxiety: a prospective study. *American journal of hypertension*. 2001 Jul;14(7 Pt 1):660-4.
25. Farney RJ, Walker BS, Farney RM, Snow GL, Walker JM. The STOP-Bang equivalent model and prediction of severity of obstructive sleep apnea: relation to polysomnographic measurements of the apnea/hypopnea index. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*. 2011 Oct 15;7(5):459-65B.
26. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *The British journal of clinical psychology / the British Psychological Society*. 2005 Jun;44(Pt 2):227-39.
27. Galletti F, Agabiti-Rosei E, Bernini G, Boero R, Desideri G, Fallo F, Mallamaci F, Morganti A, Castellano M, Nazzaro P, Trimarco B, Strazzullo P, Group M-GPS. Excess dietary sodium and inadequate potassium intake by hypertensive patients in Italy: results of the MINISAL-SIIA study program. *Journal of hypertension*. 2014 Jan;32(1):48-56.
28. Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Bohm M, Christiaens T, Cifkova R, De Backer G, Dominiczak A, Galderisi M, Grobbee DE, Jaarsma T, Kirchhof P, Kjeldsen SE, Laurent S, Manolis AJ, Nilsson PM, Ruilope LM, Schmieder RE, Sirnes PA, Sleight P, Viigimaa M, Waeber B, Zannad F, Task Force M. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Journal of hypertension*. 2013 Jul;31(7):1281-357.
29. Torlasco C, Calvanese C, Faini A, Santini F, Borghi C, Parati G. 5c.02: Prevalence of Hypertension and Other Cardiovascular Risk Factors in Participants in the 2014 Hypertension World Day Campaign in Italy. *Journal of hypertension*. 2015 Jun;33 Suppl 1:e68.
30. Tocci G, Muiesan ML, Parati G, Agabiti Rosei E, Ferri C, Virdis A, Pontremoli R, Mancia G, Borghi C, Volpe M, Working Group of the Italian Society of H, Foundation S. Trends in Prevalence, Awareness, Treatment, and Control of Blood Pressure Recorded From 2004 to 2014 During World Hypertension Day in Italy. *Journal of clinical hypertension*. 2015 Oct 13.
31. Ingelfinger JR. Clinical practice. The child or adolescent with elevated blood pressure. *The New England journal of medicine*. 2014 Jun 12;370(24):2316-25.
32. Gupta-Malhotra M, Banker A, Shete S, Hashmi SS, Tyson JE, Barratt MS, Hecht JT, Milewicz DM, Boerwinkle E. Essential hypertension vs. secondary hypertension among children. *American journal of hypertension*. 2015 Jan;28(1):73-80.
33. He FJ, Li J, Macgregor GA. Effect of longer-term modest salt reduction on blood pressure. *The Cochrane database of systematic reviews*. 2013;4:CD004937.
34. Le-Ha C, Beilin LJ, Burrows S, Huang RC, Oddy WH, Hands B, Mori TA. Oral contraceptive use in girls and alcohol consumption in boys are associated with increased blood pressure in late adolescence. *European journal of preventive cardiology*. 2013 Dec;20(6):947-55.
35. Johnson HM, Olson AG, LaMantia JN, Kind AJ, Pandhi N, Mendonca EA, Craven M, Smith

- MA. Documented lifestyle education among young adults with incident hypertension. *Journal of general internal medicine*. 2015 May;30(5):556-64.
36. Liu K, Daviglius ML, Loria CM, Colangelo LA, Spring B, Moller AC, Lloyd-Jones DM. Healthy lifestyle through young adulthood and the presence of low cardiovascular disease risk profile in middle age: the Coronary Artery Risk Development in (Young) Adults (CARDIA) study. *Circulation*. 2012 Feb 28;125(8):996-1004.
37. Vriz O, Piccolo D, Cozzutti E, Milani L, Gelisio R, Pegoraro F, Garavelli G, D'Este D, Palatini P. The effects of alcohol consumption on ambulatory blood pressure and target organs in subjects with borderline to mild hypertension. HARVEST Study Group. *American journal of hypertension*. 1998 Feb;11(2):230-4.
38. Chiva-Blanch G, Arranz S, Lamuela-Raventos RM, Estruch R. Effects of wine, alcohol and polyphenols on cardiovascular disease risk factors: evidences from human studies. *Alcohol and alcoholism*. 2013 May-Jun;48(3):270-7.
39. Ho AK, Thorpe CT, Pandhi N, Palta M, Smith MA, Johnson HM. Association of anxiety and depression with hypertension control: a US multidisciplinary group practice observational study. *Journal of hypertension*. 2015 Aug 7.
40. Chomistek AK, Chiuve SE, Eliassen AH, Mukamal KJ, Willett WC, Rimm EB. Healthy lifestyle in the primordial prevention of cardiovascular disease among young women. *Journal of the American College of Cardiology*. 2015 Jan 6;65(1):43-51.