

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The quotient set of k-generalised Fibonacci numbers is dense in \mathbb{Q}_p

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1647070> since 2017-08-22T18:50:16Z

Published version:

DOI:10.1017/S0004972716001118

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

THE QUOTIENT SET OF k -GENERALIZED FIBONACCI NUMBERS IS DENSE IN \mathbb{Q}_p

CARLO SANNA

Abstract

The quotient set of $A \subseteq \mathbb{N}$ is defined as $R(A) := \{a/b : a, b \in A, b \neq 0\}$. Using algebraic number theory in $\mathbb{Q}(\sqrt{5})$, Garcia and Luca proved that the quotient set of Fibonacci numbers is dense in the p -adic numbers \mathbb{Q}_p , for all prime numbers p . For any integer $k \geq 2$, let $(F_n^{(k)})_{n \geq -(k-2)}$ be the sequence of k -generalized Fibonacci numbers, defined by the initial values $0, 0, \dots, 0, 1$ (k terms) and such that each term afterwards is the sum of the k preceding terms. We use p -adic analysis to generalize Garcia and Luca's result, by proving that the quotient set of k -generalized Fibonacci numbers is dense in \mathbb{Q}_p , for any integer $k \geq 2$ and any prime number p .

2010 *Mathematics subject classification*: primary 11B39; secondary 11B37, 11B05.

Keywords and phrases: Fibonacci numbers, k -generalized Fibonacci numbers, p -adic numbers, density.

1. Introduction

Given a set of nonnegative integers A , the *quotient set* of A is defined as

$$R(A) := \{a/b : a, b \in A, b \neq 0\}.$$

The question of when $R(A)$ is dense in \mathbb{R}^+ is a classical topic and has been studied by many researchers. Strauch and Tóth [15] proved that if A has lower asymptotic density at least equal to $1/2$ then $R(A)$ is dense in \mathbb{R}^+ (see also [1]). Bukor, Šalát, and Tóth [3] showed that if $A \cup B$ is a partition of \mathbb{N} then at least one of $R(A)$ or $R(B)$ is dense in \mathbb{R}^+ . Moreover, the density of $R(\mathbb{P})$ in \mathbb{R}^+ , where \mathbb{P} is the set of prime numbers, is a well-known consequence of the Prime Number Theorem [10].

On the other hand, the analog question of when $R(A)$ is dense in the p -adic numbers \mathbb{Q}_p , for some prime number p , has been studied only recently [7, 8]. Let $(F_n)_{n \geq 0}$ be the sequence of Fibonacci numbers, defined by $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$, for all integers $n > 1$. Using algebraic number theory in the field $\mathbb{Q}(\sqrt{5})$, Garcia and Luca [8] proved the following result.

THEOREM 1.1. *For any prime p , the quotient set of Fibonacci numbers is dense in \mathbb{Q}_p .*

One of the many generalizations of the Fibonacci numbers is the sequence of k -generalized Fibonacci numbers $(F_n^{(k)})_{n \geq -(k-2)}$, also called *Fibonacci k -step sequence*,

Fibonacci k -sequence, or *k -bonacci sequence*. For any integer $k \geq 2$, the sequence $(F_n^{(k)})_{n \geq -(k-2)}$ is defined by

$$F_{-(k-2)}^{(k)} = \dots = F_0^{(k)} = 0, F_1^{(k)} = 1,$$

and

$$F_n^{(k)} = F_{n-1}^{(k)} + F_{n-2}^{(k)} + \dots + F_{n-k}^{(k)},$$

for all integers $n > 1$.

Usually, the study of the arithmetic properties of the k -generalized Fibonacci numbers is more difficult than that of Fibonacci numbers. Indeed, for $k \geq 3$ the sequence of k -generalized Fibonacci numbers lacks several nice properties of the sequence of Fibonacci numbers, like: being a strong divisibility sequence [13, p. 9], having a Primitive Divisor Theorem [17], and having a simple formula for its p -adic valuation [11, 14].

We give the following generalization of Theorem 1.1.

THEOREM 1.2. *For any integer $k \geq 2$ and any prime number p , the quotient set of the k -generalized Fibonacci numbers is dense in \mathbb{Q}_p .*

It seems likely that Theorem 1.2 could be extended to other linear recurrences over the integers. However, in our proof we use some specific features of the k -generalized Fibonacci numbers sequence. Therefore, we leave the following open question to the interested readers:

QUESTION 1.3. *Let $(S_n)_{n \geq 0}$ be a linear recurrence of order $k \geq 2$ satisfying*

$$S_n = a_1 S_{n-1} + a_2 S_{n-2} + \dots + a_k S_{n-k},$$

for all integers $n \geq k$, where $a_1, \dots, a_k, S_0, \dots, S_{k-1} \in \mathbb{Z}$, with $a_k \neq 0$.

For which prime numbers p is the quotient set of $(S_n)_{n \geq 0}$ dense in \mathbb{Q}_p ?

Clearly, without loss of generality, one can suppose that $\gcd(S_0, \dots, S_{k-1}) = 1$. Also, it seems reasonable assuming that $(S_n)_{n \geq 0}$ is nondegenerate, which in turn implies that $(S_n)_{n \geq 0}$ is definitely nonzero [5, §2.1]. Finally, a necessary condition for $(S_n)_{n \geq 0}$ to be dense in \mathbb{Q}_p is that $(v_p(S_n))_{n \geq 0, S_n \neq 0}$ is unbounded. This is certainly the case if $S_0 = 0$ and $p \nmid a_k$ (since $p \nmid a_k$ implies that $(S_n)_{n \geq 0}$ is periodic modulo p^h , for any positive integer h [5, §3.1]), so this could be an useful additional hypothesis.

2. Proof of Theorem 1.2

From now on, fix an integer $k \geq 2$ and a prime number p . In light of Theorem 1.1, we can suppose $k \geq 3$. Let

$$f_k(X) = X^k - X^{k-1} - \dots - X - 1$$

be the characteristic polynomial of the k -generalized Fibonacci numbers sequence.

It is known [16, Corollary 3.4] that f_k is separable. Let K be the splitting field of f_k over \mathbb{Q}_p and let $\alpha_1, \dots, \alpha_k \in K$ be the k distinct roots of f_k . We have [4, Theorem 1]

$$F_n^{(k)} = \sum_{i=1}^k c_i \alpha_i^n, \quad (2.1)$$

for all integers $n \geq 0$, where

$$c_i := \frac{\alpha_i - 1}{(k+1)\alpha_i^2 - 2k\alpha_i}, \quad (2.2)$$

for $i = 1, \dots, k$.

Now we shall interpolate a subsequence of $(F_n^{(k)})_{n \geq 0}$ by an analytic function over \mathbb{Z}_p . This is a classical method in the study of linear recurrences, which goes back at least to the proof of the Skolem–Mahler–Lech theorem [5, Theorem 2.1].

We refer the reader to [9, Ch. 4–6] for the p -adic analysis used hereafter. Let \mathcal{O}_K be the valuation ring of K ; e and f be the ramification index and the inertial degree of K over \mathbb{Q}_p , respectively; and π be a uniformizer of K .

Since $f_k(0) = -1$, we have that each α_i ($i = 1, \dots, k$) is a unit of \mathcal{O}_K , so that $|\alpha_i|_p = 1$. Hence, in particular, $\alpha_i \not\equiv 0 \pmod{\pi}$. Thus, since $\mathcal{O}_K/\pi\mathcal{O}_K$ is a finite field of p^f elements, we obtain that $\alpha_i^{p^f-1} \equiv 1 \pmod{\pi}$. Now pick any positive integer s such that $p^s \geq e + 1$. Since $|\pi|_p = p^{-1/e}$, we have $\pi^{p^s} \equiv 0 \pmod{p\pi}$, and, in turn, it follows that $\alpha_i^t \equiv 1 \pmod{p\pi}$, where $t := p^s(p^f - 1)$. At this point,

$$|\alpha_i^t - 1|_p \leq |p\pi|_p = p^{-1-1/e} < p^{-1/(p-1)}, \quad (2.3)$$

for $i = 1, \dots, k$.

Now let \log_p and \exp_p denote the p -adic logarithm and the p -adic exponential functions, respectively. Thanks to (2.3) we have that

$$\alpha_i^t = \exp_p(\log_p(\alpha_i^t)),$$

for $i = 1, \dots, k$, which together with (2.1) implies that $F_{nt}^{(k)} = G(n)$ for all integer $n \geq 0$, where

$$G(z) := \sum_{i=1}^k c_i \exp_p(z \log_p(\alpha_i^t)),$$

is an analytic function over \mathbb{Z}_p .

Let $r > 0$ be the radius of convergence of the Taylor series of $G(z)$ at $z = 0$, and let $\ell \geq 0$ be an integer. On the one hand, the radius of convergence of the Taylor series of $G(p^\ell z)$ at $z = 0$ is $p^\ell r$. On the other hand,

$$G(p^\ell z) = \sum_{i=1}^k c_i \exp_p(p^\ell z \log_p(\alpha_i^t)) = \sum_{i=1}^k c_i \exp_p(z \log_p(\alpha_i^{p^\ell t})).$$

Therefore, taking s sufficiently large, we can assume $r > 1$.

In particular, we have

$$G(z) = \sum_{j=0}^{\infty} \frac{G^{(j)}(0)}{j!} z^j, \quad (2.4)$$

for all $z \in \mathbb{Z}_p$.

Now we shall prove that $G'(0) \neq 0$. For the sake of contradiction, assume that

$$G'(0) = \sum_{i=1}^k c_i \log_p(\alpha_i^t) = 0.$$

Since $f_k(0) = -1$ and t is even, we have $\alpha_1^t \cdots \alpha_k^t = 1$, so that

$$\log_p(\alpha_k^t) = -\log_p(\alpha_1^t) - \cdots - \log_p(\alpha_{k-1}^t),$$

and consequently

$$\sum_{i=1}^{k-1} (c_i - c_k) \log_p(\alpha_i^t) = 0. \quad (2.5)$$

We need the following lemma [6, Lemma 1], which is a special case of a general result of Mignotte [12] on Pisot numbers.

LEMMA 2.1. *The roots $\alpha_1, \dots, \alpha_{k-1}$ are multiplicatively independent, that is, $\alpha_1^{e_1} \cdots \alpha_{k-1}^{e_{k-1}} = 1$ for some integers e_1, \dots, e_{k-1} if and only if $e_1 = \cdots = e_{k-1} = 0$.*

Thanks to Lemma 2.1, we know that $\alpha_1^t, \dots, \alpha_{k-1}^t$ are multiplicatively independent. Hence, $\log_p(\alpha_1^t), \dots, \log_p(\alpha_{k-1}^t)$ are linearly independent over \mathbb{Z} . Then by [2, Theorem 1] we get that $\log_p(\alpha_1^t), \dots, \log_p(\alpha_{k-1}^t)$ are linearly independent over the algebraic numbers, hence (2.5) implies

$$c_1 = c_2 = \cdots = c_k. \quad (2.6)$$

At this point, from (2.2) and (2.6), it follows that $\alpha_1, \dots, \alpha_k$ are all roots of the polynomial

$$c_1(k+1)X^2 - (2c_1k+1)X + 1,$$

but that is clearly impossible, since $k \geq 3$. Hence, we have proved that $G'(0) \neq 0$.

Taking $z = 1$ in (2.4), we find that $v_p(G^{(j)}(0)/j!) \rightarrow +\infty$, as $j \rightarrow +\infty$. In particular, there exists an integer $\ell \geq 0$ such that $v_p(G^{(j)}(0)/j!) \geq -\ell$, for all integers $j \geq 0$. As a consequence of this, and since $G(0) = F_0^{(k)} = 0$, taking $z = mp^h$ in (2.4) we get that

$$G(mp^h) = G'(0)mp^h + O(p^{2h-\ell}),$$

for all integers $m, h \geq 0$. Therefore, for $h > h_0 := \ell + v_p(G'(0))$, we have

$$\frac{G(mp^h)}{G(p^h)} - m = \frac{G'(0)mp^h + O(p^{2h-\ell})}{G'(0)p^h + O(p^{2h-\ell})} - m = \frac{O(p^{h-\ell})}{G'(0) + O(p^{h-\ell})} = O(p^{h-h_0}),$$

that is,

$$\lim_{h \rightarrow +\infty} \left| \frac{G(mp^h)}{G(p^h)} - m \right|_p = 0.$$

In conclusion, we have proved that

$$\lim_{v \rightarrow +\infty} \left| \frac{F_{mp^v(p^f-1)}^{(k)}}{F_{p^v(p^f-1)}^{(k)}} - m \right|_p = 0,$$

for all integers $m \geq 0$. In other words, the closure (with respect to the p -adic topology) of the quotient set of k -generalized Fibonacci numbers contains the nonnegative integers \mathbb{N} .

The next easy lemma is enough to conclude.

LEMMA 2.2. *Let $A \subseteq \mathbb{N}$. If the closure of $R(A)$ contains \mathbb{N} , then $R(A)$ is dense in \mathbb{Q}_p .*

PROOF. Let C be the closure of $R(A)$ as a subspace of \mathbb{Q}_p . Since \mathbb{N} is dense in \mathbb{Z}_p , we have $\mathbb{Z}_p \subseteq C$. Moreover, the inversion $\iota : \mathbb{Z}_p^\times \rightarrow \mathbb{Q}_p : x \rightarrow x^{-1}$ is continuous and, obviously, sends nonzero elements of $R(A)$ to $R(A)$, hence $\iota(\mathbb{Z}_p) \subseteq C$. Finally, $\mathbb{Q}_p = \mathbb{Z}_p \cup \iota(\mathbb{Z}_p)$, thus $C = \mathbb{Q}_p$ and $R(A)$ is dense in \mathbb{Q}_p . \square

The proof of Theorem 1.2 is complete.

Acknowledgement

The author thanks the anonymous referee for his careful reading of the paper.

References

- [1] B. Brown, M. Dairyko, S. R. Garcia, B. Lutz, and M. Someck. ‘Four quotient set gems’. *Amer. Math. Monthly* **121** (7) (2014), 590–599.
- [2] A. Brumer. ‘On the units of algebraic number fields’. *Mathematika* **14** (1967), 121–124.
- [3] J. Bukor, T. Šalát, and J. T. Tóth. ‘Remarks on R -density of sets of numbers’. *Tatra Mt. Math. Publ.* **11** (1997), 159–165. Number theory (Liptovský Ján, 1995).
- [4] G. P. B. Dresden and Z. Du. ‘A simplified Binet formula for k -generalized Fibonacci numbers’. *J. Integer Seq.* **17** (4) (2014), Article 14.4.7, 9.
- [5] G. Everest, A. van der Poorten, I. Shparlinski, and T. Ward. *Recurrence sequences, Mathematical Surveys and Monographs*, Volume 104 (American Mathematical Society, Providence, RI, 2003).
- [6] C. Fuchs, C. Hutle, F. Luca, and L. Szalay. ‘Diophantine triples and k -generalized Fibonacci sequences’. *Bull. Malays. Math. Sci. Soc.* (2016), 1–17.
- [7] S. R. Garcia, Y. X. Hong, F. Luca, E. Pinsky, E. Schechter, and A. Starr. ‘ p -adic quotient sets’ <https://arxiv.org/abs/1607.07951>.
- [8] S. R. Garcia and F. Luca. ‘Quotients of Fibonacci numbers’. *Amer. Math. Monthly* (to appear).
- [9] F. Q. Gouvêa. *p -adic numbers*. Universitext (Springer-Verlag, Berlin, 1997), 2nd edn. An introduction.
- [10] D. Hobby and D. M. Silberger. ‘Quotients of primes’. *Amer. Math. Monthly* **100** (1) (1993), 50–52.
- [11] T. Lengyel. ‘The order of the Fibonacci and Lucas numbers’. *Fibonacci Quart.* **33** (3) (1995), 234–239.

- [12] M. Mignotte. ‘Sur les conjugués des nombres de Pisot’. *C. R. Acad. Sci. Paris Sér. I Math.* **298** (2) (1984), 21.
- [13] P. Ribenboim. *My numbers, my friends* (Springer-Verlag, New York, 2000). Popular lectures on number theory.
- [14] C. Sanna. ‘The p -adic valuation of Lucas sequences’. *Fibonacci Quart.* **54** (2) (2016), 118–124.
- [15] O. Strauch and J. T. Tóth. ‘Asymptotic density of $A \subset \mathbf{N}$ and density of the ratio set $R(A)$ ’. *Acta Arith.* **87** (1) (1998), 67–78.
- [16] D. A. Wolfram. ‘Solving generalized Fibonacci recurrences’. *Fibonacci Quart.* **36** (2) (1998), 129–145.
- [17] M. Yabuta. ‘A simple proof of Carmichael’s theorem on primitive divisors’. *Fibonacci Quart.* **39** (5) (2001), 439–443.

CARLO SANNA, Department of Mathematics,
Università degli Studi di Torino, Turin, Italy
e-mail: carlo.sanna.dev@gmail.com