

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Old Yellow Enzyme homologues in *Mucor circinelloides*: Expression profile and biotransformation

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1648851> since 2017-10-09T16:53:07Z

Published version:

DOI:10.1038/s41598-017-12545-7

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Old Yellow Enzyme homologues in *Mucor circinelloides*: expression profile and biotransformation.

Alice Romagnolo^a, Federica Spina^a, Anna Poli^a, Sara Risso^a, Bianca Serito^a, Michele Crotti^b, Daniela Monti^c, Elisabetta Brenna^b, Luisa Lanfranco^a and Giovanna Cristina Varese^{a*}

^a Department of Life Sciences and Systems Biology, University of Turin, viale P. A. Mattioli 25, 10125 Turin, Italy

^b Department of Chemistry, Materials and Chemical Engineering “G. Natta”, Politecnico di Milano, via L. Mancinelli 7, 20131 Milan, Italy

^c Istituto di Chimica del Riconoscimento Molecolare, CNR, Via M. Bianco 9, 20131 Milan, Italy

*Corresponding author (G.C. Varese)

Tel.: +39 0116705984;

Fax: +39 0116705962

E-mail address: cristina.varese@unito.it

Postal address: viale P.A. Mattioli 25, 10125 Turin, Italy

Abstract

The reduction of C=C double bond, a key reaction in organic synthesis, is mostly achieved by traditional chemical methods. Therefore, the search for enzymes capable of performing this reaction is rapidly increasing. Old Yellow Enzymes (OYEs) are flavin-dependent oxidoreductases, initially isolated from *Saccharomyces pastorianus*. In this study, the presence and activation of putative OYE enzymes was investigated in the filamentous fungus *Mucor circinelloides*, which was previously found to mediate C=C reduction. Following an *in silico* approach, using *S. pastorianus* OYE1 aminoacidic sequence as template, ten putative genes were identified in the genome of *M. circinelloides*. A phylogenetic analysis revealed a high homology of McOYE1-9 with OYE1-like proteins while McOYE10 showed similarity with thermophilic-like OYEs. The activation of *mcoyes* was evaluated during the transformation of three different model substrates. Cyclohexenone, α -methylcinnamaldehyde and methyl cinnamate were completely reduced in few hours and the induction of gene expression, assessed by qRT-PCR, was generally fast, suggesting a substrate-dependent activation. Eight genes were activated in the tested conditions suggesting that they may encode for active OYEs. Their expression over time correlated with C=C double bond reduction.

Keywords

α,β -unsaturated compounds, C=C double bond reduction, gene expression, *Mucor circinelloides*, Old Yellow Enzymes.

1. Introduction

The reduction of C=C double bonds is a key reaction in organic chemistry but it is usually carried out by metal catalysts with a strong impact on the technical and economic feasibility of the process [1, 2]. For instance, Yang et al. [3] reported that toxic traces of heavy metals remained in the reaction products and needed to be removed before pharmaceutical use.

Since the major challenges of bulk and fine chemicals synthesis are the reduction of the environmental impact and process costs, biocatalysis became one of the most intriguing alternative to traditional processes. The use of microorganisms or their enzymes has recently found room in the industrial production of pharmaceuticals, flavors, aromas, etc. [2, 4]. The biological reduction of activated C=C double bonds may be carried out by flavin-dependent oxidoreductases, namely ene reductases (ERs), belonging to the Old Yellow Enzyme (OYE) family (EC 1.6.99.1) [5]. They catalyze the asymmetric hydrogenation of C=C double bond conjugated with electron withdrawing groups (EWGs) in the presence of NAD(P)H as cofactor [2]. In contrast with heavy metals, which are capable of mediating *cis*-hydrogenation, OYEs can catalyze this reaction *trans*-fashion with high stereo-selectivity [1].

The reactions catalyzed by OYEs are very interesting and have strong application outcomes. Robinson and Panaccione [6] showed the involvement of OYE homologues involvement in the biosynthetic pathway of ergot alkaloids, commonly used to treat disorders such as Alzheimer's disease, dementia, type 2 diabetes, and hyperprolactinemia or to induce labor and reduce bleeding (lysergic acid-derived drugs). OYE1 from *S. pastorianus* transformed methyl 2-hydroxymethylacrylate in (*R*)-3-hydroxy-2-methylpropanoate, known as "Roche-Ester", which is a chiral building block for the synthesis of vitamins (vitamin E) [4]. Some fragrance compounds (muscone), antibiotics (rapamycin), and natural products have been obtained by OYE-

mediated reduction [2]. The 12-oxophytodienoate reductase enzymes (OPRs, EC 1.3.1.42), OYE homologues from plants, are involved in the biosynthesis of jasmonic acid, which is implicated in the regulation of plant responses to abiotic and biotic stresses as well as plant growth and development [7]. Pentaerythritol tetranitrate reductase (PETNR) from *Enterobacter cloacae* successfully degraded tri nitro toluene (TNT) [8].

OYEs have been ubiquitously described in yeasts, bacteria, animals and plants, and recently in filamentous fungi [9]. Fungi are perfect candidates to set up biocatalysis processes: they combine operative versatility to simple growth conditions and they are a well-known enzymatic machinery [1, 2, 4, 10]. For instance, a homologue of OYE has been discovered in *Aspergillus fumigatus* and *Claviceps purpurea* and associated to the ergot biosynthesis [6, 11]. To date, most of the literature evidences focused on Ascomycetes and Basidiomycetes [9, 12] but the presence of OYE homologue within Zygomycota phylum has never been assessed.

Despite the potential application in several biotechnological fields, microorganisms and enzymes are still scarcely used in manufacturing processes, mostly due to the lack of suitable biocatalysts. Novel enzymatic activities with strong catalytic potential could be achieved with traditional functional screening or advanced molecular approaches [2, 4].

Genome-wide analysis is a useful tool to identify OYEs homologues among the available fungal genomes. For instance, Nizam et al. [9] by analysing 60 Ascomycota and Basidiomycota genomes identified 424 OYEs homologues and provided a first classification of these enzymes within the fungal kingdom. They also explored the evolutionary significance of fungal OYEs. Unfortunately, this data can be considered just a first step, and the actual capability of strains to transform target compounds by reducing C=C double bond need further validation.

In this work, we aimed to fill the lack of information about the occurrence of OYE in fungi belonging to the Zygomycota phylum. *Mucor circinelloides* was selected due to its ability of converting several substrates [13]. Despite those interesting results, the enzymatic pattern responsible for the reactions has never been investigated before. The availability of *M. circinelloides* complete genome sequence (Joint Genome Institute, JGI: <http://jgi.doe.gov>) allowed a genome-mining approach to investigate the presence of putative OYEs homologues.

2. Results

2.1 Identification of putative OYEs in the genome of *M. circinelloides*

In order to identify OYE encoding genes in the filamentous fungus *M. circinelloides*, a BlastP analysis (Basic Local Alignment Search Tool, NCBI, USA) on the complete genome of *M. circinelloides* using *S. pastorianus* OYE1 as query was performed. Ten putative sequences were retrieved and named McOYE1-McOYE10 (Table 1). The 10 amino acid sequences and the amino acid sequence of OYE1 were aligned to evaluate sequence similarities (Table 1). Nine McOYEs showed a similarity with *S. pastorianus* OYE1 of about 40 % while McOYE10 showed a lower similarity (25.33 %; Table 1).

Table 1

Putative OYE homologues of *M. circinelloides* - McOYE1, McOYE2, McOYE3, McOYE4, McOYE5, McOYE6, McOYE7, McOYE8, McOYE9 and McOYE10 - with sequence ID according to JGI database and identity percentage with *S. pastorianus* OYE1.

McOYE	Sequence ID	ID matrix (%) with OYE1	ID matrix (%) with McOYE1	ID matrix (%) with McOYE2	ID matrix (%) with McOYE3	ID matrix (%) with McOYE4	ID matrix (%) with McOYE5	ID matrix (%) with McOYE6	ID matrix (%) with McOYE7	ID matrix (%) with McOYE8	ID matrix (%) with McOYE9	ID matrix (%) with McOYE10
1	160302	44.14	97.50	89.60	70.40	75.00	67.80	66.10	60.60	67.00	58.40	37.00
2	137297	43.99	89.60	97.50	69.70	75.30	68.70	66.10	61.10	67.00	59.50	36.40
3	177510	42.19	71.20	70.50	96.40	76.10	63.30	63.60	57.10	61.60	60.50	40.70
4	155592	41.30	73.40	73.70	73.60	100.00	65.30	65.70	57.80	64.80	60.10	41.60
5	110873	43.41	65.80	66.70	60.40	65.30	100.00	64.50	61.60	65.90	59.10	39.10
6	144573	42.30	64.40	64.30	61.10	64.90	64.60	100.00	67.70	66.10	58.20	38.20
7	153280	41.80	63.70	64.20	62.10	62.20	66.30	68.80	90.50	60.50	56.60	45.10
8	76836	42.19	65.70	65.70	59.50	64.80	65.90	66.10	56.70	100.00	59.20	-

9	134845	38.19	56.90	57.90	57.90	59.50	57.70	57.50	54.90	58.50	97.00	-
10	152500	25.33	35.90	35.20	39.20	41.60	39.10	40.30	48.40	41.60	41.80	100.0

120

121

122 Three conserved domains typical of OYEs were found in all the 10 sequences: the FMN
123 binding site, the active site and the substrate binding site (Supp. Fig. 1).

124 Specific primer pairs were designed on the nucleotide sequences of the 10 putative
125 *mcoyes* (Supp. Table 1) and tested by conventional PCR on genomic DNA. Amplicons
126 of the expected size (about 200 bp) were obtained (Supp. Fig. 2). PCR products were
127 sequenced confirming the specificity of the primers pairs and the authenticity of the
128 DNA sequences.

129 A phylogenetic analysis was performed by implementing sequence data analyzed by
130 Nizam et al. [9], who divided OYEs proteins into three groups: Class I, Class II and
131 Class III. Nine out of 10 McOYEs clustered together in Class I showing a specie-
132 specific clade whereas McOYE10 was located within Class II (Fig. 1).

133 2.2 Biotransformation of conventional substrates and gene expression

134 The expression profile of the 10 putative OYEs homologues was monitored on the RNA
135 extracted from the mycelium grown in liquid culture during the biotransformation of
136 three conventional substrates presenting different EWGs. For each substrate, data on
137 biotransformation and gene expression pattern are presented. Since the substrate was
138 dissolved in dimethyl sulfoxide (DMSO), *mcoyes* activation was also evaluated in the
139 presence of this solvent to exclude artifacts. None of the genes was activated in the
140 presence of DMSO (data not shown). The 10 genes showed a basal activity in the
141 absence of substrates (data not shown).

142 2.2.1 Cyclohexenone (CE)

143 *M. circinelloides* completely reduced the substrate CE into cyclohexanol within 24 h;
144 the reaction process is well known: first an OYE reduces the C=C double bond of CE

producing cyclohexanone, then the keto group is reduced by an alcohol dehydrogenase (ADH) into cyclohexanol (Fig. 2 A) [13]. As shown in Fig. 3 A, the reaction began 30 min after the addition of CE to the medium and at 3.5 h the C=C double bond was completely reduced producing cyclohexanone which was continuously converted in its corresponding alcohol, cyclohexanol.

The transcripts level of the 10 *mcoye* homologues was monitored both in the presence and absence of CE at 30 min, 1 h, 2 h and 5 h (Fig. 4 A). With the exception of *mcoye7* and *mcoye8* that did not show activation upon CE exposure (data not shown), all the other genes were activated within the first two hours. In particular *mcoye2*, *mcoye1* and *mcoye10* displayed a fast and strong induction of gene expression: 730, 111 and 76 fold compared to the control sample without CE at 1 h and at 30 min for *mcoye10* (Fig. 4 A). *Mcoye4* and *mcoye5* showed an activation of 30-50 fold, while for *mcoye3*, *mcoye6* and *mcoye9* the induction of gene expression compared to the control sample remained below 20 fold at the different time points. Noteworthy, expression levels of these genes decreased to the control values after 5 h.

A clear relation between *mcoyes* expression profile and the biotransformation of CE was observed (Fig. 3 A). *Mcoye2* transcript levels were strongly induced at the beginning of the reductive process reaching the maximum at 1 h when 20 % of CE had been converted into cyclohexanone.

2.2.2 α -Methylcinnamaldehyde (MCA)

M. circinelloides completely reduced the C=C double bond of MCA within 20 h. Both the C=C double bond (OYE) and the aldehydic group (ADH) of the substrate were reduced (Fig. 3 B). One hour after MCA addition, α -methylcinnamyl alcohol represented 50 % of the substrates in the culture medium. The concentration of this first product increased until 8 h and then dropped down before 20 h. The production rate of

170 the saturated alcohol was constant, starting from 2 h until 20 h, when it was the
 171 remaining metabolite detected (Fig. 3 B).
 172 According to literature data [2], OYEs are able to catalyze the reduction of the C=C
 173 double bonds of unsaturated aldehydes, whereas they are usually inactive on allylic
 174 alcohols. In the case of *M. circinelloides*, α -methylcinnamyl alcohol seemed to be the
 175 intermediate of the conversion of MCA into the corresponding saturated alcohol. Thus,
 176 α -methylcinnamyl alcohol was added directly to *M. circinelloides* cultures, and indeed
 177 its conversion into α -methyldihydrocinnamyl alcohol was observed to be complete after
 178 48 h. In order to elucidate this reduction pathway, the dideuterated α -methylcinnamyl
 179 alcohol, showing two deuterium atoms linked to the carbon atom bearing the OH group,
 180 was prepared (Figure 2 B). This compound was submitted to bioreduction with *M.*
 181 *circinelloides* and a monodeuterated saturated alcohol was recovered. The formation of
 182 this compound could be explained only admitting the formation of the unsaturated
 183 aldehyde as an intermediate, because the two deuterium atoms should have been
 184 preserved in the direct reduction of the starting allylic alcohol. The alcohol
 185 dehydrogenases, which are present in the fermentation medium, catalyse the oxidation
 186 of the allylic alcohol to the unsaturated aldehyde, which is easily reduced by ERs and
 187 removed from the equilibrium. Then, the saturated aldehyde is further reduced by
 188 ADHs to afford the corresponding saturated alcohol. The intermediate aldehydes did not
 189 accumulate in the reaction medium and it was not possible to detect them during the
 190 reaction course by GC/MS analysis. On the basis of these results, the reaction sequence
 191 shown in Figure 2 C can be hypothesized for MCA.
 192 As for CE, the transcripts level of the 10 putative *mcoyes* was monitored both in the
 193 presence and absence of MCA at 30 min, 1 h, 2 h and 4 h (Fig. 4 B). *Mcoye2* showed
 194 the highest gene activation level with about 2,880 fold compared to the control sample,
 195 followed by *mcoye1* that displayed 1,860 fold induction (Fig. 4 B). An induction of

196 gene expression of about 500 fold was observed for *mcoye4*, *mcoye5* and *mcoye10*.
197 Remarkably, for these 5 genes the highest activation was reached 1 h after the addition
198 of the substrate. For *mcoye3* and *mcoye6* the highest induction levels (138 and 243,
199 respectively) were observed at 2 h. *Mcoye9* displayed moderate gene activation (about
200 14 fold) only at 4 h, *mcoye7* and *mcoye8* did not show activation, as it was observed
201 with CE (data not shown).

202 In this case too, a relation between *mcoyes* gene expression activation and MCA
203 biotransformation was observed (Fig. 3 B). The strongest activation of *mcoye1* in
204 presence of MCA was reached at 1 h, well before the beginning of the C=C reduction,
205 represented by the formation of the saturated alcohol.

206 2.2.3 Methyl cinnamate (MCI)

207 The MCI substrate was completely reduced by *M. circinelloides* within 66 h; both the
208 C=C double bond and the ester group were reduced producing cinnamyl alcohol and
209 phenylpropanol. The exact reaction profile is unknown; however, the one reported in
210 Figure 2 D can be hypothesized on the basis of what has been observed for MCA,
211 starting from the enzymatic hydrolysis of the ester moiety followed by the biocatalysed
212 reduction of the COOH group to primary alcohol. As shown in Fig. 3 C, MCI decreased
213 slightly but constantly until 66 h, when all the substrate was transformed. The detected
214 amount of cinnamyl alcohol was never more than 20 %; also the level of
215 phenylpropanol remained low (< 20 %) until 9 h, after which the concentration reached
216 100 % within 66 h.

217 The transcripts level of the 10 putative *mcoyes* was analyzed in presence and absence of
218 MCI at 1, 2, 3 and 6 h (Fig. 4 C). *Mcoye1* was the most induced gene (about 300 fold).
219 An activation of about 60 fold was observed for *mcoye2*, *mcoye5*, and *mcoye10*, while
220 *mcoye3*, *mcoye4*, *mcoye6*, and *mcoye9* showed 20 fold induction. *Mcoye1* and *mcoye6*
221 displayed the fastest activation, with the maximum within the first 2 h; after that time

point, their transcripts level rapidly decreased. *Mcoye10* showed a peak between 2 and 3 h (Fig. 4 C). Also in this case, neither *mcoye7* nor *mcoye8* were activated upon substrate exposure (data not shown).

A relation between OYE activation and the biotransformation of MCI was observed (Fig. 3 C). The transcription of *mcoye1* started early, when the substrate was still the only detectable compound in the reaction mixture.

3. Discussion

Hydrogenation of C=C double bonds is an important reaction in several manufacturing processes for the production of bulk and fine chemicals; researchers and industries are moving towards more sustainable approaches as biocatalysis and in recent years, several research groups have focused on the identification of OYEs homologues to be exploited in different processes [1]. In the last few years the attention was given to OYEs from filamentous fungi; nevertheless only few studies report their occurrence and their physiological role in this group of organisms [9, 13].

An *in silico* approach allowed to identify in the genome of the zygomycete fungus *M. circinelloides* 10 gene sequences that shared similarity and conserved domains with known OYEs. The presence of multiple OYE genes appears to be a common feature not only among Ascomycetes and Basidiomycetes [7, 12] but also in Zygomycetes: indeed with a similar approach we found from 4 to 10 putative OYE sequences within some of the completely sequenced genomes (Suppl. Table 2).

A phylogenetic analysis grouped the McOYEs in two classes: nine proteins (McOYE1-9) were placed in Class I, including most of the OYE1-like proteins [5, 9, 12], while McOYE10 clustered with Class II. Genome sequence data allowed to hypothesize that a number of Class I McOYEs are located within the same chromosome; this information may suggest duplication events for some of these genes, as suggested by Corrochano et al. [14]. Class II gathers OYEs originally identified from different thermophilic bacteria

[5, 9]; however, the recent work by Nizam et al. [9, 12] demonstrated that a number of sequences, although not yet characterized, from filamentous fungi (Ascomycota and Basidiomycota) also clusters within Class II. To the best of our knowledge, this is the first report of an OYE homologue from a Zygomycota belonging to this Class. The fungal enzymatic activity was analyzed in the presence of three different substrates while previous works considered only one substrate or a series of compounds belonging to the same chemical class [15]. *M. circinelloides* showed a strong enzymatic activity being able to completely reduce the C=C double bond of the three substrates. CE was converted very fast (3.5 h), followed by MCA (20 h) and MCI (66 h), suggesting an increasing recalcitrance of the molecules. These results are in line with those obtained by Gatti et al. [2], who demonstrated that the carbonyl moiety acts as a strong activator, while the ester group is a weak EWG. Being able to convert compounds with different EWGs, *M. circinelloides* was very versatile; during the biotransformation the EWG influenced only the timing of the reaction; the ester group of MCI was the weakest EWG as the reaction was accomplished in 66 h. The reduction of α,β -unsaturated ketones has been extensively studied using either the whole microorganism or the purified enzymes [5, 15, 16]. Generally CE is a well reduced compound; in fact *M. circinelloides* completely reduced the C=C double bond (100 %) in only 3.5 h. Comparable yields were achieved with other filamentous fungi: a previous study, which examined 28 filamentous fungi for the reduction of three different conventional compounds, showed that CE was the easiest to reduce for almost all the fungi (96.4 %); in particular, 19 fungi completely reduced this molecule [13]. Stueckler et al. [7] reported that purified OYE1 (*S. pastorianus*) reduced 92 % of CE and purified YqjM (OYE from *Bacillus subtilis*) reduced 85 % of CE. The reduction of α -substituted cinnamaldehydes is very important at industrial level [2]. Aldehyde is considered a good EWG and MCA was completely reduced within 20 h;

274 Fardelone et al. [17] obtained comparable yields using a commercial strain of *S.*
275 *cerevisiae* in the biotransformation of cinnamaldehyde derivatives. Other authors
276 reported that MCA is not always an easily reduced compound. For instance, Goretti et
277 al. [18] analyzed different non conventional yeasts in the reduction of MCA and found
278 that only *Kazachstania spenceroum* was able to convert this substrate with a yield of 60
279 %. Romagnolo et al. [13] reported that, among 19 fungi tested, only two, belonging to
280 the *Mucor* genus were able to completely convert the C=C double bond of this
281 substrate.

282 The bioreduction of MCI and its derivatives is not frequently reported in the literature,
283 suggesting a possible recalcitrance of this molecule to OYE-mediated
284 biotransformation. A biotransformation study performed on 7 bacterial, yeast and plant
285 OYEs homologues showed a conversion rate of MCI < 1 % [19]. Therefore, the ability
286 of *M. circinelloides* to completely reduce MCI is remarkable, since unsaturated esters
287 with no other EWG are rarely converted by OYEs.

288 BlastP analysis, using OYE1 of *S. pastorianus* as query, allowed the identification of 10
289 putative genes coding for OYEs, confirmed by PCR amplification and sequencing. The
290 high versatility found in the reduction of different compounds by *M. circinelloides* may
291 depend on its enzymatic pattern and on the possibility to activate distinct genes
292 specifically in the presence of different molecules or in defined environmental
293 conditions. In a recent paper, Nizam et al. [9] performed a genome-wide analysis on
294 available genomes of filamentous fungi: 60 species were investigated leading to the
295 identification of 424 OYE homologues. Surprisingly, some species were shown to
296 possess up to 22 OYEs homologues in their genome, while, in other microorganisms the
297 number of OYEs homologues number was more exiguous: only two homologues are
298 present in *S. cerevisiae*, while there are four in *Shewanella oneidensis* [20, 21].

299 Gene activation upon exposure to CE and MCA was extremely high (i.e. up to 2,900
300 fold for *mcoye2* in presence of MCA) and occurred soon after substrate addition. Nizam
301 et al. [9, 12] monitored the expression profile of 6 OYEs homologues from the
302 *Ascochyta rabiei* in two different conditions reporting an increase of 80 fold in
303 transcript levels during plant infection and a weaker activation during oxidative stress.
304 Among the 10 genes identified in *M. circinelloides*, *mcoye1* and *mcoye2* showed the
305 highest degree of gene activation (70-2,900 fold), followed by *mcoye4*, *mcoye5* and
306 *mcoye10* (20-800 fold). *Mcoye3*, *mcoye6* and *mcoye9* were poorly activated, while
307 transcripts of *mcoye7* and *mcoye8* were never activated in each condition. On the basis
308 of these results it seems reasonable to conclude that 8 out of 10 putative OYEs
309 homologues are rapidly activated in response to the substrates addition.

310 A relation between the biotransformation of each substrate and the expression profile of
311 the eight putative OYEs homologues has been observed. Generally, the transcript levels
312 reached the maximum peak before the beginning of the C=C double bond reduction. For
313 example, during CE analysis, the maximum peak of expression of *mcoye2* was reached
314 after 1 h when 20 % of substrate was reduced.

315 The biological role of these enzymes as well as their cell localization is still an open
316 question. By *in silico* analysis Nizam et al. [9], found that the majority of the OYE
317 homologues were allegedly located in the cytoplasm and in the cytoskeleton, although
318 some of them were associated to other cell compartments such as nucleus, peroxisomes,
319 plasma membrane. Only three OYE seemed to be extracellular. A preliminary
320 experiment carried out on *M. circinelloides* during the biotransformation of CE, showed
321 that ene reductase activity was detected only in presence of cell debris indicating that
322 these enzymes may be intracellular (data not shown); further and deeper experiments
323 are needed to confirm this hypothesis.

Studies are in progress to analyze the secondary and tertiary structure of these enzymes by *in silico* approaches [22]. In order to purify and catalytically characterize McOYEs, efforts will concentrate on the production of the homologues of *M. circinelloides* by heterologous expression systems.

4. Materials and methods

4.1 Fungal strain

Mucor circinelloides 277.49 was obtained from CBS (CBS-KNAW fungal biodiversity centre) and was selected due to its capability of reducing C=C double bonds [13]. The strain is preserved as MUT 44 at the *Mycotheca Universitatis Taurinensis* (MUT), Department of Life Sciences and Systems Biology, University of Turin.

4.2 Chemicals

CE, MCA and MCI were purchased from Sigma-Aldrich. Stock solutions of 500 mM of each substrate were prepared in DMSO (Sigma-Aldrich).

(*E*)-2-methyl-3-phenylprop-2-en-1,1-*d*₂-1-ol (dideuterated α -methylcinnamyl alcohol) was prepared by reduction of ethyl (*E*)-2-methyl-3-phenylacrylate (0.50 g, 2.6 mmol) with DIBAL-D (7.9 mmol, 0.7 M in toluene) in THF. After the usual work-up, the dideuterated compound was obtained (0.41 g, 2.3 mmol, 89 %). ¹H NMR (CDCl₃, 400 MHz): δ = 7.39 - 7.19 (5H, m, aromatic hydrogens), 6.53 (1H, q, *J* = 1.5 Hz, CH=C), 1.91 (3H, d, *J* = 1.5 Hz, CH₃); GC-MS (EI) *t*_R = 14.1 min: *m/z* (%) = 150 (*M*⁺, 92), 107 (68), 91 (100). 2-Methyl-3-phenylpropan-1-*d*-1-ol (monodeuterated α -methyldihydrocinnamyl alcohol) was isolated from the reaction medium and characterized by NMR and GC/MS analysis: ¹H NMR (CDCl₃, 400 MHz): δ = 7.37 – 7.13 (5H, m, aromatic hydrogens), 3.45 (1H, m, *CHDOH*), 2.75 (1H, dd *J* = 13.5 and 6.4 Hz, *CHHPh*), 2.43 (1H, dd *J* = 13.5 and 8.0 Hz, *CHHPh*), 1.97 (1H, m, *CHCH*₃), 0.92 (3H, s, *CH*₃); GC-MS (EI) *t*_R = 12.6 min: *m/z* (%) = 151 (*M*⁺, 10), 133 (23), 118 (27), 91 (100).

4.3 Genome mining and phylogenetic analyses

BlastP analysis was performed on the complete genome of *M. circinelloides* strain 277.49 (Joint Genome Institute, JGI: <http://jgi.doe.gov>) using the sequence of OYE1 of *Saccharomyces pastorianus* (UniProtKB accession no. **Q02899**) as query. Primer pairs for qRT-PCR assays were designed by using Primer 3 (<http://primer3.ut.ee/>) (Supp. Table 1). Total genomic DNA was extracted from the mycelium grown in MEA liquid medium (20 g/l glucose, 20 g/l malt extract, 2 g/l peptone) for 24 h using the CTAB method [23]. Oligonucleotides were tested by conventional PCR on genomic DNA. The PCR mixture included distilled water, PCR buffer (10 X), 1 mM deoxynucleotide triphosphates (dNTPs), 10 mM of each primer, 0.5 U of DNA polymerase (Taq DNA polymerase, Qiagen) and 100 ng of genomic DNA in a total volume of 20 µl. Amplifications were performed using a T100 Thermal Cycler (BIORAD). For the validation of *mcoye1* F-R, *mcoye2* F-R, *mcoye3* F-R, *mcoye5* F-R, *mcoye6* F-R, *mcoye9* F-R e *mcoye10* F-R, the amplification protocol was as follows: 95 °C (5 min), 34 cycles of 95 °C (40 sec), 60 °C (50 sec) and 72 °C (50 sec), 72 °C (8 min). For the detection of *mcoye4* F-R, *mcoye7* F-R e *mcoye8* F-R the amplification protocol was as follows: 95 °C (5 min), 34 cycles 95 °C (40 sec), 56 °C (50 sec) and 72 °C (50 sec), 72 °C (8 min). PCR products were loaded on a 1.5 % agarose electrophoresis gel stained with ethidium bromide; the molecular weight marker used was the GelPilot 1 kb Plus Ladder (cat. no. 239095, Qiagen). Products were purified and sequenced at MacroGen (The Netherlands). Newly generated sequences were analyzed using Sequencher 5.4 (Gene Code Corporation). To perform the phylogenetic analyses, over 400 OYEs aminoacidic sequences of fungi were aligned with MUSCLE (<http://www.ebi.ac.uk/Tools/msa/muscle/>) using default conditions for gap openings and gap extension penalties and trimmed by TrimAl (v 1.2) (<http://trimal.cgenomics.org>) with the AUTOMATED 1 setting. The analysis was

performed using two approaches. First, a phylogenetic tree was derived by Bayesian Inference (BI) implemented in MrBayes (v 3.2.2) (<http://mrbayes.sourceforge.net>) under a mixed amino acid substitution model. The alignment was run over 10 million generations with two independent runs each containing four Markov Chains Monte Carlo (MCMC) and sampling frequency of every 300 iterations. The first 2,500 trees were discarded as “burn-in” (25 %). Using the Sumt function of MrBayes a consensus tree was generated and posterior probabilities were estimated. In a second approach, Maximum Likelihood (ML) was performed using RAxML GUI (v 1.5 b) [24] with WAG+I+G model. Statistical reliability was determined by Bootstrap analysis. All the phylogenetic trees were visualized using FigTree (v 1.4) (<http://tree.bio.ed.ac.uk/software/figtree>).

4.4 Biotransformation by whole cell system

A conidia suspension of *M. circinelloides* was made from pre-growth mycelium in MEA solid medium (same composition of MEA liquid with the addition of 20 g/l of agar). 10⁶ conidia were inoculated in 100 ml flasks containing 40 ml of MEA liquid medium. Flasks were incubated at 25 °C in agitation. After 2 days, substrates were added (5 mM final concentration), each cultural line was run in triplicate. In addition, biotic controls (in absence of substrates) were set up. According to previous results (unpublished data), the conversion of CE, MCA and MCI was followed for 24 h, 48 h and 7 d, respectively. Every 2 h, 1 ml of broth and 100 mg of biomass were collected to perform chemical analysis and RNA extraction, respectively. The mycelium was frozen in liquid nitrogen and stored at - 80 °C until the analysis. At any collection time point, pH and glucose content were measured. The concentration of reducing sugars was obtained following the reaction with 3,5-dinitrosalicylic acid assay (DNS) [25], using a modified protocol as described by Spina et al. [26]. At each

time point and at the end of the experiment, fungal biomasses were separated from the culture medium by filtration and dried in oven at 60 °C for 24 h to calculate the dry weight.

4.5 Chemical analyses

Samples taken at the different time points were extracted by two-phase separation using 0.4 ml of methyl *t*-butyl ether (MTBE) as solvent; the organic phase was dried over anhydrous Na₂SO₄ and analyzed by GC/MS.

GC/MS analyses were performed on an Agilent HP 6890 gas chromatograph equipped with a 5973 mass detector and an HP-5-MS column (30 m × 0.25 mm × 0.25 µm, Agilent), employing the following temperature program: 60 °C (1 min) / 6 °C min⁻¹ / 150 °C (1 min) / 12 °C min⁻¹ / 280 °C (5 min). The end products of the

biotransformations were identified by GC/MS analysis, using authentic commercial

samples as reference compounds: i) cyclohexenone $t_R = 5.40$ min m/z 96 (M^+ , 33), 81 (19), 68 (100); cyclohexanone $t_R = 4.65$ min m/z 98 (M^+ , 47), 83 (13), 55 (100);

cyclohexanol $t_R = 4.45$ min m/z 100 (M^+ , 2), 82 (35), 57 (100), ii) α -

methylcinnamaldehyde $t_R = 14.7$ min m/z 146 (M^+ , 64), 145 (100), 117 (79), 91 (43);

α -methylcinnamyl alcohol $t_R = 15.5$ min m/z 148 (M^+ , 50), 115 (63), 91 (100); α -

methyldihydrocinnamyl alcohol $t_R = 13.7$ min m/z 150 (M^+ , 12), 117 (62), 91 (100); iii)

methyl cinnamate $t_R = 16.03$ min m/z 162 (M^+ , 58), 131 (100), 103 (72); cinnamyl

alcohol $t_R = 12.80$ min m/z 134 (M^+ , 53), 115 (65), 92 (100) ; phenylpropanol $t_R = 12.36$

min m/z 136 (M^+ , 21), 117 (100), 91 (84).

4.6 RNA extraction, first strand cDNA synthesis and quantitative Real-Time PCR experiments

The extraction of RNA was performed from about 100 mg of fungal biomass using the RNeasy Plant Mini Kit (Qiagen). Quantity and quality of RNA samples were checked spectrophotometrically (Tecan Infinite 200, i-control software). After DNase treatment

(TURBO DNA-free, Ambion), RNA quality has been tested again and for all the samples, the ratios of absorbance 260/280 were between 1.8 and 2.2. Subsequently they were processed to obtain cDNA with the use of the Super-Script II Reverse Transcriptase (Invitrogen), following instructions.

qRT-PCR were performed with an iCycler iQTM Real-Time PCR Detection System (BIORAD); reactions were carried out in a final volume of 15 µl by using iTaq Universal SYBR GREEN Supermix (BIORAD), specific primers (3 µM; Table 1) and cDNA . For the detection of *mcoye1*, *mcoye2*, *mcoye3*, *mcoye5*, *mcoye6*, *mcoye9* and *mcoye10*, the amplification protocol was as follows: 95 °C (1.5 min), 40 cycles of 95 °C (15 sec), 60 °C (30 sec) and 72 °C (50 sec), 72 °C (8 min). For the detection of *mcoye4*, *mcoye7* and *mcoye8* the amplification protocol was as follows: 95 °C (1.5 min), 40 cycles 95 °C (15 sec), 56 °C (30 sec) and 72 °C (50 sec), 72 °C (8 min). The *M. circinelloides* β-actin encoding gene was used as internal control [27]. The relative expression was calculated using the $2^{-\Delta\Delta C_t}$ method [28]. One-way ANOVA and Tukey's tests ($p < 0.05$) were performed to assess the statistical significance of the gene expression data (IBM SPSS Statistics for Macintosh, Version 22.0).

4.7 Availability of materials and data

Authors confirm that all relevant data are included in the article and its supplementary information file.

References

452 [1] Stuermer, R., Hauer, B., Hall, M. & Faber, K. Asymmetric bioreduction of activated
 453 C=C bonds using enoate reductases from the old yellow enzyme family. *Curr. Opin.*
 454 *Chem. Biol.* 11, 203–213 (2007).

455 [2] Gatti, F. G., Parmeggiani, F. & Sacchetti, A. Synthetic strategies based on C=C
 456 bioreductions for the preparation of biologically active molecules. In: *Synthetic*
 457 *methods for biologically active molecules: exploiting the potential of bioreductions*, 1st
 458 ed., Wiley WHC, Weinheim; 2013.

459 [3] Yang, J. W., Fonseca, M. T. H., Vignola, N. & List, B. Metal-free, organocatalytic
 460 asymmetric transfer hydrogenation of α,β -unsaturated aldehydes. *Angew. Chem. Intern.*
 461 *Ed.* 44, 108–110 (2005).

462 [4] Soetaert, W. & Vandamme, E. J. The scope and impact of industrial biotechnology.
 463 In: *Industrial biotechnology: sustainable growth and economic success*, 1st ed. Wiley-
 464 VHC, Weinheim; 2010.

465 [5] Toogood, H. S., Gardiner, J. M. & Scrutton, N. S. Biocatalytic reductions and
 466 chemical versatility of the old yellow enzyme family of flavoprotein oxidoreductases.
 467 *ChemCatChem* 2, 892-914 (2010).

468 [6] Robinson, S. L. & Panaccione, D. G. Diversification in ergot alkaloids in natural and
 469 modified fungi. *Toxins* 7, 201-218 (2015).

470 [7] Stueckler, C., Reiter, T. C., Baudendistel, N. & Faber, K. Nicotinamide-independent
 471 asymmetric bioreduction of C=C bonds via disproportionation of enones catalyzed by
 472 enoate reductases. *Tetrahedron* 66, 663-667 (2010).

473 [8] Huang, M., Hu, H., Ma, L., Zhou, Q., Yu, L. & Zheng, S. Carbon-carbon double-
 474 bond reductases in nature. *Drug. Metab. Rev.* 46, 362-378 (2015).

475 [9] Nizam, S., Verma, S., Borah, N. N., Gazara, R. K. & Verma, P. K. (a)
 476 Comprehensive genome-wide analysis reveals different classes of enigmatic old yellow
 477 enzyme in fungi. *Sci. Rep.* 4, 4013 (2014).

478 [10] Brandl, J. & Andersen, M. R. Current state of genome-scale modelling in
 479 filamentous fungi. *Biotechnol. Lett.* 37, 1131-1139 (2015).

480 [11] Cheng, J. Z., Coyle, C. M., Panaccione, D. G. & O'Connor, S. E. A role for Old
 481 Yellow Enzyme in ergot alkaloids biosynthesis. *J. Am. Chem. Soc.* 132, 1776–1777
 482 (2010).

483 [12] Nizam, S., Gazara, R. K., Verma, S., Singh, K. & Verma, P. K. Comparative
 484 structural modeling of six Old Yellow Enzymes (OYEs) from the necrotrophic fungus
 485 *Ascochyta rabiei*: insight into novel OYE Classes with differences in cofactor binding,
 486 organization of active site residues and stereopreferences. *Plos One* 9 (2014).

487 [13] Romagnolo, A., Spina, F., Brenna, E., Crotti, M., Parmeggiani, F. & Varese, G. C.
 488 Identification of fungal ene-reductase activity by means of a functional screening. *Fung.*
 489 *Biol.* 119, 487-493 (2015).

490 [14] Corrochano, L. M., Kuo, A., Marcel-Houben, M., Polaino, S., Salamov, A., et al.
 491 Expansion of signal transduction pathways in fungal by extensive genome duplication.
 492 *Curr. boil.* 26, 1577-1584 (2016).

493 [15] Zagozda, M. & Plenkiewicz, J. Enantioselective reduction of α,β -unsaturated
 494 ketones by *Geotrichum candidum*, *Mortierella isabellina* and *Rhodotorula rubra* yeast.
 495 *Tetrahedron: Asymm.* 17, 1958-1962 (2006).

496 [16] Hall, M., Stueckler, C., Hauer, C., Stuermer, R., Friedrich, T., Breuer, M., Kroutil,
 497 W. & Faber, K. Asymmetric bioreduction of activated C=C bonds using *Zymomonas*
 498 *mobilis* NCR enoate reductase and old yellow enzymes OYE 1-3 from yeasts. *Eur. J.*
 499 *Org. Chem.* 1511-1516 (2008).

500 [17] Fardelone, L. C., Rodrigues, A. R. & Moran, P. J. S. Baker's yeast mediated
 501 asymmetric reduction of cinnamaldehyde derivatives. *J. Mol. Catal. B: Enz.* 29, 41-45
 502 (2003).

503 [18] Goretti, M., Ponzoni, C., Caselli, E., Marchegiani, E., Cramarossa, M. R.,
504 Turchetti, B., Forti, L. & Buzzini, P. Bioreduction of α,β -unsaturated ketones and
505 aldehydes by non-conventional yeast (NCY) whole-cell. *Biores. Technol.* 102, 3993-
506 3998 (2011).

507 [19] Tasnadi, G., Winkler, C. K., Clay, D., Sultana, N., Fabian, W. M. F., Hall, M.,
508 Ditrich, K. & Faber, K. A substrate-driven approach to determine reactivities of α,β -
509 unsaturated carboxylic esters towards asymmetric bioreduction. *Chem. Eur. J.* 18,
510 10362-10367 (2012).

511 [20] Niino, Y. S., Chakraborty, S., Brown, B. J. & Massey, V. A new Old Yellow
512 Enzyme of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 27, 1983-1991 (1995).

513 [21] Brigé, A., Van Den Hemel, D., Carpentier, W., De Smet, L. & Van Beeumen, J. J.
514 Comparative characterization and expression analysis of the four Old Yellow Enzyme
515 homologues from *Shewanella oneidensis* indicate differences in physiological function.
516 *Biochem. J.* 394, 335-344 (2006).

517 [22] Catucci, G., Romagnolo, A., Spina, F., Varese, G. C., Gilardi, G. & Di Nardo, G.
518 Enzyme-substrate matching in biocatalysis: *in silico* studies to predict substrate
519 preference of ten putative ene-reductases from *Mucor circinelloides* MUT 44. *J. Mol.*
520 *Catal. B: Enz.* 131, 94-100 (2016).

521 [23] Gardes, M. & Bruns, T. D. ITS Primer with Enhanced Specificity for
522 Basidiomycetes-Application to the Identification of Mycorrhizae and Rusts. *Mol. Ecol.*
523 2, 113-118 (1993).

524 [24] Stamatakis, A. RAxML Version 8: A tool for Phylogenetic Analysis and Post-
525 Analysis of Large Phylogenies. *Bioinformatics* 30, 1312-1313 (2014).

526 [25] Miller, G. L. Use of DNS reagent for the measurement of reducing sugar. *Anal.*
527 *Chem.* 31, 426-428 (1959).

- [26] Spina, F., Fidaleo, M., Nanni, A., Romagnolo, A. & Varese, G. C. Fungal Laccases production using tomato-based medium: a factorial design approach. *Environ. Eng. Manag. J.* 14, 1743-1750 (2015).
- [27] Lee, S. C., Li, A., Calo, S. & Heitman, J. Calcineurin plays key roles in the dimorphic transition and virulence of the human pathogenic zygomycete *Mucor circinelloides*. *Plos. Path.* 9, 1-20 (2013).
- [28] Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using realtime quantitative PCR and the $2^{-\Delta\Delta C_t}$ method. *Methods* 25, 402–408 (2001).

Author Contribution Statement


A.R., F.S.: wrote the manuscript
A.P.: performed phylogenetic analysis
A.R., F.S., S.R., B.S.: performed lab experiments
L.L.: gene expression experimental design
M.C., D.M., E.B.: performed chemical analysis and data curation
G.C.V: project administration and supervision
All authors reviewed the manuscript.

Acknowledgements

This project was partially supported by Fondazione CRT (Torino, Italy). The authors thank Fondazione San Paolo (Turin, Italy) for the economic support of the PhD program of Alice Romagnolo and Simone Belmondo for the help in the first phases of gene expression analyses.
The authors declare no competing financial interests.

554

555 **Figure and figure legends**



556

557 **Figure 1.** Evolutionary relationship of deduced OYE proteins based on Bayesian
558 inference analysis of the structure-based amino acid sequence alignment. The numbers
559 at the nodes indicates Bayesian posterior probabilities. The phylogenetic tree was
560 implemented from Nizam et al., 2014 [9].




Figure 2. Reaction profiles of (A) CE, (B, C) MCA and (D) MCI biotransformations.






Figure 3. Graph combining the data of expression of (A) *mcoy2* in presence of CE (bars) with the biotransformation data of CE (lines); (B) *mcoy1* in presence of MCA (bars) with the biotransformation data of MCA (lines); (C) *mcoy1* in presence of MCI

567 (bars) with the biotransformation data of MCI (lines). Data are the averages \pm standard
568 deviations (error bars) of the results of at least three different biological replicates.


A



B



C



570 **Figure 4.** Gene expression of OYE homologues in presence of (A) CE, (B) MCA and
571 (C) MCI during the time course experiments. The relative gene expression was
572 calculated with the $2^{-\Delta\Delta C_t}$ method according to Livak & Schmittgen [27] using the β -
573 actin as housekeeping gene [26] and the control (non treated) as reference sample.
574 Different letters indicate statistically significant difference ($p < 0.05$, ANOVA and
575 Tukey's tests) for each gene at the different time points.