

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Evaluation of the use of sulfur dioxide and glutathione to prevent oxidative degradation of malvidin-3-monoglucoside by hydrogen peroxide in the model solution and real wine

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1651784> since 2017-11-14T15:07:48Z

Published version:

DOI:10.1016/j.foodres.2017.06.010

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin.

Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

FOOD RESEARCH INTERNATIONAL, 99 (Pt 1), 2017, pp: 454-460

DOI: 10.1016/j.foodres.2017.06.010

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>):

<https://doi.org/10.1016/j.foodres.2017.06.010>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1651784>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

iris-AperTO

Manuscript Number: FOODRES-D-17-00823R1

Title: EVALUATION OF THE USE OF SULFUR DIOXIDE AND GLUTATHIONE TO PREVENT OXIDATIVE DEGRADATION OF MALVIDIN-3-MONOGLUCOSIDE BY HYDROGEN PEROXIDE IN THE MODEL SOLUTION AND REAL WINE.

Article Type: Research Paper

Keywords: anthocyanins;
wine pigments;
oxidation;
sulfur dioxide;
glutathione.

Corresponding Author: Dr. angelita gambuti,

Corresponding Author's Institution: University of Napoli Federico II

First Author: angelita gambuti

Order of Authors: angelita gambuti; Luigi Picariello; Luca Rolle; Luigi Moio

Abstract: In this study the oxidative degradation by hydrogen peroxide of native grape anthocyanin was studied in model solutions and in red wines added with increasing concentration of sulfur dioxide and glutathione (GSH). The presence of hydrogen peroxide and metal ions in traces allowed to investigate the possibility to use GSH to prevent Fenton reaction in wine conditions. Two different pH of wine were considered: 3.20 and 3.80. The protective effect of sulfur dioxide on malvidin 3-monoglucoside degradation was higher at lower pH in model solution. No effect of pH on sulfur dioxide action towards the native anthocyanin in real wine was detected. Surprisingly GSH determined an increase in the degradation of malvidin 3-monoglucoside regardless of pH. Therefore, GSH is not effective in prevent native anthocyanins loss due to the Fenton reaction during red wine aging.

UNIVERSITA' DEGLI STUDI DI NAPOLI " FEDERICO II"

Prof. Angelita Gambuti
Dipartimento di Agraria
via Università n°100 - 80055 – Portici (Napoli), ITALIA

TeleFax: +39 0825 1913305
e-mail: angelita.gambuti@unina.it

April, 3th 2017

Dear Editor,

I would like to submit for publication this paper about **“EVALUATION OF THE USE OF SULFUR DIOXIDE AND GLUTATHIONE TO PREVENT OXIDATIVE DEGRADATION OF MALVIDIN-3-MONOGLUCOSIDE BY HYDROGEN PEROXIDE IN MODEL SOLUTION AND REAL WINE.”**

The originality of the work lies into the fact that, for the first time, the role of glutathione as an alternative to sulfur dioxide to prevent oxidative loss of native pigments of red wine in strong oxidative conditions was evaluated.

In last decades concerns over the ability of sulfur dioxide to induce severe allergic reactions have created a great need for its reduction or replacement in wine and glutathione has been proposed as an alternative antioxidant. However only its ability to prevent the loss of some aroma compounds has been showed and its effect on other important compounds such as pigments of wine is not clear.


Data obtained in this study showed that, in presence of hydrogen peroxide, glutathione accelerates pigments degradation. These results could have important implications into the use of glutathione to produce red wines with reduced content of sulfur dioxide. This may help in regulating the use of this compound as additive in wine production.

Instructions for authors have been carefully followed.

I hope that our editing satisfies the standards required and that the manuscript can be submitted to the attention of the referees.

I keep waiting for your answer.

Yours sincerely


Prof. Angelita Gambuti

RESPONSE TO FIRST REVIEW - LIST OF CHANGES

Dear Anderson de Souza Sant'Ana,

in response to each of the comments mentioned in the letter received on Mon, 10 Apr 2017, we changed the manuscript as follow (answers as colored text).

Sincerely yours,

Angelita Gambuti

a) Use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point).

We named the file Highlights. Each research highlight was represented with bullet points and was maximum of 85 characters, including spaces, per bullet point.

b) Define better the aims of the MS at the end of INTRODUCTION section

We better defined the aim of the MS (lines 75-87).

c) Take care of FRI style in guide for authors. For example:

21. Gambuti, A., Han, G., Peterson, A. L., & Waterhouse, A. L. (2015). Sulfur dioxide and glutathione alter the outcome of microoxygenation. American Journal of Enology and Viticulture,ajev-2015.

Should be: 21. Gambuti, A., Han, G., Peterson, A. L., & Waterhouse, A. L. (2015). Sulfur dioxide and glutathione alter the outcome of microoxygenation. American Journal of Enology and Viticulture, 66, 411-423.

We considered FRI style: lines 334-335, 387-389, 419-420, 460-463.

*Highlights (for review)

- The possibility to use GSH to prevent Fenton reaction in wine has been evaluated.
- GSH determined an increase in the degradation of malvidin 3-monoglucoside.
- The preventive action of SO₂ in red wine does not depends on pH.
- GSH is not effective in prevent anthocyanins loss during red wine aging.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

EVALUATION OF THE USE OF SULFUR DIOXIDE AND GLUTATHIONE TO PREVENT
OXIDATIVE DEGRADATION OF MALVIDIN-3-MONOGLUCOSIDE BY HYDROGEN
PEROXIDE IN THE MODEL SOLUTION AND REAL WINE.

*Gambuti Angelita**^a, *Picariello Luigi*^a, *Rolle Luca*^b, *Moio Luigi*^a

^aDipartimento di Agraria - Università degli Studi di Napoli Federico II - Sezione di “Scienze della Vigna e del Vino” SVV. Viale Italia (angolo via Perrottelli), 83100, Avellino.

^bUniversità di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy.

* Corresponding author: angelita.gambuti@unina.it, Tel.: +3908251913305; Fax: +390825784678.

Running title: Glutathione and Sulfur Dioxide affect the degradation of wine native pigments by hydrogen peroxide.

ABSTRACT

In this study the oxidative degradation by hydrogen peroxide of native grape anthocyanin was studied in model solutions and in red wines added with increasing concentration of sulfur dioxide and glutathione (GSH). The presence of hydrogen peroxide and metal ions in traces allowed to investigate the possibility to use GSH to prevent Fenton reaction in wine conditions. Two different pH of wine were considered: 3.20 and 3.80. The protective effect of sulfur dioxide on malvidin 3-monoglucoside degradation was higher at lower pH in model solution. No effect of pH on sulfur dioxide action towards the native anthocyanin in real wine was detected. Surprisingly GSH determined an increase in the degradation of malvidin 3-monoglucoside regardless of pH. Therefore, GSH is not effective in prevent native anthocyanins loss due to the Fenton reaction during red wine aging.

Keywords: anthocyanins, wine pigments, oxidation, sulfur dioxide, glutathione.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

During production and aging of red wine a change of wine color is observed due to the involvement of wine pigments, the anthocyanins, in numerous reactions with other compounds present in solution and with compounds deriving from oxygen action. The shift of wine color from red to yellow hue is an indication of aging but also of oxidative spoilage of red wine and it is essentially due to the action of oxygen determining a loss of native anthocyanins not balanced by the formation of more stable red pigments. The chemical oxidation of wine is triggered by the oxidation of polyphenols to quinones while oxygen is reduced to hydrogen peroxide in presence of trace metals such as iron and copper (Danilewicz, 2011). Wine polyphenols involved in this starting phase contain at least two vicinal hydroxyls. Because wine native red pigments, such as malvidin 3-monoglucoside, contain isolated phenolic hydroxyl groups and need higher potential to be oxidized (Kilmartin, Zou & Waterhouse, 2001), they are not involved in the first steps of wine oxidation. Their loss is due to the action of hydrogen peroxide H₂O₂ that, in the second step of wine oxidation, in presence of ferrous or cuprous species, reacts in the Fenton reaction to give the destructive oxidant radicals (Elias, Andersen, Skibsted & Waterhouse, 2009). A direct reaction of malvidin 3-monoglucoside with H₂O₂ has been reported (Sondheimer & Kertesz, 1952; Ozkan, Yemencioğlu, Asefi & Cemeroglu, 2002) but the main reason of their loss is linked to their involvement in complex reactions triggered by reactive carbonyls produced by the action of radicals produced by Fenton reaction such as acetaldehyde from oxidation of ethanol and glyoxylic acid from oxidation of tartaric acid (Es-Safi, Fulcrand, Cheynier & Moutounet, 1999; Es Safi, Cheynier & Moutounet, 2003; He et al., 2012).

Oxidation is a long-standing problem in wine industry and sulfur dioxide SO₂ is the generally used chemical to control it. SO₂ acts as antioxidant in three ways, scavenging hydrogen peroxide, reacting with ortho-quinones acting as sacrificial nucleophiles and binding carbonyl compounds produced by Fenton reaction (Adachi et al., 1979; Danilewicz & Wallbridge, 2010). However, concerns over its ability to induce severe allergic reactions have created a great need for its reduction or replacement and, as a consequence, regulatory restrictions has been established by World Health Organization (WHO) and International Organization of Vine and Wine (OIV). Moreover, its excessive use in winemaking can determine a distinctive irritating odor in wine.

In last decades, the tripeptide glutathione (GSH) has been proposed in winemaking as alternative antioxidant to decrease the use of SO₂ (Kritzinger, Bauer & Du Toit, 2012). This is

1
2
3 why it has been recently authorized by OIV (International Organization of Vine and Wine) in
4 must (maximum up to 20 mg/L) but it is still not admitted by EC as wine additive. Low
5 concentration (20 mg/L) of GSH protected against loss of esters, terpenes (Roussis,
6 Lambropoulos & Tzimas, 2007) and volatile thiols during bottle storage (Ugliano et al.,
7 2011). At higher concentrations (180 mg/L) it delayed the oxidative browning limiting the
8 formation of yellow xanthylum cation pigments in white wines (Roussis, Lambropoulos &
9 Tzimas, 2007; Sonni, Clark, Prenzler, Riponi, & Scollary, 2011; Bouzanquet, Barril, Clark,
10 Dias & Scollary, 2012;). Recently a moderate protective effect of GSH (30 mg/L) on native
11 anthocyanins during micro-oxygenation (MOx) has been showed (Gambutu, Han, Peterson &
12 Waterhouse, 2015). Its anti-oxidant activity in wine is mainly due to the abilities to reduce
13 back *o*-quinone compounds (Cheynier & Van Hulst, 1988; Nikolantonaky & Waterhouse,
14 2012) and it can also bind wine reactive aldehydes (Sonni, et al., 2011).

15
16
17 GSH also prevent cellular damage owing to its hydrogen peroxide scavenging activity
18 (Winterbourn & Metodiewa, 1999) but the same activity in wine has not been evaluated.
19 Among wine features, pH is one of main wine parameter influencing wine oxidation because
20 it affects the formation of new stable polymeric pigments from native anthocyanins
21 (Kountoudakis et al., 2011; Pechamat, Zeng, Jourdes, Ghidossi, & Teissedre, 2014;) and SO₂
22 forms in hydroalcoholic solution. However, no information on the effect pH on the
23 antioxidant activity of GSH in wine conditions has been reported.

24
25
26 The aim of the present study was to investigate the potentiality of GSH as an alternative to
27 SO₂ to scavenge hydrogen peroxide, to inhibit Fenton reaction and to prevent grape native
28 anthocyanins loss during wine aging. With this purpose the oxidative degradation of malvidin
29 3-monoglucoside was studied in model solutions and in red wines treated with hydrogen
30 peroxide and added with increasing concentration of sulfur dioxide and glutathione. The
31 experiments were carried out at two different wine pH: 3.20 and 3.80.

32
33
34 In addition, GSH prevent cellular damage owing to its hydrogen peroxide scavenging activity
35 (Winterbourn & Metodiewa, 1999) but the same activity in wine has not been evaluated.

36
37
38 In this study, to better understand if GSH can be proposed as an alternative to SO₂ to scavenge
39 hydrogen peroxide, inhibit Fenton reaction and prevent grape native anthocyanins loss during
40 wine aging, the oxidative degradation of malvidin 3 monoglucoside was studied in model
41 solutions and in red wines treated with hydrogen peroxide and added with increasing
42 concentration of sulfur dioxide and glutathione.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

~~Moreover, because pH is one of main wine feature affecting wine oxidation owing to its role on the formation of new stable polymeric pigments from native anthocyanins (Kountoudakis et al., 2011; Pechamat, Zeng, Jourdes, Ghidossi, & Teissedre, 2014;) and on SO₂ forms in hydroalcoholic solution, the experiments were carried out at two different wine pH: 3.20 and 3.80.~~

2. MATERIALS AND METHODS

2.1 Experimental trial. Oxidation reactions were performed in 10 mL reagent bottles. The bottles were purged with nitrogen and placed in darkness at 20 °C. In the first experiment the effect of antioxidants and pH was evaluated in model solutions. All solutions contained malvidin 3-monoglucoside 100 mg/L (203 mM), ethanol (12% v/V) and tartaric acid (q. s.). The pH was adjusted adding NaOH. The oxidation Ox was performed adding hydrogen peroxide at a concentration of 39.2 mg/L of O₂ eq (1.225 mM) to trigger Fenton reaction (Elias and Waterhouse, 2010). Because this reaction involves hydrogen peroxide and metal ions at concentration (< 0.2 µM) much lower than expected in all commercial water supplies (Clark, Prenzler & Scollary, 2007) the occurrence of Fenton reaction in our experimental conditions is guaranteed. Six oxidized (Ox) samples were obtained: Ox, adding only hydrogen peroxide; Ox+SO₂ low, adding hydrogen peroxide and 37.4 mg/L of SO₂ (0.584 mM); Ox+SO₂ high, adding hydrogen peroxide and 202 mg/L of SO₂ (3.16 mM); Ox+GSH, adding hydrogen peroxide and 30 mg/L of glutathione (0.098 mM); Ox+SO₂ low+GSH, adding hydrogen peroxide, 37.4 mg/L of SO₂ (0.584 mM) and 30 mg/L of glutathione (0.098 mM); Ox+SO₂high+GSH, adding hydrogen peroxide, 202 mg/L of SO₂ (3.16 mM) and glutathione (0.098 mM). Samples were prepared at two pH: 3.20 and 3.80 and monitored after 0, 16 and 72 hours of incubation at 20 °C. Dilution coefficient were considered to compare treated sample with control one. In the second experiment the same treatments were performed on a red wine produced in 2013 (*Vitis vinifera* L. Casavecchia). The base parameters were: ethanol content 13.60 ± 0.07 % v/V, pH 3.80 ± 0.03, residual sugars 1.72 ± 0.06 g/L. The pH of wine was adjusted to pH 3.20 by adding tartaric acid. All samples (model solutions and wines) were prepared in duplicate. On each replicate two analyses were performed to have a datum from the mean of four values.

2.2 Reagents and standards. Solvents of HPLC-gradient grade and all other chemicals of analytical reagent grade were purchased from Sigma (Milan, Italy). The solutions were prepared in deionized water produced by a Purelab Classic system (Elga Labwater, Marlow,

1
2
129 UK). About standards for calibration curves, syringic acid was purchase from Sigma-Aldrich
130 (Milan, Italy) whereas malvidin-3-glucoside chloride was supplied by Extrasynthèse (Genay,
131 France). For identification purposes, anthocyanin standards (delphinidin-3-glucoside chloride,
132 malvidin-3-glucoside chloride, petunidin chloride, peonidin-3-glucoside chloride, and
133 cyanidin-3-glucoside chloride) were purchased from Extrasynthèse.

10
134 **2.3 Methods.** HPLC separation of anthocyanins and for the determination of syringic acid were
135 carried out according to the OIV Compendium of International Methods of Analysis of Wine
136 and Musts (2017). Analyses were performed in a HPLC Shimadzu LC10 ADVP apparatus
137 (Shimadzu Italy, Milan), consisting of a SCL-10AVP system controller, two LC-10ADVP
138 pumps, a SPD-M 10 AVP detector, and an injection system full Rheodyne model 7725
139 (Rheodyne, Cotati, CA) equipped with a 50 μ L loop. A Waters Spherisorb column (250 x 4.6
140 mm, 4 μ m particles diameter) with pre-column was used. Twenty μ L of wine or calibration
141 standards were injected onto the column. Detection was performed by monitoring the
142 absorbance signals at 518 nm. All the samples were filtered through 0.45 μ m, Durapore
143 membrane filters (Sigma Aldrich, Milan, Italy) into glass vials and immediately injected into
144 the HPLC system. The HPLC solvents were: solvent A: water/formic acid/acetonitrile
145 (87:10:3) v/v; solvent B: water/formic acid/acetonitrile (40:10:50) v/v. The gradient used was:
146 zero-time conditions 94 % A and 6 % B, after 15 min the pumps were adjusted to 70 % A and
147 30 % B, at 30 min to 50 % A and 50 % B, at 35 min to 40 % A and 60 % B, at 41 min, end of
148 analysis, to 94 % A and 6 % B. After 10-min equilibrium period the next sample was injected.
149 The flow rate was 0.80 mL/min. For calibration the external standard method was used: the
150 calibration curve was plotted for the malvidin-3-monoglucoside (Extrasynthese, Lyon,
151 France) on the basis of peak area and the concentration was expressed as mg/L of malvidin-3-
152 monoglucoside. The calibration curve for the identification and determination of syringic acid
153 was prepared starting from a stock solution containing 5 mg/L of syringic acid (Sigma-
154 Aldrich, Milan, Italy). All the analyses were made in duplicate on each experimental replicate.

155 **2.4 Statistical Analyses.** Quantitative data of the wines were compared using Fisher's least
156 significant differences (LSDs) procedure. Multifactorial ANOVA with third-order interactions
157 was used to evaluate the relationships among factors. Differences of $p < 0.05$ were considered
158 significant. These analyses were performed using XLSTAT (Addinsoft, XLSTAT Version
159 2013.6.04). All data are means of four values (2 experimental replicates X 2 analytical
160 replicates).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. RESULTS AND DISCUSSION

3.1 Model solution experiments

In aqueous solution anthocyanins exist in different forms in equilibrium depending on the pH: the reddish-pink flavylium salt predominates at lower pH values while the carbinol pseudo-base and chalcone are the main species at pH higher than 7 (Brouillard & Dubois, 1977). Concerning the range of pH typical of red wines it has been reported that at pH of 3.4–3.6, 20–25% of anthocyanins are in the colored flavylium forms, whereas at pH of 4.0, only 10% of anthocyanins are in such ionized state (Jackson, 2008). In the first part of this study a model solution containing malvidin 3-monoglucoside was analyzed by detection at 518 nm after chromatographic separation and oxidation by means of H₂O₂ addition. A degradation of malvidin 3-monoglucoside after the addition of hydrogen peroxide occurred. Lopes et al. (2007) showed that under wine pH malvidine 3-O-glucoside was degraded to 2,4,6-trihydroxybenzaldehyde, syringic acid, and the 8-β-D-glucopyranosyl-2,4-dihydroxy-6-oxocyclohexa-2,4-dienyl acetic acid (anthocyanone A). These last molecules result from the degradation of malvidin 3-O-glucoside, formed by Baeyer-Villiger-type oxidation triggered by hydrogen peroxide. The degradation peaks (peaks 1-5 in Fig. 1) detected during chromatographic run are the same previously reported by Lopes and colleagues (2007). In this study we evaluated the concentrations of malvidin 3-monoglucoside and that of the main degradation product, the syringic acid (peak 4 in the chromatogram). As expected by the chemistry of anthocyanins in aqueous solution, a lower content of malvidin 3-monoglucoside was observed at higher pH (Fig. 2). After the addition of hydrogen peroxide a dramatic loss of malvidin 3-monoglucoside has been detected at both pH; the loss is slightly enhanced increasing the pH. It is well known that fruit native anthocyanins are susceptible to be destroyed by H₂O₂ (Sondheimer & Kertesz, 1952; Ozkan, Yemencioğlu, Asefi & Cemeroglu, 2002) but the effect of pH seems to be in disagreement with a previous study where flavilum cation, which is dominant at lower pH, has been showed to react 4.3 ± 0.4 times faster with hydrogen peroxide molecules than the neutral pseudo-base (Thompson, Spiro & Griffith, 1996) (Fig. 2). However, Zhang, Duan, Ji and Pang (2000) found, in agreement with our results, a less degradation of litchi anthocyanins by H₂O₂ at lower pH. The discrepancy between Thompson, Spiro and Griffith (1996) results and our study can be related to the fact that in solution the direct chemical degradation of malvidin 3-monoglucoside is not the only reaction triggered by hydrogen peroxide. It is likely that part of radicals produced by Fenton

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

reaction are quenched by tartaric acid (Elias & Waterhouse, 2010) more than from malvidin 3-monoglucoside in solution at pH 3.20.

At pH 3.20 an antioxidant effect of SO₂ is detected regardless the concentration used. In contrast, at higher pH, the protective activity against oxidative degradation was detected only at higher SO₂ concentration. Probably at these concentrations SO₂ not only quenches hydrogen peroxide limiting Fenton reaction but, also binds malvidin 3-monoglucoside. To understand these results the chemistry of SO₂ in aqueous solution has to be considered. As well-known SO₂ exists in different forms, molecular SO₂(g) in equilibrium with bisulfite ion, in turn in equilibrium with bound SO₂. In the model solution under investigation bound SO₂ is related to the reaction of bisulfite ion with carbonyls deriving from oxidation of tartaric acid and ethanol (glyoxylic acid and acetaldehyde) as well as with malvidin 3-O glucoside. The extend of combination and the rate of binding is slower the lower the pH (Amerine & Joslin, 1970). In addition, lower is the pH more the equilibria are shifted towards the molecular SO₂. Therefore, at lower pH both phenomena determine a higher presence of bisulfite ions dissociating from bound SO₂ and capable to react with hydrogen peroxide and act as quenching compounds (Danilewicz, 2007).

Surprisingly at pH 3.80 low concentration of SO₂ determined a loss of malvidin 3-monoglucoside (Fig. 2). It is possible that in these conditions SO₂ did not protect the ethanol but promoted its oxidation, which is in agreement with the fact that a substantial proportion of SO₂ is oxidized to produce highly oxidizing SO radicals such as the peroxomonosulfate radical (Danilewicz, 2007). A past study of Connick and Zhang (1986) showing that the reaction of formation of peroxomonosulfate radical is enhanced at higher pH, confirm this hypothesis. The bleaching of anthocyanin solutions due to pH and SO₂ should be also considered (Brouillard, & El Hage Chahine, 1980) but this reaction is secondary with respect to the dominant kinetic of reaction between SO₂ and H₂O₂ in presence of trace metals and organic acids (Breytenbach, van Pareen, Pienaar, & van Eldik, 1994) and with respect to direct degradation of malvidin 3-Oglucoside by hydrogen peroxide when in excess of SO₂.

To understand if GSH may fulfil the antioxidant roles of SO₂ in wine conditions, such as to bind aldehyde compounds and to scavenge hydrogen peroxide, GSH and a combination of both antioxidant (SO₂ and GSH) were used. An increase in the degradation of malvidin 3-monoglucoside was detected when GSH was used alone and in combination with low concentration of SO₂ at both pH. The scavenging effect of SO₂ was instead dominant when its concentration was high and, in this case, no significant activity of GSH was detected. These

1
2
227 results seem in disagreement with previous results obtained during micro-oxygenation of
228 wine (Gambuti, Han, Peterson, & Waterhouse, 2015) where a little protective effect of GSH
5
229 against anthocyanins degradation was detected. The reason of the different behavior can be
7
230 found in the fact that in the present experiment only the anti-Fenton activity of GSH was
8
231 determined while in the previous study the oxidation was obtained adding directly oxygen (in
10
232 a micro-oxygenation experiment). It is well known that mechanism of oxidation of wine can
11
233 be separated in two parts, the first one regulated by quinones chemistry and leading to the
13
234 reduction of O₂ and production of H₂O₂, the second one regulated by Fenton reaction and
14
235 giving the high reactive radicals. The present results suggest that the antioxidant action of
16
236 GSH at wine pH is mainly due to its action on quinones chemistry and not on hydrogen
17
237 peroxide scavenging activity. The small loss of malvidin 3-monoglucoside may be due to the
19
238 formation, in a strong oxidant medium, of oxidized glutathione GSSG that may acts as
20
239 oxidant even if this activity has been reported only in living systems and in presence of
22
240 enzymes (Ceballos-Picot et al., 1996). Recently it has been showed that, at higher GSH/SO₂
23
241 ratio, in the presence of oxygen, GSH gives glutathione disulfide (GSSG), and GSSG reacts
25
242 with SO₃H⁻ to provide S-sulfonated glutathione (GSSO₃H) (Arapitsas et al., 2016). This
26
243 mechanism was favored in wine stored with a larger amount of oxygen. In our experimental
28
244 condition it could be assumed that a large amount of GSH was transformed into its sulfonated
29
245 analogue, thus simultaneously depleting the concentration of the two major wine antioxidants,
31
246 SO₂ and GSH. Further investigation can help to elucidate the reason of the disappear of
32
247 malvidin 3-monoglucoside in presence of GSH at lower pH.
34

248 The possible action of GSH as oxidant is confirmed by data on syringic acid (Table 1). This
36
249 molecule derive from breakdown of malvidin-3-O-monoglucoside (Lopes et al., 2007). It has
38
250 been detected after photodegradation (Maccarone, Ferrigno, Longo & Rapisarda, 1987) and
39
251 enzymatic and thermal degradation of anthocyanins (Piffaut, Kader, Girardin & Metche,
41
252 1994). It origins, together with carboxyaldehyde from the ring opening of 2,4,6-
42
253 trihydroxychalcone and proton transfer and rearrangements in the acidic aqueous medium.
44
254 Syringic acid concentrations detected after oxidation of model solutions containing GSH are
45
255 higher than in Ox solution indicating that, under strong oxidative stress, GSH never acts as
47
256 oxidant. Moreover, these results confirmed the possibility to use syringic acid as marker of
48
257 malvidin-3-O-glucoside oxidation. Another chemical antioxidant used in winemaking, the
50
258 ascorbic acid, showed a similar behavior (Iacobucci & Sweeny, 1983). Hence only proper
51
259 concentration of sulfur dioxide is effective in prevent Fenton reaction and, at the best of our
53
260 knowledge, no other molecule can exercise the same effect.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3.2 Red wine experiments

The degradation of malvidin 3-monoglucoside by hydrogen peroxide is lower in wine than in model solution (Tab. 2 and Tab. 3). This result can be easily explained considering that in wine very dangerous radicals produced by Fenton reaction react with a greater number of compounds in solution and not only with native anthocyanins, therefore it is likely that the occurrence of other competitive reactions decreased the malvidin degradation in wine. In contrast with results obtained in model solution, in wine the lower the pH was the higher and faster pigment oxidation. This is in agreement with previous results obtained on wine (Pechamat, Zeng, Jourdes, Ghidossi & Teissedre, 2014) and it is probably due to the occurrence of reactions involving malvidin 3-monoglucoside, the glyoxylic acid produced by tartaric acid oxidation and, other flavanols present in wine and not in model solution (Es-Safi, Cheynier & Moutounet, 2003).

Among native anthocyanins, malvidin-3-acetylglucoside and malvidin-3-monoglucoside are less degraded by hydrogen peroxide than delphinidin-3-monoglucoside and peonidin-3-monoglucoside. This results is due to O-methylation of these molecules that results in a higher stability of anthocyanidin molecule while, the existence of hydroxyl groups makes the molecule more sensitive to oxidation (Mazza & Francis 1995). In agreement with previous studies (Bakker & Timberlake, 1997; Morata, Calderón, González, Gómez-Cordovés & Suárez, 2007), a lower degradation of vitisin B has been even observed as expected because pyranoanthocyanidins are more stable. As expected a positive action of SO₂ against malvidin 3-monoglucoside and all anthocyanins degradation at both pH was detected and it resulted highly correlated with concentration. In contrast with model solution never an oxidant activity of SO₂ was detected. Departure from model solution in the activity of SO₂ in real wine has been also recently observed by Danilewicz (2016). These results are not surprising and indicate that in real wine ethanol, anthocyanins and tartaric acid are not the only target of radicals produced by Fenton but other compounds such as other phenolics, malic acid, glyceraldehyde and volatile compounds compete with them (Elias & Waterhouse, 2010). These results showed that, as suggested by Danilewicz (2007), the oxidant activity of SO₂ is prevented by the radical scavenging action of polyphenols in real wine and its ability to scavenge H₂O₂ is dominant (Table 2 and Table 3). About the GSH, as observed in model solution, a slight oxidative action was observed when it was used alone or in combination with high concentration of SO₂ to contrast hydrogen peroxide action. Thus, also in real wine, GSH is not capable to inhibit Fenton reaction and/or to bind efficiently carbonyls as SO₂

Field Code Changed

1
2
294 while it seems that, in presence of hydrogen peroxide it can instead contribute to oxidation
3
4
295 and/or reacts with flavonoids. Considering the protective role played by GSH towards varietal
5
296 thiols (Nikolantonaky & Waterhouse, 2012) and previous results obtained during micro-
7
297 oxygenation (Gambut, Han, Peterson & Waterhouse, 2015) this study furnish serious
8
298 evidences that the action of this tripeptide on wine may be limited only to its effect on
10
299 quinones chemistry but not on complementary antioxidant actions of SO₂. Future work must
11
300 be conducted in real wine to understand the reasons of the slight loss of malvidin 3-
13
301 monoglucoside observed in presence of an excess of hydrogen peroxide and GSH.
14

15
302 ANOVA analysis of Ox samples showed that all of the variables tested in this study had, in
16
303 model solution, the ability to significantly affect the degradation of malvidin 3-
18
304 monoglucoside by hydrogen peroxide while, in red wine, pH showed no significant effect
19
205 (Table 4). This last result seems in contrast with data reported by Pechamat, Zeng, Jourdes,
21
306 Ghidossi and Teissedre (2014) who observed a higher formation of
22
307 pyranomalvidin–procyanidin dimers in oxygenated red wines at lower pH. However, we use
24
308 very strong oxidative conditions and, the effect of pH on the loss of monomeric anthocyanins
25
309 during MOx changes with time (Kontoudakis et al., 2011). As expected SO₂ has the most
27
310 significant effect in preventing malvidin 3-monoglucoside loss (significant at p < 0.001). A
28
311 significant effect of GSH was observed and it resulted in a slight loss of anthocyanin in
30
312 presence of hydrogen peroxide. The interaction between pH and SO₂ was significant only in
31
313 model solution while interactions between SO₂ and GSH were significant in real wine leading
33
314 to a slight minor anthocyanin preservation. These results partially agree with a recent study
35
315 showing no protective activity of GSH to prevent white wines oxidation after one year of
36
316 aging in bottles (Panero, Motta, Petrozziello, Guaita & Bosso, 2015).
38

39 40 418 4. CONCLUSIONS 42

43
44 The protective effect of sulfur dioxide on malvidin 3-monoglucoside degradation was, as
45
46 expected, higher at lower pH in model solution. The use of GSH alone determined an increase
47
48 in the degradation of malvidin 3-monoglucoside regardless of pH in model solution and in
49
50 real wine. Results obtained in this study showed that the possibility to use GSH to prevent
51
52 anthocyanins oxidation is not linked to its capability to quench hydrogen peroxide but only, in
53
54 the first steps of oxidation, to act on quinones chemistry and limit the reduction of oxygen to
55
56 hydrogen peroxide. When in wine is present hydrogen peroxide GSH is not able to scavenge
57
58 it and contrast Fenton reaction nor alone and not in combination with SO₂ at concentration
59
60
61
62
63
64
65

1
2 usually proposed during winemaking. Taking into account these results, the use of this
3
4 tripeptide as an alternative to SO₂ has to be revised and the chemistry of action of these
5
6 compounds in wine conditions better understood.
7

8 9 10 **5. REFERENCES**

- 11
12 1. Adachi, T., Nonogi, H., Fuke, T., Ikuzawa, M., Fujita, K., Izumi, T., ... & Iwaida, M. (1979).
13 On the combination of sulphite with food ingredients (aldehydes, ketones and sugars).
14 II. *Zeitschrift für Lebensmittel-Untersuchung und Forschung*, 168(3), 200-205.
15
16 2. Amerine, M. A., & Joslyn, M. A. (1970). *Table wines: the technology of their production*
17 (2nd ed.). University of California Press. Berkeley, CA, (Chapter 2).
18 ~~2. Amerine, M. A., & Joslyn, M. A. (1970). *Table wines: the technology of their production (Vol.*~~
19 ~~*2).*~~
20
21 3. Arapitsas, P., Ugliano, M., Perenzoni, D., Angeli, A., Pangrazzi, P., & Mattivi, F. (2016).
22 Wine metabolomics reveals new sulfonated products in bottled white wines, promoted by
23 small amounts of oxygen. *Journal of Chromatography A*, 1429, 155-165.
24
25 4. Bouzanquet, Q., Barril, C., Clark, A. C., Dias, D. A., & Scollary, G. R. (2012). A novel
26 glutathione-hydroxycinnamic acid product generated in oxidative wine conditions. *Journal of*
27 *Agricultural and Food Chemistry*, 60(49), 12186-12195.
28
29 5. Breytenbach, L., van Pairen, W., Pienaar, J. J., & van Eldik, R. (1994). The influence of
30 organic acids and metal ions on the kinetics of the oxidation of sulfur (IV) by hydrogen
31 peroxide. *Atmospheric Environment*, 28(15), 2451-2459.
32
33 6. Brouillard, R., & Dubois, J. E. (1997). Mechanism of the structural transformation of
34 anthocyanins in acid media. *J. Am. Chem. Soc.*, 99, 1359-1364.
35
36 7. Brouillard, R., & El Hage Chahine, J. M. (1980). Chemistry of anthocyanin pigments. 6.
37 Kinetic and thermodynamic study of hydrogen sulfite addition to cyanin. Formation of a
38 highly stable Meisenheimer-type adduct derived from a 2-phenylbenzopyrylium salt. *Journal*
39 *of the American Chemical Society*, 102(16), 5375-5378.
40
41 8. Ceballos-Picot, I., Witko-Sarsat, V., Merad-Boudia, M., Nguyen, A. T., Thévenin, M.,
42 Jaudon, M. C., ... & Descamps-Latscha, B. (1996). Glutathione antioxidant system as a
43 marker of oxidative stress in chronic renal failure. *Free Radical Biology and Medicine*, 21(6),
44 845-853.
45
46 9. Cheynier, V. F., & Van Hulst, M. W. (1988). Oxidation of trans-caftaric acid and 2-S-
47 glutathionylcaftaric acid in model solutions. *Journal of Agricultural and Food*
48 *Chemistry*, 36(1), 10-15.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

10. Clark, A. C., Prenzler, P. D., & Scollary, G. R. (2007). Impact of the condition of storage of tartaric acid solutions on the production and stability of glyoxylic acid. *Food Chemistry*, 102(3), 905-916.

11. Connick, R. E., & Zhang, Y. X. (1996). Kinetics and mechanism of the oxidation of HSO₃-by O₂. 2. the manganese (II)-catalyzed reaction. *Inorganic Chemistry*, 35(16), 4613-4621.

12. Danilewicz, J. C. (2007). Interaction of sulfur dioxide, polyphenols, and oxygen in a wine-model system: Central role of iron and copper. *American Journal of Enology and Viticulture*, 58(1), 53-60.

13. Danilewicz, J. C. (2016). Reaction of oxygen and sulfite in wine. *American Journal of Enology and Viticulture*, 67(1), 13-17.

14. Danilewicz, J. C., & Wallbridge, P. J. (2010). Further studies on the mechanism of interaction of polyphenols, oxygen, and sulfite in wine. *American Journal of Enology and Viticulture*, 61(2), 166-175.

15. Danilewicz, J. C., (2011). Mechanism of Autoxidation of Polyphenols and Participation of Sulfite in Wine: Key Role of Iron. *American Journal of Enology and Viticulture*, 62, 319-328.

16. Elias, R. J. & Waterhouse, A. L. (2010). Controlling the Fenton Reaction in Wine. *Journal of Agricultural and Food Chemistry*, 58, 1699-1707.

17. Elias, R. J., Andersen, M. L., Skibsted, L. H., & Waterhouse, A. L. (2009). Identification of Free Radical Intermediates in Oxidized Wine Using Electron Paramagnetic Resonance Spin Trapping. *Journal of Agricultural and Food Chemistry*, 57, 4359-4365.

18. Es-Safi, N. E., Cheynier, V., & Moutounet, M. (2003). Implication of phenolic reactions in food organoleptic properties. *Journal of Food Composition and Analysis*, 16(5), 535-553.

19. Es-Safi, N. E., Fulcrand, H., Cheynier, V., & Moutounet, M. (1999). Studies on the acetaldehyde-induced condensation of (-)-epicatechin and malvidin 3-O-glucoside in a model solution system. *Journal of Agricultural and Food Chemistry*, 47, 2096-2102.

20. Finley, J. W., Wheeler, E. L., & Witt, S. C. (1981). Oxidation of glutathione by hydrogen peroxide and other oxidizing agents. *Journal of Agricultural and Food Chemistry*, 29, 404-7.

21. Gambuti, A., Han, G., Peterson, A. L., & Waterhouse, A. L. (2015). Sulfur dioxide and glutathione alter the outcome of microoxygenation. *American Journal of Enology and Viticulture*, 66, 411-423.

~~21. Gambuti, A., Han, G., Peterson, A. L., & Waterhouse, A. L. (2015). Sulfur dioxide and glutathione alter the outcome of microoxygenation. *American Journal of Enology and Viticulture*, ajev-2015.~~

- 1
2
3
394 22. He, F., Liang, N.-N., Mu, L., Pan, Q.-H., Wang, J., Reeves, M. J., & Duan, C.-Q. (2012).
395 Anthocyanins and Their Variation in Red Wines I. Monomeric Anthocyanins and Their Color
396 Expression. *Molecules*, 17, 1571-1601.
- 397 23. Iacobucci, G. A., & Sweeny, J. G. (1983). The chemistry of anthocyanins, anthocyanidins and
398 related flavylum salts. *Tetrahedron*, 39(19), 3005-3038.
- 399 24. Jackson, R. S. (2008). *Wine science: principles and applications*.
- 400 25. Kilmartin, P. A., Zou, H., & Waterhouse, A. L. (2001). A cyclic voltammetry method suitable
401 for characterizing antioxidant properties of wine and wine phenolics. *Journal of Agricultural
402 and Food Chemistry*, 49(4), 1957-1965.
- 403 26. Kontoudakis, N., González, E., Gil, M., Esteruelas, M., Fort, F., Canals, J. M., & Zamora, F.
404 (2011). Influence of wine pH on changes in color and polyphenol composition induced by
405 micro-oxygenation. *Journal of Agricultural and Food Chemistry*, 59(5), 1974-1984.
- 406 27. Kritzing, E. C., Bauer, F. F., & Du Toit, W. J. (2012). Role of glutathione in winemaking: a
407 review. *Journal of Agricultural and Food Chemistry*, 61(2), 269-277.
- 408 28. Lopes, P., Richard, T., Saucier, C., Teissedre, P. L., Monti, J. P., & Glories, Y. (2007).
409 Anthocyanone A: A quinone methide derivative resulting from malvidin 3-O-glucoside
410 degradation. *Journal of agricultural and food chemistry*, 55(7), 2698-2704.
- 411 29. Maccarone, E., Ferrigno, V., Longo, L. L. & Rapisarda, P. (1987). Effects of light on
412 anthocyanins. Kinetics and photodegradation products in acidic aqueous solutions. *Annali di
413 Chimica*, 77, 499-508.
- 414 30. Morata, A., Calderón, F., González, M. C., Gómez-Cordovés, M. C., & Suárez, J. A. (2007).
415 Formation of the highly stable pyranoanthocyanins (vitisins A and B) in red wines by the
416 addition of pyruvic acid and acetaldehyde. *Food Chemistry*, 100(3), 1144-1152.
- 417 31. Nikolantonaki, M., & Waterhouse, A. L. (2012). A method to quantify quinone reaction rates
418 with wine relevant nucleophiles: a key to the understanding of oxidative loss of varietal
419 thiols. *Journal of Agricultural and Food Chemistry*, 60(34), 8484-8491.
- 420 ~~32. OIV Compendium of International Methods of Wine and Must Analysis. (2017).
421 <http://www.oiv.int/> Accessed 10.01.17.~~
- 422 ~~32. OIV Compendium of International Methods of Wine and Must Analysis. Off. Int. Vigne Vin.
423 Paris 2017, URL (<http://www.oiv.int/>).~~
- 424 33. Özkan, M., Yemenicioglu, A., Asefi, N., & Cemeroglu, B. (2002). Degradation kinetics of
425 anthocyanins from sour cherry, pomegranate, and strawberry juices by hydrogen
426 peroxide. *Journal of Food Science*, 67(2), 525-529.
- 52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

34. Panero, L., Motta, S., Petrozziello, M., Guaita, M., & Bosso, A. (2015). Effect of SO₂, reduced glutathione and ellagitannins on the shelf life of bottled white wines. *European Food Research and Technology*, 240(2), 345-356.

35. Pechamat, L., Zeng, L., Jourdes, M., Ghidossi, R., & Teissedre, P. L. (2014). Occurrence and formation kinetics of pyranomalvidin-procyanidin dimer pigment in merlot red wine: Impact of acidity and oxygen concentrations. *Journal of Agricultural and Food Chemistry*, 62(7), 1701-1705.

36. Piffaut, B., Kader, F., Girardin, M., & Metche, M. (1994). Comparative degradation pathways of malvidin 3, 5-diglucoside after enzymatic and thermal treatments. *Food Chemistry*, 50(2), 115-120.

37. Roussis, I. G., Lambropoulos, I., & Tzimas, P. (2007). Protection of volatiles in a wine with low sulfur dioxide by caffeic acid or glutathione. *American Journal of Enology and Viticulture*, 58, 274-278.

38. Sondheimer, E., & Kertesz, Z. I. (1952). The kinetics of the oxidation of strawberry anthocyanin by hydrogen peroxide. *Journal of Food Science*, 17(1-6), 288-298.

39. Sonni, F., Clark, A. C., Prenzler, P. D., Riponi, C., & Scollary, G. R. (2011). Antioxidant action of glutathione and the ascorbic acid/glutathione pair in a model white wine. *Journal of Agricultural and Food Chemistry*, 59(8), 3940-3949.

40. Sonni, F., Moore, E. G., Clark, A. C., Chinnici, F., Riponi, C., & Scollary, G. R. (2011). Impact of glutathione on the formation of methylmethine-and carboxymethine-bridged (+)-catechin dimers in a model wine system. *Journal of Agricultural and Food Chemistry*, 59(13), 7410-7418.

41. Thompson, K. M., Spiro, M., & Griffith, W. P. (1996). Mechanism of bleaching by peroxides. Part 4.-Kinetics of bleaching of malvin chloride by hydrogen peroxide at low pH and its catalysis by transition-metal salts. *Journal of the Chemical Society, Faraday Transactions*, 92(14), 2535-2540.

42. Ugliano, M., Kwiatkowski, M., Vidal, S., Capone, D., Siebert, T., Dieval, J. B., ... & Waters, E. J. (2011). Evolution of 3-mercaptohexanol, hydrogen sulfide, and methyl mercaptan during bottle storage of Sauvignon blanc wines. Effect of glutathione, copper, oxygen exposure, and closure-derived oxygen. *Journal of Agricultural and Food Chemistry*, 59(6), 2564-2572.

43. Von Elbe, J. H., & Schwartz, S. J. (1996). Colorants. *Food Chemistry*, 3, 651-723.

44. Winterbourn, C. C., & Metodiewa, D. (1999). Reactivity of biologically important thiol compounds with superoxide and hydrogen peroxide: Free Radical Biology and Medicine, 27, 322-328.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

45. Zhang, Z. Q., Duan, X. W., Ji, Z. L., & Pang, X. Q. (2000, June). Influence of pH and active oxygen on the stability of anthocyanins from litchi pericarp. Paper presented at the I International Symposium on Litchi and Longan, Guangzhou, China 16-19 June (pp. 339-342). ISHS: Acta Horticulturae.

~~45. Zhang, Z. Q., Duan, X. W., Ji, Z. L., & Pang, X. Q. (2000, June). Influence of pH and active oxygen on the stability of anthocyanins from litchi pericarp. In *I International Symposium on Litchi and Longan* 558 (pp. 339-342).~~

Formatted: Indent: Left: 1.27 cm, No bullets or numbering

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

Figure captions

Fig. 1. HPLC chromatogram recorded at 280 nm representing the malvidin 3-O-glucoside degradation over 72 hours of reaction with hydrogen peroxide. Compounds 1–5 correspond to breakdown products.

Fig. 2 Effect of SO₂ and GSH on the degradation of malvidin 3-monoglucoside by hydrogen peroxide in model solution. Ox: sample added with hydrogen peroxide 39.2 mg/L of O₂ eq; Ox+SO₂ low: Ox + 37.4 mg/L of SO₂; Ox+SO₂ high: Ox + 202 mg/L of SO₂ (3.16 mM); Ox+GSH: Ox+ 30 mg/L of glutathione (0.098 mM); Ox+SO₂ low+GSH: Ox + 37.4 mg/L of SO₂ and 30 mg/L of glutathione (0.098 mM); Ox+SO₂high+GSH: Ox + 202 mg/L of SO₂ and 30 mg/L of glutathione.




Fig. 1. HPLC chromatogram recorded at 280 nm representing the malvidin 3-O-glucoside degradation over 72 hours of reaction with hydrogen peroxide. Compounds 1–5 correspond to breakdown products.




Fig. 2 Effect of SO₂ and GSH on the degradation of malvidin 3-monoglucoside by hydrogen peroxide in model solution. Ox: sample added with hydrogen peroxide 39.2 mg/L of O₂ eq; Ox+SO₂ low: Ox + 37.4 mg/L of SO₂; Ox+SO₂ high: Ox + 202 mg/L of SO₂ (3.16 mM); Ox+GSH: Ox+ 30 mg/L of glutathione (0.098 mM); Ox+SO₂ low+GSH: Ox + 37.4 mg/L of SO₂ and 30 mg/L of glutathione (0.098 mM); Ox+SO₂high+GSH: Ox + 202 mg/L of SO₂ and 30 mg/L of GSH. Oxidative treatments (a, b, c, ...) and oxidation time (A, B) sharing the same letters are not significantly different.

Table 1. Effect of SO₂ and GSH on the formation of syringic acid (microgram/L), a breakdown compound of malvidin 3-O-glucoside detected at 15.25 min at 280 nm (peak 4) during HPLC run under oxidative conditions over time. Oxidative treatments (a, b, c, ...) and oxidation time (A, B) sharing the same letters are not significantly different.

pH 3.2			
	0 h	16 h	72 h
Control	8 ± 1 C	47 ± 1 eB	107 ± 5 dA
OX		224 ± 14 dB	528 ± 17 aA
OX+SO ₂ low		266 ± 16 cB	487 ± 3 cA
OX+SO ₂ high		24 ± 0 fA	24 ± 1 eA
OX+GSH		349 ± 11 bB	504 ± 4 bA
OX+SO ₂ low+GSH		381 ± 8 aB	492 ± 6 bcA
OX+SO ₂ high+GSH		22 ± 1 fB	30 ± 1 eA
pH 3.8			
Control	9 ± 1 C	43 ± 2 eB	94 ± 1 eA
OX		208 ± 14 dB	461 ± 1 dA
OX+SO ₂ low		260 ± 12 cB	526 ± 1 aA
OX+SO ₂ high		30 ± 5 efA	27 ± 0 gA
OX+GSH		378 ± 7 aB	519 ± 2 bA
OX+SO ₂ low+GSH		330 ± 6 dB	503 ± 4 cA
OX+SO ₂ high+GSH		24 ± 0 fB	36 ± 1 fA

Table 2. Effect of SO₂ and GSH on the degradation of monomeric anthocyanins by hydrogen peroxide in red wine at pH 3.2. Ox: wine added with hydrogen peroxide 39.2 mg/L of O₂ eq; Ox+SO₂ low: Ox + 37.4 mg/L of SO₂; Ox+SO₂ high: Ox + 202 mg/L of SO₂ (3.16 mM); Ox+GSH: Ox+ 30 mg/L of glutathione (0.098 mM); Ox+SO₂ low+GSH: Ox + 37.4 mg/L of SO₂ and 30 mg/L of glutathione (0.098 mM); Ox+SO₂high+GSH: Ox + 202 mg/L of SO₂ and 30 mg/L of glutathione.

	Df3glc (mg/L)		Pn3glc (mg/L)		Mv3glc (mg/L)		Vit.B (mg/L)		Mv3acglc (mg/L)		Tot of mon. anth.* (mg/L)	
Control	61.9 ± 3.6	A	39.5 ± 0.3	A	649.5 ± 7.2	A	11.6 ± 0.0	A	139.2 ± 1.8	A	997.1 ± 14.3	A
Control16h	62.7 ± 1.9	aA	38.8 ± 2.9	aA	652.0 ± 14.1	abA	11.7 ± 0.0	aA	136.8 ± 5.2	aA	1000.5 ± 22.2	aA
Control 72h	57.6 ± 0.3	bA	39.2 ± 0.3	abA	640.3 ± 1.4	aA	9.8 ± 2.9	aA	136.4 ± 3.3	aA	977.0 ± 7.8	aA
Ox 16h	28.6 ± 0.1	cA	17.6 ± 0.4	cA	349.6 ± 4.3	dA	9.4 ± 0.0	dB	77.2 ± 2.7	cA	537.4 ± 7.8	cA
Ox 72h	25.5 ± 0.1	eB	16.2 ± 0.1	dB	313.7 ± 5.8	cB	9.5 ± 0.0	aA	69.7 ± 2.4	cB	487.0 ± 8.6	dB
Ox+SO ₂ low 16h	37.3 ± 0.1	bA	23.5 ± 0.1	bA	428.8 ± 1.0	aA	9.8 ± 0.0	aB	92.5 ± 0.6	bcA	660.5 ± 0.8	bA
Ox+SO ₂ low 72h	34.3 ± 0.3	dB	21.5 ± 0.4	cB	397.5 ± 8.3	cB	9.9 ± 0.0	aA	86.1 ± 3.9	bB	613.2 ± 13.6	aB
Ox+SO ₂ high 16h	64.0 ± 3.3	aA	39.0 ± 0.5	aA	665.0 ± 6.9	cA	11.8 ± 0.1	cA	139.5 ± 9.4	aA	1009.0 ± 21.0	aA
Ox+SO ₂ high 72h	65.7 ± 0.2	aA	40.6 ± 0.2	aA	660.7 ± 4.4	aA	11.5 ± 0.1	eA	142.2 ± 1.6	aA	1012.6 ± 6.4	cA
Ox+GSH 16h	27.2 ± 0.6	bA	18.8 ± 0.6	cA	354.9 ± 6.5	dA	8.8 ± 0.0	eB	77.7 ± 3.1	cA	546.0 ± 11.0	cA
Ox+GSH 72h	23.5 ± 2.8	eA	15.6 ± 2.6	dB	297.2 ± 37.0	cB	9.6 ± 0.6	aA	64.3 ± 9.1	cB	459.7 ± 55.9	dB
Ox+SO ₂ low+GSH 16h	35.2 ± 1.0	cA	23.8 ± 0.0	bA	425.0 ± 0.2	cA	9.3 ± 0.1	dB	101.4 ± 13.9	bcA	662.5 ± 12.6	bA
Ox+SO ₂ low+GSH 72h	34.5 ± 1.4	dA	22.7 ± 1.3	bA	412.0 ± 24.9	bA	9.7 ± 0.0	aA	89.5 ± 4.9	bA	635.8 ± 33.1	cA
Ox+SO ₂ high+GSH 16h	61.5 ± 1.1	aA	38.3 ± 1.1	aA	636.7 ± 15.1	bA	10.9 ± 0.3	bA	139.5 ± 9.2	aA	979.4 ± 24.0	aA
Ox+SO ₂ high+GSH 72h	42.1 ± 0.3	cB	37.3 ± 0.4	cA	621.6 ± 2.1	aA	11.1 ± 0.4	aA	144.5 ± 0.8	aA	943.9 ± 1.4	aA

Dp3glc = delphinidin 3-glucoside, Pn3glc = peonidin 3-monoglucoside, Mv3glc = malvidin 3-glucoside, Vit.B=vitisin B, Mv3acglc = malvidin 3-(6^{II}-acetyl)-glucoside, *Sum of monomeric anthocyanins. Oxidative treatments (a, b, c, ...) and oxidation time (A, B) sharing the same letters are not significantly different.

Table 3. Effect of SO₂ and GSH on the degradation of monomeric anthocyanins by hydrogen peroxide in red wine at pH 3.8. Ox: wine added with hydrogen peroxide 39.2 mg/L of O₂ eq; Ox+SO₂ low: Ox + 37.4 mg/L of SO₂; Ox+SO₂ high: Ox + 202 mg/L of SO₂ (3.16 mM); Ox+GSH: Ox+ 30 mg/L of glutathione (0.098 mM); Ox+SO₂ low+GSH: Ox + 37.4 mg/L of SO₂ and 30 mg/L of glutathione (0.098 mM); Ox+SO₂high+GSH: Ox + 202 mg/L of SO₂ and 30 mg/L of GSH.

	Dp3glc (mg/L)	Pn3glc (mg/L)	Mv3glc (mg/L)	Vit.B (mg/L)	Mv3acglc (mg/L)	Tot of mon. anth.* (mg/L)
Control	56.1 ± 1.7 AB	38.4 ± 0.8 A	627.7 ± 22.0 A	12.0 ± 0.5 A	136.5 ± 10.2 A	962.4 ± 37.5 A
Control16h	57.6 ± 0.5 bA	36.0 ± 0.3 bB	629.0 ± 7.6 bA	12.1 ± 0.1 aA	137.6 ± 18.9 aA	967.7 ± 28.6 aA
Control 72h	53.9 ± 0.1 cB	35.7 ± 0.8 bB	609.3 ± 0.4 bA	11.9 ± 0.1 aA	131.1 ± 1.2 bA	940.7 ± 0.5 bA
Ox 16h	33.3 ± 0.5 eB	18.2 ± 0.4 dA	385.3 ± 6.9 eB	9.1 ± 0.1 eA	85.0 ± 4.4 bcA	596.5 ± 14.9 cA
Ox 72h	29.2 ± 1.4 eC	17.0 ± 1.6 eA	344.9 ± 3.0 eC	8.8 ± 0.0 eA	75.2 ± 0.1 dB	533.5 ± 6.5 dB
Ox+SO ₂ low 16h	39.3 ± 0.6 cA	23.0 ± 0.8 cA	443.1 ± 9.1 cA	9.8 ± 0.0 dA	96.0 ± 4.1 bA	685.1 ± 14.2 bA
Ox+SO ₂ low 72h	36.1 ± 1.6 dA	22.3 ± 0.0 cA	407.2 ± 0.7 dB	9.5 ± 0.0 dB	86.3 ± 0.1 cB	630.1 ± 2.6 cB
Ox+SO ₂ high 16h	61.9 ± 0.1 aA	39.6 ± 0.7 aA	651.8 ± 3.8 aA	11.8 ± 0.1 bA	139.9 ± 2.5 aA	992.1 ± 7.1 aA
Ox+SO ₂ high 72h	63.7 ± 0.2 aA	39.0 ± 0.1 aB	636.3 ± 4.2 aA	11.3 ± 0.1 bB	137.6 ± 1.7 aA	975.8 ± 6.7 aA
Ox+GSH 16h	29.7 ± 0.5 fA	17.8 ± 1.8 dA	358.2 ± 6.6 cA	8.4 ± 0.1 gA	77.4 ± 1.6 cA	551.6 ± 10.4 dA
Ox+GSH 72h	26.4 ± 1.4 fB	16.7 ± 0.4 eA	323.9 ± 7.4 eB	8.4 ± 0.0 gA	70.9 ± 2.7 eB	500.6 ± 12.7 eB
Ox+SO ₂ low+GSH 16h	34.5 ± 0.9 dA	19.8 ± 0.3 cA	401.2 ± 5.2 cA	9.2 ± 0.0 dA	87.0 ± 0.4 bcA	616.9 ± 5.5 bA
Ox+SO ₂ low+GSH 72h	36.2 ± 0.7 dA	23.2 ± 1.1 dB	434.2 ± 5.3 cB	9.9 ± 0.0 dB	94.7 ± 1.4 cB	670.2 ± 5.6 cB
Ox+SO ₂ high+GSH 16h	60.8 ± 0.1 aA	38.5 ± 0.9 aA	633.0 ± 9.8 bA	11.3 ± 0.2 cA	136.9 ± 4.3 aA	968.2 ± 16.0 aA
Ox+SO ₂ high+GSH 72h	60.1 ± 0.8 bA	37.4 ± 0.9 abA	607.9 ± 14.2 bA	10.3 ± 0.3 cB	129.8 ± 1.7 bA	932.2 ± 14.6 bA

Dp3glc = delphinidin 3-glucoside, Pn3glc = peonidin 3-monoglucoside, Mv3glc = malvidin 3-glucoside, Vit.B=vitisin B, Mv3acglc = malvidin 3-(6^{II}-acetyl)-glucoside, *Sum of monomeric anthocyanins. Oxidative treatments (a, b, c, ...) and oxidation time (A, B) sharing the same letters are not significantly different.

Table 4. F values and significance of variables SO₂, GSH and pH for malvidin 3-monoglucoside degradation by hydrogen peroxide.

	Model solution		Red wine	
	F	Pr > F	F	Pr > F
pH	24.24	0.000	2.887	0.124
SO₂	4558.60	< 0.0001	1548.140	< 0.0001
GSH	18.46	0.000	6.563	0.031
pH x SO₂	5.75	0.028	2.558	0.144
pH x GSH	0.73	0.405	0.178	0.683
SO₂ x GSH	2.44	0.137	10.828	0.009

