

On the interpretation of the atmospheric neutrino data in terms of flavor changing neutrino interactions

This content has been downloaded from IOPscience. Please scroll down to see the full text.

JHEP07(2000)006

(<http://iopscience.iop.org/1126-6708/2000/07/006>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 130.237.122.245

This content was downloaded on 09/09/2015 at 01:50

Please note that [terms and conditions apply](#).

On the interpretation of the atmospheric neutrino data in terms of flavor changing neutrino interactions

Nicolao Fornengo, M. Concha Gonzalez-Garcia, and Jose W. Furtado Valle

Instituto de Física Corpuscular - C.S.I.C.
Departamento de Física Teòrica, Universitat of València
46100 Burjassot, València, Spain
E-mail: fornengo@to.infn.it, concha@flamenco.ific.uv.es,
valle@flamenco.ific.uv.es

ABSTRACT: Flavour changing (FC) neutrino-matter interactions have been proposed as a solution to the atmospheric neutrino anomaly. Here we perform the analysis of the full set of the recent 52 kTy SUPER-KAMIOKANDE atmospheric neutrino data, including the zenith angle distribution of the contained events as well as the higher energy upward-going stopping and through-going muon events. Our results show that the FC mechanism can describe the full data sample with a $\chi^2_{\min} = 44/33$ d.o.f. which is acceptable only at the 90.5% confidence level, while the $\nu_\mu \rightarrow \nu_\tau$ oscillation scenario gives a $\chi^2_{\min} = 23.5/33$ d.o.f. which is valid at the 11% CL. The combined analysis confines the amount of FC to be either close to maximal or to the level of about (10–50)%.

KEYWORDS: Solar and Atmospheric Neutrinos, Beyond Standard Model, Neutrino Physics.

Contents

1. Introduction	1
2. Massless neutrino evolution with FC interaction	2
3. Fitting the data to the FC hypothesis	3
4. Conclusions	10

1. Introduction

Neutrinos produced as decay products in hadronic showers from cosmic ray collisions with nuclei in the upper atmosphere [1] have been observed by several detectors [2]–[7]. Although the absolute fluxes of atmospheric neutrinos are largely uncertain, the expected ratio (μ/e) of the muon neutrino flux $(\nu_\mu + \bar{\nu}_\mu)$ over the electron neutrino flux $(\nu_e + \bar{\nu}_e)$ is robust, since it largely cancels out the uncertainties associated with the absolute flux. In fact, this ratio has been calculated [1] with an uncertainty of less than 5% over energies varying from 0.1 GeV to 100 GeV. In this resides our confidence on the long-standing atmospheric neutrino anomaly.

Although the first iron-calorimeter detectors in Fréjus [2] and NUSEX [3] reported a value of the double ratio, $R(\mu/e) = (\mu/e)_{\text{data}}/(\mu/e)_{\text{MC}}$, consistent with one, all the water Cerenkov detectors KAMIOKANDE [4], IMB [5] and SUPER-KAMIOKANDE [6] have measured $R(\mu/e)$ significantly smaller than one. Moreover, the SOUDAN-2 collaboration, also using an iron-calorimeter, reported a small value of $R(\mu/e)$ [7], showing that the so-called atmospheric neutrino anomaly was not a feature of water Cerenkov detectors.

Recent SUPER-KAMIOKANDE high statistics observations [6] indicate that the deficit in the total ratio $R(\mu/e)$ is due to the number of neutrinos arriving in the detector at large zenith angles. Although e -like events do not present any compelling evidence of a zenith-angle dependence, the μ -like event rates are substantially suppressed at large zenith angles.

The $\nu_\mu \rightarrow \nu_\tau$ as well as the $\nu_\mu \rightarrow \nu_s$ [8, 9] oscillation hypothesis provides a very good explanation for this smaller-than-expected ratio, which is also simple and well-motivated theoretically. This led the SUPER-KAMIOKANDE collaboration to conclude that their data provide good evidence for neutrino oscillations and neutrino

masses [10]. However, alternative explanations to the atmospheric neutrino data have been proposed in the literature including the possibility of neutrino decay [11], the violation of relativity principles [12, 13] or the violation of CPT symmetry [14]. These explanations, however, have been challenged by the precise data of SUPER-KAMIOKANDE on upward going muon events [15] which allows to study the energy dependence of the neutrino survival (or disappearance) probability [16, 17]. Based on such observations, both the possibility of an explanation of the anomaly in terms of neutrino decay [18] as well as the violation of relativity principles or the violation of CPT symmetry [17], have been disfavoured.

In ref. [19] an alternative explanation of the atmospheric neutrino data in terms of FC neutrino-matter interactions [20] was proposed, and it was shown that even if neutrinos have vanishing masses and/or the vacuum mixing angle is negligible, FC neutrino matter interactions could account for the SUPER-KAMIOKANDE results on contained events providing an excellent description to the data, statistically as good as neutrino oscillations. The validity of this explanation was first questioned in ref. [16] where the authors presented arguments against the FC neutrino-matter interaction solution on the basis of a fit to the up-going muons data from SUPER-KAMIOKANDE.

In this paper we re-analyze the possibility of explaining the atmospheric neutrino anomaly by means of $\nu_\mu \rightarrow \nu_\tau$ conversion induced by flavour-changing neutrino-matter interaction which can be effective during the neutrino propagation in the Earth. We extend the analysis of ref. [19] to the new set of SUPER-KAMIOKANDE data by including also the up-going muon samples.

2. Massless neutrino evolution with FC interaction

In our phenomenological approach we assume that the evolution equations which describe the $\nu_\mu \rightarrow \nu_\tau$ transitions in matter may be written as

$$i \frac{d}{dr} \begin{pmatrix} \nu_\mu \\ \nu_\tau \end{pmatrix} = \sqrt{2} G_F \begin{pmatrix} 0 & \epsilon_\nu n_f(r) \\ \epsilon_\nu n_f(r) & \epsilon'_\nu n_f(r) \end{pmatrix} \begin{pmatrix} \nu_\mu \\ \nu_\tau \end{pmatrix}, \quad (2.1)$$

where $\nu_a \equiv \nu_a(r)$ ($a = \mu, \tau$) are the probability amplitudes to find these neutrinos at a distance r from their creation position, $\sqrt{2} G_F n_f(r) \epsilon_\nu$ is the $\nu_\mu + f \rightarrow \nu_\tau + f$ forward scattering amplitude and $\sqrt{2} G_F n_f(r) \epsilon'_\nu$ is the difference between the $\nu_\tau - f$ and $\nu_\mu - f$ elastic forward scattering amplitudes, with $n_f(r)$ being the number density of the fermions which induce such processes.

The parameters ϵ and ϵ' contain the information about FC neutrino interactions. Such FC interactions may be accompanied by neutrino mass [21] but this need not be the case [22, 23]. One description would be to parametrize directly the FC interactions in terms of an effective four-fermion hamiltonian. This could, for instance, arise by renormalization effects from the unification scale down to the electroweak

scale in, say, supergravity models [23]. An alternative more phenomenological way is to consider the existence of a tree-level FC process $\nu_\alpha + f \rightarrow \nu_\beta + f$ where f is an elementary fermion (charged lepton or quark). The interaction can be mediated by a scalar or vector boson of mass m and the neutrino-fermion coupling is generically denoted by $g_{\alpha f}$ (α is a flavour index) and can be written as

$$\epsilon'_\nu = \frac{|g_{\tau f}|^2 - |g_{\mu f}|^2}{4m^2\sqrt{2}G_F} \quad \text{and} \quad \epsilon_\nu = \frac{g_{\tau f} \cdot g_{\mu f}}{4m^2\sqrt{2}G_F}. \quad (2.2)$$

Since we are assuming vanishing neutrino masses, the anti-neutrino transitions $\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$ are governed by the same evolution matrix given in eq. (2.1). For the sake of simplicity, we consider $\epsilon_{\bar{\nu}} = \epsilon_\nu$ and $\epsilon'_{\bar{\nu}} = \epsilon'_\nu$, which implies that we have only two free parameters in the analysis. Moreover, we set our normalization on these parameters by assuming that the relevant neutrino interaction in the Earth is only with down-type quarks. One could also assume that the incoming atmospheric neutrino has FC interactions off-electrons or equivalently, due to charge neutrality, off-up-type quarks. For simplicity, in the present analysis we consider only the case of interactions on down-type quarks.

We have calculated the transition probabilities of $\nu_\mu \rightarrow \nu_\tau$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$) as a function of the zenith angle by numerically solving the evolution equation using the density distribution in [24] and a realistic chemical composition with proton/neutron ratio 0.497 in the mantle and 0.468 in the core [25].

3. Fitting the data to the FC hypothesis

We have then used these probabilities to compute, as a function of the two parameters, ϵ_ν and ϵ'_ν , the theoretically expected numbers of events for the four sets of data reported by SUPER-KAMIOKANDE: sub-GeV, multi-GeV, stopping muons and through-going muons. The expected number of contained events are computed by convoluting the probability with the corresponding neutrino fluxes (for which we use the Bartol calculations [1]) and interaction cross sections and taking into account the experimental efficiencies as detailed in ref. [8]. For the up-going muon samples we obtain the effective muon fluxes for both stopping and through-going muons by convoluting the probabilities with the corresponding muon fluxes produced by the neutrino interactions with the Earth. We include the muon energy loss during propagation both in the rock and in the detector according to [26, 27] and we take into account also the effective detector area for both types of events, stopping and through-going. We compute the effective area using the simple geometrical picture given in ref. [28]. Our final results show good agreement with the full MC simulation of the SUPER-KAMIOKANDE collaboration in the Standard Model case (see the thick solid line in figure 3.)




Figure 1: Allowed regions for ϵ_ν and ϵ'_ν in the FC massless-neutrino scenario for the different SUPER-KAMIOKANDE data sets: (a) sub-GeV, (b) multi-GeV, (c) stopping muons and (d) through-going muons. The best fit points for each case are indicated by stars. The shaded area refers to the 90% C.L. while the contours stand for 95% and 99% C.L.

In our statistical analysis we adopted the technique [8, 29] of fitting separately the angular distributions of the μ - and e -like contained events (N_μ^i and N_e^i , i stands for sub-GeV and multi-GeV) and the up-going muon fluxes (Φ_μ^j , $j = \text{stopping, through-going}$). The expected number of events have been compared with the recent 52kTy data reported by the SUPER-KAMIOKANDE collaboration [30] and the allowed regions in the $(\epsilon_\nu, \epsilon'_\nu)$ plane have been determined from a χ^2 fit. In constructing the χ^2 function, we explicitly take into account the correlation of errors, both of theoretical and experimental origin. Details on the definition of the correlation matrix for contained events can be found in ref. [8], while the definition of the sources of errors and their correlations for the up-going muons fluxes are given in ref. [31, 29]. Here we simply summarize that we consider the overall normalization of the up-going




Figure 2: Expected up-down asymmetry in Super-Kamiokande for the FC-neutrino scenario as a function of the muon momentum for fully contained μ -like events, compared to the Super-Kamiokande experimental data.

muon fluxes to be affected by an uncertainty of 20% but in order to account for the uncertainties in the primary cosmic ray flux spectrum we allow a 5% variation in the ratio between muon events in different energy samples. We further introduce a 10% theoretical error in the ratio of electron-type to muon-type events of the different samples. Other important source of theoretical uncertainty arises from the neutrino interaction cross section which at SUPER-KAMIOKANDE ranges from 10–15%. Uncertainties in the ratio between different angular bins are treated, similarly to ref. [29], by allowing a variation of 5% times the difference between the mean bin cosines. With our definition we obtain, for instance, $\chi^2_{SM} = 122/(35 \text{ d.o.f.})$ which means that the SM has a CL of 10^{-11} ! Using this same χ^2 function for the case of oscillations we obtained allowed regions for masses and mixing angles very similar to those obtained by the SUPER-KAMIOKANDE collaboration both for contained events as well as for upward going muons [8].

In figure 1 we show the contours of the regions allowed by the SUPER-KAMIOKANDE data. The different panels of the figure refer to the fits performed over the different sets of data separately: (a) sub-GeV; (b) multi-GeV; (c) stopping muons; (d) through-going muons. The shaded areas are the regions allowed at 90% C.L., while the dashed and dotted contours refer to 95 and 99 % C.L., respectively. The condition used to determine the allowed regions is: $\chi^2 = \chi^2_{\min} + \Delta\chi^2$ where $\Delta\chi^2 = 4.6, 6.0, 9.2$ for 90, 95 and 99 % C. L., respectively.

The allowed regions for the contained events are, as expected, similar to the ones obtained in ref. [19]. The individual best fits now improve with respect to the analysis of the old data: $\chi^2_{\min} = 2.4/(8 \text{ d.o.f.})$ for the sub-GeV data ($\epsilon_\nu = 0.196$ and $\epsilon'_\nu = 0.013$) and to $\chi^2_{\min} = 6.4/(8 \text{ d.o.f.})$ for the multi-GeV sample ($\epsilon_\nu = 0.689$ and




Figure 3: Allowed regions for ϵ_ν and ϵ'_ν for the combination of the SUPER-KAMIOKANDE data sets: (a) the binned contained events are combined with total (unbinned) up-going events; (b) binned contained and up-going events. The best fit points for each case are indicated by stars. The shaded area refers to the 90% C.L. while the contours stand for 95% and 99% C.L.

$\epsilon'_\nu = 0.284$). The combination of the two sets of contained events leads to allowed regions which are analogous to the ones reported in the ref. [19] and which are not reproduced again here. The best fit point corresponds to $\epsilon_\nu = 0.95$ and $\epsilon'_\nu = 0.084$ with $\chi^2_{\min} = 9.3/(18 \text{ d.o.f.})$. The goodness of the fit to the contained events in the FC-neutrino interaction scenario can be understood since the suppression of the expected event rates for contained events is the same for sub-GeV and multi-GeV samples. The SUPER-KAMIOKANDE collaboration has also measured the energy dependence of the up-down asymmetry for contained events [10] and this clearly indicates a strong energy dependence of the asymmetry for muon-like events in the momentum range $0.2 \text{ GeV} < p_\mu < 2 \text{ GeV}$. The asymmetry is consistent with zero at low momentum but significantly deviates from the expectation in the SM at higher momenta. One may naively expect that since the FC conversion mechanism is energy-independent it could be in contradiction with this measurement. However, one must notice that the average angle between the directions of the final-state lepton and the incoming neutrino ranges from 70° at 200 MeV to 20° at 1.5 GeV, so that at low momenta the possible asymmetry of the neutrino flux is largely washed out. In figure 2 we plot, together with the SUPER-KAMIOKANDE data, the momentum behaviour of the asymmetry in the FC-neutrino interaction scenario calculated for the best fit point to the contained event sample. As seen in the figure the agreement is excellent.

Data	d.o.f.	χ_{minFC}^2	χ_{minOsc}^2
sub-GeV	8	2.4	2.4
multi-GeV	8	6.4	6.3
contained	18	9.3	8.8
stopping- μ	3	1.	1.3
through-going- μ	8	10.3	10.4
contained + total up- μ	10	20.4	9.6
contained + angular up- μ	33	44.	23.5

Table 1: χ_{min}^2 obtained for several data combinations in the framework of FC- ν interactions as compared to the case of the neutrino vacuum-oscillation scenario.

In figure 1, we also show the regions which are allowed by the up-going muons samples of SUPER-KAMIOKANDE. Panel (c) stands for stopping muons and panel (d) for the through-going sample. In the case of stopping muons, we see that, analogously to the contained events, the allowed region lies in the sector of the plane where the average survival probability is of the order of a half, which is what appears to be needed for explaining the data. Instead, in the case of through-going muons, the experimental data do not show such a strong reduction with respect to the theoretical calculations, and therefore the allowed region lies in the upper-left corner of the parameter space, which refers to a smaller transition probability. In both cases, the best fit point for each individual sample is good: $\chi_{min}^2 = 1/(3 \text{ d.o.f.})$ for stopping muons ($\epsilon_\nu = 0.756$ and $\epsilon'_\nu = 0.196$) and $\chi_{min}^2 = 10.3/(8 \text{ d.o.f.})$ for the through-going case ($\epsilon_\nu = 0.081$ and $\epsilon'_\nu = 0.260$). Both for the contained and for the up-going events, the best fits for each individual sample have the same level of statistical confidence as compared to the oscillation interpretation of the atmospheric neutrino data. This is shown in table 1, where we report the best fit values we obtain for the different data sets in the case of the FC- ν interactions scenario and in the case of the neutrino oscillation scenario[31].

The allowed regions can be qualitatively understood in the approximation of constant matter density. The conversion probability in this case is

$$P(\nu_\mu \rightarrow \nu_\tau) = \frac{4\epsilon_\nu^2}{4\epsilon_\nu^2 + \epsilon'_\nu{}^2} \sin^2\left(\frac{1}{2}\eta L\right), \tag{3.1}$$

where $\eta = \sqrt{4\epsilon_\nu^2 + \epsilon'_\nu{}^2} \sqrt{2}G_F n_f$. For $n_f = n_d \approx 3n_e$ and $\epsilon'_\nu < \epsilon_\nu$, the oscillation length in matter is given by

$$L_{osc} = \frac{2\pi}{\eta} \approx 1.2 \times 10^3 \left[\frac{2 \text{ mol/cc}}{n_e} \right] \left[\frac{1}{\epsilon_\nu} \right] \text{ km}. \tag{3.2}$$

From eq. (3.1) one can see that in order to have a relatively large transition probability, as required by the contained events and, also, by the stopping muons

events, the FC parameters are required to be in the region $\epsilon'_\nu \lesssim \epsilon_\nu$ and $\eta \gtrsim \pi/R_\oplus$. This last condition leads to a lower bound on ϵ_ν . The island in figure 1b corresponds to $\eta \sim \pi/R_\oplus$.

The combination of the different data sets in a single χ^2 -analysis is shown in figure 3. Panel (a) shows the combination of the full angular distribution of contained events with the total (unbinned) event rate of stop and through-going muons data, while panel (b) refers to the combination of all the angular distributions, including that of through-going muon events. In figure 3a the information brought by the higher energy data is effective at the normalization level, since no information about their angular dependence is included. In this case the allowed region is still relatively large although the description is already worse than in the oscillation case as can be seen by comparing the corresponding χ^2_{\min} (20.4/(10 d.o.f.) for FC as compared to 9.6/(10 d.o.f.) for the oscillation scenario). This worsening is due to the fact that in the FC scenario the transition probability is energy independent while the data shows a smaller conversion for the higher energy through-going muon events. As seen in figure 3b, when the angular information of both stopping and through-going muons is included in the data analysis, the description becomes even worse, mainly due to the angular distribution of the through-going data set. The allowed regions now form a set of isolated small 'islands'. The best fit point corresponds to $\epsilon_\nu = 0.57$ and $\epsilon'_\nu = 0.45$ which is acceptable only at the 90.5% confidence level ($\chi^2_{\min} = 44/(33$ d.o.f.)).

The behavior of the allowed regions can be understood by observing figure 4 where we show the angular distributions for the four cases: (a) sub-GeV; (b) multi-GeV; (c) stopping muons; (d) through-going muons. We show the distributions for the best fit point obtained from the combination of contained events with the total number of upward going muons $P_1 = (\epsilon_\nu, \epsilon'_\nu) = (0.17, 0.28)$ and for the best fit point obtained from the analysis of the full data set $P_2 = (0.57, 0.45)$. Although both points give a similar normalization to the up-going muon data samples, point P_1 gives a better description to the angular dependence of the contained events, but it does not describe well the zenith angle distribution of the through-going muon events. As commented above, such point correspond to an effective FC-oscillation length of the order of the Earth radius. In this case we can see the imprints of the ‘‘oscillatory’’ sine behavior in the expected angular distribution of the up-going muon events. Such behavior, however, does not appear to be present in the SUPER-KAMIOKANDE data, leading to a worse overall fit. In the case of multi-GeV contained events, this oscillatory behavior is averaged out due to the smaller angular resolution in the data and point P_1 can give a good description of the data. On the other hand, point P_2 gives a worse description of the contained events but fits better the shape of upward going muon data, with the exception of the last three angular bins of the through-going sample, where it does not produce a sufficient amount of through-going muons at angles $0 < \theta < 20^\circ$ below the horizon.




Figure 4: Best-fit zenith angle distributions in the massless-neutrino FC scenario. The thick-solid lines correspond to the calculation in absence of new physics. The dotted lines correspond to the best fit point obtained by the analysis of the contained events combined with total (unbinned) up-going events. The thin-solid line is for the best fit point of the combined analysis of contained and up-going muon events. The 52 kTy SUPER-KAMIOKANDE data are indicated by crosses.

4. Conclusions

In summary, in this paper we have re-analyzed the possibility of explaining the atmospheric neutrino anomaly by means of $\nu_\mu \rightarrow \nu_\tau$ conversion induced by flavour-changing neutrino-matter interaction which can be effective during the neutrino propagation in the Earth. We extend the analysis of ref. [19] to the new set of SUPER-KAMIOKANDE data by including also the up-going muon samples. Our results show that that flavour changing ν_μ -matter interactions are able to describe the full set of data of SUPER-KAMIOKANDE on atmospheric neutrinos with a probability of 9.6 %. Thus they are ruled out as an explanation at the 90 %CL. The agreement between the data and the calculated events for the SUPER-KAMIOKANDE detector is good for the individual sets of data collected by SUPER-KAMIOKANDE, with a confidence level as good as for the oscillation hypothesis. However, when the data are combined together, in particular once the upward-going muon zenith-angle distribution is included in the analysis, the ν_μ oscillation provides a much better description. The worsening of the fit which occurs when the through-going muons sample is included is partly due to the fact that in the FC scenario the transition probability is energy independent while the data shows a smaller conversion for the higher energy up-through-going muon events. The ensuing result is that the expectations from FC-neutrino interaction for neutrinos arriving mainly at angles above 20 degrees below the horizon do not reproduce the experimental data. The accumulation of additional up-going events should therefore enable one to impose new model independent limits on the strength of the ϵ and ϵ' parameters, as can be foreseen by looking at figure 1(d).

For the time being it is cautious to say that, unaccompanied by mass-related oscillations, the FC hypothesis is disfavored by the data. However, they could still be there at some level, even if the data would admit a very good interpretation in terms of standard $\nu_\mu \rightarrow \nu_\tau$ oscillations. This is theoretically not an ad hoc assumption, since in many theoretical models neutrino masses naturally co-exist with FC-neutrino interactions. Another argument against FC neutrino conversions as the only source of explanation of the atmospheric anomaly is the required magnitude of the FC couplings: although it is technically possible to accommodate it (for example in models with broken R-parity [32]) these neutrino FC couplings would be accompanied by too large lepton flavour violating τ decays in models which respect an $SU_L(2)$ symmetry in FC couplings [33]. Last, but not least, one virtue of the above FC mechanism is that it can be also tested at future Long Baseline experiments. From eq. (3.1), using $n_e \sim 2$ mol/cc, we can predict that for $\epsilon \simeq \epsilon' \sim 1$ (0.1) the planned K2K experiment [34] should obtain $P(\nu_\mu \rightarrow \nu_\tau) \sim 0.35$ (0.004) while for MINOS [35] one finds $P(\nu_\mu \rightarrow \nu_\tau) \sim 0.75$ (0.04).

Acknowledgments

We thank R. Vazquez for providing us with the program to compute the muon energy loss. We are also grateful to S. Nussinov for pointing us out the limits from the neutrino neutral current measurements. M.C. G-G is grateful to the Instituto de Fisica Teorica from UNESP and to the CERN Theory division for their kind hospitality during her visits. This work was supported by Spanish DGICYT under grant PB98-0693 and by the European Union TMR network ERBFMRXCT960090.

References

- [1] L.V. Volkova, *Energy spectra and angular distributions of atmospheric neutrinos*, *Sov. J. Nucl. Phys.* **31** (1980) 784;
M. Honda, T. Kajita, K. Kasahara and S. Midorikawa, *Calculation of the flux of atmospheric neutrinos*, *Phys. Rev.* **D 52** (1995) 4985 [[hep-ph/9503439](#)];
V. Agrawal, T.K. Gaisser, P. Lipari and T. Stanev, *Atmospheric neutrino flux above 1 GeV*, *Phys. Rev.* **D 53** (1996) 1314 [[hep-ph/9509423](#)];
T.K. Gaisser and T. Stanev, *Path length distributions of atmospheric neutrinos*, *Phys. Rev.* **D 57** (1998) 1977 [[astro-ph/9708146](#)].
- [2] FREJUS. collaboration, K. Daum, *Determination of the atmospheric neutrino spectra with the Frejus detector*, *Z. Physik* **C 66** (1995) 417.
- [3] NUSEX collaboration, *Experimental study of atmospheric neutrino flux in the Nusex experiment*, *Europhys. Lett.* **8** (1989) 611.
- [4] KAMIOKANDE-II collaboration, *Observation of a small atmospheric ν_μ/ν_e ratio in kamiokande*, *Phys. Lett.* **B 280** (1992) 146; *Atmospheric ν_μ/ν_e ratio in the multigeV energy range*, *Phys. Lett.* **B 335** (1994) 237.
- [5] R. Becker-Szendy et al., *The ν_μ/ν_e content of the atmospheric flux*, *Phys. Rev.* **D 46** (1992) 3720.
- [6] SUPER-KAMIOKANDE collaboration, *Measurement of a small atmospheric ν_μ/ν_e ratio*, *Phys. Lett.* **B 433** (1998) 9 [[hep-ex/9803006](#)];
Study of the atmospheric neutrino flux in the multi-GeV energy range, *Phys. Lett.* **B 436** (1998) 33 [[hep-ex/9805006](#)].
- [7] SOUDAN-2 collaboration, *The atmospheric neutrino flavor ratio from a 3.9 fiducial kiloton-year exposure of SOUDAN 2*, *Phys. Lett.* **B 449** (1999) 137 [[hep-ex/9901024](#)].
- [8] M.C. Gonzalez-Garcia, H. Nunokawa, O.L. G. Peres, T. Stanev and J.W.F. Valle, *Update on atmospheric neutrinos*, *Phys. Rev.* **D 58** (1998) 033004 [[hep-ph/9801368](#)];
M.C. Gonzalez-Garcia, H. Nunokawa, O.L. G. Peres and J.W. F. Valle, *Active-active and active-sterile neutrino oscillation solutions to the atmospheric neutrino anomaly*, *Nucl. Phys.* **B 543** (1999) 3 [[hep-ph/9807305](#)];

- For an updated global analysis of the atmospheric neutrino data in terms of neutrino oscillations, see N. Fornengo, M.C. Gonzalez-Garcia and J.W.F. Valle, *Updated global analysis of the atmospheric neutrino data in terms of neutrino oscillations*, *Nucl. Phys. B* **580** (2000) 58 [[hep-ph/0002147](#)].
- [9] R. Foot, R.R. Volkas and O. Yasuda, *Comparing and contrasting the $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ solutions to the atmospheric neutrino problem with SUPER-KAMIOKANDE data*, *Phys. Rev. D* **58** (1998) 013006 [[hep-ph/9801431](#)];
 O. Yasuda, *Three flavor neutrino oscillation analysis of the SUPER-KAMIOKANDE atmospheric neutrino data*, *Phys. Rev. D* **58** (1998) 091301 [[hep-ph/9804400](#)];
 G.L. Fogli, E. Lisi, A. Marrone and G. Scioscia, *SUPER-KAMIOKANDE atmospheric neutrino data, zenith distributions and three-flavor oscillations*, *Phys. Rev. D* **59** (1999) 033001 [[hep-ph/9808205](#)];
 E.K. Akhmedov, A. Dighe, P. Lipari and A.Y. Smirnov, *Atmospheric neutrinos at SUPER-KAMIOKANDE and parametric resonance in neutrino oscillations*, *Nucl. Phys. B* **542** (1999) 3 [[hep-ph/9808270](#)].
- [10] SUPER-KAMIOKANDE collaboration, *Evidence for oscillation of atmospheric neutrinos*, *Phys. Rev. Lett.* **81** (1998) 1562 [[hep-ex/9807003](#)];
 See also the talk by K. Scholberg, for SUPERKAMIOKANDE collaboration, *Atmospheric neutrinos at SUPER-KAMIOKANDE*, [hep-ex/9905016](#).
- [11] V. Barger, J.G. Learned, S. Pakvasa and T.J. Weiler, *Neutrino decay as an explanation of atmospheric neutrino observations*, *Phys. Rev. Lett.* **82** (1999) 2640 [[astro-ph/9810121](#)].
- [12] M. Gasperini, *Testing the principle of equivalence with neutrino oscillations*, *Phys. Rev. D* **38** (1988) 2635;
 J. Pantaleone, A. Halprin and C.N. Leung, *Neutrino mixing due to a violation of the equivalence principle*, *Phys. Rev. D* **47** (1993) 4199 [[hep-ph/9211214](#)];
 A. Halprin, C.N. Leung and J. Pantaleone, *A possible violation of the equivalence principle by neutrinos*, *Phys. Rev. D* **53** (1996) 5365 [[hep-ph/9512220](#)].
- [13] S. Coleman and S.L. Glashow, *Cosmic ray and neutrino tests of special relativity*, *Phys. Lett. B* **405** (1997) 249 [[hep-ph/9703240](#)];
 S.L. Glashow, A. Halprin, P.I. Krastev, C.N. Leung and J. Pantaleone, *Comments on neutrino tests of special relativity*, *Phys. Rev. D* **56** (1997) 2433 [[hep-ph/9703454](#)].
- [14] S. Coleman and S.L. Glashow, *High-energy tests of Lorentz invariance*, *Phys. Rev. D* **59** (1999) 116008 [[hep-ph/9812418](#)].
- [15] SUPERKAMIOKANDE collaboration, *Measurement of the flux and zenith-angle distribution of upward through-going muons by SUPER-KAMIOKANDE*, *Phys. Rev. Lett.* **82** (1999) 2644 [[hep-ex/9812014](#)].
- [16] P. Lipari and M. Lusignoli, *On exotic solutions of the atmospheric neutrino problem*, *Phys. Rev. D* **60** (1999) 013003 [[hep-ph/9901350](#)].

- [17] G.L. Fogli, E. Lisi, A. Marrone and G. Scioscia, *Testing violations of special and general relativity through the energy dependence of $\nu_\mu \leftrightarrow \nu_\tau$ oscillations in the SUPER-KAMIOKANDE atmospheric neutrino experiment*, *Phys. Rev. D* **60** (1999) 053006 [[hep-ph/9904248](#)].
- [18] G.L. Fogli, E. Lisi, A. Marrone and G. Scioscia, *SUPER-KAMIOKANDE data and atmospheric neutrino decay*, *Phys. Rev. D* **59** (1999) 117303 [[hep-ph/9902267](#)].
- [19] M.C. Gonzalez-Garcia, M.M. Guzzo, P.I. Krastev, H. Nunokawa, O.L.G. Peres, V. Pleitez, J.W.F. Valle and R. Zukanovich Funchal, *Atmospheric neutrino observations and flavor changing interactions*, *Phys. Rev. Lett.* **82** (1999) 3202 [[hep-ph/9809531](#)].
- [20] J.W.F. Valle, *Resonant oscillations of massless neutrinos in matter*, *Phys. Lett. B* **199** (1987) 432;
M. Fukugita and T. Yanagida, *Neutrino oscillation in matter induced by charged scalar particles*, *Phys. Lett. B* **206** (1988) 93;
M.M. Guzzo, A. Masiero and S.T. Petcov, *On the MSW effect with massless neutrinos and no mixing in the vacuum*, *Phys. Lett. B* **260** (1991) 154;
E. Roulet, *Mikheyev-Smirnov-Wolfenstein effect with flavor-changing neutrino interactions*, *Phys. Rev. D* **44** (1991) 935;
V. Barger, R.J. N. Phillips and K. Whisnant, *Solar neutrino solutions with matter enhanced flavor changing neutral current scattering*, *Phys. Rev. D* **44** (1991) 1629;
P.I. Krastev and J.N. Bahcall, *Finite solutions to the solar neutrino problem*, [hep-ph/9703267](#);
S. Bergmann, *The solar neutrino problem in the presence of flavor-changing neutrino interactions*, *Nucl. Phys. B* **515** (1998) 363 [[hep-ph/9707398](#)];
E. Ma and P. Roy, *Neutrino oscillation data versus three light neutrino flavors*, *Phys. Rev. Lett.* **80** (1998) 4637 [[hep-ph/9706309](#)].
- [21] J. Schechter and J.W.F. Valle, *Neutrino masses in $SU(2) \times U(1)$ theories*, *Phys. Rev. D* **22** (1980) 2227.
- [22] R.N. Mohapatra and J.W. F. Valle, *Neutrino mass and baryon number nonconservation in superstring models*, *Phys. Rev. D* **34** (1986) 1642;
D. Wyler and L. Wolfenstein, *Massless neutrinos in left-right symmetric models*, *Nucl. Phys. B* **218** (1983) 205.
- [23] L.J. Hall, V.A. Kostelecky and S. Raby, *New flavor violations in supergravity models*, *Nucl. Phys. B* **267** (1986) 415;
Y. Okada, *Flavor physics in the supergravity model*, [hep-ph/9809297](#).
- [24] A.M. Dziewonski and D.L. Anderson, *Preliminary Reference Earth Model (PREM)*, *Phys. Earth and Planet Interiors*, **25** (1981) 297.
- [25] J.N. Bahcall and P.I. Krastev, *Does the sun appear brighter at night in neutrinos?*, *Phys. Rev. D* **56** (1997) 2839 [[hep-ph/9706239](#)].

- [26] W. Lohmann, R. Kopp and R. Voss, *Energy loss of muons in the energy range 1–10000 GeV*, CERN Yellow Report EP/85-03.
- [27] E. Zas, F. Halzen, R.A. Vazquez, *High-energy neutrino astronomy: horizontal air shower arrays versus underground detectors*, *Astropart. Phys.* **1** (1993) 297.
- [28] P. Lipari and M. Lusignoli, *Comparison of $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ oscillations as solutions of the atmospheric neutrino problem*, *Phys. Rev.* **D 58** (1998) 073005 [[hep-ph/9803440](#)].
- [29] G.L. Fogli and E. Lisi, *On the atmospheric neutrino anomaly and its statistical significance*, *Phys. Rev.* **D 52** (1995) 2775 [[hep-ph/9504287](#)];
G.L. Fogli, E. Lisi, A. Marrone and G. Scioscia, *SUPER-KAMIOKANDE atmospheric neutrino data, zenith distributions and three-flavor oscillations*, *Phys. Rev.* **D 59** (1999) 033001 [[hep-ph/9808205](#)].
- [30] M. Nakahata, *Recent results from SUPER-KAMIOKANDE*, talk at the *Sixth international workshop on topics in astroparticle and underground physics, TAUP99*, Paris, September 1999.
- [31] N. Fornengo, M.C. Gonzalez-Garcia and J.W. F. Valle, *Updated global analysis of the atmospheric neutrino data in terms of neutrino oscillations*, *Nucl. Phys.* **B 580** (2000) 58 [[hep-ph/0002147](#)].
- [32] L.J. Hall and M. Suzuki, *Explicit R-parity breaking in supersymmetric models*, *Nucl. Phys.* **B 231** (1984) 419;
G.G. Ross and J.W. F. Valle, *Supersymmetric models without R-parity*, *Phys. Lett.* **B 151** (1985) 375;
J. Ellis, G. Gelmini, C. Jarlskog, G.G. Ross and J.W.F. Valle, *Phenomenology of supersymmetry with broken R-parity*, *Phys. Lett.* **B 150** (1985) 142.
- [33] S. Bergmann, Y. Grossman and D.M. Pierce, *Can lepton flavor violating interactions explain the atmospheric neutrino problem?*, *Phys. Rev.* **D 61** (2000) 053005 [[hep-ph/9909390](#)].
- [34] C. Yanagisawa et al., in *Physics beyond the standard model: from theory to experiment*, I. Antoniadis, L.E. Ibañez and J.W.F. Valle eds., World Scientific, Singapore 1998.
- [35] D. Michael et al., *Nucl. Phys.* **66** (*Proc. Suppl.*) (1998) 432.