

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Anti-leukemic activity of microRNA-26a in a chronic lymphocytic leukemia mouse model

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1654404> since 2017-12-06T13:20:36Z

Published version:

DOI:10.1038/onc.2017.269

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

L D'Abundo, E Callegari, A Bresin, A Chillemi, B K Elamin, P Guerriero, X Huang, E Saccenti, E M A A Hussein, F Casciano, P Secchiero, G Zauli, G A Calin, G Russo, L J Lee, C M Croce, G Marcucci, S Sabbioni, F Malavasi & M Negrini

Paper: Anti-leukemic activity of microRNA-26a in a chronic lymphocytic leukemia mouse model

ONCOGENE, 36(47), 2017, pp: 6617–6626

DOI: 10.1038/onc.2017.269

The publisher's version is available at:

<https://doi.org/10.1038/onc.2017.269>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1654404>

Anti-leukemic activity of microRNA-26a in a chronic lymphocytic leukemia mouse model

Lucilla D'Abundo^{1**}, *Elisa Callegari*^{1**}, *Antonella Bresin*^{1,2}, *Antonella Chillemi*^{3,4}, *Bahaeldin K. Elamin*^{5,6}, *Paola Guerriero*¹, *Xiaomeng Huang*^{7,8,9}, *Elena Saccenti*¹⁰, *Enaam M.A.A. Hussein*^{1,11}, *Fabio Casciano*¹², *Paola Secchiero*¹², *Giorgio Zauli*¹², *George A. Calin*¹³, *Giandomenico Russo*², *Ly James Lee*^{7,8}, *Carlo Maria Croce*⁹, *Guido Marcucci*¹⁴, *Silvia Sabbioni*¹⁵, *Fabio Malavasi*^{3,4}, *Massimo Negrini*^{1*}

- 1 Department of Morphology, Surgery and Experimental Medicine, University of Ferrara, Ferrara, Italy
- 2 Laboratorio di Oncologia Molecolare, Istituto Dermopatico dell'Immacolata, IDI-IRCCS, Rome, Italy
- 3 Laboratory of Immunogenetics, Department of Medical Sciences, University of Torino, Torino 10126, Italy
- 4 Laboratorio di Immunologia dei Trapianti, Città della Salute e della Scienza, Torino 10126, Italy
- 5 Microbiology Department, Faculty of Medical laboratory Sciences, University of Khartoum, Sudan
- 6 Microbiology Department, College of Medicine, University of Bisha, Saudi Arabia
- 7 Nanoscale Science and Engineering Center for Affordable Nanoengineering of Polymeric Biomedical Devices, The Ohio State University, Columbus, Ohio, USA
- 8 Department of Chemical and Biomolecular Engineering, The Ohio State University, Columbus, Ohio, USA
- 9 Comprehensive Cancer Center, The Ohio State University, Columbus, Ohio, USA
- 10 Hematology Section, Department of Medical Sciences, University of Ferrara, University Hospital Arcispedale S. Anna, Ferrara, Italy
- 11 Hematology and Immunohematology Department, Faculty of Medical laboratory Sciences, University of Khartoum, Sudan
- 12 Department of Morphology, Surgery, Experimental Medicine and LTTA Centre, University of Ferrara, Ferrara, Italy
- 13 Department of Experimental Therapeutics, Division of Cancer Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX
- 14 Gehr Family Center For Leukemia Research Hematologic Malignancies Institute City of Hope, Duarte, CA, 90010, USA
- 15 Department of Life Sciences and Biotechnology, University of Ferrara, Ferrara, Italy

** LD'A and EC contributed equally to this work

***Corresponding Authors**

Massimo Negrini, PhD
University of Ferrara
Dept of Surgery, Morphology and Experimental Medicine
Via Luigi Borsari, 46
44121 Ferrara, Italy
Phone: +39 0532 455 399
Email: ngm@unife.it

Lucilla D'Abundo, PhD
University of Ferrara
Dept of Surgery, Morphology and Experimental Medicine
Via Luigi Borsari, 46
44121 Ferrara, Italy
Phone: +39 0532 455 871
Email: dbnlll@unife.it

Running Title: Anti-leukemic activity of microRNA-26a in CLL

Abstract

The dysregulation of microRNAs (miRNAs) plays an important role in the pathogenesis of chronic lymphocytic leukemia (CLL). The E μ -*TCL1* transgenic mouse, which develops a leukemia similar to the aggressive form of human B-CLL, is a valuable model for testing novel therapeutic approaches. Here, we adopted this model to investigate the therapeutic effect of miR-26a, miR-130a, and anti-miR-155.

To improve delivery of miRNA molecules into CLL cells, we developed novel anti-CD38-conjugated lipid nanoparticles, which were highly effective in delivering miRNA molecules into leukemic cells. Short and long-term experiments showed that miR-26a, miR-130a, and anti-miR-155 increased the levels of apoptosis after *in vitro* or *in vivo* treatment. In particular, *miR-26a* was the most effective in reducing leukemic cell expansion. At the end of long-term treatment, apoptosis was readily detectable by analysis of cleaved PARP and caspase-7. This effect could be directly related to miR-26a, as shown by the significant downregulation of its proven targets, the cyclin-dependent kinase 6 and Mcl1.

This study provides the first evidence that miR-26a could elicit an *in vivo* anti-leukemic activity through an increase in cell apoptosis. It also provides a novel *in vivo* approach that improves the efficiency and specificity of miRNA delivery into CD38+ leukemic cells.

Keywords: miRNA, CLL, Nanoparticles, CD38, miRNA delivery.

1 Introduction

2 Chronic lymphocytic leukemia (CLL) is the most common B-cell malignancy in Western countries.
3 CLL lymphocytes resemble memory B cells with a mature phenotype (1). This leukemia is
4 characterized by the accumulation of CD5+ B lymphocytes in the blood, spleen, liver, lymph nodes,
5 and bone marrow. Despite its morphological homogeneity, CLL is a clinically heterogeneous
6 disease (2-4) with a number of adverse factors that affect the clinical course, including stage(5);
7 CD38 positivity; unmutated variable region of the immunoglobulin heavy chain gene (*IGHV*) (6);
8 ZAP70 positivity (7); karyotype aberrations (8); and TP53, NOTCH1, SF3B1, and other gene
9 mutations (9, 10). Despite the progress in chemoimmunotherapy, CLL remains an incurable
10 disease (11).

11 To help decipher the pathogenic mechanisms of disease and evaluate the efficacy and
12 mechanisms of action of novel therapies, researchers have developed mouse models that
13 resemble human CLL (12, 13). The E μ -*TCL1* transgenic (*TCL1*-tg) mouse develops a leukemia that is
14 highly similar to the aggressive form of human CLL (14-16). An important feature of *TCL1*
15 overexpression is that it exhibits 100% disease penetrance. At the age of 16–20 months, these
16 mice show a B-CLL-like disease characterized by spleno- and hepatomegaly associated with high
17 counts of white blood cells. Leukemias are characterized by clonal expansion of B cells with a
18 B220+/IgM+/CD5+ immunophenotype, unmutated *IGHV*, increased proliferation, and enhanced
19 AKT phosphorylation. Tumor cells in *TCL1*-tg mice have wild-type (WT) p53 and initially respond to
20 fludarabine treatment (17). Notably, leukemic cells from a *TCL1*-tg donor can be transplanted by
21 intraperitoneal (IP) or intravenous injection into syngeneic WT or immunodeficient mice (e.g.,
22 SCID) to accelerate the disease course and generate genetically homogeneous populations of
23 leukemic cells, which better allows for systematic comparison of novel therapies.

24 MicroRNAs (miRNAs) play a central role in the pathogenesis of CLL. They regulate gene expression
25 at the post-transcriptional level by targeting messenger RNA for degradation or translational
26 inhibition and thus can modulate several biological processes. As a consequence of their
27 deregulation, miRNAs may act as oncogenes or tumor suppressors (18). The discovery of the
28 involvement of miRNAs in human cancer originated from studies of CLL (19). miR-15a and miR-16-
29 1 were found within the minimal region of deletion at chromosome 13q, the most frequent
30 genetic alteration found in human CLL. Deletion of miR-15a/miR-16-1 in a knockout mouse model
31 was shown to confer predisposition to the development of an indolent form of leukemia similar to
32 human CLL (20). BCL2 was a target of these miRNAs; thus, their loss eliminated control over BCL2
33 expression (21). It is notable that venetoclax, a highly active and recently approved drug against
34 CLL, was designed to inhibit BCL2 activity (22). A prognostic signature consisting of a panel of
35 aberrantly expressed miRNAs has been identified in CLL (23). Other important miRNAs in human
36 CLL include miR-34a, which, being an effector of the p53 protein, is strongly downregulated in
37 17p-CLL cases, or miR-181b, which is downregulated during disease progression (24-26). In a more
38 recent comprehensive study, we found that a number of miRNAs are deregulated in CLL (27).
39 miRNAs were differentially expressed between CLL cells and mature antigen-experienced B cells,
40 or correlated to specific pathological features (*IGHV* somatic mutations or specific cytogenetic
41 aberrations) or clinical parameters, such as time to first treatment (27). These studies established
42 the groundwork from which miRNA-based therapies, either by restoring or repressing miRNA
43 activity, could be designed and tested. We have already investigated miR-181b as a potential
44 therapeutic molecule against leukemias of the *TCL1*-tg mouse model (28). The study proved that
45 this miRNA induced a measurable anti-leukemic effect. However, an improvement in delivery
46 methods was suggested.

1 Here, we used the E μ -*TCL1FL* mouse model (15) to test the anti-leukemic activity of various miRNA
2 mimics or anti-miRNA oligonucleotides (AMOs). To increase *in vivo* delivery, we developed a
3 lipopolyplex formulation that included anti-CD38 conjugated lipid nanoparticles (CD38-NPs), which
4 considerably increased the specificity for hCD38+ leukemic cells of this mouse model. The choice
5 of CD38 as a molecule to target CLL cells was guided by its known importance in CLL (29), as well
6 as its structural characteristics (30) and the evidence of its antibody-mediated internalization (31).

7

8 **Results**

9 **Leukemic cells of the E μ -*TCL1FL* transgenic mouse are CD38+.** Leukemic cells of the E μ -*TCL1FL*
10 transgenic mouse or transplanted into syngeneic FVB mice are characterized by B220+/CD5dim
11 surface markers (28). Here, we evaluated the expression of CD38. To this end, the splenic
12 lymphocyte population from transgenic or transplanted FVB mice with a high level of disease was
13 first sorted on the basis of physical parameters, and the B220+/CD5dim leukemic population was
14 identified; finally, the lymphocytes within the B220+/CD5dim gate were analyzed for CD38
15 expression, showing that more than 98% of the cells were positive (**Figure 1**). This analysis not only
16 confirmed that the *TCL1*-tg model exhibits an immunophenotype typical of an aggressive form of
17 B-CLL characterized by high expression of CD38, but it also provided a way to specifically target
18 leukemic cells.

19 **Lipid nanoparticles conjugated with anti-CD38 antibodies can efficiently deliver miRNAs into CLL**
20 **cells.** On the basis of this evidence, we developed lipid nanoparticles conjugated with anti-CD38
21 antibodies (CD38-NPs) to improve the efficiency of *in vivo* miRNA delivery into leukemic cells. The
22 lipid NP-miRNAs were synthesized as described (32).

23 We investigated the efficiency of CD38-NPs both *in vitro* and *in vivo*. *In vitro*, we tested whether
24 CD38-NPs could deliver mature miRNAs into murine *TCL1*-tg leukemic cells. We transfected cells
25 with miR-181b mimics, miRNA that we had previously investigated (28). Twenty-four hours after
26 transfection, cells were collected for RNA extraction and the levels of miR-181b measured
27 (**Supplementary Figure 1**). These *in vitro* results indicated that CD38-NPs could efficiently deliver
28 small RNA into leukemic cells and were approximately 2-fold more efficient than unconjugated
29 NPs.

30 The *TCL1*-tg mouse represented an excellent model for testing *in vivo* delivery efficiency. We
31 quantitatively investigated the delivery of 100 μ g of miR-181b in different organs of FVB-*TCL1*
32 transplanted mice. Mice were treated when they exhibited a lymphocyte disease burden of 30-
33 50% as the percentage of leukemic cells in peripheral blood. We tested CD38-NPs, non-conjugated
34 NPs, or Jet-PEI, a formulation used in our previous study (28), in 4 mice for each condition. The
35 nanoparticles were administered by IP injection. After 24 hours, mice were sacrificed and organs
36 collected. Total RNA was isolated from liver, splenocytes, heart, kidneys, thyroid, lungs, muscle,
37 stomach, intestine, brain, and bladder and the expression levels of miR-181b were quantified by
38 digital PCR (**Figure 2A**). Spleen was the most efficiently targeted organ by CD38-NPs, followed by
39 liver. Compared with the physiological levels in these organs, the amount of miR-181b increased
40 more than 300-fold following CD38-NP-mediated delivery. In splenocytes, CD38-NPs induced an
41 18-fold increase compared with Jet-PEI and a 3-fold increase compared with non-conjugated NPs
42 (**Figure 2B**). All other organs exhibited increased expression levels of miR-181b, generally at lower
43 levels and independent to the type of NP.

44 The greater delivery mediated by CD38-NPs in spleen and liver of diseased animals was likely
45 attributable to the presence of a high number of CD38+ cells. This conclusion was supported by

1 the finding that no significant difference of miR-181b in spleen and liver was observed between
2 CD38-NPs and non-conjugated NPs delivery efficacy when experiment was performed in FVB WT
3 mice (**Supplementary Figure 2**). These findings indicated that CD38-NPs were capable of
4 improving efficiency of delivery and specificity of targeting for CD38+ CLL cells.

5 **miR-26a, miR-130a, and anti-miR-155 are selected as potential anti-leukemic molecules.** After
6 the delivery efficacy of CD38-NPs was verified *in vivo* and *in vitro*, we tested the potential anti-
7 leukemic effects of a group of selected miRNAs/anti-miRNAs.

8 From the results of our previous microRNAome study in human CLL (27), we selected a group of
9 miRNAs that were deregulated in human CLL. Herein, we tested miRNA mimics miR-15, miR-16,
10 miR-26a, miR-125a, miR-130a, and miR-34a or AMOs of miR-21, miR-155, and miR-130a. A
11 scrambled RNA oligo was used as a negative control.

12 First, we investigated the *in vitro* biological activity on leukemic splenocytes isolated from the
13 spleen of E μ -*TCL1FL* transgenic mice. Leukemic splenocytes were transfected by the use of CD38-
14 NPs and 48 hours after transfection, apoptotic activity was assessed by Annexin V assay (**Figure**
15 **3A, supplementary figure 3**). The highest apoptotic activity was detected in splenocytes treated
16 with miR-26a, miR-125a, anti-miR-155 and miR-130a ($p < 0.05$). The efficiency of transfection of
17 mimics/AMOs was assessed and confirmed in all the samples by analyzing the expression levels of
18 each miRNA (**Figure 3B**).

19 Next, for *in vivo* studies, we first evaluated if the CD38-NPs could also mediate a more effective
20 functional outcome than non-conjugated NPs. Given the high pro-apoptotic activity demonstrated
21 *in vitro*, we selected the miR-26a for measuring *in vivo* biological effects. 100 μ g of miR-26a
22 mimics was delivered into mice with a high disease burden (30-50% TCL1+ cells in peripheral
23 blood) by using either CD38-NPs or non-conjugated NPs (3 mice in each group). As control, 3 mice
24 with a similar disease burden were treated with a scrambled RNA oligo. After 48 hours from
25 treatment (IP injection), mice were sacrificed and splenocytes isolated to analyze the biological
26 effects. As shown in **Figure 4**, in comparison with non-conjugated NPs, the use of CD38-NPs
27 induced a more effective down-regulation CDK6, a target of miR26a, and a clearer activation of
28 apoptosis, detectable by cleavage of PARP.

29 Based on these results, we evaluated a number of miRNA mimics/AMOs for their *in vivo* biological
30 activity using CD38-NPs as delivery vehicles. Transplanted mice with a high disease burden (30-
31 50% TCL1+ cells in peripheral blood) were treated with 100 μ g of different miRNA mimics or AMOs
32 (at least 3 mice in each group) by using CD38-NPs; as control, 4 mice with a similar disease burden
33 were treated with scrambled miRNA. After 48 hours of a single treatment (IP injection), mice were
34 sacrificed and splenocytes isolated to analyze the effects of miRNAs on apoptosis by the Annexin V
35 assay. As shown in **Figure 5A**, treatment with miR-26a or miR-130a mimics or with anti-miR-155
36 oligonucleotides induced a significant increase in apoptotic cells compared with control ($p < 0.05$),
37 while several other miRNAs did not exhibit significant differences. Notably, CD38-NP-Empty
38 treated mice exhibited a negligible effect in comparison with control ($p = 0.6$). The efficiency of
39 delivery of mimic/AMOs in splenocytes was confirmed by quantitative PCR in all samples (**Figure**
40 **5B**). These results revealed the potential pro-apoptotic activity of miR-26a, miR-130a, and anti-
41 miR-155 against leukemic cells.

42 **miR-26a reduces the expansion of leukemic cells of E μ -*TCL1* mice.** Given the observed pro-
43 apoptotic effects after short-term treatments *in vitro* and *in vivo*, we sought to assess the
44 potential anti-leukemic activity of miR-26a, miR-130a, or anti-miR-155 after 3 weeks of treatment.
45 We performed the experiments by using FVB WT mice transplanted with E μ -*TCL1FL* leukemic cells

1 as we have previously described (28). Syngeneic transplantations were performed in 6-week-old
2 mice by IP injection of 5×10^5 splenocytes, collected from an E μ -*TCL1*FL transgenic mouse with
3 advanced disease. We monitored the progression of leukemia in transplanted syngeneic mice over
4 time by quantitatively measuring the increase of the human *TCL1* transgene, using a biomarker of
5 transplanted leukemic cells, in DNA isolated from peripheral blood with digital PCR.

6 Mice were enrolled for treatment at about 8 weeks after transplantation, when leukemic cells
7 reached 15-30% of peripheral blood lymphocytes. At this stage, mice were assigned to 5 groups of
8 treatment with the same schedule of administration (IP injection of 100 μ g of miRNA mimics or
9 AMOs 3 times a week for 3 weeks): (1) miR-26a, (2) miR-130a, (3) anti-miR-155, (4) scrambled
10 oligo control, and (5) CD38-NP-Empty nanoparticles. At the beginning of treatment, each group of
11 mice exhibited the same or a very similar arithmetic mean and standard deviation of leukemic
12 burden.

13 Before starting the treatments and 48 hours after the last treatment, we measured by digital PCR
14 and FACS analysis the percentage of B220+/CD5+ leukemic cells present in the blood. Results of
15 the two assays were in agreement. The differences between percentages of leukemic cells before
16 and after treatment are shown in **Figure 6**. miR-26a treatment was the most effective in reducing
17 the accumulation/expansion of leukemic cells in blood. The other molecules also induced a
18 reduction of the leukemic cell burden in blood, albeit to a significantly less extent (**Figure 6**).

19 **miR-26a promotes apoptosis in CLL cells.** Given the significant anti-leukemic effect of miR-26a, we
20 investigated the effect of miR-26a on leukemic splenocytes after 3 weeks of *in vivo* treatment.

21 The activation of apoptosis was clearly detectable by analysis of poly(ADP-ribose) polymerase
22 (PARP) and caspase-7: the appearance of the 85-kD fragment (cleaved PARP), together with the
23 strong increase in cleavage of caspase-7 were seen only in mice treated with CD38-NP-miR-26a,
24 thus indicating the induction of apoptosis (**Figure 7A**).

25 To confirm that the apoptotic effect could be attributable to miR-26a activity, we analyzed the
26 levels of two of its targets, Mcl1 and cyclin-dependent kinase 6 (Cdk6). Treatment with CD38-NP-
27 miR-26a induced their downregulation, thus confirming the molecular activity of the delivered
28 miR-26a (**Figure 7B**).

29

30 Discussion

31 The aim of this study was to demonstrate that miRNA-based therapies may have a place in the
32 treatment of CLL, a human leukemia that is still incurable despite recent multiple successes. Our
33 approach was based on evidence related to aberrant expressions of miRNA in CLL and on
34 numerous observations (19, 23, 25, 27, 33).

35 To address the working hypothesis, we adopted the E μ -*TCL1* mouse model, which develops a
36 leukemia that is highly similar to an aggressive form of human CLL (15, 16). Using this model, we
37 previously reported the anti-leukemic activity of miR-181b mimics (28). Here, we assayed miR-15a,
38 miR-16-1, miR-26a, miR-125a, miR-130a, miR-34a, ant-miR-130a, anti-miR-21, and anti-miR-155.
39 Among these, miR-26a, miR-130a, and anti-miR-155 induced apoptosis in leukemic cells both *in*
40 *vitro* and *in vivo* in short-term assays. On the basis of these results, we tested the ability of miR-
41 26a, miR-130a, and anti-miR-155 to counteract the *in vivo* accumulation/expansion of leukemic
42 cells of the E μ -*TCL1* mouse model.

43 miR-155 is a well-known oncomiR (34-39). In human CLL, a high level of miR-155 was associated
44 with various adverse prognostic factors (27, 40-43). Overexpression of miR-155 in transgenic mice

1 was shown to induce polyclonal B-cell expansion (34). The use of anti-miR-155 molecules has been
2 reported to significantly decrease *in vivo* tumor growth of BCWM1 cells derived from a patient
3 with Waldenstrom macroglobulinemia (44). The role of miR-130a in tumorigenesis is, on the other
4 hand, controversial. Various reports have indicated either tumor-suppressive (45-51) or oncogenic
5 (52-56) activity, depending on the experimental settings. In human CLL, miR-130a was shown to
6 repress a survival autophagic pathway by targeting ATG2B and DICER1 (57). No animal models are
7 available to examine the physiological or pathological function of miR-130a. Similarly, contrasting
8 results of dysregulated miR-26a have been reported in different tumor types (58-60). For example,
9 it was shown that miR-26a could facilitate glioblastoma formation *in vivo* (58, 61), but it was also
10 shown that low miR-26a expression conferred a shorter overall survival in liver cancer patients
11 (62) and that its delivery through an adeno-associated viral vector could achieve a therapeutic
12 effect on a MYC-induced liver cancer mouse model (59). In human CLL, a tumor suppressive
13 function of miR-26a was supported by evidence that its low expression was associated with a
14 shortened time from diagnosis to first treatment (27), a clinical feature associated with poor
15 prognosis.

16 Despite the limited evidence linking miR-26a to CLL pathogenesis, this miRNA exhibited the
17 strongest activity against the accumulation/expansion of leukemic cells by enhancing apoptosis.
18 Several studies have previously documented that miR-26a is able to target and down-modulate
19 several protein-coding gene targets, including CDK6, cyclins D2/E2, and Mcl-1, in different tumor
20 cells (59, 63-65). Here, we found that a strong apoptotic effect was detectable in the spleen of
21 treated mice and proof of miR-26a activity was confirmed by the downregulation of two of its
22 known targets, Mcl1 and Cdk6. The concomitant downregulation of the anti-apoptotic Mcl1 and
23 the cell cycle-promoting Cdk6 proteins suggests not only that miR-26a can promote apoptosis, but
24 that it can also inhibit CLL cell proliferation.

25 A second important result of this study was the development of nanoparticles conjugated with
26 anti-CD38 antibodies that could efficiently deliver miRNA or AMOs into CD38+ cells, which
27 typically characterize the most aggressive forms of CLL (29) and other hematological diseases such
28 as myeloma. In fact, efficient, specific, and safe delivery of miRNA mimics or AMOs is a major
29 challenge in miRNA-based therapeutic applications (66, 67).

30 Currently, methods for miRNA systemic delivery are mainly designed to target the liver (68, 69)
31 and they may use either viral (59, 70, 71) or non-viral systems (72-76). Significant therapeutic
32 effects in murine liver cancer models have been described (59, 70, 77). More difficult, however, is
33 the systemic delivery of miRNA mimics to other organs or tissues. Regarding hematopoietic cells,
34 *in vivo* delivery of antagomiR-126 was achieved in acute myeloid leukemia subpopulations by
35 using lipopolyplex nanoparticles conjugated with transferrin or antibody (anti-CD45.2) (72). The
36 chemical and physical characteristics of this formulation were shown to bypass hepatic uptake and
37 ultimately achieve better delivery to hematopoietic organs (32). An analogous system based on
38 stable nucleic acid lipid vesicles conjugated with transferrin was developed to target multiple
39 myeloma cells expressing transferrin receptors (78, 79).

40 To develop a method for *in vivo* delivery to CLL cells, here we produced lipid nanoparticles, based
41 on the formulation by Huang *et al* (32), conjugated with an anti-CD38 antibody, whose antigen is
42 present on the surface of leukemic cells of E μ -*TCL1* mouse model. Delivery to leukemic cells was
43 superior to that of either non-conjugated nanoparticles or JetPEI, the nanosystem that we used in
44 our previous report (28). The presence of the anti-CD38 antibody conferred specificity of action
45 and also led to internalization of the nanoparticle complex in targeted spleen cells. The
46 internalization was previously shown by using a panel of rat and mouse-anti CD38 mAbs (31). The

1 results indicated that the adopted methodology is efficient in transferring a cargo of different
2 miRNAs to leukemic cells. This competence was maintained *in vivo*. Spleen and liver were the most
3 efficiently targeted organs by CD38-NPs. A small increase of miR-181b was also detectable in other
4 organs, especially when non-conjugated nanoparticles were used. Since diseased mice are
5 characterized by spleno- and hepatomegaly, these results strongly suggested that the CD38-NPs
6 could increase delivery efficiency to these organs because they accumulate CD38+ CLL cells. The
7 characteristics of CD38 targeting in terms of safety were recently confirmed *in vivo* in mAb-
8 mediated therapy of human myeloma (80).

9 The main result of this study is that miR-26 exhibits a clear apoptotic effect on leukemic cells and a
10 significant reduction in leukemic cell expansion, suggesting its consideration for future CLL
11 therapy, possibly in combination with presently used therapeutic approaches. Moreover, we
12 provide a novel delivery method, which improves the efficiency and specificity of miRNA delivery
13 into CD38+ CLL cells. In humans, this approach could help target not only CLL cells in lymphatic
14 organs, but also other CD38+ B-cell malignancies, including various types of lymphoma, plasma
15 cell-derived neoplasms, and acute myeloid leukemia of the t(6;9) subtype (81-86).

16

17 **Materials and Methods**

18 **Cell cultures and transfections.** Mouse splenocytes were freshly isolated from diseased mice by
19 using the procedure described by Bresin et al.(28). Transient transfections were performed with
20 100 nM pre-miR (Ambion) or single-stranded anti-miRNA (IDT) or scrambled negative controls
21 complexed with lipidic nanoparticles (NPs or CD38-NPs). Cells were harvested 48 hours after
22 transfection to evaluate apoptosis and miR expression.

23 **Mice and syngeneic transplantation.** The *TCL1*-tg mouse model used for these experiments has
24 been previously described (15). Breeding pairs were provided to our group as a generous gift from
25 C. M. Croce (Ohio State University). FVB WT mice were obtained from Charles River Laboratories.
26 Mice had ad libitum access to water and a pellet diet. The animal room was maintained at 23°C on
27 a 12-h light/12-h dark cycle. At the age of 6 weeks, the WT FVB female mice were transplanted by
28 IP injection of 5×10^5 lymphocytes isolated from the spleen of an adult *TCL1*-tg mouse with
29 established leukemia. The engraftment of leukemic cells and the progression of disease were
30 monitored over time by the expression of B220/CD5 using flow cytometry analysis (FACS) or
31 absolute quantification of the human *TCL1* transgene by digital PCR of peripheral blood DNA. All
32 experiments with mice were conducted according to the 2010/63/EU directive of the European
33 Parliament and Council. The protocol for animal experimentation was approved by the Italian
34 Ministry of Health (approval n. 40-2014 PR released on November 6, 2014).

35 **Anti-CD38 monoclonal antibody.** The anti-CD38 antibody is a rat monoclonal antibody (mAb)
36 specific for mouse CD38 antigen. It was produced in the laboratory of one of the authors (FM,
37 University of Turin). The NIMR-5 clone was expanded *in vitro* in Iscove's Modified Dulbecco's
38 Medium and purified by high pressure liquid chromatography. The purified IgG was then sterilized
39 by 0.22- μ m filtration (Millipore Polyethersulfone Millex-GP Syringe Filter Unit, radiosterilized) and
40 detoxified by Detoxi-Gel Endotoxin Removing Gel (Thermo Scientific) (87). Specificity was
41 confirmed by reacting the purified mAb with the murine X63.Ag8 myeloma and analyzing binding
42 by indirect immunofluorescence (87).

43 **Preparation of nanoparticles.** The lipid components of the nanoparticles were 1,2-dioleoyl-sn-
44 glycerol-3-phosphoethanolamine (DOPE), 1,2-dimyristoyl-sn-glycerol, methoxypolyethylene glycol
45 (MW ~2000; DMG-PEG; Avanti Polar Lipids, Alabaster, AL), and linoleic acid (Sigma-Aldrich, St.

1 Louis, MO). The molar ratio of DOPE:linoleic acid:DMG-PEG was 50:48:2. The preparation of empty
2 nanoparticles was performed as previously described (32). The anti-CD38 antibody was conjugated
3 with 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[maleimide (polyethylene glycol)-2000]
4 (DSPE-PEG2000 maleimide) according to the method described in a previous study (72) and was
5 then post-inserted into the surface of lipopolyplex nanoparticles to form the CD38-NP-miRs.

6 ***In vivo* treatment with mimics or AMOs.** The synthetic mimics and anti-miRNAs for *in vivo* delivery
7 were purchased from Axolab (Germany). For short-term experiments, the FVB mice (at least 3
8 mice in each group) were enrolled for treatment when the disease reached 30-50% of *TCL1*-
9 positive cells in peripheral blood, and they were treated once with 100 µg of specific single-strand
10 mimics/anti-miRNA and sacrificed after 48 hours. For long-term experiments, the mice (at least 6
11 mice in each group) were enrolled for treatment when the disease reached 15-30% of *TCL1*-
12 positive cells in peripheral blood, and they were treated with 100 µg of the specified molecules 3
13 times a week for 3 weeks. For long-term treatments, the levels of disease were measured by FACS
14 and droplet digital PCR (ddPCR) analysis the day before the start of treatment and 48 hours after it
15 ended.

16 **Flow cytometry.** Blood samples (20 µL) were placed in a tube containing 0.5 M EDTA as an
17 anticoagulant. Erythrocytes were lysed by treatment with ammonium chloride (0.8%) and EDTA
18 (0.1 mM) (Sigma). Cells were incubated with specific antibodies for 10 min in ice, washed, and
19 analyzed by flow cytometry in a FACSCalibur flow cytometer (Becton Dickinson, San Jose,
20 California, USA) by using FlowJo software (TreeStar, Ashland, OR). Data for 5×10^4 cells within the
21 lymphocyte light-scatter gate were collected. In these assays, color compensation was performed
22 before cell acquisition by using the MACS Comp Bead Kit, anti-rat Igk (Miltenyi Biotec, Gladbach,
23 Germany). During analysis, gates were set by using a Fluorescence Minus One control strategy.
24 Leukemic cells were identified as B220+/CD5dim cells, normal B lymphocytes as B220+/CD5- cells,
25 and T lymphocytes as B220-/CD5+ cells. The following antibodies were used: FITC rat anti-mouse
26 CD5 (Cat. 553020, BD Pharmingen) and PeCy5 rat anti-mouse CD45R/B220 (Cat. 553091, BD
27 Pharmingen). CD38 expression on the surface of leukemic splenocytes was detected by triple
28 staining using a PE rat anti-mouse CD38 antibody (Cat. 130-103-008, Miltenyi Biotec) together
29 with the FITC rat anti-mouse CD5 and the PeCy5 rat anti-mouse CD45R/B220 antibody indicated
30 above.

31 **Analysis of apoptosis.** The apoptotic effect of mimic or anti-miR molecules on murine splenocytes
32 treated *in vitro* or *in vivo* was assessed by the Muse™ Annexin V and Dead Cell Assay kit (Cat.
33 MCH100105, Merck) according to manufacturer's protocol. To count viable cells, we used the
34 Muse Count & Viability Assay kit (Cat. MHC100102, Merck).

35 **RNA and DNA extraction.** Total RNA and DNA were isolated from cells or tissues by using the
36 Maxwell Rapid Sample Concentrator (RSC) Instrument (Promega) with the Maxwell RSC miRNA
37 Tissue Kit and the Maxwell RSC Blood DNA Kit.

38 **Reverse transcription and ddPCR.** The ddPCR method was used to measure the expression level
39 of miRNAs. The reverse transcription reaction was performed on 5 ng of total RNA by using the
40 TaqMan miRNA Reverse Transcription assay. After appropriate dilution, 1 µL of the cDNA was used
41 for amplification in a 20-µL reaction volume containing ddPCR Supermix for Probes (Bio-Rad) and
42 the TaqMan miRNA PCR probe set. Droplets generation, cycling conditions for TaqMan assays and
43 the counting of positive droplets were performed according to procedures described by Miotto et
44 al. (88). To normalize the relative abundance of miRNAs, we used the Taqman Assays for RNAs U6
45 or SNO412 (Applied Biosystems).

1 **Monitoring leukemic cells by ddPCR.** A ddPCR approach was used to measure the human *TCL1*
2 transgene in DNA extracted from peripheral blood cells of FVB transplanted mice. We used
3 primers/probes specific for the human *TCL1* transgene (forward 5'-CTCTGGCTCTTGCTTCTTAG-3';
4 reverse 5'-CACCCGTAACCTGTAACCTATC-3'; probe--/56-FAM/TCGTGTATT/Zen/TGGACGAGAAGCA
5 GCA/3IABkFQ/) and primers/probes specific for the mouse *Gapdh* gene as an endogenous
6 reference (forward 5'-GGTGTGAACCACGAGAAATA-3'; reverse 5'-CTCATGGCAGGGTAAGATAAG-3';
7 probe--/5HEX/ACAAC TTTG/Zen/GCATTGTGGAAGGGC/3IABkFQ/). The probes for human *TCL1* and
8 mouse *Gapdh* were conjugated with FAM and HEX, respectively. The ddPCR assay was performed
9 on 10 ng of genomic DNA, using the same procedures as described above.

10 **Western blot analysis.** Splenocytes of treated and control mice were suspended in
11 radioimmunoprecipitation assay (RIPA) buffer (Sigma-Aldrich), supplemented with protease and
12 phosphatase inhibitors (Sigma-Aldrich), according to the manufacturer's protocol. Protein
13 concentrations were measured with the Bradford assay (Bio-Rad). Equal amounts (30 µg) of
14 protein extracts from all samples were applied to SDS-PAGE electrophoresis and then transferred
15 to a PVDF membrane (Bio-Rad). The membrane was incubated with primary antibodies as follows:
16 anti-CDK6 (rabbit, Cat. sc177, Santa Cruz), anti-Mcl-1 (Rabbit, Cat.5453 Cell Signalling) , anti-PARP
17 (rabbit, Cat. 9542, Cell Signalling) , anti-caspase-7 (rabbit, Cat. 9492, Cell Signaling) 1:1000. β-
18 Tubulin H235 (rabbit, Cat. sc9104, Santa Cruz) , , was used as a normalized control. The membrane
19 was incubated with anti-rabbit IgG, HRP-linked antibody (Cat. 7074, Cell Signaling). For signal
20 detection, ClarityTM Western ECL Substrate (Cat. 170-5060, Bio-Rad) was used according to the
21 manufacturer's instructions. Digital images were acquired with Chemidoc (BioRad). Signals were
22 quantified by ImageJ software and protein expression levels normalized according to β-tubulin
23 expression.

24

25 **Conflict of Interest:** The authors declare no competing financial interests.

26

27 **Acknowledgements**

28 This work was supported by funds from the Italian Association for Cancer Research (AIRC Special
29 program 5xmille n.9980) and the University of Ferrara to MN, from the US National Cancer
30 Institute grant CA19-7706 to CMC, from FIRB Programme of the Ministry of University and
31 Research (Rome, Italy), Fondazione Cassa di Risparmio di Torino and Fondazione Ricerca Molinette
32 Torino to FM.

33 **Supplementary Information** accompanies the paper on the *Oncogene* website
34 (<http://www.nature.com/onc>)

1 Reference

- 2 1. Chiorazzi N, Rai KR, Ferrarini M. Chronic lymphocytic leukemia. *The New England journal of*
3 *medicine*. 2005;352(8):804-15.
- 4 2. Zenz T, Mertens D, Kuppers R, Dohner H, Stilgenbauer S. From pathogenesis to treatment of chronic
5 lymphocytic leukaemia. *Nature reviews Cancer*. 2010;10(1):37-50.
- 6 3. Chiorazzi N, Hatzi K, Albesiano E. B-cell chronic lymphocytic leukemia, a clonal disease of B
7 lymphocytes with receptors that vary in specificity for (auto)antigens. *Annals of the New York Academy of*
8 *Sciences*. 2005;1062:1-12.
- 9 4. Grever MR, Lucas DM, Dewald GW, Neuberg DS, Reed JC, Kitada S, et al. Comprehensive
10 assessment of genetic and molecular features predicting outcome in patients with chronic lymphocytic
11 leukemia: results from the US Intergroup Phase III Trial E2997. *Journal of clinical oncology : official journal*
12 *of the American Society of Clinical Oncology*. 2007;25(7):799-804.
- 13 5. Binet JL, Auquier A, Dighiero G, Chastang C, Piguët H, Goasguen J, et al. A new prognostic
14 classification of chronic lymphocytic leukemia derived from a multivariate survival analysis. *Cancer*.
15 1981;48(1):198-206.
- 16 6. Damle RN, Wasil T, Fais F, Ghiotto F, Valetto A, Allen SL, et al. Ig V gene mutation status and CD38
17 expression as novel prognostic indicators in chronic lymphocytic leukemia. *Blood*. 1999;94(6):1840-7.
- 18 7. Wiestner A, Rosenwald A, Barry TS, Wright G, Davis RE, Henrickson SE, et al. ZAP-70 expression
19 identifies a chronic lymphocytic leukemia subtype with unmutated immunoglobulin genes, inferior clinical
20 outcome, and distinct gene expression profile. *Blood*. 2003;101(12):4944-51.
- 21 8. Dohner H, Stilgenbauer S, Benner A, Leupolt E, Krober A, Bullinger L, et al. Genomic aberrations and
22 survival in chronic lymphocytic leukemia. *The New England journal of medicine*. 2000;343(26):1910-6.
- 23 9. Landau DA, Tausch E, Taylor-Weiner AN, Stewart C, Reiter JG, Bahlo J, et al. Mutations driving CLL
24 and their evolution in progression and relapse. *Nature*. 2015;526(7574):525-30.
- 25 10. Rossi D, Brusca A, Spina V, Rasi S, Khiabanian H, Messina M, et al. Mutations of the SF3B1
26 splicing factor in chronic lymphocytic leukemia: association with progression and fludarabine-
27 refractoriness. *Blood*. 2011;118(26):6904-8.
- 28 11. Cuneo A, Cavazzini F, Ciccone M, Daghia G, Sofritti O, Saccetti E, et al. Modern treatment in chronic
29 lymphocytic leukemia: impact on survival and efficacy in high-risk subgroups. *Cancer medicine*.
30 2014;3(3):555-64.
- 31 12. Simonetti G, Bertilaccio MT, Ghia P, Klein U. Mouse models in the study of chronic lymphocytic
32 leukemia pathogenesis and therapy. *Blood*. 2014;124(7):1010-9.
- 33 13. Bresin A, D'Abundo L, Narducci MG, Fiorenza MT, Croce CM, Negrini M, et al. TCL1 transgenic
34 mouse model as a tool for the study of therapeutic targets and microenvironment in human B-cell chronic
35 lymphocytic leukemia. *Cell death & disease*. 2016;7:e2071.
- 36 14. Bichi R, Shinton SA, Martin ES, Koval A, Calin GA, Cesari R, et al. Human chronic lymphocytic
37 leukemia modeled in mouse by targeted TCL1 expression. *Proceedings of the National Academy of Sciences*
38 *of the United States of America*. 2002;99(10):6955-60.
- 39 15. Efanov A, Zanesi N, Nazaryan N, Santanam U, Palamarchuk A, Croce CM, et al. CD5+CD23+ leukemic
40 cell populations in TCL1 transgenic mice show significantly increased proliferation and Akt phosphorylation.
41 *Leukemia*. 2010;24(5):970-5.
- 42 16. Yan XJ, Albesiano E, Zanesi N, Yancopoulos S, Sawyer A, Romano E, et al. B cell receptors in TCL1
43 transgenic mice resemble those of aggressive, treatment-resistant human chronic lymphocytic leukemia.
44 *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(31):11713-8.
- 45 17. Johnson AJ, Lucas DM, Muthusamy N, Smith LL, Edwards RB, De Lay MD, et al. Characterization of
46 the TCL-1 transgenic mouse as a preclinical drug development tool for human chronic lymphocytic
47 leukemia. *Blood*. 2006;108(4):1334-8.
- 48 18. Negrini M, Ferracin M, Sabbioni S, Croce CM. MicroRNAs in human cancer: from research to
49 therapy. *Journal of cell science*. 2007;120(Pt 11):1833-40.

- 1 19. Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, et al. Frequent deletions and down-
2 regulation of micro- RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. Proceedings of
3 the National Academy of Sciences of the United States of America. 2002;99(24):15524-9.
- 4 20. Klein U, Lia M, Crespo M, Siegel R, Shen Q, Mo T, et al. The DLEU2/miR-15a/16-1 cluster controls B
5 cell proliferation and its deletion leads to chronic lymphocytic leukemia. Cancer cell. 2010;17(1):28-40.
- 6 21. Cimmino A, Calin GA, Fabbri M, Iorio MV, Ferracin M, Shimizu M, et al. miR-15 and miR-16 induce
7 apoptosis by targeting BCL2. Proceedings of the National Academy of Sciences of the United States of
8 America. 2005;102(39):13944-9.
- 9 22. Roberts AW, Davids MS, Pagel JM, Kahl BS, Puvvada SD, Gerecitano JF, et al. Targeting BCL2 with
10 Venetoclax in Relapsed Chronic Lymphocytic Leukemia. The New England journal of medicine.
11 2016;374(4):311-22.
- 12 23. Calin GA, Ferracin M, Cimmino A, Di Leva G, Shimizu M, Wojcik SE, et al. A MicroRNA signature
13 associated with prognosis and progression in chronic lymphocytic leukemia. The New England journal of
14 medicine. 2005;353(17):1793-801.
- 15 24. Fabbri M, Bottoni A, Shimizu M, Spizzo R, Nicoloso MS, Rossi S, et al. Association of a
16 microRNA/TP53 feedback circuitry with pathogenesis and outcome of B-cell chronic lymphocytic leukemia.
17 JAMA : the journal of the American Medical Association. 2011;305(1):59-67.
- 18 25. Visone R, Veronese A, Rassenti LZ, Balatti V, Pearl DK, Acunzo M, et al. miR-181b is a biomarker of
19 disease progression in chronic lymphocytic leukemia. Blood. 2011;118(11):3072-9.
- 20 26. He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, et al. A microRNA component of the p53 tumour
21 suppressor network. Nature. 2007;447(7148):1130-4.
- 22 27. Negrini M, Cutrona G, Bassi C, Fabris S, Zagatti B, Colombo M, et al. microRNAome expression in
23 chronic lymphocytic leukemia: comparison with normal B cell subsets and correlations with prognostic and
24 clinical parameters. Clinical cancer research : an official journal of the American Association for Cancer
25 Research. 2014;20(15):4141-53.
- 26 28. Bresin A, Callegari E, D'Abundo L, Cattani C, Bassi C, Zagatti B, et al. miR-181b as a therapeutic agent
27 for chronic lymphocytic leukemia in the Emicro-TCL1 mouse model. Oncotarget. 2015;6(23):19807-18.
- 28 29. Malavasi F, Deaglio S, Damle R, Cutrona G, Ferrarini M, Chiorazzi N. CD38 and chronic lymphocytic
29 leukemia: a decade later. Blood. 2011;118(13):3470-8.
- 30 30. Malavasi F, Deaglio S, Funaro A, Ferrero E, Horenstein AL, Ortolan E, et al. Evolution and function of
31 the ADP ribosyl cyclase/CD38 gene family in physiology and pathology. Physiological reviews.
32 2008;88(3):841-86.
- 33 31. Chillemi A, Zaccarello G, Quarona V, Ferracin M, Ghimenti C, Massaia M, et al. Anti-CD38 antibody
34 therapy: windows of opportunity yielded by the functional characteristics of the target molecule. Mol Med.
35 2013;19:99-108.
- 36 32. Huang X, Schwind S, Yu B, Santhanam R, Wang H, Hoellerbauer P, et al. Targeted delivery of
37 microRNA-29b by transferrin-conjugated anionic lipopolyplex nanoparticles: a novel therapeutic strategy in
38 acute myeloid leukemia. Clinical cancer research : an official journal of the American Association for Cancer
39 Research. 2013;19(9):2355-67.
- 40 33. Calin GA, Pekarsky Y, Croce CM. The role of microRNA and other non-coding RNA in the
41 pathogenesis of chronic lymphocytic leukemia. Best practice & research Clinical haematology.
42 2007;20(3):425-37.
- 43 34. Costinean S, Zanesi N, Pekarsky Y, Tili E, Volinia S, Heerema N, et al. Pre-B cell proliferation and
44 lymphoblastic leukemia/high-grade lymphoma in E(mu)-miR155 transgenic mice. Proceedings of the
45 National Academy of Sciences of the United States of America. 2006;103(18):7024-9.
- 46 35. O'Connell RM, Rao DS, Chaudhuri AA, Boldin MP, Taganov KD, Nicoll J, et al. Sustained expression of
47 microRNA-155 in hematopoietic stem cells causes a myeloproliferative disorder. The Journal of
48 experimental medicine. 2008;205(3):585-94.
- 49 36. Tagawa H, Ikeda S, Sawada K. Role of microRNA in the pathogenesis of malignant lymphoma.
50 Cancer science. 2013;104(7):801-9.

- 1 37. Mattiske S, Suetani RJ, Neilsen PM, Callen DF. The oncogenic role of miR-155 in breast cancer.
2 Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer
3 Research, cosponsored by the American Society of Preventive Oncology. 2012;21(8):1236-43.
- 4 38. Zanesi N, Pekarsky Y, Trapasso F, Calin G, Croce CM. MicroRNAs in mouse models of lymphoid
5 malignancies. Journal of nucleic acids investigation. 2010;1(1):36-40.
- 6 39. Garzon R, Croce CM. MicroRNAs in normal and malignant hematopoiesis. Current opinion in
7 hematology. 2008;15(4):352-8.
- 8 40. Cui B, Chen L, Zhang S, Mraz M, Fecteau JF, Yu J, et al. MicroRNA-155 influences B-cell receptor
9 signaling and associates with aggressive disease in chronic lymphocytic leukemia. Blood. 2014;124(4):546-
10 54.
- 11 41. Visone R, Rassenti LZ, Veronese A, Taccioli C, Costinean S, Aguda BD, et al. Karyotype-specific
12 microRNA signature in chronic lymphocytic leukemia. Blood. 2009;114(18):3872-9.
- 13 42. Rossi S, Shimizu M, Barbarotto E, Nicoloso MS, Dimitri F, Sampath D, et al. microRNA fingerprinting
14 of CLL patients with chromosome 17p deletion identify a miR-21 score that stratifies early survival. Blood.
15 2010;116(6):945-52.
- 16 43. Vargova K, Curik N, Burda P, Basova P, Kulvait V, Pospisil V, et al. MYB transcriptionally regulates
17 the miR-155 host gene in chronic lymphocytic leukemia. Blood. 2011;117(14):3816-25.
- 18 44. Zhang Y, Roccaro AM, Rombaoa C, Flores L, Obad S, Fernandes SM, et al. LNA-mediated anti-miR-
19 155 silencing in low-grade B-cell lymphomas. Blood. 2012;120(8):1678-86.
- 20 45. Zhu X, Zhao H, Lin Z, Zhang G. Functional studies of miR-130a on the inhibitory pathways of
21 apoptosis in patients with chronic myeloid leukemia. Cancer gene therapy. 2015;22(12):573-80.
- 22 46. Acunzo M, Visone R, Romano G, Veronese A, Lovat F, Palmieri D, et al. miR-130a targets MET and
23 induces TRAIL-sensitivity in NSCLC by downregulating miR-221 and 222. Oncogene. 2012;31(5):634-42.
- 24 47. Pan Y, Wang R, Zhang F, Chen Y, Lv Q, Long G, et al. MicroRNA-130a inhibits cell proliferation,
25 invasion and migration in human breast cancer by targeting the RAB5A. International journal of clinical and
26 experimental pathology. 2015;8(1):384-93.
- 27 48. Li Y, Challagundla KB, Sun XX, Zhang Q, Dai MS. MicroRNA-130a associates with ribosomal protein
28 L11 to suppress c-Myc expression in response to UV irradiation. Oncotarget. 2015;6(2):1101-14.
- 29 49. Fujita Y, Kojima T, Kawakami K, Mizutani K, Kato T, Deguchi T, et al. miR-130a activates apoptotic
30 signaling through activation of caspase-8 in taxane-resistant prostate cancer cells. The Prostate.
31 2015;75(14):1568-78.
- 32 50. Li B, Huang P, Qiu J, Liao Y, Hong J, Yuan Y. MicroRNA-130a is down-regulated in hepatocellular
33 carcinoma and associates with poor prognosis. Med Oncol. 2014;31(10):230.
- 34 51. Boll K, Reiche K, Kasack K, Morbt N, Kretzschmar AK, Tamm JM, et al. MiR-130a, miR-203 and miR-
35 205 jointly repress key oncogenic pathways and are downregulated in prostate carcinoma. Oncogene.
36 2013;32(3):277-85.
- 37 52. Shen S, Guo X, Yan H, Lu Y, Ji X, Li L, et al. A miR-130a-YAP positive feedback loop promotes organ
38 size and tumorigenesis. Cell research. 2015;25(9):997-1012.
- 39 53. Lee SH, Jung YD, Choi YS, Lee YM. Targeting of RUNX3 by miR-130a and miR-495 cooperatively
40 increases cell proliferation and tumor angiogenesis in gastric cancer cells. Oncotarget. 2015;6(32):33269-
41 78.
- 42 54. Chen J, Yan D, Wu W, Zhu J, Ye W, Shu Q. MicroRNA-130a promotes the metastasis and epithelial-
43 mesenchymal transition of osteosarcoma by targeting PTEN. Oncology reports. 2016;35(6):3285-92.
- 44 55. Jiang H, Yu WW, Wang LL, Peng Y. miR-130a acts as a potential diagnostic biomarker and promotes
45 gastric cancer migration, invasion and proliferation by targeting RUNX3. Oncology reports.
46 2015;34(3):1153-61.
- 47 56. Liu L, Nie J, Chen L, Dong G, Du X, Wu X, et al. The oncogenic role of microRNA-130a/301a/454 in
48 human colorectal cancer via targeting Smad4 expression. PloS one. 2013;8(2):e55532.
- 49 57. Kovaleva V, Mora R, Park YJ, Plass C, Chiramel AI, Bartenschlager R, et al. miRNA-130a targets
50 ATG2B and DICER1 to inhibit autophagy and trigger killing of chronic lymphocytic leukemia cells. Cancer
51 research. 2012;72(7):1763-72.

- 1 58. Kim H, Huang W, Jiang X, Pennicooke B, Park PJ, Johnson MD. Integrative genome analysis reveals
2 an oncomir/oncogene cluster regulating glioblastoma survivorship. *Proceedings of the National Academy of*
3 *Sciences of the United States of America*. 2010;107(5):2183-8.
- 4 59. Kota J, Chivukula RR, O'Donnell KA, Wentzel EA, Montgomery CL, Hwang HW, et al. Therapeutic
5 microRNA delivery suppresses tumorigenesis in a murine liver cancer model. *Cell*. 2009;137(6):1005-17.
- 6 60. Sander S, Bullinger L, Klapproth K, Fiedler K, Kestler HA, Barth TF, et al. MYC stimulates EZH2
7 expression by repression of its negative regulator miR-26a. *Blood*. 2008;112(10):4202-12.
- 8 61. Huse JT, Brennan C, Hambardzumyan D, Wee B, Pena J, Rouhanifard SH, et al. The PTEN-regulating
9 microRNA miR-26a is amplified in high-grade glioma and facilitates gliomagenesis in vivo. *Genes &*
10 *development*. 2009;23(11):1327-37.
- 11 62. Ji J, Shi J, Budhu A, Yu Z, Forgues M, Roessler S, et al. MicroRNA expression, survival, and response
12 to interferon in liver cancer. *The New England journal of medicine*. 2009;361(15):1437-47.
- 13 63. Yang X, Liang L, Zhang XF, Jia HL, Qin Y, Zhu XC, et al. MicroRNA-26a suppresses tumor growth and
14 metastasis of human hepatocellular carcinoma by targeting interleukin-6-Stat3 pathway. *Hepatology*.
15 2013;58(1):158-70.
- 16 64. Zhu Y, Lu Y, Zhang Q, Liu JJ, Li TJ, Yang JR, et al. MicroRNA-26a/b and their host genes cooperate to
17 inhibit the G1/S transition by activating the pRb protein. *Nucleic acids research*. 2012;40(10):4615-25.
- 18 65. Gao J, Li L, Wu M, Liu M, Xie X, Guo J, et al. MiR-26a inhibits proliferation and migration of breast
19 cancer through repression of MCL-1. *PloS one*. 2013;8(6):e65138.
- 20 66. Zhang Y, Wang Z, Gemeinhart RA. Progress in microRNA delivery. *Journal of controlled release :*
21 *official journal of the Controlled Release Society*. 2013;172(3):962-74.
- 22 67. Chen Y, Gao DY, Huang L. In vivo delivery of miRNAs for cancer therapy: Challenges and strategies.
23 *Advanced drug delivery reviews*. 2014;81:128-41.
- 24 68. Krutzfeldt J, Rajewsky N, Braich R, Rajeev KG, Tuschl T, Manoharan M, et al. Silencing of microRNAs
25 in vivo with 'antagomirs'. *Nature*. 2005;438(7068):685-9.
- 26 69. Elmen J, Lindow M, Schutz S, Lawrence M, Petri A, Obad S, et al. LNA-mediated microRNA silencing
27 in non-human primates. *Nature*. 2008;452(7189):896-9.
- 28 70. Callegari E, Elamin BK, D'Abundo L, Falzoni S, Donvito G, Moshiri F, et al. Anti-tumor activity of a
29 miR-199-dependent oncolytic adenovirus. *PloS one*. 2013;8(9):e73964.
- 30 71. Amendola M, Passerini L, Pucci F, Gentner B, Bacchetta R, Naldini L. Regulated and multiple miRNA
31 and siRNA delivery into primary cells by a lentiviral platform. *Molecular therapy : the journal of the*
32 *American Society of Gene Therapy*. 2009;17(6):1039-52.
- 33 72. Dorrance AM, Neviani P, Ferenchak GJ, Huang X, Nicolet D, Maharry KS, et al. Targeting leukemia
34 stem cells in vivo with antagomiR-126 nanoparticles in acute myeloid leukemia. *Leukemia*.
35 2015;29(11):2143-53.
- 36 73. Li Z, Rana TM. Therapeutic targeting of microRNAs: current status and future challenges. *Nature*
37 *reviews Drug discovery*. 2014;13(8):622-38.
- 38 74. Masotti A, Miller MR, Celluzzi A, Rose L, Micciulla F, Hadoke PW, et al. Regulation of angiogenesis
39 through the efficient delivery of microRNAs into endothelial cells using polyamine-coated carbon
40 nanotubes. *Nanomedicine : nanotechnology, biology, and medicine*. 2016;12(6):1511-22.
- 41 75. Taccioli C, Garofalo M, Chen H, Jiang Y, Tagliazucchi GM, Di Leva G, et al. Repression of Esophageal
42 Neoplasia and Inflammatory Signaling by Anti-miR-31 Delivery In Vivo. *Journal of the National Cancer*
43 *Institute*. 2015;107(11).
- 44 76. Ghosh R, Singh LC, Shohet JM, Gunaratne PH. A gold nanoparticle platform for the delivery of
45 functional microRNAs into cancer cells. *Biomaterials*. 2013;34(3):807-16.
- 46 77. Jia XQ, Cheng HQ, Qian X, Bian CX, Shi ZM, Zhang JP, et al. Lentivirus-mediated overexpression of
47 microRNA-199a inhibits cell proliferation of human hepatocellular carcinoma. *Cell biochemistry and*
48 *biophysics*. 2012;62(1):237-44.
- 49 78. Di Martino MT, Campani V, Misso G, Gallo Cantafio ME, Gulla A, Foresta U, et al. In vivo activity of
50 miR-34a mimics delivered by stable nucleic acid lipid particles (SNALPs) against multiple myeloma. *PloS*
51 *one*. 2014;9(2):e90005.

- 1 79. Scognamiglio I, Di Martino MT, Campani V, Virgilio A, Galeone A, Gulla A, et al. Transferrin-
2 conjugated SNALPs encapsulating 2'-O-methylated miR-34a for the treatment of multiple myeloma.
3 BioMed research international. 2014;2014:217365.
- 4 80. van de Donk NW, Moreau P, Plesner T, Palumbo A, Gay F, Laubach JP, et al. Clinical efficacy and
5 management of monoclonal antibodies targeting CD38 and SLAMF7 in multiple myeloma. *Blood*.
6 2016;127(6):681-95.
- 7 81. Ibrahim S, Keating M, Do KA, O'Brien S, Huh YO, Jilani I, et al. CD38 expression as an important
8 prognostic factor in B-cell chronic lymphocytic leukemia. *Blood*. 2001;98(1):181-6.
- 9 82. Vaisitti T, Audrito V, Serra S, Buonincontri R, Sociali G, Mannino E, et al. The enzymatic activities of
10 CD38 enhance CLL growth and trafficking: implications for therapeutic targeting. *Leukemia*. 2015;29(2):356-
11 68.
- 12 83. Perez-Galan P, Dreyling M, Wiestner A. Mantle cell lymphoma: biology, pathogenesis, and the
13 molecular basis of treatment in the genomic era. *Blood*. 2011;117(1):26-38.
- 14 84. Bende RJ, Smit LA, Bossenbroek JG, Aarts WM, Spaargaren M, de Leval L, et al. Primary follicular
15 lymphoma of the small intestine: alpha4beta7 expression and immunoglobulin configuration suggest an
16 origin from local antigen-experienced B cells. *The American journal of pathology*. 2003;162(1):105-13.
- 17 85. Colomo L, Loong F, Rives S, Pittaluga S, Martinez A, Lopez-Guillermo A, et al. Diffuse large B-cell
18 lymphomas with plasmablastic differentiation represent a heterogeneous group of disease entities. *The*
19 *American journal of surgical pathology*. 2004;28(6):736-47.
- 20 86. Ansari MQ, Dawson DB, Nador R, Rutherford C, Schneider NR, Latimer MJ, et al. Primary body
21 cavity-based AIDS-related lymphomas. *American journal of clinical pathology*. 1996;105(2):221-9.
- 22 87. Horenstein AL, Durelli I, Malavasi F. Purification of clinical-grade monoclonal antibodies by
23 chromatographic methods. *Methods Mol Biol*. 2005;308:191-208.
- 24 88. Miotto E, Saccenti E, Lupini L, Callegari E, Negrini M, Ferracin M. Quantification of circulating
25 miRNAs by droplet digital PCR: comparison of EvaGreen- and TaqMan-based chemistries. *Cancer*
26 *epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research,*
27 *cosponsored by the American Society of Preventive Oncology*. 2014;23(12):2638-42.

28

FIGURES LEGENDS

Figure 1. In the Eμ-TCL1FL TG mouse, the leukemic population (B220+/CD5dim) is CD38+. Cells were isolated from the spleen of an adult Eμ-TCL1FL transgenic mouse (upper panels) or a transplanted FVB-TCL1 mouse (lower panels). The left panels show the lymphocyte population, the middle panels the B220+/Cd5dim leukemic population. The right panels show the CD38+ lymphocytes within the B220+/CD5dim population: more than 98% of the cells are positive.

Figure 2. Bio-distribution of miR-181 after the use of CD38-NPs as an *in vivo* delivery agent. (A) Bio-distribution 24 hours after intraperitoneal injection of miR-181b mimics (100 μg) delivered into a transplanted FVB-TCL1 mouse with a high level of disease by the use of CD38-NP, regular nanoparticles (NPs), or Jet-PEI. **(B)** Highlights of splenocytes and liver. Treatment with CD38-NPs induces the most significant increase in the miR-181b level in splenocytes compared with other tissues or methods of delivery (** p < 0.001, * p < 0.05, respectively), likely reflecting the high number of CD38+ leukemic lymphocytes present in this organ. The results are normalized to SNO412 expression.

Figure 3. Pro-apoptotic activity of different miRNA mimics/AMOs in leukemic splenocytes *in vitro*. **(A)** The apoptotic activity of miRNAs/anti-miRNAs in leukemic splenocytes was evaluated by the Muse Cell Analyzer using the Muse Annexin V & Dead Cell Assay 48 hours after transfection. Statistical assessment of each miRNA effect is referred to the Scramble control. * = p < 0.05 (t-test). **(B)** The level of each miRNA after transfection was assessed by ddPCR at the time when apoptosis measured. miRNA levels were normalized on SNO412 and non-transfected cells (UNT).

Figure 4. CD38-NPs mediate an *in vivo* more effective functional outcome than non-conjugated NPs. Forty-eight hours after a single injection of miR-26a mimics (100 ug) with CD38-NPs or non-conjugated NPs, mice with established leukemia were sacrificed, splenocytes were isolated and analyzed for RNA and protein expression. CD38-NP-miRNA26a produced a higher level of miR-26a than NP-miR26a. The results are shown as relative expression of miR-26a normalized to SNO412 expression, as measured by ddPCR. As a result, the treatment with CD38-NPs could mediate more effective functional outcomes as revealed by a clearer apoptotic effect (cleavage of PARP) and a stronger down regulation of a miR-26a target (CDK6).

Figure 5. Apoptotic activity of miRNA/anti-miRNA molecules after a single *in vivo* treatment of mice with established leukemia. Mice with established leukemia were sacrificed 48 hours after a single treatment with miRNA/anti-miRNA molecules. Splenocytes were isolated to evaluate apoptosis. **(A)** Using an Annexin V assay, we observed a significant increase in the percentage of apoptotic cells 48 hours after treatment with CD38-NP-miR-26a (p = 0.013), CD38-NP-miR-130a (p = 0.03), or CD38-NP-anti-miR-155 (p = 0.002). **(B)** At the same time, the relative expression of miR-16, miR-26a, miR-155, miR-130a, miR-34a, and miR-21 was measured in splenocytes by using ddPCR. The results are shown as the relative expression level, normalized to SNO412 expression.

Figure 6. miR-26a exhibits the strongest activity against Eμ-TCL1FL leukemic cells. Percentages of circulating leukemic cells were measured just before starting treatments and 48 hours after their completion. The differences between these points were acquired and plotted for each mouse enrolled in the study. miR-26a exhibited the strongest effect on expansion of leukemic burden, as shown in the graph and by the p-values in the associated table. P-values < 0.05 are highlighted by a grey background. N = number of mice, SD = standard deviation.

1
2
3
4
5
6
7
8
9
10
11

Figure 7. Modulation of pro-apoptotic proteins and direct targets of miR-26a *in vivo*. Forty-eight hours after the end of long-term treatments, mice were sacrificed and the spleen collected. Proteins were extracted from splenocytes of CD38-NP-miR-26a-treated and control mice for Western blot analysis. **(A)** Cleaved PARP and cleaved caspase-7 are evidence of apoptosis and were clearly visible in the samples treated with miR-26a, but not in the controls. **(B)** The direct molecular activity of miR-26a was demonstrated by showing that two of its known targets, Mcl1 and Cdk6, were down-modulated in the samples treated with miR-26a, but not in the control.




Figure 1



A**B**

Figure 2




Figure 3




Figure 4




A**B**

Figure 5



	Leukemia Burden Difference post-pre			t-tests				
	N	Mean	SD	mR-26a	miR-130a	anti-miR-155	Ctrl	CD38-NP-empty
mR-26a	10	23.1	3.2	1.00E+00	8.30E-03	1.00E-01	2.60E-03	2.00E-04
miR-130a	7	34.5	1.7	8.30E-03	1.00E+00	5.32E-01	2.49E-01	1.36E-02
anti-miR-155	6	32.0	3.7	1.00E-01	5.32E-01	1.00E+00	1.83E-01	2.61E-02
Ctrl	9	38.6	3.0	2.60E-03	2.49E-01	1.83E-01	1.00E+00	2.53E-01
CD38-NP-empty	6	43.6	2.5	2.00E-04	1.36E-02	2.61E-02	2.53E-01	1.00E+00

Figure 6




Figure 7