AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Immunolocalization of H⁺-ATPase and IRT1 enzymes in N₂-fixing common bean nodules subjected to iron deficiency

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1655386> since 2018-01-15T10:30:19Z

Published version:

DOI:10.1016/j.jplph.2011.10.003

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Slatni T1, Dell'Orto M, Ben Salah I, Vigani G, Smaoui A, Gouia H, Zocchi G, Abdelly C..
**Immunolocalization of H(+)-ATPase and IRT1 enzymes in N(2)-fixing common bean
nodules subjected to iron deficiency.** Journal of Plant Physiology . volume169 fascicolo 3,
anno 2012, pagg 242-248-11. Doi: [10.1016/j.jplph.2011.10.003](https://doi.org/10.1016/j.jplph.2011.10.003)

The publisher's version is available at:

<http://www.sciencedirect.com/science/article/pii/S017616171100424X?via%3Dihub>

When citing, please refer to the published version.

Link to this full text:

https://ac.els-cdn.com/S017616171100424X/1-s2.0-S017616171100424X-main.pdf?_tid=5209bfbf-f9d6-11e7-a3ab-00000aacb35d&acdnat=1516008635_57434167abd1ec6da56827018640fa89

This full text was downloaded from iris-Aperto: <https://iris.unito.it/>

Abstract

The demand for iron in leguminous plants increases during symbiosis, as the metal is utilised for the synthesis of various Fe-containing proteins in both plant and bacteroids. However, the acquisition of this micronutrient is problematic due to its low bioavailability at physiological pH under aerobic conditions. Induction of root Fe(III)-reductase activity is necessary for Fe uptake and can be coupled to the rhizosphere acidification capacity linked to the H⁺-ATPase activity. Fe uptake is related to the expression of a Fe²⁺ transporter (IRT1). In order to verify the possible role of nodules in the acquisition of Fe directly from the soil solution, the localization of H⁺-ATPase and IRT1 was carried out in common bean nodules by immuno-histochemical analysis. The results showed that these proteins were particularly abundant in the central nitrogen-fixing zone of nodules, around the periphery of infected and uninfected cells as well as in the vascular bundle of control nodules. Under Fe deficiency an over-accumulation of H⁺-ATPase and IRT1 proteins was observed especially around the cortex cells of nodules. The results obtained in this study suggest that the increase in these proteins is differentially localized in nodules of Fe-deficient plants when compared to the Fe-sufficient condition and cast new light on the possible involvement of nodules in the direct acquisition of Fe from the nutrient solution.

Abbreviations

BMbacteroid membrane CZcortex zone Ex Cexternal cortex FC-Rferric chelate reductase I Cinfected cell In Cinternal cortex IRT1iron regulated transporter 1 IZinfected zone LHleghemoglobin NRnitrogenase PBMperibacteroid membrane PBSperibacteroid space Ssiderophore UCuninfected cell VBvascular bundle

Keywords

Fe absorption Immunodetection Iron transporter *Phaseolus vulgaris* Symbiosis

Introduction

Iron is an important nutrient in N₂-fixing legume nodules. Iron supplied to the nodule is used by the plant for the synthesis of leghemoglobin (LH), while in the bacteroid it is used as an essential cofactor for the bacterial N₂-fixing enzyme, nitrogenase (NR), and Fe-containing proteins of the electron transport chain. *In situ*, bacteroids are always surrounded by a membrane of plant origin (Verma et al., 1978). This so-called peribacteroid membrane (PBM) physically isolates the microbial symbiont from the cytoplasm of the host cell to form the symbiosome, delimiting a peribacteroid space (PBS). The supply of Fe to the bacteroids requires initial transport across the plant-derived PBM. The main metabolic exchange that occurs between plant and bacteroids is related to the reduced carbon (usually malate) supplied by the plant to the bacteroids for N₂-fixation (Udvardi et al., 1988). Specific transport mechanisms responsible for this exchange have been identified (Bergersen and Turner, 1990). However, the bacteroids are dependent on the plant for all micronutrients, and transporters for these must also exist on the PBM. The mechanisms involved in bacteroid Fe acquisition within the nodule have been investigated at the biochemical level and three activities were identified (Day et al., 2001). Fe³⁺ is transported across the PBM complexed with organic acids such as citrate and accumulates in the PBS (Le Vier et al., 1996; Moreau et al., 1995), where it is bound to siderophore (S)-like compounds (Wittenberg et al., 1996). Fe(III)-chelate reductase (FC-R) activity has been measured on isolated PBM and Fe³⁺ uptake into symbiosomes was dependent on the presence of NADH (Le Vier et al., 1996). However, Fe²⁺ is also readily

transported across the PBM and has been found to be the favoured form of Fe taken up by bacteroids (Moreau et al., 1998). One of the proteins involved in this transport has been identified by Kaiser et al. (2003) in soybean nodules. This is a *Glycine max* divalent metal transporter 1 (GmDMT1) homologue of the NRAMP/Dmt1 family of divalent metal ion transporters. In the same work, the localization and the activity of GmDMT1 found in soybean nodules suggest that the protein is involved in Fe²⁺ transport and Fe homeostasis in symbiotic N₂-fixing nodules. In addition, another transporter was identified and characterized as a symbiotic zinc transporter (GmZip1), which does not transport Fe²⁺ (Moreau et al., 2002). Some work reported that ferrous Fe was taken up by symbiosomes more efficiently than the ferric form, because in the microaerobic conditions and the slightly acidic pH of nodules *in situ*, Fe²⁺ should be more stable (Appleby, 1984). Consequently, it is possible that ferrous Fe is the main form of iron present *in vivo* (Moreau et al., 1998). Concerning the genetic approach of Fe regulation in plants, many works focused in the regulation of the expression of genes involved in Fe acquisition, transport and homeostasis in plant, among these genes are the transcription factor *AtFIT*, the ferric reductase *AtFRO*, the iron transporter *AtIRT1* and the H⁺-ATPase *AtAHA* (Colangelo and Guerinot, 2004; Walker and Connolly, 2008; Buckhout et al., 2009; García et al., 2011). Nevertheless, little information is available concerning the Fe-related genes involved in Fe acquisition and transport in root-nodules of legumes. Hakoyama et al. (2009) showed that the gene of nicotianamine synthase (NAS) was expressed exclusively in the vascular bundles (VBs) of *Lotus japonicas* nodules during the late state of nodule development and was involved in the intracellular Fe translocation in nodules.

Generally, upon sensing Fe limitation, strategy I plants such as common bean, induce a coordinated set of responses that, taken together, allow the plant to maximize Fe mobilization and uptake from the soil. Under symbiotic N₂-fixing condition, in which the demand for Fe increases, the preferential allocation of Fe to nodules and the Fe-use efficiency for nodules growth and functions were found to be the base of tolerance of some common bean cultivars (Krouma et al., 2008; Slatni et al., 2008, 2009). This adaptive mechanism includes the induction of three activities localized at the PM of roots cells and PBM and BM of symbiosome cells: (i) a proton pump acidifies the rhizosphere or PBS, thus driving more Fe into solution, (ii) a FC-R converts Fe³⁺-chelates to Fe²⁺ and (iii) a Fe²⁺ transporter moves Fe across the plasma or PBM into cells (Slatni et al., 2011). In spite of several studies clearly elucidating the biochemical and molecular aspects involved in the regulation of the enzymes cited above in roots, little is known about their involvement in Fe uptake and mobilization in nodules and the role of these organs in enhancing Fe acquisition. Previous results focused to verify the ability of nodules to acquire Fe directly from the soil solution were obtained and showed, by detection both “*in vivo*” and “*in vitro*”, the stimulation of FC-R activity in nodulated roots and detached nodules of Fe-deficient plants (Slatni et al., 2009, 2011). Two common bean cultivars were used in these studies, Coco blanc, sensitive and Flamingo, tolerant to Fe deficiency (Slatni et al., 2008). The performance of Flamingo in Fe deficiency is related to (i) its ability to maintain a better symbiotic N₂ fixation linked to a better Fe allocation to nodules and a significant Fe use efficiency for nodule growth and N₂ fixation, and (ii) an important ammonium assimilation linked to the maintenance of an important GS activity and particularly to a significant induction of GDH activity (Slatni et al., 2008). The same studies suggested that the nodules of tolerant cultivar contributed to the rhizosphere acidification and to the reduction of apoplastic Fe³⁺. It is also noteworthy that the activity of FC-R under Fe deficiency was higher in nodules than in roots (about two fold), suggesting that under this condition nodules play an important role in Fe³⁺ reduction and Fe uptake and consequently in Fe supply to the bacteroids (Slatni et al., 2011). Like in roots, transporters

homologues to iron regulated transporter (IRT1) should be present in nodules. The amount of this protein was increased under Fe deficiency also in nodules enabling them to acquire Fe directly from the soil nutrient solution (Slatni et al., 2011). Additionally, Slatni et al. (2011) showed that also H⁺-ATPase activity and protein level were increased under Fe deficiency in nodules of both cultivars, mainly in Flamingo. From the data obtained in this previous work we might now hypothesise that, when common bean plants grow with low Fe availability, nodules acquire the capacity to take up Fe not only from the plant cytosol, but also directly from the nutrient solution. The aim of the present work was: to provide more evidence about the implication of different tissues of nodules in Fe availability and uptake by immuno-histochemical analysis of H⁺-ATPase and IRT1 in common bean nodules.

Materials and methods

Plant materials and growth conditions

Two common bean cultivars, Coco blanc, sensitive and Flamingo, tolerant to Fe deficiency (Slatni et al., 2008), were inoculated with *Rhizobium tropici* CIAT 899 originating from International Center of Tropical Agriculture, Colombia, maintained in the “Laboratory of legumes” (CBBC, Hammam-Lif, Tunisia) and grown with the following N-free nutrient solution (Hewitt, 1966): KH₂PO₄ (1.60 mM), MgSO₄ (1.50 mM), K₂SO₄ (1.50 mM), CaSO₄ (3.50 mM), H₃BO₃ (4 µM), MnSO₄ (4 µM), ZnSO₄ (1 µM), CuSO₄ (1 µM), CoCl₂ (0.12 µM), (Na)₆(Mo)O₂₄ (0.12 µM) added with 5 µM Fe as K–Fe–EDTA and 1 mM urea as starter N needed for growth during the first two weeks. The nutrient solution was aerated with a flow of filtered air. After two weeks of pre-treatment (appearance of nodules), plants were separated into two plots; the first one did not receive Fe (Fe-deficient plants), while the second one received 45 µM Fe(III)EDTA (Fe-sufficient plants). Nodules from three-week-old plants were harvested and used for the analysis.

Fixing, embedding and sectioning

Nodules were detached from roots and fixed overnight at 4 °C either in ethanol–acetic acid (3:1 (v/v), for immunolocalization) or in 100 mM sodium phosphate buffer (pH 7.0) containing 2% paraformaldehyde (w/v) and 2.5% glutaraldehyde, then dehydrated through an ethanol–tertiary butanol series and embedded in paraffin (Paraplast plus, Sigma) as described by Villalba et al. (1991). Serial sections of 7 µm were cut with a microtome. Sections were mounted on polylysine-treated slides, deparaffinised in xylene and rehydrated through an ethanol series.

Immunolocalization

Immunological detection in tissue sections fixed as above described was performed as in Parets-Soler et al. (1990) with some modifications. Sections were incubated for 30 min at room temperature in 3% H₂O₂, then blocked for 1 h in TBS (150 mM NaCl, 25 mM Tris–HCl, pH 7.6) with 2% BSA (w/v). Polyclonal antibodies raised against a IRT1 synthetic peptide (a kind gift from Dr. Connolly; Connolly et al., 2002), and the central domain of *Arabidopsis thaliana* PM H⁺-ATPase (a kind gift from Dr. Serrano, Universidad Politécnica, Valenci2a, Spain) were used. Antibodies were diluted 1:1000 in TBS with 0.5% BSA and incubated overnight at 4 °C with the sections. After 3× 5 min washes in TBS, the sections were incubated 2 h at room temperature with a biotinylated secondary antibody (anti-rabbit IgG biotin conjugate developed against goat, Sigma) diluted 1:1000 in TBS with 0.5% BSA. After 3× 5 min washes in TBS the sections were incubated 30 min in the ExtrAvidin-Peroxidase (ExtrAvidin Peroxidase Staining Kit, Sigma). After 3× 5 min washes in TBS the sections were incubated 5–10 min in 0.05 M acetate buffer pH 5.0 containing 5% dimethylformamide, 0.04% 3-amino-9-ethylcarbazole (AEC) and 0.015% H₂O₂ and finally washed in distilled water.

Results

According to the microscopy observation of nodules, three major parts can be recognized at the histological level in the control section treated without primary antibodies (Fig. 1): first, the central part of the nodules or infected zone (IZ) (Fig. 1A) containing the infected cells (IC) and interconnected aggregates of smaller, uninfected cells (UCs) (Fig. 1B); second, the nodule parenchyma or internal cortex (In C) surrounding the central tissue and including the VBs (Fig. 1A and C) and third the external cortex (Ex C) (Fig. 1A and D). Fig. 1 shows the control sections which appear not stained, indicating that a specific reaction attributable to the secondary antibody and/or extravidin–peroxidase complex does not occur during the immunolocalization experiments.

[Download full-size image](#)

Fig. 1. (A)–(D) Control sections of common bean nodule tissues treated without polyclonal antibodies. CZ: cortex zone; Ex C: external cortex; IC: infected cell; In C: internal cortex; IZ: infected zone; UC: uninfected cell; VB: vascular bundle.

Iron-sufficient (+Fe) and Fe-deficient (–Fe) nodule sections from both Flamingo and Coco blanc were subjected to immunoreaction with antibodies raised against H⁺-ATPase and IRT1 in order to localize the expression of these proteins.

In +Fe nodules of the tolerant cultivar (Flamingo) H⁺-ATPase protein is expressed in the VBs (Fig. 2A and B) and the IZ (Fig. 2C), but not in the Ex C (Fig. 2D). In –Fe nodules a considerable signal relative to H⁺-ATPase appeared in IZ (Fig. 2E and G) and also in the cortex zone (CZ), both internal and external (In C, Ex C) (Fig. 2H) but less pronounced in the VB (Fig. 2E and F) compared to +Fe nodules. In the sensitive genotype (Coco blanc),

similarly to what observed in Flamingo, H⁺-ATPase protein was mainly expressed in VB and IZ (Fig. 2J and K) but not in Ex C of nodules (Fig. 2L) of +Fe plants, but, conversely to Flamingo, in -Fe nodules H⁺-ATPase did not show any significant difference compared with the +Fe condition. Indeed, the signal intensity was similar in both the VB and Ex C between +Fe (Fig. 2J and L) and -Fe (Fig. 2N and P) conditions, while a slightly decreased signal was detected in the IZ of -Fe nodules (Fig. 2M and O).

[Download full-size image](#)

Fig. 2. (A)–(P) Immunolocalization of H⁺-ATPase in common bean nodule tissues of (i) Flamingo. (A)–(D) sections of Fe sufficient nodules. (E)–(H) sections of Fe deficient nodules and of (ii) Coco blanc. (I)–(L) sections of Fe sufficient nodules. (M)–(P) sections of Fe deficient nodules treated with polyclonal antibody. Ex C: external cortex; IC: infected cell; In C: internal cortex; UC: uninfected cell; VB: vascular bundle.

Concerning IRT1, the protein is particularly expressed in VB (Fig. 3A and B) and IZ (Fig. 3A and C) of +Fe Flamingo nodules, while in -Fe ones an increased signal was detected in the IZ (Fig. 3G) and also in cortex cells (Fig. 3H) but decreased in VB (Fig. 3F) respect

to the +Fe ones. In +Fe Coco blanc nodules, IRT1 antibodies evidenced a weak signal in the VB (Fig. 3J) which slightly increased in the -Fe (Fig. 3N), while in the Ex C no appreciable difference in the signal intensity between the two conditions was observed (Fig. 3L and P). Moreover, the IZ of the -Fe nodules shows an increase in IRT1 signal (compare Fig. 3O with K).

[Download full-size image](#)

Fig. 3. (A)–(P) Immunolocalization of IRT1 in common bean nodule tissues of (i) Flamingo. (A)–(D) sections of Fe sufficient nodules. (E)–(H) sections of Fe deficient nodules and of (ii) Coco blanc. (I)–(L) sections of Fe sufficient nodules. (M)–(P) sections of Fe deficient nodules treated with polyclonal antibody. Ex C: external cortex; IC: infected cell; In C: internal cortex; UC: uninfected cell; VB: vascular bundle.

In general two main conclusions might be drawn: (i) the signal relative to the proteins studied in this work was less intense in the sensitive cultivar than in the tolerant one, (ii) in -Fe treated plants H⁺-ATPase and IRT1 proteins were significantly detected in CZ of the tolerant cultivar nodules (Figs. 2H and 3H), but not in the sensitive one (Figs. 2P and 3P). In addition, enhanced signal of these proteins was also detected in the IZ of -Fe nodules

of Flamingo, while in Coco the same is true only for IRT1 (Figs. 2C, G, K, and O and 3C, G, K, and O).

Discussion

In the present study the expression and the histological distribution of H⁺-ATPase and IRT1 proteins in nodules of Fe-sufficient and Fe-deficient roots from two bean cultivars differently tolerant to Fe deficiency was compared. The results suggest that an increase in these proteins occurs in nodules of -Fe plants compared to +Fe ones, especially in the tolerant cultivar, Flamingo. In nodules of +Fe plants these proteins are expressed particularly in IZ and VB, while nodules of -Fe plants are characterized by high proteins accumulation in IZ and in the cortex, which is involved in the nutrient absorption and exchange with the external medium.

The accumulation of H⁺-ATPase and IRT1 proteins in cortex cells of Fe-deficient nodules of Flamingo suggests their participation in the acidification of the rhizosphere around the nodule and in the Fe uptake directly from the soil solution. Considerable signal of these proteins was also detected in the IZ of -Fe Flamingo nodules. This finding suggests that H⁺-ATPase might participate in maintaining low pH in the apoplast of IZ, contributing to the solubilisation of Fe compounds. Consequently, Fe³⁺ reduction via PBM is facilitated, since FC-R activity is favoured by acidic pH in the apoplast. Several studies showed that a H⁺-ATPase pumps protons into the PBS and generates a $\Delta\psi$ across the PBM (Udvardi and Day, 1997; Day et al., 2001). Moreover, the bacteroid respiratory electron chain pumps protons out of the bacteroid into the PBS, resulting in additional acidification of the PBS. Thus, the pH of the PBS, determined by the relative activities of proton pumps and counter-ion movements, could be up to two pH units more acidic than the plant cytosol (Udvardi and Day, 1997). The nodule cortex is the interface between the nodule and its environment and probably, under some severe condition like Fe deficiency, it performs essential functions such as nutrient uptake. Basing on the presented data we may suggest that the driving force for the movement of nutrients across the nodule cortex is provided by the H⁺-ATPase located on the PM of cortex cells.

On the other hand, the determination of protein level of IRT1 in nodules by western blot analysis (Slatni et al., 2011), showed a great accumulation of this protein in both -Fe Flamingo and -Fe Coco blanc with respect to the relative control conditions. In this work we confirmed these results but we also observed that IRT1 localized in the cortex of nodules from Fe-deficient plants, suggesting that IRT1 is directly involved in Fe²⁺ transport from the soil solution, across the Ex C of nodules, to the bacteroid interior. Two different behaviours can be raised in this work concerning the allocation of Fe to nodules under Fe limiting condition. The nodules of the tolerant cultivar, Flamingo, in addition to Fe supply through xylem flux from roots, depend by large part, if not exclusively, on their own ability to acquire Fe directly from nutrient solution; thus, Fe is rapidly available for nodule tissues and bacteroids to support the synthesis of Fe-dependent proteins, i.e. NR and LH. This in part explains the greater efficiency of Flamingo to fix N₂ under limited Fe availability (Slatni et al., 2008). On the contrary, under the same condition, the nodule of Fe-deficient plants of the sensitive cultivar, Coco blanc, depends on Fe supply through xylem flux from roots via VB and when this source is blocked or decreases the availability of Fe, the nodule is rapidly affected by Fe deficiency. Overall, in this work the comparison between tolerant and susceptible cultivars allowed to identify the traits conferring efficiency in Fe uptake/supply to nodules. Basing on our results (data from this work; Slatni et al., 2009, 2011) and also on studies focalized on Fe uptake and transport to bacteroids (Le Vier et al., 1996; Udvardi and Day, 1997; Moreau et al., 1998), we propose a model summarising Fe acquisition and transport in nodules (Fig. 4).

[Download full-size image](#)

Fig. 4. Model of Fe acquisition and transport in infected and external cortex cells of common bean nodules, of tolerant plant, under Fe sufficient (+Fe) (A) or Fe deficient (-Fe) (B) conditions. At the level of infected cells Fe deficiency induced the Fe uptake system (FC-R, IRT1, H⁺-ATPase) both at the PBM and the plasma membrane levels compared to +Fe condition. As well, at the level of the external cortex cells, Fe deficiency induced the Fe uptake system compared to +Fe condition, suggesting that -Fe nodules are able to acquire Fe directly from soil nutrient solution. The colored enzyme in the figure means an activated enzyme compared with the gray scale enzyme. LH: leghemoglobin; NR: nitrogenase; PBM: peribacteroid membrane; PBS: peribacteroid space; S: siderophore. (For interpretation of the references to color in this figure caption, the reader is referred to the web version of the article.)

In the presence of adequate Fe supply, plants deliver Fe to nodules via the VBs. Iron

coming from roots reaches the cytoplasm of nodule cells either as Fe²⁺ or Fe³⁺ form. Fe³⁺ can be transported across the PBM complexed with organic acids such as citrate and accumulates in the PBS (Le Vier et al., 1996; Moreau et al., 1995), where it is bound to S-like compounds by the bacteroids (Wittenberg et al., 1996) (Fig. 4A). On the other hand, the PBM possesses a FC-R capable to reduce Fe³⁺ to Fe²⁺. In Fe-deficient condition (Fig. 4B), we suggest that nodules are able to acquire Fe directly from soil nutrient solution. This ability implies the induction, in the Ex C of nodule, of (a) H⁺-ATPase which, acidifying the rhizosphere, increases the availability of Fe and creates a favourable pH condition for (b) FC-R activity which is increased in external cells of nodules (Slatni et al., 2009, 2011), and (c) IRT1 to transport Fe²⁺ from the rhizosphere into cytoplasm of nodule cells. Once Fe²⁺ has reached the cytoplasm of nodule cells, it is transported into the PBS through the same mechanism as in Fe-sufficient nodules (i.e. by IRT1 of the PBM). In addition to the direct absorption of iron by the cortex cells of nodules to support the iron requirement in the bacteroids, the xylem way from the root to the nodules of Fe uptake and transport is also present.

The main results of the present study suggest that under Fe deficiency, similarly to what occurs at the PBM, the nodule cortex shows a higher protein level of H⁺-ATPase and IRT1. At the whole, the data suggest that (i) these proteins play a critical role in the regulation of Fe uptake in nodules; (ii) nodules, especially from Flamingo roots, could be able to acquire Fe directly from the nutrient solution, particularly upon Fe-limitation. This would confer a considerable advantage to this cultivar, allowing the N₂ fixation process to take place even in such a stress-inducing condition.

Acknowledgment

This work was supported by grants from the Tunisian Ministry of Higher Education and Scientific Research (LR10CBBC02).

References

Appleby, 1984

C.A. Appleby

Leghemoglobin and rhizobium respiration

Annu Rev Plant Physiol, 35 (1984), pp. 443-478

[CrossRef](#)

Bergersen and Turner, 1990

F.J. Bergersen, G.L. Turner

Bacteroids from soybean root nodules: accumulation of poly-β-hydroxybutyrate during the supply of malate and succinate in relation to N₂ fixation in flow chamber reactions with oxyleghaemoglobin

Proc Soc Lond B, 238 (1990), pp. 295-320

[CrossRefView Record in Scopus](#)

Buckhout et al., 2009

T.J. Buckhout, T.J.W. Yang, W. Schmidt

Early iron-deficiency-induced transcriptional changes in Arabidopsis roots as revealed by microarray analyses

BMC Genom, 10 (2009), p. 147

[CrossRef](#)

Colangelo and Guerinot, 2004

E.P. Colangelo, M.L. Guerinot

The essential basic helix-loop-helix protein FIT1 is required for the iron deficiency response

Plant Cell, 16 (2004), pp. 3400-3412

[CrossRefView Record in Scopus](#)

Connolly et al., 2002

E.L. Connolly, J.P. Fett, M.L. Guerinot

Expression of the IRT1 metal transporter is controlled by metals at the levels of transcript and protein accumulation

Plant Cell, 14 (2002), pp. 1347-1357

[CrossRefView Record in Scopus](#)

Day et al., 2001

D.A. Day, P.S. Poole, S.D. Tyerman, L. Rosendahl

Ammonia and amino acid transport across symbiotic membranes in nitrogen-fixing legume nodules

Cell Mol Life Sci, 58 (2001), pp. 61-71

[CrossRefView Record in Scopus](#)

García et al., 2011

M.J. García, J. García María, Suárez Vicente, J. Romera Francisco, Alcántara Esteban, Pérez-Vicente Rafael

A new model involving ethylene, nitric oxide and Fe to explain the regulation of Fe acquisition genes in Strategy I plants

Plant Physiol Biochem, 49 (2011), pp. 537-544

[ArticlePDF \(918KB\)View Record in Scopus](#)

Hakoyama et al., 2009

T. Hakoyama, H. Watanabe, J. Tomita, A. Yamamoto, S. Sato, Y. Mori, *et al.*

Nicotianamine synthase specifically expressed in root nodules of *Lotus japonicas*

Planta, 230 (2009), pp. 309-317

[CrossRefView Record in Scopus](#)

Hewitt, 1966

E.J. Hewitt

Sand and water culture methods used in the study of plant nutrition

(2nd ed.), Commonwealth Bureau of Horticulture Technical Communications (1966)

p. 22

Kaiser et al., 2003

B.N. Kaiser, S. Moreau, J. Castelli, R. Thomson, A. Lambert, S. Bogliolo, *et al.*

The soybean NRAMP homologue, GmDMT1, is a symbiotic divalent metal transporter capable of ferrous iron transport

Plant J, 35 (2003), pp. 295-304

[CrossRefView Record in Scopus](#)

Krouma et al., 2008

A. Krouma, T. Slatni, C. Abdelly

Differential tolerance to lime-induced chlorosis of N₂ fixing common bean (*Phaseolus vulgaris* L.)

Symbiosis, 46 (2008), pp. 137-143

[View Record in Scopus](#)

Le Vier et al., 1996

K. Le Vier, D. Day, M.L. Guerinot

Iron uptake by symbiosomes from soybean root nodules

Plant Physiol, 111 (1996), pp. 893-900

[CrossRefView Record in Scopus](#)

Moreau et al., 1998

S. Moreau, D.A. Day, A. Puppo

Ferrous iron is transported across the peribacteroid membrane nodules

Planta, 207 (1998), pp. 83-87

[CrossRefView Record in Scopus](#)

Moreau et al., 2002

S. Moreau, R.M. Thomson, B.N. Kaiser, B. Trevaskis, M.L. Guerinot, M.K. Udvardi, *et al.*

GmZIP1 encodes a symbiosis-specific zinc transporter in soybean

J Biol Chem, 277 (2002), pp. 4738-4746

[CrossRefView Record in Scopus](#)

Moreau et al., 1995

S. Moreau, J.M. Meyer, A. Puppo

Uptake of iron by symbiosomes and bactéroïdes from soybean nodules

FEBS Lett, 361 (1995), pp. 225-228

[ArticlePDF \(442KB\)CrossRefView Record in Scopus](#)

Parets-Soler et al., 1990

A. Parets-Soler, J.M. Pardo, S. Serrano

Immunocytolocalization of plasma membrane H⁺-ATPase

Plant Physiol, 93 (1990), pp. 1654-1658

[CrossRefView Record in Scopus](#)

Slatni et al., 2008

T. Slatni, A. Krouma, S. Aydi, C. Chaiffi, H. Gouia, C. Abdelly

Growth, nitrogen fixation and ammonium assimilation in common bean (*Phaseolus vulgaris* L.) subjected to iron deficiency

Plant Soil, 312 (2008), pp. 49-57

[CrossRefView Record in Scopus](#)

Slatni et al., 2009

T. Slatni, A. Krouma, H. Gouia, C. Abdelly

Importance of ferric-chelate reductase activity and acidification capacity in root nodules of N₂-fixing Common bean (*Phaseolus vulgaris* L.) subjected to iron deficiency

Symbiosis, 47 (2009), pp. 35-42

[CrossRefView Record in Scopus](#)

Slatni et al., 2011

T. Slatni, G. Vigani, I. Ben salah, S. Kouas, M. Dell'Orto, H. Gouia, *et al.*

Metabolic changes of iron uptake in N₂-fixing common bean nodules during iron deficiency

Plant Sci, 181 (2011), pp. 151-158

[ArticlePDF \(689KB\)View Record in Scopus](#)

Udvardi and Day, 1997

M.K. Udvardi, D. Day

Metabolite transport across symbiotic membranes of legume nodules

Annu Rev Plant Physiol Plant Mol Biol, 48 (1997), pp. 493-523

[CrossRefView Record in Scopus](#)

Udvardi et al., 1988

M.K. Udvardi, G.D. Price, P.M. Gresshoff, D.A. Day

A dicarboxylate transporter on the peribacteroid membrane of soybean nodules

FEBS Lett, 231 (1988), pp. 36-40

[ArticlePDF \(416KB\)CrossRefView Record in Scopus](#)

Verma et al., 1978

D.P.S. Verma, V. Kazazian, V. Zogbi, A.K. Bal

Isolation and characterization of the membrane envelope enclosing the bacteroids in soybean root nodules

J Cell Biol, 78 (1978), pp. 919-936

[CrossRefView Record in Scopus](#)

Villalba et al., 1991

J.M. Villalba, M. Lützelschwab, R. Serrano

Immunocytochemical localization of plasma membrane H⁺-ATPase in maize coleoptiles and enclosed leaves

Planta, 185 (1991), pp. 458-461

[View Record in Scopus](#)

Walker and Connolly, 2008

E.L. Walker, E.L. Connolly

Time to pump iron: iron-deficiency-signaling mechanisms of higher plants

Curr Opin Plant Biol, 11 (2008), pp. 530-535

[ArticlePDF \(355KB\)View Record in Scopus](#)

Wittenberg et al., 1996

J.B. Wittenberg, B.A. Wittenberg, D.A. Day, M.K. Udvardi, C.A. Appleby

Siderophore-bound iron in the peribacteroid space of soybean root nodules

Plant Soil, 178 (1996), pp. 161-169

[CrossRefView Record in Scopus](#)

Copyright © 2011 Elsevier GmbH. All rights reserved.