AperTO - Archivio Istituzionale Open Access dell'Università di Torino

EXCI-CEST: Exploiting pharmaceutical excipients as MRI-CEST contrast agents for tumor imaging

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1658308> since 2018-01-19T14:36:40Z

Published version:

DOI:10.1016/j.ijpharm.2017.04.040

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Dario Livio Longo, Fatima Zzahra Moustaghfir, Alexandre Zerbo, Lorena Consolino,
Annasofia Anemone, Martina Bracesco and Silvio Aime

EXCI-CEST: Exploiting pharmaceutical excipients as MRI-CEST contrast agents for
tumor imaging

In INTERNATIONAL JOURNAL OF PHARMACEUTICS 2017, 525, 275

The definitive version is available at:

DOI: 10.1016/j.ijpharm.2017.04.040

23 **Title page**

24 *Title*

25 **EXCI-CEST: exploiting pharmaceutical excipients as MRI-CEST contrast agents for tumor**
26 **imaging**

27

28 *Authors:*

29 Dario Livio Longo^{1,*} Fatima Zzahra Moustaghfir^{2,3}, Alexandre Zerbo², Lorena Consolino²,
30 Annasofia Anemone², Martina Bracesco² and Silvio Aime²

31 ¹Istituto di Biostrutture e Bioimmagini (IBB), Consiglio Nazionale delle Ricerche (CNR), Via
32 Nizza 52, 10126, Torino, Italy

33 ²Dipartimento di Biotecnologie Molecolari e Scienze per la Salute, Università degli Studi di Torino,
34 Via Nizza 52, 10126, Torino, Italy

35

36 *Corresponding author:*

37 Dario Livio Longo, Istituto di Biostrutture e Bioimmagini, (CNR) c/o Centro di Biotecnologie
38 Molecolari, Via Nizza 52, 10126, Torino, Italy

39 Phone: +39-011-6706473, Fax: +39-011-6706487, email: dario.longo@unito.it.

40

41 *Present address:*

42 ³ Fatima Zzahra Moustaghfir: MSD Belgium, Lynx Binnenhof 5, 1200, Brussels, Belgium

43

44 **Abstract**

45 Chemical Exchange Saturation Transfer (CEST) approach is a novel tool within magnetic resonance
46 imaging (MRI) that allows visualization of molecules possessing exchangeable protons with water.
47 Many molecules, employed as excipients for the formulation of finished drug products, are
48 endowed with hydroxyl, amine or amide protons, thus can be exploitable as MRI-CEST contrast
49 agents. Their high safety profiles allow them to be injected at very high doses. Here we investigated
50 the MRI-CEST properties of several excipients (ascorbic acid, sucrose, N-acetyl-D-glucosamine,
51 meglumine and 2-pyrrolidone) and tested them as tumor-detecting agents in two different murine
52 tumor models (breast and melanoma cancers). All the investigated molecules showed remarkable
53 CEST contrast upon i.v. administration in the range 1-3 ppm according to the type of mobile proton
54 groups. A marked increase of CEST contrast was observed in tumor regions up to 30 min post
55 injection. The combination of marked tumor contrast enhancement and lack of toxicity make these
56 molecules potential candidates for the diagnosis of tumors within the MRI-CEST approach.

57

58

59 **Keywords**

60 Excipients, MRI, CEST, tumor, imaging, chemical exchange saturation transfer

61

62 **Chemical Compounds studied in this article**

63 Ascorbic acid (PubChem CID: 54670067);

64 Meglumine (PubChem CID: 8567);

65 Sucrose (PubChem CID: 5988);

66 N-acetyl-D-glucosamine (PubChem CID: 439174);

67 2-Pyrrolidone (PubChem CID: 12025);

68

69 **Abbreviations**

70 MRI: Magnetic Resonance Imaging

71 CEST: Chemical Exchange Saturation Transfer

72 i.v.: intravenous

73

74 **1. Introduction**

75 Medicines could not be made without the use of pharmaceutical excipients that contribute notably
76 to guarantee efficacy and safety of the final pharmaceutical product (Casas et al., 2015). Moreover,
77 excipients perform multiple functions, besides completing the formulation volume, such as
78 improving bioavailability, administration and acceptance of the treatment by the patient (Loftsson,
79 2015; Narayan, 2011; Wening and Breitzkreutz, 2011). Another fundamental characteristic of
80 excipients is their pharmacological and toxicological inactivity that allows them to be used at high
81 doses (Abrantes et al., 2016). Several natural products, simple substances and mixtures are
82 commonly used in formulating medicines, with chemical structures that vary from small molecules
83 to polymers.

84 Interestingly, most, if not all of these molecules, possess exchangeable protons (hydroxyl, amine,
85 amide groups) that can be potentially detected by chemical exchange saturation transfer (CEST)
86 magnetic resonance imaging (MRI) (van Zijl and Yadav, 2011; Vinogradov et al., 2013). This
87 technique enables the indirect visualization of molecules via magnetization transfer between
88 exchangeable protons and bulk water protons. By applying a selective radiofrequency irradiation to
89 the mobile protons, the induced saturation is transferred to the bulk water protons, thus inducing a
90 reduction of the water signal (Liu et al., 2013). Several natural molecules and polymers (glucose,

glycogen, glycosaminoglycan, sialic acid, gelatin) have already been exploited as MRI-CEST contrast agents, since these molecules have precedence of use with human exposure (Chan et al., 2012; Jin et al., 2017; Liang et al., 2015; Ling et al., 2008; Shinar et al., 2014; van Zijl et al., 2007; Walker-Samuel et al., 2013). Also metabolites, drugs and polypeptides/proteins have been investigated to demonstrate their capability to generate contrast within this approach (Bar-Shir et al., 2015; Bar-Shir et al., 2014; Cai et al., 2015; Haris et al., 2012; Li et al., 2016; Liu et al., 2016; Longo et al., 2014a; McMahon et al., 2008; Zaiss et al., 2013). Moreover, several diamagnetic CEST agents have been proposed as exogenous probes for tumor imaging (Geraldès and Laurent, 2009). However, diamagnetic molecules require high doses to discriminate their contrast from direct water saturation and from endogenous magnetization transfer effects, due to the small chemical shift difference from water.

These considerations limit the effective use of exogenous molecules as CEST agents to those possessing low *in vivo* toxicity. According to these considerations, researchers firstly turned their attention to already clinically approved contrast agents, such as iodinated contrast media, exploiting their high safety profile and FDA approval (Aime et al., 2005b; Longo et al., 2011). Consequently, radiographic agents have been exploited for assessing tumor microenvironment properties, including perfusion (Anemone et al., 2017; Longo et al., 2016b), acidosis (Chen et al., 2015; Jones et al., 2015; Longo et al., 2016a; Longo et al., 2014b; Sun et al., 2014) and for assessing renal functionality (Longo et al., 2013; Longo et al., 2017).

Pharmaceutical excipients have attracted our interest since they can be used at very high dose due to their well-known safety profiles. In addition, excipients do not have any pharmacological effects, in contrast to active pharmaceutical ingredients. Ideally, a MRI-CEST contrast agent should possess good solubility and high safety profile, it should accumulate enough in the region of interest to produce contrast; afterwards, it should be excreted through the kidneys without long-term accumulation (Aime et al., 2005a; Sherry et al., 2009). The present investigation reports the MRI-

116 CEST properties of several pharmaceutical excipients (sucrose, N-acetyl-D-glucosamine, ascorbic
117 acid, meglumine and 2-pyrrolidone), as novel, biocompatible MRI contrast agents for molecular
118 imaging of tumors. We describe the *in vitro* MRI-CEST contrast enhancing properties and the *in*
119 *vivo* investigation of these molecules in two murine tumor models.

120

121 **2. Methods**

122 *2.1 Materials*

123 All chemicals (Sucrose, N-acetyl-D-glucosamine, Meglumine, 2-pyrrolidone, Ascorbic acid) were
124 purchased from Sigma-Aldrich (Sigma Aldrich, Milan, Italy).

125

126 *2.2 In vitro MRI CEST acquisition*

127 Phantoms containing vials of phosphate buffer solution of Sucrose, N-acetyl-D-glucosamine,
128 Meglumine, 2-pyrrolidone or ascorbic acid were prepared at a concentration of 30mM and titrated
129 over a range 6-7.4 pH units. CEST-MRI experiments were performed on a vertical 7 T MRI scanner
130 Bruker Avance 300 (Bruker, Ettlingen, Germany) using a fast spin-echo sequence with centric
131 encoding after presaturation pulses varying in power (1.5, 2.0, 3.0 and 6.0 μ T) for 5 or 7s at 37°C.
132 A modified RARE sequence including a magnetization transfer module was used to acquire CEST-
133 weighted images from -10 to +10 ppm with increments of 0.1 ppm around the water resonance.

134

135 *2.3 Cell lines for xenograft tumor models*

136 TS/A cells, derived from a metastasizing mouse cell line, originated from a mammary
137 adenocarcinoma which arose spontaneously in a BALB/c female, were grown in RPMI 1640
138 medium supplemented with 10% fetal bovine serum (FBS), 100U/mL Penicillin with 100 μ g/mL
139 Streptomycin (Pen/Strep) and 2mM L-Glutamine (Nanni et al., 1983). B16-F10 cells, an

140 established murine melanoma cell line, were cultured in DMEM supplemented with 10% FBS, 100
141 $\mu\text{g/ml}$ penicillin and 100 $\mu\text{g/ml}$ streptomycin. B16-F10 cells were obtained from American Type
142 Culture Collection (ATCC).

143

144 *2.4 Animal experiments*

145 6-old-week female BALB/c mice (n=5 for each molecule) were inoculated subcutaneously with 2.5
146 $\times 10^5$ TS/A cells in 100 μL of PBS on both flanks and 6-old-week male C57BL/6 mice (n=5 for
147 each molecule) were inoculated subcutaneously with 3×10^5 B16-f10 cell in 100 μL of PBS on both
148 flanks. BALB/c and C57BL/6 mice (Charles River Laboratories Italia S.r.l., Calco Italia) were
149 maintained under specific pathogen free conditions in the animal facility of the Molecular
150 Biotechnology Center, University of Turin, and treated in accordance with the EU guidelines
151 (EU2010/63). All in vivo studies were conducted according to approved procedures of the
152 Institutional Animal Care and Use Committee of the University of Torino.

153 Before imaging, mice were anaesthetized with a mixture of tiletamine/zolazepam (Zoletil 100;
154 Vibac, Milan, Italy) 20mg/kg and xylazine (Rompun; Bayer, Milan, Italy) 5mg/kg and during the
155 acquisition their breath rate was monitored throughout in vivo MRI experiments using a respiratory
156 probe. Cannulation of the lateral tail vein with a catheter was exploited for intravenous injection of
157 the investigated molecules.

158

159 *2.5 In vivo MRI CEST acquisition and analysis*

160 A Bruker 7T Avance 300 MRI scanner (Bruker Biospin, Ettlingen, Germany) equipped with a 30
161 mm ^1H quadrature coil was used to scan mammary adenocarcinoma (TS/A cell line) and melanoma
162 (B16-f10 cell line) tumor bearing mice 15 days post-inoculation. After the scout image acquisition,
163 T_{2w} anatomical images were acquired with a Fast Spin Echo sequence and the same geometry was

164 used for the following CEST experiments. CEST images were acquired with a single shot FSE
165 sequence with centric encoding (TR: 6000 ms, TE: 4.0 ms) after a CW-RF presaturation pulse of B_1
166 = 1.5 μ T x 5s from a single axial slice with high in-plane resolution of 234 μ m (FOV 3 cm, MTX
167 96, zero filled to 128, slice thickness 1.5mm) with 55 frequency offsets unevenly spaced in the
168 range ± 10 ppm. Each investigated molecule was administrated intravenously at the dose of 1.2 g/kg
169 b.w. with a single bolus of 100 μ L followed by continuous infusion at a rate of 500 μ L/h and CEST
170 images were acquired before and every 10 minutes up to 30 minutes.

171 CEST images were analyzed using homemade scripts implemented in MATLAB (The Mathworks,
172 Inc, Natick, MA). The Z-spectra were interpolated, on a voxel-by-voxel basis, by smoothing
173 splines, B_0 -shift corrected and saturation transfer efficiency (ST%) was measured by punctual
174 analysis at 1.2 ppm (Terreno et al., 2009). For in vivo images, difference contrast maps (Δ ST%)
175 were calculated by subtracting the ST contrast after each molecules injection from the ST contrast
176 before the injection on a per voxel basis. Extravasation fraction of each molecule was calculated as
177 the percentage of pixels showing a Δ ST% above the threshold (2%) in the manually-defined tumor
178 region of interest.

179

180 *2.6 Statistical analysis*

181 Calculations were performed using GraphPad Prism (GraphPad Software, La Jolla, CA, USA)
182 software package; data are presented as mean $\pm$ SD unless otherwise stated. Statistical significance
183 was established at $P < 0.05$.

184

185 **3. Results**

186 *3.1 In vitro characterization of CEST properties*

187 The investigated molecules differ for the types of mobile protons, i.e. belonging to the hydroxyl,
188 amide and amine groups, and for the number of exchanging protons (Figure 1). The ability to yield
189 CEST contrast in the MR images is shown in Figure 2, where the contrast efficiency (ST effect) is
190 plotted as a function of the chemical shift. Sucrose, ascorbic acid, meglumine and N-acetyl-D-
191 glucosamine show CEST contrast peaking at ca. 0.7-1.2 ppm, due to the presence of hydroxyl
192 protons. 2-Pyrrolidone showed a small CEST contrast at 0.7 ppm downfield from water, due to the
193 presence of a cyclic amide (lactam) group. Both meglumine and N-acetyl-D-glucosamine showed,
194 in addition to the less shifted hydroxyl protons, a second broad CEST contrast peak between 2 and
195 3.5 ppm, due to amine and amide protons, respectively. As shown in Figure 2, when keeping
196 constant the saturation field strength ($1.5 \mu\text{T} \times 7\text{s}$), the CEST contrast showed a significant
197 dependence with pH. The hydroxyl mobile protons of sucrose showed higher CEST contrast values
198 at lower pH values (Figure 3A). Conversely, the hydroxyl protons of N-acetyl-D-glucosamine,
199 meglumine and ascorbic acid showed a steady increase of the CEST contrast at high pH values, at
200 all the investigated saturation field strengths ($1.5 - 6 \mu\text{T}$, Figures 3B, D and E). The CEST contrast
201 of 2-pyrrolidone was almost independent from pH (Figure 3C). Sucrose, meglumine, ascorbic acid
202 and N-acetyl-D-glucosamine showed in vitro a ST efficiency close to 1% per 1 mM concentration
203 (irradiation pulse of $1.5 \mu\text{T} \times 5\text{s}$, pH 7, $T=37^\circ\text{C}$; Figure 3F). The highest CEST contrast efficiency
204 observed for sucrose (1.5 % ST per 1 mM concentration) likely depends on the large number of
205 mobile protons (8 $-\text{OH}$ protons), whereas 2-pyrrolidone, having only one exchanging proton,
206 showed the lowest CEST contrast.

207

208 3.2 *In vivo* CEST detection in tumor murine models

209 Two cancer xenograft models, TS/A (highly metastatic mouse breast cancer cells) and B16 (mouse
210 melanoma cancer cells) were used for *in vivo* experiments. CEST agents were administered at the
211 same dose by intravenous injection through the tail vein. A pronounced increase in the CEST

contrast in both TS/A and B16 tumor models for all the investigated molecules, with an average ΔST increase in the range 2- 6% in comparison to the pre-contrast ST values was observed (Figure 4). Sucrose showed a slightly higher CEST contrast in the B16 model ($\Delta ST=4.1 \pm 0.7\%$) in respect to TS/A tumors ($\Delta ST=2.7 \pm 0.5\%$) at all the investigated time points (Figure 4A). N-Acetyl-D-glucosamine slightly raised the CEST effect from the baseline with no difference between the two tumor models ($\Delta ST=2.1 \pm 0.5\%$ and $2.5 \pm 0.5\%$ for B16 and TS/A, respectively, Figure 4B). The CEST signal of meglumine increased by $4.1 \pm 1.0\%$ for the B16 tumors and by $2.5 \pm 0.4\%$ for the TS/A tumors (Figure 4C), at 10 min post injection ($P<0.05$). Ascorbic acid showed the highest enhancement in TS/A tumors ($\Delta ST=5.4 \pm 1.1\%$) in comparison to the B16 model ($\Delta ST=2.7 \pm 0.7\%$), with statistically significant difference already 10 min after the i.v. administration ($P<0.05$, Figure 4D). For all the investigated molecules the CEST contrast measured in tumors persisted up to 30 min after the administration. Representative CEST contrast maps overlaid on anatomical images show the differential enhancement pattern among the investigated excipients within the investigated two murine tumor models (Figure 5). In particular, ΔST images showed a marked increase in CEST contrast in B16 and TS/A tumors for meglumine and ascorbic acid, respectively. The percentage of the tumor pixels where the CEST effect was detectable (ΔST higher than 2%) is dependent on both the molecules and the tumor model (Figure 6). In particular, CEST contrast increase was higher for the B16 model than for the TS/A one. Sucrose was detected in 40-60% of the tumor area according to the tumor model, whereas all the other molecules showed a detection fraction lower than 50% of the whole tumor volume.

232

233 4. Discussion

234 This study demonstrated that several excipients can be exploited as MRI-CEST contrast agents for
235 tumor detection in mice. A moderate to marked increase in CEST contrast in the tumor region was
236 detected up to 30 min following intravenous administration, according to the exploited agents or the

237 investigated tumor models. Meglumine and ascorbic acid yielded the highest contrast enhancement
238 ($\Delta ST > 5\%$) in B16 and TS/A models, respectively (Figure 4). Conversely, lower CEST
239 enhancements were measured for sucrose and N-acetyl-D-glucosamine. The potential of glucose
240 and its derivatives to act as MRI-CEST agents for tumor imaging has already been demonstrated
241 (Chan et al., 2012; Walker-Samuel et al., 2013; Xu et al., 2015). However, for glucose, the main
242 drawback was associated to its rapid metabolism once entered in the tumor cells, with consequent
243 reduction of CEST contrast capabilities. For this reason, glucose analogs, such as 2-deoxy-glucose
244 (2DG) and 3-oxy-methyl-glucose (3OMG) have been proposed as they showed superior contrast
245 efficiency owing to the reduced metabolic conversion in the case of 2DG (Nasrallah et al., 2013;
246 Rivlin et al., 2013) or to the lack of metabolic transformation in the case of 3OMG (Rivlin et al.,
247 2014). As a consequence, such derivatives provide an improved and long-lasting CEST contrast in
248 mice carrying xenograft tumors. On the other hand, the safety of these compounds has still to be
249 demonstrated, since the high concentrations ($> 1\text{--}1.5\text{ g/kg b.w.}$) required to generate enough CEST
250 contrast might limit their use to laboratory animals. An analogous limitation may be envisaged for
251 the recently reported CEST agents based on salicylic acid (a metabolite of aspirin), although their
252 very large chemical shift difference (unusual for diamagnetic CEST agents) holds promise for
253 applications at lower magnetic fields (Lesniak et al., 2016; Yang et al., 2013). Conversely, the
254 herein investigated excipients hold a very high safety profile, considering that they are used at high
255 dosages to provide suitable formulations for drugs.

256 In contrast to N-acetyl-D-glucosamine and ascorbic acid that can be metabolized within the body,
257 sucrose (when injected i.v.) and meglumine are rapidly excreted unchanged in the urine, with no
258 evidence for metabolism (Heeg et al., 1977). Clearly, this represents a great advantage in
259 comparison to metabolizable probes that do not accumulated in the extracellular extravascular space
260 of tumors. Furthermore, metabolic products cannot provide enough CEST contrast as the original
261 molecule, with a following decrease of their observed CEST contribution. This may partly explain

the non-optimal CEST contrast in tumors upon N-acetyl-D-glucosamine administration, in comparison to the other excipients, despite the high *in vitro* CEST contrast efficiency. Similar findings for N-acetyl-D-glucosamine were observed by Navon group, who investigated the CEST properties of this molecule and of glucosamine as alternatives to glucose analogs (Rivlin and Navon, 2016). In our study, the observed increase in CEST contrast in tumors with N-Acetyl-D-Glucosamine was lower ($\Delta ST = 2\text{-}3\%$) than what previously reported ($\Delta ST = 6\text{-}7\%$), albeit a similar dosage was administered. These observations may be accounted by the use of different tumor cell lines or by the lower saturation pulse power that has been applied in our experimental protocol ($1.5\ \mu\text{T}$ vs $2.4\ \mu\text{T}$).

Meglumine showed distinct contrast enhancement capabilities between TS/A and B16 tumors. Meglumine is not internalized inside cells, therefore differences in CEST contrast enhancements are only dependent on the accumulation within the extracellular extravascular space, hence reflecting different vascularization properties. As such, this molecule can be considered an extracellular-fluid agent analogous to the clinically approved Gadolinium-based contrast agents (Morana et al., 2013) or to Iodine-containing X-ray systems (Rutten and Prokop, 2007). Thus, meglumine may be an attractive candidate to be used as MRI-CEST contrast agents for tumor imaging with remarkable contrast efficiency.

5. Conclusions

The herein investigated excipients show remarkable MRI-CEST properties as demonstrated by the *in vivo* visualization of tumors in two murine models. The extremely good safety profile of the excipients provides support to the view that these systems may be considered reliable candidates for clinical translation.

286 **Acknowledgements:**

287 The study was supported by the EU Horizon 2020 GLINT project (grant #667510).

288

289 **References**

- 290 Abrantes, C.G., Duarte, D., Reis, C.P., 2016. An Overview of Pharmaceutical Excipients: Safe or Not Safe? J.
291 Pharm. Sci. 105, 2019-2026.
- 292 Aime, S., Botta, M., Terreno, E., 2005a. Gd(III)-based contrast agents for MRI. Advances in Inorganic
293 Chemistry - Including Bioinorganic Studies, Vol 57 57, 173-237.
- 294 Aime, S., Calabi, L., Biondi, L., De Miranda, M., Ghelli, S., Paleari, L., Rebaudengo, C., Terreno, E., 2005b.
295 Iopamidol: Exploring the potential use of a well-established x-ray contrast agent for MRI. Magn.
296 Reson. Med. 53, 830-834.
- 297 Anemone, A., Consolino, L., Longo, D.L., 2017. MRI-CEST assessment of tumour perfusion using X-ray
298 iodinated agents: comparison with a conventional Gd-based agent. Eur. Radiol. 27, 2170-2179.
- 299 Bar-Shir, A., Liang, Y., Chan, K.W., Gilad, A.A., Bulte, J.W., 2015. Supercharged green fluorescent proteins as
300 bimodal reporter genes for CEST MRI and optical imaging. Chem Commun (Camb) 51, 4869-4871.
- 301 Bar-Shir, A., Liu, G., Chan, K.W., Oskolkov, N., Song, X., Yadav, N.N., Walczak, P., McMahon, M.T., van Zijl,
302 P.C., Bulte, J.W., Gilad, A.A., 2014. Human protamine-1 as an MRI reporter gene based on chemical
303 exchange. ACS Chem Biol 9, 134-138.
- 304 Cai, K., Singh, A., Poptani, H., Li, W., Yang, S., Lu, Y., Hariharan, H., Zhou, X.J., Reddy, R., 2015. CEST signal at
305 2ppm (CEST@2ppm) from Z-spectral fitting correlates with creatine distribution in brain tumor.
306 NMR Biomed. 28, 1-8.
- 307 Casas, M., Aguilar-de-Leyva, A., Caraballo, I., 2015. Towards a rational basis for selection of excipients:
308 Excipient Efficiency for controlled release. Int. J. Pharm. 494, 288-295.

309 Chan, K.W., McMahon, M.T., Kato, Y., Liu, G., Bulte, J.W., Bhujwala, Z.M., Artemov, D., van Zijl, P.C., 2012.
 310 Natural D-glucose as a biodegradable MRI contrast agent for detecting cancer. *Magn. Reson. Med.*
 311 68, 1764-1773.

312 Chen, L.Q., Randtke, E.A., Jones, K.M., Moon, B.F., Howison, C.M., Pagel, M.D., 2015. Evaluations of Tumor
 313 Acidosis Within In Vivo Tumor Models Using Parametric Maps Generated with AcidoCEST MRI. *Mol*
 314 *Imaging Biol.*

315 Geraldles, C.F., Laurent, S., 2009. Classification and basic properties of contrast agents for magnetic
 316 resonance imaging. *Contrast Media Mol. Imaging* 4, 1-23.

317 Haris, M., Nanga, R.P., Singh, A., Cai, K., Kogan, F., Hariharan, H., Reddy, R., 2012. Exchange rates of creatine
 318 kinase metabolites: feasibility of imaging creatine by chemical exchange saturation transfer MRI.
 319 *NMR Biomed.* 25, 1305-1309.

320 Heeg, J.F., Born, G.S., Kessler, W.V., Shaw, S.M., Lange, W.E., 1977. Absorption, distribution, and excretion
 321 of 14C-meglumine in rats and dogs after administration of 14C-meglumine salicylate. *J. Pharm. Sci.*
 322 66, 96-99.

323 Jin, T., Nicholls, F.J., Crum, W.R., Ghuman, H., Badylak, S.F., Modo, M., 2017. Diamagnetic chemical
 324 exchange saturation transfer (diaCEST) affords magnetic resonance imaging of extracellular matrix
 325 hydrogel implantation in a rat model of stroke. *Biomaterials* 113, 176-190.

326 Jones, K.M., Randtke, E.A., Howison, C.M., Cardenas-Rodriguez, J., Sime, P.J., Kottmann, M.R., Pagel, M.D.,
 327 2015. Measuring Extracellular pH in a Lung Fibrosis Model with acidoCEST MRI. *Mol Imaging Biol*
 328 17, 177-184.

329 Lesniak, W.G., Oskolkov, N., Song, X., Lal, B., Yang, X., Pomper, M., Laterra, J., Nimmagadda, S., McMahon,
 330 M.T., 2016. Salicylic Acid Conjugated Dendrimers Are a Tunable, High Performance CEST MRI
 331 NanoPlatform. *Nano Lett* 16, 2248-2253.

332 Li, Y., Chen, H., Xu, J., Yadav, N.N., Chan, K.W., Luo, L., McMahon, M.T., Vogelstein, B., van Zijl, P.C., Zhou,
 333 S., Liu, G., 2016. CEST theranostics: label-free MR imaging of anticancer drugs. *Oncotarget* 7, 6369-
 334 6378.

335 Liang, Y., Bar-Shir, A., Song, X., Gilad, A.A., Walczak, P., Bulte, J.W., 2015. Label-free imaging of gelatin-
336 containing hydrogel scaffolds. *Biomaterials* 42, 144-150.

337 Ling, W., Regatte, R.R., Navon, G., Jerschow, A., 2008. Assessment of glycosaminoglycan concentration in
338 vivo by chemical exchange-dependent saturation transfer (gagCEST). *Proc. Natl. Acad. Sci. U. S. A.*
339 105, 2266-2270.

340 Liu, G., Song, X., Chan, K.W., McMahon, M.T., 2013. Nuts and bolts of chemical exchange saturation
341 transfer MRI. *NMR Biomed.* 26, 810-828.

342 Liu, H., Jablonska, A., Li, Y., Cao, S., Liu, D., Chen, H., Van Zijl, P.C., Bulte, J.W., Janowski, M., Walczak, P., Liu,
343 G., 2016. Label-free CEST MRI Detection of Citicoline-Liposome Drug Delivery in Ischemic Stroke.
344 *Theranostics* 6, 1588-1600.

345 Loftsson, T., 2015. Excipient pharmacokinetics and profiling. *Int. J. Pharm.* 480, 48-54.

346 Longo, D.L., Bartoli, A., Consolino, L., Bardini, P., Arena, F., Schwaiger, M., Aime, S., 2016a. In Vivo Imaging
347 of Tumor Metabolism and Acidosis by Combining PET and MRI-CEST pH Imaging. *Cancer Res.* 76,
348 6463-6470.

349 Longo, D.L., Busato, A., Lanzardo, S., Antico, F., Aime, S., 2013. Imaging the pH evolution of an acute kidney
350 injury model by means of iopamidol, a MRI-CEST pH-responsive contrast agent. *Magn. Reson. Med.*
351 70, 859-864.

352 Longo, D.L., Cutrin, J.C., Michelotti, F., Irrera, P., Aime, S., 2017. Noninvasive evaluation of renal pH
353 homeostasis after ischemia reperfusion injury by CEST-MRI. *NMR Biomed.* doi: 10.1002/nbm.3720

354 Longo, D.L., Dastru, W., Digilio, G., Keupp, J., Langereis, S., Lanzardo, S., Prestigio, S., Steinbach, O., Terreno,
355 E., Uggeri, F., Aime, S., 2011. Iopamidol as a responsive MRI-chemical exchange saturation transfer
356 contrast agent for pH mapping of kidneys: In vivo studies in mice at 7 T. *Magn. Reson. Med.* 65,
357 202-211.

358 Longo, D.L., Di Gregorio, E., Abategiovanni, R., Ceccon, A., Assfalg, M., Molinari, H., Aime, S., 2014a.
359 Chemical exchange saturation transfer (CEST): an efficient tool for detecting molecular information
360 on proteins' behaviour. *Analyst* 139, 2687-2690.

361 Longo, D.L., Michelotti, F., Consolino, L., Bardini, P., Digilio, G., Xiao, G., Sun, P.Z., Aime, S., 2016b. In Vitro
 362 and In Vivo Assessment of Nonionic Iodinated Radiographic Molecules as Chemical Exchange
 363 Saturation Transfer Magnetic Resonance Imaging Tumor Perfusion Agents. *Invest. Radiol.* 51, 155-
 364 162.

365 Longo, D.L., Sun, P.Z., Consolino, L., Michelotti, F.C., Uggeri, F., Aime, S., 2014b. A general MRI-CEST
 366 ratiometric approach for pH imaging: demonstration of in vivo pH mapping with iobitridol. *J. Am.*
 367 *Chem. Soc.* 136, 14333-14336.

368 McMahon, M.T., Gilad, A.A., DeLiso, M.A., Berman, S.M., Bulte, J.W., van Zijl, P.C., 2008. New "multicolor"
 369 polypeptide diamagnetic chemical exchange saturation transfer (DIACEST) contrast agents for MRI.
 370 *Magn. Reson. Med.* 60, 803-812.

371 Morana, G., Cugini, C., Scatto, G., Zanato, R., Fusaro, M., Dorigo, A., 2013. Use of contrast agents in
 372 oncological imaging: magnetic resonance imaging. *Cancer Imaging* 13, 350-359.

373 Nanni, P., de Giovanni, C., Lollini, P.L., Nicoletti, G., Prodi, G., 1983. TS/A: a new metastasizing cell line from
 374 a BALB/c spontaneous mammary adenocarcinoma. *Clin. Exp. Metastasis* 1, 373-380.

375 Narayan, P., 2011. Overview of drug product development. *Curr Protoc Pharmacol* Chapter 7, Unit 7 3 1-29.

376 Nasrallah, F.A., Pages, G., Kuchel, P.W., Golay, X., Chuang, K.H., 2013. Imaging brain deoxyglucose uptake
 377 and metabolism by glucoCEST MRI. *J. Cereb. Blood Flow Metab.* 33, 1270-1278.

378 Rivlin, M., Horev, J., Tsarfaty, I., Navon, G., 2013. Molecular imaging of tumors and metastases using
 379 chemical exchange saturation transfer (CEST) MRI. *Sci Rep* 3, 3045.

380 Rivlin, M., Navon, G., 2016. Glucosamine and N-acetyl glucosamine as new CEST MRI agents for molecular
 381 imaging of tumors. *Sci Rep* 6, 32648.

382 Rivlin, M., Tsarfaty, I., Navon, G., 2014. Functional molecular imaging of tumors by chemical exchange
 383 saturation transfer MRI of 3-O-Methyl-D-glucose. *Magn. Reson. Med.* 72, 1375-1380.

384 Rutten, A., Prokop, M., 2007. Contrast agents in X-ray computed tomography and its applications in
 385 oncology. *Anticancer Agents Med Chem* 7, 307-316.

386 Sherry, A.D., Caravan, P., Lenkinski, R.E., 2009. Primer on gadolinium chemistry. *J. Magn. Reson. Imaging* 30,
387 1240-1248.

388 Shinar, H., Battistel, M.D., Mandler, M., Lichaa, F., Freedberg, D.I., Navon, G., 2014. Sialo-CEST: chemical
389 exchange saturation transfer NMR of oligo- and poly-sialic acids and the assignment of their
390 hydroxyl groups using selective- and HSQC-TOCSY. *Carbohydr. Res.* 389, 165-173.

391 Sun, P.Z., Longo, D.L., Hu, W., Xiao, G., Wu, R., 2014. Quantification of iopamidol multi-site chemical
392 exchange properties for ratiometric chemical exchange saturation transfer (CEST) imaging of pH.
393 *Phys. Med. Biol.* 59, 4493-4504.

394 Terreno, E., Stancanello, J., Longo, D., Castelli, D.D., Milone, L., Sanders, H.M., Kok, M.B., Uggeri, F., Aime,
395 S., 2009. Methods for an improved detection of the MRI-CEST effect. *Contrast Media Mol. Imaging*
396 4, 237-247.

397 van Zijl, P.C., Jones, C.K., Ren, J., Malloy, C.R., Sherry, A.D., 2007. MRI detection of glycogen in vivo by using
398 chemical exchange saturation transfer imaging (glycoCEST). *Proc. Natl. Acad. Sci. U. S. A.* 104, 4359-
399 4364.

400 van Zijl, P.C., Yadav, N.N., 2011. Chemical exchange saturation transfer (CEST): what is in a name and what
401 isn't? *Magn. Reson. Med.* 65, 927-948.

402 Vinogradov, E., Sherry, A.D., Lenkinski, R.E., 2013. CEST: from basic principles to applications, challenges
403 and opportunities. *J. Magn. Reson.* 229, 155-172.

404 Walker-Samuel, S., Ramasawmy, R., Torrealdea, F., Rega, M., Rajkumar, V., Johnson, S.P., Richardson, S.,
405 Goncalves, M., Parkes, H.G., Arstad, E., Thomas, D.L., Pedley, R.B., Lythgoe, M.F., Golay, X., 2013. In
406 vivo imaging of glucose uptake and metabolism in tumors. *Nat. Med.* 19, 1067-1072.

407 Wening, K., Breitzkreutz, J., 2011. Oral drug delivery in personalized medicine: unmet needs and novel
408 approaches. *Int. J. Pharm.* 404, 1-9.

409 Xu, X., Chan, K.W., Knutsson, L., Artemov, D., Xu, J., Liu, G., Kato, Y., Lal, B., Larterra, J., McMahon, M.T., van
410 Zijl, P.C., 2015. Dynamic glucose enhanced (DGE) MRI for combined imaging of blood-brain barrier
411 break down and increased blood volume in brain cancer. *Magn. Reson. Med.* 74, 1556-1563.

412 Yang, X., Song, X., Li, Y., Liu, G., Ray Banerjee, S., Pomper, M.G., McMahon, M.T., 2013. Salicylic acid and
413 analogues as diaCEST MRI contrast agents with highly shifted exchangeable proton frequencies.
414 Angew. Chem. Int. Ed. Engl. 52, 8116-8119.

415 Zaiss, M., Kunz, P., Goerke, S., Radbruch, A., Bachert, P., 2013. MR imaging of protein folding in vitro
416 employing nuclear-Overhauser-mediated saturation transfer. NMR Biomed. 26, 1815-1822.

417

418

419 **Figure Legends**

420 **Figure 1.** Chemical structures and molecular weights (g/mol) of the investigated excipients.

421 **Figure 2.** CEST contrast curves for the investigated molecules (sucrose, black; N-acetyl-D-
422 glucosamine, red; 2-Pyrrolidone, grey; meglumine, green; ascorbic acid, blue) obtained at
423 concentration of 30 mM at pH 6 (A) and pH 7.4 (B) using a saturation power level of 1.5 μ T with
424 duration of 7s at 7T and 37°C.

425 **Figure 3.** CEST contrast dependence on pH measured in the range of 6-7.4 pH units at different B_1
426 levels (saturation power from 1.5 μ T to 6 μ T for 5s, 7T, 37°C) for sucrose (A), N-acetyl D-
427 glucosamine (B), 2-Pyrrolidone (C), meglumine (D) and ascorbic acid (E).

428 **Figure 4.** Box-plots showing averaged mean Δ ST increase (calculated as ST post -ST pre injection)
429 in TS/A (grey bars) and B16 (black bar) tumor regions using $B_1 = 1.5\mu$ Tx5s on a 7T MRI scanner.
430 CEST contrast observed after i.v. administration of sucrose (A), N-acetyl-D-glucosamine (B),
431 meglumine (C) and ascorbic acid (D) at a dose of 1.2 g/kg b.w. was observed at 10, 20 and 30
432 minutes post-injection.

433 **Figure 5.** Representative Δ ST tumor maps superimposed on anatomical images showing CEST
434 contrast increase for sucrose, N-acetyl-D-glucosamine, meglumine and ascorbic acid 20 min after
435 i.v. administration using $B_1 = 1.5 \mu$ T x 5s in B16 (top row) and TS/A (bottom row) tumors.

436 **Figure 6.** Box-plot showing extravasation fraction calculated as percentage of pixels with ΔST
437 higher than 2% for sucrose, N-acetyl-D-glucosamine, meglumine and ascorbic acid 20 min after i.v
438 administration using $B_1 = 1.5 \mu T \times 5s$ for B16 (A) and TS/A (B) murine tumors.