

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

### Cytogenetic characterization of an hybrid goat-sheep by GTG-banding

**This is the author's manuscript**

*Original Citation:*

Cytogenetic characterization of an hybrid goat-sheep by GTG-banding / Pauciullo, A; Knorr, C; Perucatti, A; Pieper, SD; Iannuzzi, A; Iannuzzi, L; Erhardt, G. - In: CHROMOSOME RESEARCH. - ISSN 0967-3849. - 23(2015), pp. 127-127.

*Availability:*

This version is available <http://hdl.handle.net/2318/1662763> since 2018-03-18T21:18:18Z

*Published version:*

DOI:10.1007/s10577-015-9476-6

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

Pauciullo A., Knorr C., Perucatti A., Stölzl A., Pieper D., Iannuzzi A., Iannuzzi L., Erhardt G.  
Cytogenetic characterization of an hybrid goat-sheep by GTG-banding. *Chromosome Research*  
(2015) 23 (Suppl. 1): S127

DOI: 10.1007/s10577-015-9476-6

**The publisher's version is available at:**

<https://link.springer.com/article/10.1007%2Fs10577-015-9476-6>

**When citing, please refer to the published version.**

**Link to this full text:**

<https://link.springer.com/article/10.1007%2Fs10577-015-9476-6>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

## Cytogenetic characterization of an hybrid goat-sheep by GTG-banding

Pauciullo A.<sup>1,2,3,\*</sup>, Knorr C.<sup>4</sup>, Perucatti A.<sup>2</sup>, Stölzl A.<sup>5</sup>, Pieper D.<sup>6</sup>, Iannuzzi A.<sup>2</sup>, Iannuzzi L.<sup>2</sup>, Erhardt G.<sup>1</sup>

<sup>1</sup> Institute for Animal Breeding and Genetics, Justus-Liebig-University Ludwigstraße 21b, D-35390 Giessen, Germany

<sup>2</sup> National Research Council (CNR), ISPAAM, Laboratory of Animal Cytogenetics and Gene Mapping, Naples, Italy

<sup>3</sup> Department of Agricultural, Forest and Food Sciences, University of Torino, Grugliasco, Italy

<sup>4</sup> Department for Animal Science, Biotechnology and Reproduction of Livestock, Georg-August-University, Burckhardtweg 2, D-37077 Göttingen, Germany

<sup>5</sup> Department for Animal Science, Livestock Production Systems, Georg-August-University, Albrecht-Thaer-Weg 3, D-37075 Göttingen, Germany

<sup>6</sup> Mariengarten 3, D-37124 Rosdorf, Germany

\* Email: [alfredo.pauciullo@agr.uni-giessen.de](mailto:alfredo.pauciullo@agr.uni-giessen.de)

Phone: +4906419937620

Fax: +4906419937629

In mammals interspecies hybridization occurs rarely under natural conditions. Sheep (*Ovis aries*, 2n=54) and goats (*Capra hircus*, 2n=60) are genetically different and do not readily interbreed, although information of such interspecies matings exists. Cytogenetic incompatibility is one of the causes of embryo death due to incorrect chromosome pairing during the zygote formation and/or aneuploidy occurrence during the zygote division. In this study we report a preliminary cytogenetic characterization of a healthy female hybrid goat-sheep born in a flock under natural conditions.

Peripheral blood sample cultures were performed to get normal lymphocyte cell cultures. After the fixation, a part of the obtained metaphases was stained with a 5% Giemsa solution, whereas a part was treated with 0.05% of trypsin solution and Giemsa staining to obtain the GTG-banding. Conventional and G-banded karyotypes were arranged respectively.

All cells that were observed showed an intermediate karyotype between sheep and goat with 57 chromosomes in total. The G-banding karyotype revealed the presence of 3 metacentric and 54 acrocentric chromosomes. The autosomes involved in the hybrid combination were CHI1,3, CHI2,8 and CHI5,11 corresponding to the metacentric chromosomes OAR1, OAR2 and OAR3. The sex chromosomes were correctly arranged and no further morphological differences were evidenced by classical cytogenetic investigation. Further molecular cytogenetic and genetic analysis are in progress to clarify the recombination events occurred in this rare interspecies hybrid.