AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Sausage fermentation and starter cultures in the era of molecular biology methods

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1680016> since 2021-03-10T10:01:05Z

Published version:

DOI:10.1016/j.ijfoodmicro.2018.04.038

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Sausage fermentation and starter cultures in the era of molecular biology methods**

2

3 Irene Franciosa, Valentina Alessandria, Paola Dolci, Kalliopi Rantsiou, Luca Cocolin*

4

5 University of Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Largo Braccini 2,
6 10095 Grugliasco, Torino, Italy

7

8 *corresponding author: phone +39 011 670-8553, fax +39 011 670-8549, email:
9 lucasimone.cocolin@unito.it

10

Highlights

- Starter cultures in fermented sausages are mixtures of lactobacilli, pediococci and coagulase negative cocci
- Their correct use guarantees improved safety and quality
- The ecology of fermented sausage is very complex and starters have to be able to compete with native microbiota
- *Lactobacillus*, *Staphylococcus*, *Debaryomyces* and *Penicillium* are the main genera involved in sausage fermentation
- Next generation sequencing approaches allow for more in depth studies of the microbial ecology and functionality during fermentation

Abstract

Fermented sausages have a long tradition originating from Europe and they constitute a significant part of the Mediterranean diet. This kind of products has a specific microbiota that is typical of the region or area where they are produced. Therefore, in order to protect the traditional aspect of these products, it is essential to understand the microbial ecology during fermentation by studying the dynamic changes that occur and to select autochthonous starter cultures that can be used in the production. In this paper we summarize the state of the art concerning the selection and use of starter cultures and ecology aspects of naturally fermented sausages. We pay particular attention to the application of bacteriocinogenic strains as they could provide an additional tool in the prevention of foodborne pathogens as well as enhancing the competitiveness of the starter organisms. Microbial ecology of fermented sausages has been determined by traditional microbiological methods, but the introduction in food microbiology of new molecular techniques complements the studies carried out so far and allows scientists to overcome the limitations of traditional methods. Next Generation Sequencing (NGS) techniques represent a change in the way microbiologists address ecology and diversity in foods. Indeed the application of metataxonomics and metagenomics will permit a detailed understanding of microbial ecology. A thorough knowledge of the mechanisms behind the biological processes will enhance meat fermentation control and modulation to obtain products with desired organoleptic properties.

39

40 **Keywords:** Lactic Acid Bacteria; Coagulase Negative Cocci; Starter Cultures; Bioprotection; Ecology;

41 Metataxonomics; Metagenomics

42

43 **1. Introduction**

44 In Europe, dry fermented sausages have a long tradition originating from Mediterranean countries during
45 Roman times. Processing conditions, as well as ingredients and additives, vary among the different types of
46 fermented sausages (Gardini *et al.*, 2001). In fact, ‘typical’ foods of any region or area have their own
47 peculiar characteristics that are deeply rooted in tradition and linked to the territory and which arise from the
48 use of local ingredients and specific production techniques (Aquilanti *et al.*, 2007; Casaburi *et al.*, 2007). The
49 production process begins with small pieces of meat and fat that are minced; salt and spices and in some
50 cases sugar, herbs and/or other ingredients are then added. The homogenised mixture is then stuffed into
51 casings, and undergoes fermentation and drying. European legislation, under Reg. EC 1333/2008 (and
52 subsequent modifications), allows the use of nitrate and nitrite as preservatives, unless subject to other
53 regulations for protected denomination of origin (PDO) products (Aquilanti *et al.*, 2016). The qualitative
54 characteristics of fermented sausages are known to be largely dependent on the quality of the ingredients and
55 raw materials, the specific conditions of the processing and ripening, and the composition of the microbial
56 population (Aquilanti *et al.*, 2007). Pathogenic and spoilage bacteria are inhibited; consequently, the final
57 product has an increased shelf-life (Hugas and Monfort, 1997).

58 Meat fermentations are complex microbial ecosystems in which bacteria, yeasts and molds coexist.
59 Considerable microbial diversity is observed during the fermentation process and is evidenced by the
60 presence of several species belonging to different genera, but also strains of the same species. Through
61 fermentation, highly perishable raw materials, such as meat and fat, are transformed in microbiologically
62 stable final products, characterized by a defined sensory profile, enhanced due to sodium chloride
63 supplementation and to the drying process (Cocolin *et al.*, 2011). Changes that occur during fermentation
64 and drying influence the aroma development in fermented sausages (Flores *et al.*, 2004).

65 Many typical fermented meat products are still produced with traditional technologies without selected
66 starters. However, in the modern sausage production, the use of starter cultures is becoming more frequent to
67 guarantee safety and to standardize product properties, for example consistent flavor and color and shorter
68 ripening time (Cocolin *et al.*, 2001).

69 **2. Starter cultures**

70 Starter cultures are preparations that contain actively growing or resting forms of microorganisms that with
71 their metabolic activity (Fig. 1) impart desired effects during fermentation (Hammes and Hertel, 1998).
72 Industrialized production of starter cultures is a consequence of the gradual shift in sausage production from
73 small local producers to large-scale processing plants and the increasing awareness of the risks for consumer
74 health, in view of overall process efficiency (Magistà *et al.*, 2017). The introduction of starter cultures has
75 become essential in order to shorten the ripening period, ensure colour development, enhance the flavour and
76 improve product safety, given that industrial production of fermented sausages is increasing (Lücke, 1986).
77 In fact, a starter culture should be capable of conducting the fermentation, colonizing the product and
78 dominating over other microorganisms from the beginning to the end of the process (Cocolin *et al.*, 2006).
79 On the other hand, the use of commercially available starters, mainly constituted of lactic acid bacteria and
80 coagulase negative cocci, may result in a loss of peculiar organoleptic characteristics found in spontaneously
81 fermented sausages with an impoverishment of flavor and aroma. For this reason, in several European
82 countries, the artisanal sausages that are manufactured by relying on an unknown ‘factory biota’ are
83 preferred by the consumer (Samelis *et al.*, 1994). The quality of such artisanal, spontaneously fermented
84 sausages possess distinctive characteristics and are often superior if compared to controlled fermentations,
85 inoculated with industrial starters. The principal differences between traditional and industrial fermented
86 product are summarized in the table 1.

87 This is due to the technology used, the properties of the raw material (Moretti *et al.*, 2004) and the specific
88 composition of the microbiota (Leroy *et al.*, 2006). Nonetheless, Sunesen and Stahnke (2003) reported that
89 sausages produced with commercial molds show more consistent flavor, taste, drying rate, and a more
90 uniform appearance with respect to artisan ally fermented sausages.

91 The microbial ecology of fermented sausages has become of increasing interest over the last few decades
92 given that different genera, species, and even strains, have been shown to significantly affect the sensory
93 traits of fermented sausages (Rantsiou and Cocolin, 2006). Production of artisanal sausages largely depends
94 on the skill and experience of the meat manufacturer and may be considered an art rather than a process fully
95 based on scientific and technological understanding. Meat fermentation is, in fact, a complex biological
96 phenomenon accelerated by the desirable action of certain microbes in the presence of a variety of

synergistically acting or competing species. A great variability in the quality of the products is due to traditional practices and variation in the microorganisms involved in the process. De Vuyst (2000) underlines that it is of primordial importance to investigate and analyze the influence of the environment on the performance of a starter culture before using it in a selected product. In order to protect the traditional aspects of these products and to select autochthonous starter cultures to be used, it is essential to understand the microbial dynamics during fermentation (Rantsiou and Cocolin, 2006). Therefore, a current quest is to develop indigenous starters that guarantee hygienic quality and improve the sensorial aspects of the product (Talon *et al.*, 2007). It should, however, be considered that the law allows only the use of qualified presumption of safety (QPS) in the EU, or generally recognized as safe (GRAS) in the US, microorganisms in food production. In Italy, only *Lactobacillus*, *Pediococcus*, *Micrococcus*, *Debaryomyces*, and *Staphylococcus xylosus*, *Staphylococcus simulans* and *Staphylococcus carnosus* are authorized as starters cultures for sausage production (Gazzetta Ufficiale, 1995).

2.1 Starter culture selection parameters

So far the selection of a starter culture has been based on the screening of a great number of isolates in small-scale food fermentations. A satisfactory performance of the selected starter culture in the process, and an acceptable organoleptic evaluation of the food product are the fundamental characteristics to be found. The behavior of the starter culture in relation to the environmental factors and ripening conditions encountered during a specific production needs to be carefully investigated and standardized in the selection process. It is necessary to understand the properties required and the specific technology and recipe for which a strain will be used in order to develop the ideal starter culture (Hansen, 2002). According to Holzapfel (1997), in order to improve product quality, the introduction of starter cultures in traditional small-scale fermentations should incorporate considerations as (i) rapid metabolic activities (acid production); (ii) improved and predictable fermentation processes; (iii) desirable sensory attributes; (iv) improved safety and reduced hygienic and toxicological risks. Another important factor is the interaction in mixed cultures of selected starter strains, with consideration for the behavior of these strains under defined conditions, and within the food matrix. Other aspects, which should be considered, include: (i) competitive behavior, viability and survival; (ii) antagonism against pathogens and spoilage microbes; (iv) rate of acid production; (v) organoleptic changes; (vi) primary metabolites of fermentation; (vii) degradation of

125 antinutritive factors; (vii) detoxification; (viii) probiotic features (Holzapfel, 1997). Modern approaches for
126 selection of the best strain(s) for a process integrate also technical safety and health-promoting features
127 (Holzapfel, 2002).

128 It is essential to know the autochthonous microbiota of fermented food that is to be analyzed because
129 commercial starter cultures usually originate either from substrates or from the processes in which they are
130 applied. Factors that can contribute to the selection of microbial populations typical of a fermentation
131 process are environmental conditions, back-slopping, adaptation and the repeated use of specific tools
132 (Holzapfel, 1997).

133 **3. Ecological aspects of sausage fermentation**

134 Most European fermented sausages still follow the traditional procedures in which fermentation and ripening
135 depend on the activities of heterogeneous microbial communities (Gardini *et al.*, 2001, Cocolin *et al.*, 2006).
136 Two wide groups of bacteria largely predominate, lactic acid bacteria (LAB) and the group known as either
137 coagulase-negative cocci (CNC) or (gram-positive)-catalase-positive cocci (GCC+/CPC), which includes
138 both micrococci and coagulase-negative staphylococci (CNS) (Aquilanti *et al.*, 2016). Yeasts and
139 filamentous fungi also play a relevant role, through the formation of a superficial film which exerts a
140 protective action against both excessive dehydration and the oxidation of the lipid fraction due to oxygen and
141 light (Gardini *et al.*, 2001, Cocolin *et al.*, 2006). The acidification process that is the result of the
142 fermentation of the sugars into lactic acid by LAB, plays a fundamental role to prevent spoilage and
143 pathogen outgrowth. Coagulase-negative cocci are involved in the proteolytic and lipolytic processes thus
144 playing a central role in the formation of the final organoleptic characteristics (Hammes and Hertel, 1998),
145 and contributing to nitrate reduction and color formation, as well as to prevention of rancidity. In addition,
146 the characteristic flavors and surface aspect are due to yeasts and molds (Cocolin *et al.*, 2006). LAB are more
147 numerous than CNC during fermentation and ripening, remaining more stable in the ripened products.
148 Within LAB, facultatively heterofermentative lactobacilli generally prevail and, among them, the two
149 psychrotrophic species *Lactobacillus sakei* and *Lactobacillus curvatus* are dominant. Within CNC,
150 *Staphylococcus xylosus* neatly dominates (Aquilanti *et al.*, 2016).

151 The selection of specific populations adapted to a specific environment depends on ingredient composition,
152 fermentation and maturation factors (Rantsiou *et al.*, 2005a). The production environment may have an
153 impact on the microbial ecology of the product. For example, Greppi *et al.* (2015) showed that strains
154 isolated from environmental samples were also detected either in the raw materials or in the product. This
155 finding highlights that microorganisms may enter in the production plant with the raw materials.
156 Furthermore, it underlines that the production environment is a source of continuous “inoculation”, during
157 fermentation and ripening, with strains that may have important technological characteristics and influence
158 the characteristics of the final product.

159

160 **3.1 Lactic acid bacteria**

161 The term lactic acid bacteria is used to define a large and diverse group of microorganisms. LAB may be
162 described as a group of Gram-positive, non-spore-forming cocci and rods, microaerophilic or facultative
163 anaerobes, that they produce lactic acid as the major end-product during fermentation of carbohydrates
164 (Halász, 2009).

165 *L. sakei*, *L. curvatus* and *L. plantarum* are the principal species of LAB usually found in meat and meat
166 products, including fermented sausages made with different production process (Hugas *et al.*, 1993;
167 Kittisakulnam *et al.*, 2017; Pisacane *et al.*, 2015). *L. sakei* is often isolated with the higher frequency with
168 respect to *L. curvatus*, although sometimes the opposite occurs, or they are found at similar levels; *L.*
169 *plantarum* is generally isolated with less frequency, but even in this case exceptions are found, probably due
170 to particular processing conditions. The same is also true for the members of other LAB genera, such as
171 *Weissella*, *Leuconostoc*, *Lactococcus*, and *Pediococcus* since they are in general found as minority species
172 (Aquilanti *et al.*, 2016).

173 The genus *Pediococcus*, at the current time, consists of 13 species (Haakensen *et al.*, 2009). *Pediococcus*
174 *pentosaceus* and *Pediococcus acidilactici* are the main species used in i) pediocin production, ii)
175 fermentation processes as a starter (co-culture) to avoid contamination, and iii) probiotic supplements for
176 animals and humans. *P. pentosaceus* cells are spherical arranged in tetrads. They are homofermentative, i.e.
177 produce lactic acid as sole product of hexose fermentation (Porto *et al.*, 2017).

178 Lactobacilli and pediococci are the dominant microorganisms in sausages with a short ripening time from
 179 early stages to the end of the process: this type of product has an acid flavor with little aroma. In contrast,
 180 sausages with longer maturation times contain higher numbers of lactobacilli (Demeyer *et al.*, 1986). Several
 181 studies have been conducted, employing molecular methods for species and strain identification, in order to
 182 understand the diversity and dynamics of LAB populations during fermented sausages production with long
 183 maturation times. Rantsiou *et al.* (2005a) studied the dynamics of LAB populations involved in the process
 184 of traditional fermentations performed in three countries: Hungary, Italy and Greece. In this study, 14
 185 different species of LAB were detected. The only common species for Greek, Hungarian and Italian sausages
 186 were *L. plantarum*, *L. curvatus* and *L. sakei*. Furthermore, molecular characterization of the isolates revealed
 187 a country-specific geographic distribution of LAB populations... In Pisacane *et al.* (2015) the production of
 188 Salame Mantovano using two different types of natural casing, deriving from two different portions of the
 189 pig's intestine, was studied. Community dynamics suggested that the predominant LAB species in the two
 190 types of sausages were the same.

191 Aquilanti *et al.* (2016) summarized the studies concerning the structure of LAB in Mediterranean (Northern,
 192 Central and Southern Italy, Greece, France, Spain and Portugal) traditional fermented sausages. Within the
 193 LAB population, facultatively heterofermentative lactobacilli generally prevailed and, among them, *L. sakei*
 194 and *L. curvatus* were found to be dominant in most studies. Authors underlined that there was low species
 195 variability between products of the different countries (Aquilanti *et al.* 2016).

196

197 **3.2 Coagulase–Negative Cocci**

198 *Staphylococcus* and *Kocuria* are the most representative genera of the Gram-positive Catalase-positive Cocci
 199 (GCC+) group (Morot-Bizot *et al.*, 2006). The characteristic microbiota in sausages is composed of *S.*
 200 *xylosus*, *S. saprophyticus* and *S. equorum*, but many other species have been identified such as *S. succinus*, *S.*
 201 *warneri*, *S. vitulinus*, *S. pasteurii*, *S. epidermidis*, *S. lentus* and *S. haemolyticus* (Cocolin *et al.*, 2001; Talon *et*
 202 *al.*, 2011; Mainar *et al.*, 2017). *Kocuria* species are ubiquitous and are highly adapted to their ecological
 203 niches (Kim *et al.*, 2011). In fermented sausages *Kocuria varians* and *Kocuria kristinae* were mainly found
 204 (Fischer and Schleifer, 1980); moreover, *K. varians* is often found in biofilms (Raghupathi *et al.*, 2016).

205 Although the environmental, production plant associated microbiota, can contribute to the spoilage of the
 206 meat products, ecology of *Staphylococcus* occurring in the environment of spontaneously fermented
 207 sausages has not been thoroughly studied. In fact, they showed high capacity to colonize the surfaces, the
 208 equipment and the meat products (Morot-Bizot *et al.*, 2006).

209 Iacumin *et al.* (2006) studied the ecology and dynamics of staphylococci in three different local meat
 210 producers in the North East of Italy. In all three fermentations the same species of CNC (*S. epidermidis*, *S.*
 211 *equorum*, *S. warneri*, *S. saprophyticus*, *S. xylosus*, *S. pasteurii*) took part, but in variable quantity and
 212 proportions. The study evidenced that the slaughterhouse can partly influence the microbial composition of
 213 meat and a correlation between the isolated *S. xylosus* strains and the specific plant of production exists. This
 214 confirms the hypothesis that selection of the microbiota takes place in a production plant, depending on
 215 temperature, humidity and ingredients and influences the final sensory aspect of the product. In a
 216 comparative evaluation of the CNC communities from sausages produced in Italy, France, Greece Spain and
 217 Portugal, The dominance of *S. xylosus* clearly emerged, with the exception of the sausage productions in
 218 Greece and France. In fact, the CNC diversity, between different countries, was generally higher than that
 219 recorded for LAB (Aquilanti *et al.*, 2016).

220 Finally, Quijada *et al.* (2018) analyzed five Chorizo de Leon factories in the North-West region of Castilla y
 221 Leon (Spain). The factories didn't use microbial starters and adopted similar traditional manufacturing
 222 procedures. Among the five manufactures differences in microbiota composition were observed. For the
 223 CNC, *Staphylococcus* was found in all the samples, but its distribution depended on the manufacturer. In this
 224 work again the importance of country-specific microbiota in the development of traditionally manufactured
 225 products was more evident for CNC compared to *Lactobacillus* species.

226

227 **3.3 Yeasts**

228 Spontaneous fermentations are usually characterised by the presence of yeasts, but studies on the yeast
 229 biodiversity in sausages are limited. *Debaryomyces hansenii* is the yeast species most commonly isolated
 230 according to several researches but other yeast genera have also been found, such as *Candida* spp. (Gardini
 231 *et al.*, 2001). An increase in pH and a decrease in lactic acid content in the sausages can be caused by these
 232 yeasts that contribute to the characteristics of the final product (Gardini *et al.*, 2001). Both *D. hansenii* and

233 *Candida utilis* initially proliferate in sausages and then slowly decline (Olesen and Stahnke, 2000). *C. utilis*
 234 shows a considerable potential production of several volatile compounds, such as alcohols and esters which
 235 probably derive from the amino acids isoleucine, leucine, valine and phenylalanine (Olesen and Stahnke,
 236 200). On the contrary, The primary and secondary metabolism, where lipases and proteinases are key
 237 enzymes, are the principal processes of these organisms and can produce the typical aroma of the products
 238 (Cocolin *et al.*, 2006). A yeast can be added as aroma enhancer and can also stabilise the red colour of
 239 fermented sausages (Olesen and Stahnke, 2000).

240 *Debaryomyces* spp. are extremophilic, perfect, haploid yeasts that asexually reproduce by multilateral
 241 budding, the pseudomycelium is absent, primitive or occasionally well developed. Heterogamous
 242 conjugation is the way for the sexual reproduction. In particular, *D. hansenii* is an osmo-, alo- and
 243 xerotolerant yeast (Breuer and Harms, 2006). Flores *et al.* (2004) underlined that in fermented sausages,
 244 *Debaryomyces* spp. can have important effects on the generation of volatile compounds during the ripening.
 245 The development of the typical aroma of the sausage was possible through the inhibition of the generation of
 246 lipid oxidation products and promoting the generation of ethyl esters. When *D. hansenii* was used as a starter
 247 it showed a positive effect on the development of flavour characteristics and stabilisation of the reddening
 248 reaction (Gardini *et al.*, 2001).

249 Cocolin *et al.* (2006a) employed a multiphasic approach during the fermentation of a traditional sausage
 250 produced in Northern Italy. To profile the dynamics of yeast communities present during the maturation
 251 culture-dependent and independent methods were used. Through the molecular identification by PCR-DGGE
 252 and sequencing of partial 26S rRNA encoding gene of 180 isolates, *D. hansenii* resulted to be the dominant
 253 species throughout the fermentation process. With molecular characterization, *D. hansenii* isolates displayed
 254 a change in their population density during the maturation process of the sausages.

255 Although the origin of the meat and the factory environment have been reported as factors that can cause
 256 variations on yeast populations in fermented meat products, most of the studies point towards *D. hansenii* as
 257 the most frequently and abundantly isolated yeast species (Flores *et al.*, 2015; Mendonça *et al.*, 2013).
 258

259 3.4 Filamentous fungi

260 The surface of dry-cured meat is colonized by molds able to grow on different environments and
261 substrates (Magistà *et al.*, 2017). Xerotolerant and xerophilic fungi grow preferably in an environment with
262 low water activity (a_w) and high salt concentrations as dry-cured meats. In this kind of products, fungi have
263 also an important role in the production process because they can lead to the development of specific flavors
264 and aromas, due to their lipolytic and proteolytic activities (Sonjak *et al.* 2011).

265 The genus *Penicillium* represents the major mold population of the surface mycobiota on dry-cured meat
266 products (Sonjak *et al.*, 2011). *Penicillium* is one of the most common fungi that can grow in a diverse range
267 of habitats, from soil to vegetation to air, indoor environments and various food products (Visagie *et al.*,
268 2014). Important taxonomic characters of *Penicillium* are the presence of conidiophore and cleistothecium
269 (when produced). Conidiophore branching patterns have been traditionally used in the classification of
270 *Penicillium* (Visagie *et al.*, 2014). Species of *Penicillium* have been found in fermented meat sausages to be
271 responsible for the surface colonization, most importantly *P. nalgiovense* and to a lesser extent, *P.*
272 *chrysogenum* (López-Díaz *et al.*, 2001). This layer of mold is important to the sausage since it has an
273 antioxidative effect, protecting from development of the rancidity and keeping the colour; it gives the
274 sausage its typical appearance because it allows the development of a positive microclimate at the surface for
275 preventing, for example, sticky or slimy characteristic of the surface (Visagie *et al.*, 2014).

276 4. Bioprotection

277 In fermented meat, the accumulation of particular metabolites as lactic acid, acetic acid, formic acid, ethanol,
278 ammonium, fatty acids, hydrogen peroxide, acetaldehyde and bacteriocins can inhibit the growth of
279 pathogenic and spoilage bacteria (Hugas and Monfort, 1997). The production of antimicrobial bacteriocins
280 that leads to a better preservation of the product is a characteristic of particular starter cultures (Cleveland *et*
281 *al.*, 2001). Strains of all genera of LAB have been identified as bacteriocin producers. They are important in
282 meat microbiota composition and act against bacteria closely related to the producer organisms (Lücke,
283 2000). However, many lactic acid bacteria (LAB) strains produce bacteriocins that are active towards
284 pathogens or food spoilers *in vitro*, but not *in situ*, in a meat matrix (De Vuyst, 2000).

285 Different bacteriocins produced by LAB strains could be applied in food products but, at the moment, only
286 nisin and pediocin PA-1/AcH are approved for use in food preservation (Cleveland *et al.*, 2001; Barbosa *et*
287 *al.*, 2017; Kęska *et al.*, 2017). The application of bacteriocins in meats and meat products is allowed in three
288 different modalities: (i) direct inoculation of bacteriocinogenic LAB strains as starter or protective cultures,
289 (ii) direct application of bacteriocins from cell free supernatant (CFS) as food additive and (iii) incorporation
290 of totally or partially purified bacteriocins into the packaging (Woraprayote *et al.*, 2016). Bacteriocins
291 improve a strain's competitiveness for the nutrients during fermentation, but without reducing the growth of
292 the starter organisms towards the fortuitous microbiota. (Hugas and Monfort, 1997).

293 Many strains of *Lactococcus lactis* are able to produce Nisin A that has a wide antimicrobial spectrum
294 against Gram-positive bacteria, including staphylococci, streptococci, *Listeria* spp., bacilli, and enterococci
295 (Woraprayote *et al.*, 2016). Several nisin-producing *Lact. lactis* strains isolated from fermented sausages
296 showed the potential use of lactococci in this kind of products (Castellano *et al.*, 2008).

297 The effect of commercial pure nisin from *Lact. lactis* subsp. *lactis* (Sigma-Aldrich) against *List.*
298 *monocytogenes* was evaluated in Turkish fermented sausages (sucuks). All products treated with nisin,
299 showed a reduction of *List. monocytogenes* population compared to the control (Hampikyan and Ugur,
300 2007).

301 *L. sakei* and *L. curvatus* strains are able to produce sakacins, bacteriocins that show high inhibitory activities
302 towards *List. monocytogenes*. Sakacin Q, produced by *L. curvatus* ACU-1, was used by Rivas *et al.* (2014)
303 for growth control of *List. innocua*. Cooked meat was artificially inoculated on surface during chilled storage
304 and four different forms of bacteriocin applications were tested: protective culture, cell-free supernatant
305 (CFS), mixture of both protective culture and CFS and freeze-dried reconstituted CFS. In this study, the most
306 effective one to control the pathogen growth was freeze-dried reconstituted CFS. In Barbosa *et al.* (2015)
307 meat-borne strains of *L. curvatus* have been described as the main source of antimicrobial compounds:
308 curvaticins. One of the isolates exhibiting inhibitory activity against *List. monocytogenes* ATCC 7644 was
309 identified as *L. curvatus* 54M16 and studied in detail for its antimicrobial substances (Casaburi *et al.*, 2016).

310 In the last decades, great number of *L. plantarum* strains that produce bacteriocins were isolated from
311 different matrices including meat (Kanatani and Oshimura, 1994). In literature, numerous small, heat-stable
312 plantaricins have been described but not completely well characterized (Todorov, 2009). Schillinger and

313 Lücke (1989) isolated, from different meat products, various bacteriocinogenic lactobacilli including *L.*
314 *plantarum*. Enan *et al.* (1996) showed, for example, that plantaricin UG1 is produced by *L. plantarum* UG1
315 isolated from dry sausage and that this compound had no effect on Gram-negative bacteria, but a variety of
316 Gram-positive bacteria were sensitive. An important control of *List. monocytogenes* growth has been
317 obtained by application of plantaricins or *L. plantarum* bacteriocin producing strains (Todorov, 2009).
318 Pediocins are biomolecules that can be synthesized by some LAB and present a broad spectrum of
319 antimicrobial activity against Gram-positive bacteria (Papagianni and Anastasiadou, 2009), among which
320 *List. monocytogenes* (Porto *et al.*, 2017). Kingcha *et al.* (2012) observed a significant decrease of *List.*
321 *monocytogenes* ATCC 19115 growth in Nhan, a Thai traditional fermented pork sausage, when it was
322 inoculated with *P. pentosaceus* BCC 3772 cells. The antimicrobial activity was attributed to the production,
323 by *P. pentosaceus* BCC3772, of pediocin that shows 100% amino acid identity with the commercial pediocin
324 PA-1 isoform. A correlation was observed between anti-listerial activity and *P. pentosaceus* BCC3772
325 inoculum. In addition, the authors suggested that this strain is a suitable candidate for *Listeria* control in
326 fermented pork sausage, given that no significant changes of Nahn's organoleptic properties were observed.

327 **5. Direct analysis of sausages and omics approaches**

328 Traditional microbiological methods, namely plate counts, isolation, and biochemical identification, have
329 been often used for ecological studies of spontaneously fermented sausages. With this approach only easily
330 culturable microorganisms can be detected, while the information about microorganisms that need elective
331 enrichments or that are in a sublethal or injured state (physiological condition) are lost (Rantsiou *et al.*,
332 2005b). In fact, traditional microbiological techniques do not give a correct view of microbial diversity (Justé
333 *et al.*, 2008; Stefanis *et al.*, 2016; Silvetti *et al.*, 2017). New approaches, that take advantage of molecular
334 methods and are applied directly in a sample (direct or culture independent approaches), have been
335 introduced in the field of food microbiology and food fermentation, allowing scientists to overcome the
336 limitations of classical methods (Rantsiou and Cocolin, 2006; De Filippis *et al.*, 2018). These studies may
337 also be able to identify sentinel microbes (essentially indicator microbes linked to various pathogens), which
338 could be incorporated into food safety plans (Cocolin *et al.*, 2018). The first direct or culture-independent
339 approaches were based on fingerprinting techniques such as DGGE and have been extensively applied in

340 fermented sausages ecology studies (Rantsiou and Cocolin, 2006; Aquilanti *et al.*, 2016). In recent years
341 however, food microbiota studies are based on sequencing (Cocolin and Ercolini, 2015). High-throughput
342 sequencing (HTS) techniques, undoubtedly represent a step change in the way microbiologists address
343 ecology and diversity in foods. Unlike traditional Sanger approach sequencing that could be performed on a
344 single DNA molecule, in HTS mixed nucleic acid molecules from a complex ecosystem can be sequenced
345 and therefore can lead to detailed profile of the microbial populations (identified as Operational Taxonomic
346 Units, OTU) present (Cocolin *et al.* 2018; Zhang *et al.*, 2017). In the mid 2000s HTS technologies became
347 ubiquitous in microbial ecology studies; in fact, these technologies have been used to monitor the dynamics
348 of microbial communities during fermentation of different types of foodstuffs and beverages (De Filippis *et*
349 *al.*, 2017; Fontana *et al.*, 2016).

350 Metataxonomics or amplicon sequencing, is a powerful tool that allows taxonomic characterization of
351 microbial communities, which would have been difficult if not otherwise impossible to determine using
352 traditional microbiological techniques. For bacteria, the common amplification target is various regions of
353 the 16S rRNA encoding gene, while for yeasts the 26S rRNA encoding gene has been used. Large sequence
354 databases exist for these two genes (Cocolin *et al.*, 2013). On the other hand, metagenome sequencing, or
355 shotgun metagenome sequencing, is a technique where an entire mixed microbial community DNA is
356 fragmented, prepared into a sequencing library and sequenced. Metagenomics offers the opportunity to look
357 beyond the presence/absence of taxonomically defined entities (i.e. specific organisms) and instead to
358 understand the relationships between microorganisms and their activities and functionalities in a particular
359 niche (Cocolin *et al.*, 2017). To understand and characterize the composition and function of the microbiota
360 in a food ecosystem the evolution of the massive sequencing technologies such as shotgun DNA-seq or
361 RNA-seq can help (Ferrocino *et al.*, 2018). In the field of food microbiota structure and function, the
362 identification of enzymes, pathways and mechanisms and how these operate under specific conditions may
363 perhaps be of superior value than determining the taxonomic composition of specific samples alone
364 (Santiago-Rodriguez *et al.*, 2016).

365 DNA is a chemically stable molecule, which can be found a long time after the death of a cell, RNA is more
366 sensitive to degradation, especially in environments, like foods, where enzymes, such as hydrolases, are
367 present. While DNA can give a good overview of the microorganisms that are or were present in a given

ecosystem, it cannot provide any information on what microbes are doing regarding metabolic and spoilage activities and virulence factors expression. For this reason, if the goal of the investigation is to get an insight on how the microorganism is behaving, the RNA is a better option (Cocolin *et al.*, 2013). Analysis of RNA (transcriptomics), proteins (proteomics) and metabolites (metabolomics), preferably in an integrated framework, is fundamental for a full description of microbial community inasmuch metagenomic sequencing has an important limitation: it cannot directly measure the functional activity of a community under a given set of conditions (Franzosa *et al.*, 2015). Studies have used RT-PCR-DGGE coupled with 16S rRNA-based sequencing to evaluate the diversity of metabolically active microbiota during the spontaneous fermentation of sausages (Połka *et al.*, 2015; Rebecchi *et al.*, 2015). The organoleptic characteristics of the final products are influenced by volatile organic compounds (VOCs) that are produced through the breakdown of carbohydrates, proteins and lipids. Ferrocino *et al.* (2018) focused on studying the microbiota development and functions in an Italian fermented sausage through a shotgun DNA metagenomic approach. For the first time an integrated analysis related to volatilome profile, microbiota, gene content and consumers acceptability was presented. The study displayed the evolution of those pathways over time and condition. The most prominent differences during spontaneous and inoculated fermentation, according to the analysis, involved key genes in particular pathways: pyruvate metabolism and glycolysis. For the faster metabolic activity, confirmed by the meta-metabolomics data, in this study the sensory test showed that the presence of starter cultures had a negative impact on the properties of the product. These methods allowed to detect shifts in microbiota composition through the recognition of changes in the microbial gene content and abundance. This is important for the study of complex and dynamic microbiota of food products and for food safety both to spoilage and fraud level, especially when starter cultures are used.

391

392 **6. Conclusions**

393 The use of starter cultures in food fermentations, including sausages, has allowed for a greater level of safety
394 and quality. Undoubtedly, the addition of starter strains in the meat batter for the production of sausages
395 makes the fermentation process less prone to modifications, which are responsible for products with deviated

396 organoleptic properties and potentially hazardous due to the presence of foodborne pathogens not inhibited
397 during the process. However, due to the traditional and artisanal aspects that fermented sausages often
398 possess, spontaneous fermentations are still used and represent a common practice especially in the
399 Mediterranean countries. Isolation and selection of new strains of well known LAB and CNC species from
400 those traditional products represents a possible alternative to the use of commercial starter cultures, which
401 nowadays are criticized since they may lead to flattened organoleptic characteristics.

402 The application of modern molecular methods, such as metataxonomics and metagenomics, in fermented
403 sausages will permit, in the near future, the understanding in detail of the microbial ecology and functions
404 and at the same time allow for a better comprehension of the interactions of the starter cultures with the meat
405 microbiota during sausage production. Only after a thorough knowledge of the mechanisms behind the
406 fermentation process in meat it will be possible its control and modulation to obtain products with desired
407 and expected organoleptic profiles.

408

References

- Aquilanti L., Garofalo C., Osimani A., Clementi F. (2016). Ecology of lactic acid bacteria and coagulase negative cocci in fermented dry sausages manufactured in Italy and other Mediterranean countries: an overview. *International Food Research Journal* 23(2), 429-445.
- Aquilanti L., Santarelli S., Silvestri G., Osimani A., Petruzzelli A., Clementi F. (2007). The microbial ecology of a typical Italian salami during its natural fermentation. *International Journal of Food Microbiology*, 120, 136-145.
- Barbosa A.A.T., Mantovani H.C., Jain S. (2017). Bacteriocins from lactic acid bacteria and their potential in the preservation of fruit products. *Critical reviews in biotechnology*, 37(7), 852-864.
- Barbosa M.S., Todorov S.D., Ivanova I., Chobert J.-M., Haertlé T., Franco B.D. G.M. (2015). Improving safety of salami by application of bacteriocins produced by an autochthonous *Lactobacillus curvatus* isolate. *Food Microbiol.*, 46, 254–262.
- Breuer U., Harms H. (2006). *Debaryomyces hansenii* - an extremophilic yeast with biotechnological potential. *Yeast*, 23(6), 415-437.
- Casaburi A., Aristoy M-C., Cavella S., Di Monaco R., Ercolini D., Toldra´ F., Villani F. (2007). Biochemical and sensory characteristics of traditional fermented sausages of Vallo di Diano (Southern Italy) as affected by the use of starter cultures. *Meat Science*, 76, 295–307.
- Casaburi A., Di Martino V., Ferranti P., Picariello L., Villani F. (2016). Technological properties and bacteriocins production by *Lactobacillus curvatus* 54M16 and its use as starter culture for fermented sausage manufacture. *Food Control*, 59, 31-45.
- Castellano P., Belfiore C., Fadda S., Vignolo G. (2008). A review of bacteriocinogenic lactic acid bacteria used as bioprotective cultures in fresh meat produced in Argentina. *Meat Science*, 79(3), 483-499.
- Cleveland J., Montville T.J., Nes I.F., Chikindas M.L. (2001). Bacteriocins: safe, natural antimicrobials for food preservation. *Int. J. Food. Microbiol.* 71, 1-20.
- Cocolin L., Alessandria V., Dolci P., Gorra R., Rantsiou K. (2013). Culture independent methods to assess the diversity and dynamics of microbiota during food fermentation. *Int. J. Food Microbiol.* 167, 29-43.

- Cocolin L., Dolci P., Rantsiou K. (2011). Biodiversity and dynamics of meat fermentations: The contribution of molecular methods for a better comprehension of a complex ecosystem. *Meat Science*, 89, 296-302.
- Cocolin L., Manzano M., Cantoni C., Comi G. (2001). Denaturing Gradient Gel Electrophoresis Analysis of the 16S rRNA Gene V1 Region To Monitor Dynamic Changes in the Bacterial Population during Fermentation of Italian Sausages. *Applied and Environmental Microbiology*, 5113-512.
- Cocolin L., Mataragas M., Bourdichon F., Doulgeraki A., Pilet M.-F., Jagadeesan B., Rantsiou K., Phister T. (2018). Next generation Microbiological Risk Assessment: Meta-Omics: the next need for integration. *International journal of food microbiology*.
- Cocolin L., Urso R., Rantsiou K., Cantoni C. and Comi G. (2006). Dynamics and characterization of yeasts during natural fermentation of Italian sausages. *FEMS Yeast Research* 6(5): 692-701.
- De Filippis F., Parente E., Ercolini D. (2017). Metagenomics insights into food fermentations. *Microbial biotechnology*, 10(1), 91-102.
- De Filippis F., Parente E., Ercolini D. (2018). Recent Past, Present, and Future of the Food Microbiome. *Annual review of food science and technology*, 9(1).
- De Vuyst L. (2000). Technology aspects related to the application of functional starter cultures. *Food Technology and Biotechnology*, 38(2), 105-112.
- Demeyer D.I., Verplaetse A., Gistelink M. (1986). Fermentation of meat: an integrated process. *Belgian Journal of Food Chemistry and Biotechnology*, 41, 131–140.
- El Sheikha A., Montet D. (2016). Fermented foods - artisan household technology to biotechnology era. In : Montet Didier (ed.), Ray Ramesh Chandra (ed.). *Fermented foods. Part I: biochemistry and biotechnology*. Boca Raton : CRC Press, p. 1-15. (Food Biology Series).
- Enan G.; Essaway A.A.; Uyttendaele M.; Debevere J. (1996). Antibacterial activity of *Lactobacillus plantarum* UG1 isolated from dry sausages: characterization, production, and bactericidal action of plantaricin UG1. *Int. J. Food Microbiol.*, 30: 189-215.
- Ferrocino I., Bellio A., Giordano M., Macori G., Romano A., Rantsiou K., Decastelli L., Cocolin, L. (2018). Shotgun metagenomics and volatilome profile of the microbiota of fermented sausages. *Applied and environmental microbiology*, 84(3), e02120-17.

- .
- Fischer U., Schleifer K. H. (1980). Vorkommen von staphylokokken und mikrokokken in Rohwurst. Fleischwirtschaft, 60, 1046-1049.
- Flores M., Dura M.A., Marco A., Toldra F. (2004). Effect of *Debaryomyces* spp. on aroma formation and sensory quality of dry-fermented sausages. Meat Sci 68: 439–446.
- Flores M., Corral S., Cano-García L., Salvador A., Belloch C. (2015). Yeast strains as potential aroma enhancers in dry fermented sausages. International journal of food microbiology, 212, 16-24.
- Fontana C., Bassi D., López C., Pisacane V., Otero M.C., Puglisi E., Rebecchi A., Cocconcelli P.S., Vignolo, G. (2016). Microbial ecology involved in the ripening of naturally fermented llama meat sausages. A focus on lactobacilli diversity. International journal of food microbiology, 236, 17-25.
- Franzosa E.A., Hsu T., Sirota-Madi A., Shafquat A., Abu-Ali G., Morgan X.C., Huttenhower C. (2015). Sequencing and beyond: integrating molecular 'omics' for microbial community profiling. Nature Reviews Microbiology, 13(6), 360-372.
- Gardini F., Suzzi G., Lombardi A., Galgano F., Crudele M. A., Andrighetto C., Schirone M., Tofalo R. (2001). A survey of yeasts in traditional sausages of southern Italy. FEMS Yeast Research 1(2), 161-167.
- Gazzetta Ufficiale, Serie Generale, N. 89 del 15/4/1995
- Greppi A., Ferrocino I., La Stora A., Rantsiou K., Ercolini D., Cocolin L. (2015). Monitoring of the microbiota of fermented sausages by culture independent rRNA-based approaches. International journal of food microbiology, 212, 67-75.
- Haakensen M., Dobson C.M., Hill J.E., Ziola B. (2009). Reclassification of *Pediococcus dextrinicus* (Coster and White 1964) Back 1978 (Approved Lists 1980) as *Lactobacillus dextrinicus* comb. nov., and emended description of the genus *Lactobacillus*. International journal of systematic and evolutionary microbiology, 59(3), 615-621.
- Halász A. (2009). Lactic acid bacteria. Food Quality and Standards, 3, 70-82.
- Hammes W.P., Hertel C. (1998). New developments in meat starter cultures. Meat Sci 49: S125–S138.
- Hampikyan H., Ugur M. (2007). The effect of nisin on *L. monocytogenes* in Turkish fermented sausages (sucuks). Meat science, 76(2), 327-332.

- Hansen E. B. (2002). Commercial bacterial starter cultures for fermented foods of the future. *International Journal of Food Microbiology*, 78, 119–131.
- Holzapfel W.H. (1997). Use of starter cultures in fermentation on a household scale. *Food control*, 8 (5-6), 241-258.
- Holzapfel W.H. (2002). Appropriate starter culture technologies for small-scale fermentation in developing countries. *International journal of food microbiology*, 75(3), 197-212.
- Hugas M., Garriga M., Aymerich T., Monfort J.M. (1993). Biochemical characterization of lactobacilli from dry fermented sausages. *International Journal of Food Microbiology*, 18, 107-113.
- Hugas M., Monfort J.M. (1997). Bacterial starter cultures for meat fermentation. *Food Chemistry*, 59, No. 4, 547-554.
- Iacumin L., Comi G., Cantoni C., Cocolin L. (2006). Ecology and dynamics of coagulase-negative cocci isolated from naturally fermented Italian sausages. *Systematic and applied microbiology*, 29(6), 480-486.
- Justé A., Thomma B.P.H. J., Lievens B. (2008). Recent advances in molecular techniques to study microbial communities in food-associated matrices and processes. *Food Microbiology*, 25(6), 745-761.
- Kanatani K., Oshimura M. (1994). Plasmid-associated bacteriocin production by a *Lactobacillus plantarum* strain. *Bioscience, biotechnology, and biochemistry*, 58(11), 2084-2086.
- Kęska P., Stadnik J., Zielińska D., Kołożyn-Krajewska D. (2017). Potential of bacteriocins from lab to improve microbial quality of dry-cured and fermented meat products. *Acta Scientiarum Polonorum Technologia Alimentaria*, 16(2), 119-126.
- Kim W.J., Kim Y.O., Kim D.S., Choi S.H., Kim D.W., Lee J.S., Kong H.J., Nam B.H., Kim B.S., Lee S.J., Park H.S., Chae S.H. (2011). Draft genome sequence of *Kocuria rhizophila* 7-4. *J Bacteriol* 193:4286–4287.
- Kingcha Y., Tosukhowong A., Zendo T., Roytrakuls S., Luxananil P., Chareonpornsook K., Valyasevi R., Jang S., Lee J., Jung U., Choi H.S., Suh H.J. (2012). Anti-*Listeria* activity of *Pediococcus pentosaceus* BCC 3772 and application as starter culture for Nham, a traditional fermented pork sausage. *Food Control* 25, 190–196.

Kittisakulnam S., Saetae D., Suntornsuk W. (2017). Antioxidant and antibacterial activities of spices traditionally used in fermented meat products. *Journal of Food Processing and Preservation*, 41(4).

Leroy F., De Vuyst L. (2004). Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology*, 15(2), 67-78.

Leroy F., Verluyten J., De Vuyst L. (2006). Functional meat starter cultures for improved sausage fermentation. *International Journal of Food Microbiology* 106, 270-285.

López-Díaz T.M., Santos J.A., García-López M.L., Otero A. (2001). Surface mycoflora of a Spanish fermented meat sausage and toxigenicity of *Penicillium* isolates. *International Journal of Food Microbiology*, 68(1), 69-74.

Lücke F.K. (1986). Microbiological processes in the manufacture of dry sausage and raw ham. *Fleischwirtschaft* 66, 1505-1509.

Lücke F.K. (2000). Utilization of microbes to process and preserve meat. *Meat Science*, 56(2), 105-115.

Magistà D., Susca A., Ferrara M., Logrieco A.F., Perrone G. (2017). *Penicillium* species: crossroad between quality and safety of cured meat production. *Current Opinion in Food Science*, 17, 36-40.

Mainar M.S., Stavropoulou D.A., Leroy F. (2017). Exploring the metabolic heterogeneity of coagulase-negative staphylococci to improve the quality and safety of fermented meats: a review. *International journal of food microbiology*, 247, 24-37.

Mendonça R.C., Gouvêa D.M., Hungaro H.M., Sodré A.D.F., Querol-Simon A. (2013). Dynamics of the yeast flora in artisanal country style and industrial dry cured sausage (yeast in fermented sausage). *Food control*, 29(1), 143-148.

Moretti V.A., Madoni, G., Diaferi, C., Mentast, T., Palear, M.A., Panser, S., Pirone G., Gandini G., 2004. Chemical and microbiological parameters and sensory attributes of a typical Sicilian salami ripened in different conditions. *Meat Science* 66, 845– 854.

Morot-Bizot S.C., Leroy S., Talon R. (2006). *Staphylococcal* community of a small unit manufacturing traditional dry fermented sausages. *International journal of food microbiology*, 108(2), 210-217.

Olesen P.T., Stahnke L.H. (2000). The influence of *Debaryomyces hansenii* and *Candida utilis* on the aroma formation in garlic spiced fermented sausages and model minces. *Meat Science*, 56(4), 357-368.

- Papagianni M., Anastasiadou S. (2009). Pediocins: The bacteriocins of *Pediococci*. Sources, production, properties and applications. *Microbial Cell Factories*, 8(1), 3.
- Pisacane V., Callegari M. L., Puglisi E., Dallolio G., Rebecchi A. (2015). Microbial analyses of traditional Italian salami reveal microorganisms transfer from the natural casing to the meat matrix. *International journal of food microbiology*, 207, 57-65.
- Polka J., Rebecchi A., Pisacane V., Morelli L., Puglisi E. (2015). Bacterial diversity in typical Italian salami at different ripening stages as revealed by high-throughput sequencing of 16S rRNA amplicons. *Food microbiology*, 46, 342-356.
- Porto M.C.W., Kuniyoshi T.M., de Azevedo P.O.D.S., Vitolo M., de Souza Oliveira R.P. (2017). *Pediococcus* spp.: An important genus of lactic acid bacteria and pediocin producers. *Biotechnology Advances*. *Biotechnology advances*, 35(3), 361-374.
- Quijada N.M., De Filippis F., Sanz J.J., del Camino García-Fernández M., Rodríguez-Lázaro D., Ercolini D., Hernández M. (2018). Different *Lactobacillus* populations dominate in “Chorizo de León” manufacturing performed in different production plants. *Food microbiology*, 70, 94-102.
- Raghupathi P.K., Herschend J., Røder H.L., Sørensen S.J., Burmølle M. (2016). Genome sequence of *Kocuria varians* G6 isolated from a slaughterhouse in Denmark. *Genome announcements*, 4(2), e00076-16.
- Rantsiou K., Cocolin L. (2006). New developments in the study of the microbiota of naturally fermented sausages as determined by molecular methods: a review. *International Journal of Food Microbiology*, 108, 255–267.
- Rantsiou K., Drosinos E. H., Gialitaki M., Urso R., Krommer J., Gasparik-Reichardt J., Tóth S., Metaxopoulos I., Comi G., Cocolin L. (2005a). Molecular characterization of *Lactobacillus* species isolated from naturally fermented sausages produced in Greece, Hungary and Italy. *Food Microbiology*, 22(1), 19-28.
- Rantsiou K., Urso R., Iacumin L., Cantoni C., Cattaneo P., Comi G., Cocolin L. (2005b). Culture-dependent and-independent methods to investigate the microbial ecology of Italian fermented sausages. *Applied and Environmental Microbiology*, 71(4), 1977-1986.

- Rebecchi A., Pisacane V., Miragoli F., Polka J., Falasconi I., Morelli L., Puglisi E. (2015). High-throughput assessment of bacterial ecology in hog, cow and ovine casings used in sausages production. *International journal of food microbiology*, 212, 49-59.
- Rivas F.P., Castro M.P., Vallejo M., Marguet E., Campos C.A. (2014). Sakacin Q produced by *Lactobacillus curvatus* ACU-1: functionality characterization and antilisterial activity on cooked meat surface. *Meat science*, 97(4), 475-479.
- Samelis J., Stavropoulos S., Kakouri A., Metaxopoulos J. (1994). Quantification and characterization of microbial population associated with naturally fermented Greek dry salami. *Food Microbiol.* 11, 447-460.
- Santiago-Rodriguez T. M., Cano R., Jiménez-Flores R. (2016). Potential applications of metagenomics to assess the biological effects of food structure and function. *Food & function*, 7(10), 4160-4169.
- Schillinger U.; Lücke F. (1989). Antibacterial activity of *Lactobacillus sake* isolated from meat. *Appl. Environ. Microbiol.*, 55: 1901-1906.
- Silveti T., Capra E., Morandi S., Cremonesi P., Decimo M., Gavazzi F., Giannico R., De Noni I., Brasca M. (2017). Microbial population profile during ripening of Protected Designation of Origin (PDO) Silter cheese, produced with and without autochthonous starter culture. *LWT-Food Science and Technology*, 84, 821-831.
- Sonjak S., Ličen M., Frisvad J. C., Gunde-Cimerman N. (2011). The mycobiota of three dry-cured meat products from Slovenia. *Food microbiology*, 28(3), 373-376.
- Stefanis C., Mantzourani I., Plessas S., Alexopoulos A., Galanis A., Bezirtzoglou E. (2016). Reviewing Classical and Molecular Techniques Regarding Profiling of Probiotic Character of Microorganisms. *Current Research in Nutrition and Food Science Journal*, 4(1), 27-47.
- Sunesen L. O., Stahnke L. H. (2003). Mould starter cultures for dry sausages-selection, application and effects. *Meat Science*, 65(3), 935-948.
- Talon R., Leroy S. (2011). Diversity and safety hazards of bacteria involved in meat fermentations. *Meat science*, 89(3), 303-309.
- Talon R., Leroy S., Lebert I. (2007). Microbial ecosystems of traditional fermented meat products: the importance of indigenous starters. *Meat Science*, 77(1), 55-62.

- Todorov S.D. (2009). Bacteriocins from *Lactobacillus plantarum* production, genetic organization and mode of action: produção, organização genética e modo de ação. Brazilian journal of microbiology, 40(2), 209-221.
- Visagie C.M., Houbraken J., Frisvad J.C., Hong S.B., Klaassen C.H.W., Perrone G., Seifert K.A., Varga J., Yaguchi T., Samson R.A. (2014). Identification and nomenclature of the genus *Penicillium*. Studies in Mycology, 78, 343-371.
- Woraprayote W., Malila Y., Sorapukdee S., Swetwiwathana A., Benjakul S., Visessanguan W. (2016). Bacteriocins from lactic acid bacteria and their applications in meat and meat products. Meat science, 120, 118-132.
- Zhang Q.Q., Jiang M., Rui X., Li W., Chen X.H., Dong M.S. (2017). Effect of rose polyphenols on oxidation, biogenic amines and microbial diversity in naturally dry fermented sausages. Food Control, 78, 324-330.

616 **Figure legenda**

617 Table 1. Main differences between traditional and industrial fermented food products (adapted and modified
618 from El Sheikha and Montet, 2016).

619

620 Figure 1 Summary of biochemical activities performed by principal microbial groups in fermented sausages.

621 The most frequently isolated species of each group are nominated.

622

623

Traditional fermented products	Industrial fermented products
Small-scale	Large-scale
Manual	Automated
Intensive to time	Time-sensitive
Possible exposure to contaminants	Minimal exposure to contaminants
Varying quality	Constant quality
Complex sensory attributes	Less complex sensory attributes
Attention to organoleptic characteristic of the product	Safety driven operation
Shorter shelf-life	Longer shelf-life
Large undefined microbial diversity	Reduced microbial diversity
Limited use of selected microbial cultures	Extensive use of microbial cultures

624

625

626

627