



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Fate of selected pathogens in spiked «SALAME NOSTRANO» produced without added nitrates following the application of NONIT™ technology

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1687703> since 2021-12-29T17:55:21Z

Published version:

DOI:10.1016/j.meatsci.2018.02.002

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Table 1.

Kinetic parameters ⁽¹⁾	<i>Lc. lactis</i> lactis 16	<i>Lc. lactis</i> lactis 340	<i>Lb. casei</i> casei 208	<i>E. faecium</i> UBEF-41						
Vm ($\Delta\text{pH}/\text{min}$)	-2.06x10 ⁻⁰⁴	-2.35x10 ⁻⁰⁴	-1.51x10 ⁻⁰⁴	-1.39x10 ⁻⁰⁴						
Tm (min)	3754	7564	5003	4105						
pHm	6.00	5.47	5.70	6.09						
Regression coefficients										
	estimate	se	estimate	se	estimate	se	estimate	se		
a	6.37	7.96x10 ⁻⁰⁴	6.35	7.67x10 ⁻⁰⁴	6.36	0.09x10 ⁻⁰²	6.34	4.01x10 ⁻⁰⁴		
b	1.42x10 ⁻⁰⁴	9.85x10 ⁻⁰⁷	1.87x10 ⁻⁰⁵	8.96x10 ⁻⁰⁷	-8.84x10 ⁻⁰⁵	1.81x10 ⁻⁰⁶	1.17x10 ⁻⁰⁴	4.87x10 ⁻⁰⁷		
c	-1.07x10 ⁻⁰⁷	3.58x10 ⁻¹⁰	-6.36x10 ⁻⁰⁹	3.07x10 ⁻¹⁰	4.49x10 ⁻⁰⁸	1.05x10 ⁻⁰⁹	-7.28x10 ⁻⁰⁸	1.74x10 ⁻¹⁰		
d	1.33x10 ⁻¹¹	4.80x10 ⁻¹⁴	-3.32x10 ⁻¹²	3.89x10 ⁻¹⁴	-1.80x10 ⁻¹¹	2.25x10 ⁻¹³⁴	8.44x10 ⁻¹²	2.29x10 ⁻¹⁴		
e	-5.06x10 ⁻¹⁶	2.12x10 ⁻¹⁸	2.38x10 ⁻¹⁶	1.63x10 ⁻¹⁸	1.50x10 ⁻¹⁵	1.60x10 ⁻¹⁷	-3.08x10 ⁻¹⁶	9.96x10 ⁻¹⁹		
r ² ⁽²⁾	0.998		0.999		0.998		0.999			
Kinetic parameters ⁽¹⁾	NoNit™ formulation	NoNit™ formulation + <i>S. aureus</i> 27R	NoNit™ formulation + <i>E. coli</i> CSH26 K 12	NoNit™ formulation + <i>L. innocua</i> ATCC33090	NoNit™ formulation + <i>S. Derby</i> 27					
Vm ($\Delta\text{pH}/\text{min}$)	-3.09x10 ⁻⁰⁴	-5.13x10 ⁻⁰⁴	-4.29x10 ⁻⁰⁴	-4.14x10 ⁻⁰⁴	-3.92x10 ⁻⁰⁴					
Tm (min)	2296	3262	2231	2554	2887					
pHm	5.69	5.5	5.71	5.63	5.53					
Regression coefficients										
	estimate	se	estimate	se	estimate	se	estimate	se		
a	6.33	1.50x10 ⁻⁰³	6.17	9.48x10 ⁻⁰⁴	6.21	1.15x10 ⁻⁰³	6.10	7.04x10 ⁻⁰⁴	6.24	9.70x10 ⁻⁰⁴
b	-2.13x10 ⁻⁰⁴	1.89x10 ⁻⁰⁶	1.10x10 ⁻⁰⁴	2.65x10 ⁻⁰⁶	2.41x10 ⁻⁰⁴	2.53x10 ⁻⁰⁶	3.36x10 ⁻⁰⁴	1.33x10 ⁻⁰⁶	6.05x10 ⁻⁰⁵	2.49x10 ⁻⁰⁶
c	-4.56x10 ⁻⁰⁸	7.01x10 ⁻¹⁰	-5.38x10 ⁻⁰⁹	2.18x10 ⁻⁰⁹	-3.46x10 ⁻⁰⁷	1.63x10 ⁻⁰⁹	-3.41x10 ⁻⁰⁷	7.39x10 ⁻¹⁰	-1.65x10 ⁻⁰⁷	1.88x10 ⁻⁰⁹
d	8.32x10 ⁻¹²	9.61x10 ⁻¹⁴	-5.64x10 ⁻¹¹	6.60x10 ⁻¹³	7.24x10 ⁻¹¹	3.90x10 ⁻¹³	6.30x10 ⁻¹¹	1.51x10 ⁻¹³	2.19x10 ⁻¹¹	5.23x10 ⁻¹³
e	-3.70x10 ⁻¹⁶	4.35x10 ⁻¹⁸	8.73x10 ⁻¹⁵	6.60x10 ⁻¹⁷	-4.64x10 ⁻¹⁵	3.07x10 ⁻¹⁷	-3.62x10 ⁻¹⁵	1.03x10 ⁻¹⁷	-4.93x10 ⁻¹⁶	4.80x10 ⁻¹⁷
r ² ⁽²⁾	0.998		0.999		0.999		0.9995		0.999	

⁽¹⁾ Vm is the maximum instantaneous acidification rate, Tm and pHm the time and the pH at which Vm occurred.

⁽²⁾ Significant at p<0.001.

Table 2.

<i>S. aureus</i> 27R				<i>E. coli</i> CSH26 K12				<i>L. innocua</i> ATCC 33090				<i>S. Derby</i> 27				
Time (hrs)	log cfu ml ⁻¹		log reduction		log cfu ml ⁻¹		log reduction		log cfu ml ⁻¹		log reduction		log cfu ml ⁻¹		log reduction	
	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd
0 (control)	5,73	0,63	0,00	0,00	6,96	0,24	0,00	0,00	6,20	0,26	0,00	0,00	6,45	0,24	0,00	0,00
24	5,24	0,31	-0,50	0,17	7,17	0,1	0,21	0,17	6,38	0,21	0,18	0,19	6,68	0,37	0,23	0,16
72	5,20	0,24	-0,53	0,14	7,13	0,19	0,17	0,17	6,88	0,29	0,76	0,37	6,87	0,42	0,42	0,23
120	5,80	0,55	0,41	0,44	6,80	0,38	-0,36	0,14	6,91	0,09	0,71	0,13	7,06	0,17	0,71	0,12
168	5,23	0,19	-0,51	0,11	6,61	0,41	-0,56	0,18	5,60	0,46	-0,60	0,20	6,32	0,19	-0,13	0,43
240	3,94*	0,93	-1,45*	0,68	6,28*	0,21	-0,81*	0,23	4,80*	0,12	-1,30*	0,19	4,05*	0,21	-2,18*	0,33

*: indicates significant different means ($p<0,005$) when compared to control (time 0).

Table 3.

	Commercial		NoNit™	
	mean	sd	mean	sd
<i>S. aureus</i> 27R				
day-0	4,19	0,17	4,26	0,15
day-3	5,45 ^a	0,14	4,04 ^b	0,47
day-7	5,89 ^a	0,10	3,69 ^b	0,50
day-13	5,62 ^a	0,48	3,87	0,59
day-21	6,00 ^a	0,12	3,85 ^b	0,15
day-28	6,01 ^a	0,33	3,82 ^b	0,16
<i>E. coli</i> CSH26 K-12				
day-0	4,03	0,22	4,18	0,24
day-3	4,54	0,08	4,53	0,20
day-7	5,05 ^a	0,03	4,39 ^b	0,21
day-13	4,62 ^a	0,14	3,75 ^b	0,17
day-21	3,92 ^a	0,39	2,94 ^b	0,16
day-28	3,54 ^a	0,32	1,93 ^b	0,63
<i>L. innocua</i> ATCC 33090				
day-0	4,09	0,20	4,31	0,05
day-3	5,99 ^a	0,08	4,73 ^b	0,12
day-7	5,88 ^a	0,12	4,97 ^b	0,01
day-13	5,89 ^a	0,11	4,98 ^b	0,05
day-21	5,89 ^a	0,12	5,02 ^b	0,24
day-28	5,80 ^a	0,06	4,93 ^b	0,05
<i>S. Derby</i> 27				
day-0	3,69	0,16	3,90	0,13
day-3	5,42 ^a	0,13	4,43 ^b	0,17
day-7	4,97	0,10	4,35	0,24
day-13	4,37	0,47	3,63	0,28
day-21	3,41 ^a	0,20	2,55 ^b	0,42
day-28	3,67 ^a	0,44	1,79 ^b	0,86

Table 4

		Commercial mean	sd	NoNit™ mean	sd
<i>Total mesophilic microbiota</i>					
day-0	6,69 ^a	0,02		6,34 ^b	0,06
day-3	7,87	0,07		8,00	0,13
day-7	8,50 ^a	0,11		8,23 ^b	0,08
day-13	8,50 ^a	0,10		8,24 ^b	0,07
day-21	8,40 ^a	0,10		8,10 ^b	0,07
day-28	8,48 ^a	0,06		8,06 ^b	0,12
<i>Lactococcus</i> spp.					
day-0	6,34	0,15		6,16	0,07
day-3	6,72 ^a	0,06		6,29 ^b	0,04
day-7	6,94 ^a	0,16		6,35 ^b	0,04
day-13	7,06 ^a	0,03		6,37 ^b	0,06
day-21	7,05 ^a	0,05		6,49 ^b	0,35
day-28	6,76 ^a	0,03		6,37 ^b	0,12
<i>Lactobacillus</i> spp.					
day-0	6,66 ^a	0,06		6,20 ^b	0,05
day-3	8,38 ^a	0,13		8,05 ^b	0,10
day-7	8,21	0,10		8,12	0,06
day-13	8,31	0,04		8,25	0,08
day-21	8,25	0,12		8,08	0,06
day-28	8,40 ^a	0,16		8,05 ^b	0,15
<i>Enterococcus</i> spp.					
day-0	4,14 ^a	0,23		6,11 ^b	0,04
day-3	6,76	0,26		6,98	0,06
day-7	6,64	0,39		6,82	0,20
day-13	6,24	0,03		6,58	0,19
day-21	6,49	0,31		6,75	0,15
day-28	6,47	0,29		6,66	0,08
<i>Staphylococcus</i> spp.					
day-0	4,03	0,22		3,06	0,23
day-3	4,54	0,08		3,83	0,34
day-7	5,05 ^a	0,03		3,24 ^b	0,10
day-13	4,62 ^a	0,14		3,13 ^b	0,49
day-21	3,92 ^a	0,39		1,59 ^b	1,20
day-28	3,25 ^a	0,26		0,22 ^b	0,67

Table 5

Initial batch	day of stuffing	Mean	sd	n
proteins g 100 g ⁻¹ total solids		40.44	0.35	3
fat g 100 g ⁻¹ total solids		50.32	2.56	3
ashes g 100 g ⁻¹ total solids		8.88	0.19	3
moisture g 100 g ⁻¹		58.67	1.77	3
Salami with commercial starter and pathogens	end of ripening	Mean	sd	n
proteins g 100 g ⁻¹ total solids		43.38	4.11	3
fat g 100 g ⁻¹ total solids		45.84	6.02	3
ashes g 100 g ⁻¹ total solids		9.51	0.58	3
moisture g 100 g ⁻¹		28.28	1.40	3
Salami with NoNit™ starter and pathogens	end of ripening	Mean	sd	n
proteins g 100 g ⁻¹ total solids		43.91	4.72	3
fat g 100 g ⁻¹ total solids		45.50	4.13	3
ashes g 100 g ⁻¹ total solids		9.26	1.33	3
moisture g 100 g ⁻¹		29.30	0.72	3

Table 6.

	Commercial		NoNit™	
	mean	sd	mean	sd
color intensity	4,38	0,92	4,53	1,81
color uniformity	3,38 ^a	1,30	5,00 ^b	1,65
fat/lean connection	3,63	1,19	4,47	1,64
fat/lean distribution	3,43 ^a	0,79	4,87 ^b	1,69
odour (global)	3,88	0,83	4,53	1,36
mould odor	3,00	1,51	2,47	1,46
elasticity	3,88	1,81	3,27	1,62
hardness	2,50 ^a	1,60	3,87 ^b	1,41
cohesiveness	4,63	1,92	4,93	1,34