

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Fate of selected pathogens in spiked «SALAME NOSTRANO» produced without added nitrates following the application of NONIT™ technology

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1687703> since 2021-12-29T17:55:21Z

Published version:

DOI:10.1016/j.meatsci.2018.02.002

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)






Figure 2a


16 at 10 °C



208 at 10 °C



340 at 10 °C



Enterelle at 10 °C






Figure 2b

LAB at 10 °C




LAB + E. coli at 10 °C




b

LAB + Salm at 10 °C



LAB + List at 10 °C



LAB + Staph at 10 °C




Figure 3a




Figure 3b




Figure 3c




Figure 3d




Figure 4a




Figure 4b




Figure 4c




Figure 4d




Figure 5a




Figure 5b




Figure 5c




Figure 5d




Figure 6a




Figure 6b




Figure 6c




Figure 6d




Figure 6e




Figure 7




Figure 8





Figure 9a






Figure 9b