

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Ectopic activation of cortical cell division during the accommodation of arbuscular mycorrhizal fungi

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1688618> since 2019-01-30T11:18:00Z

Published version:

DOI:10.1111/nph.15398

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Title page**

2

3 **Authors**

4 Giulia Russo¹, Gennaro Carotenuto¹, Valentina Fiorilli¹, Veronica Volpe¹, Marco Chiapello¹, Daniel
5 Van Damme^{2,3}, Andrea Genre¹

6

7 ¹Department of Life Sciences and Systems Biology, University of Torino, 10125 Torino, Italy

8 ²Ghent University, Department of Plant Biotechnology and Bioinformatics, Technologiepark 927,
9 9052 Ghent, Belgium

10 ³VIB Center for Plant Systems Biology, Technologiepark 927, 9052 Ghent, Belgium

11

12 **Corresponding author:**

13 Andrea Genre: andrea.genre@unito.it, Tel: +39 011 6705083

14

15 **Title**

16 Ectopic activation of cortical cell division during the accommodation of arbuscular mycorrhizal fungi

17

18 **Short title**

19 Cortical cell division in arbuscular mycorrhizas

20

21 Total word count for the main body: 5671

22 Word count for section Introduction: 909

23 Word count for Materials and methods: 1619

24 Word count for Results: 2049

25 Number of Figures: 5

26 Number of colored Figures: 4

27 Number of Supplementary Figures: 7

28 Number of tables: 0

29 Word count for Discussion: 1089

30

31 **ORCID IDs:**

32 Valentina Fiorilli: 0000-0001-9805-1559

33 Marco Chiapello: 0000-0001-7768-3047

34 Daniel Van Damme: 0000-0002-9385-4851

35 Andrea Genre: 0000-0001-5029-6194

36

37

1 **Summary**

- 2 • Arbuscular mycorrhizas (AM) between plants and soil fungi are widespread symbioses with a
3 major role in soil nutrient uptake.
- 4 • In this study we investigated the induction of root cortical cell division during AM colonization
5 by combining morphometric and gene expression analyses with promoter activation and protein
6 localization studies of the cell plate-associated exocytic marker TPLATE.
- 7 • Our results show that TPLATE promoter is activated in colonized cells of the root cortex where
8 we also observed the appearance of cells that are half the size of the surrounding cells. Furthermore,
9 TPLATE-GFP recruitment to developing cell plates highlighted ectopic cell division events in the
10 inner root cortex during early AM colonization. Lastly, transcripts of TPLATE, KNOLLE, and
11 CYC1 are all upregulated in the same context, alongside endocytic markers AP2A1 and CHC2,
12 known to be active during cell plate formation. This pattern of gene expression was recorded in wild-
13 type *Medicago truncatula* roots but not in a CSSP mutant where fungal colonization is blocked at the
14 epidermal level.
- 15 • Altogether, these results suggest the activation of cell division-related mechanisms by AM hosts
16 during the accommodation of the symbiotic fungus.

1 **Introduction**

2 The majority of extant plants, including both wild and domesticated crop species, is supported in its
3 uptake of soil nutrients by Glomeromycotina fungi (Lin *et al.*, 2014) - in a beneficial symbiosis called
4 arbuscular mycorrhiza (AM). The broad diffusion of this interaction among plant species is
5 considered a consequence of its ancient evolutionary origin (Bonfante & Genre, 2008). Indeed, 400-
6 million-year old plant fossils already hosted hyphal structures that closely resemble the arbuscules
7 produced by living AM fungi (Remy *et al.*, 1994; Strullu-Derrien *et al.*, 2018). Arbuscules are highly
8 branched hyphae that develop inside living cells of the root cortex (Gutjahr & Parniske, 2013). In
9 fact, arbuscules are accommodated inside a novel cell compartment enveloped by the perifungal (or
10 periarbuscular) membrane, a specialized extension of the host plasmalemma (Pumplin & Harrison,
11 2009). This symbiotic interface compartment assembles within the pre-penetration apparatus (PPA),
12 a broad, nucleus-associated cytoplasmic aggregation, where the secretory process is focused and
13 coordinated (Genre *et al.*, 2005; 2008).

14 Both PPA organization and fungal accommodation depend on the activation of a conserved signal
15 transduction pathway (Delaux *et al.*, 2015), the so-called Common Symbiotic Signaling Pathway, or
16 CSSP (see Oldroyd, 2013 for a comprehensive review) and represent the earliest visible response of
17 plant cells to AM fungi (Genre *et al.*, 2005). The CSSP is believed to transduce diffusible AM fungal
18 signals through the sequential activation of a number of proteins, starting from receptor-like kinases
19 in the plasma membrane and culminating with the nuclear localized kinase DOESN'T MAKE
20 INFECTIONS 3 (DMI3; Lévy *et al.*, 2004), which mediates the regulation of gene expression by
21 activating a set of transcription factors (Oldroyd, 2013; Zipfel and Oldroyd, 2017; Floss *et al.*, 2017).
22 Altogether, CSSP activation is essential for root colonization (Novero *et al.*, 2002; Lévy *et al.* 2004,
23 Kistner *et al.*, 2005; Parniske, 2008), as demonstrated by the block of PPA formation (Genre *et al.*,
24 2005) and epidermal penetration in *dmi3* mutants (Morandi *et al.*, 2005).

25 Biogenesis of the perifungal membrane has been described as a massive exocytic process that
26 involves cytoskeletal rearrangements as well as the preemptive accumulation of an extensive quantity
27 of secretory membranes, such as trans-Golgi network (TGN) compartments and vesicles (Genre *et*
28 *al.*, 2012). Alongside secretion-related endomembrane compartments, transmission electron
29 microscopy imaging of PPA aggregates has shown the abundant presence of multivesicular bodies -
30 or late endosomes - indicating that endocytic processes are likely also in place during interface
31 biogenesis (Genre *et al.*, 2012), even if their role has not been investigated further.

32 Such intense and targeted membrane dynamics have few analogs in plant cells and possibly the most
33 notable similarity is with cell division. This process initiates with vacuole partitioning by fusion of
34 multiple trans-vacuolar cytoplasmic bridges (phragmosome) and nuclear positioning in the cell center

1 (Smith, 2001; Lloyd & Buschmann, 2007). Phragmosome formation is followed by regular mitosis
2 and cell plate assembly through the progressive fusion of TGN-derived cytokinetic vesicles (Segui-
3 Simarro *et al.*, 2004; Smertenko *et al.*, 2017). Intriguingly, Siciliano *et al.*, (2007) reported that
4 histone H2B1 gene was induced in root segments of *Medicago* on which hyphopodium formation had
5 occurred, hinting at the possibility that cell cycle determinants are active during early AM
6 colonization. The involvement of cell division-related processes has also been proposed for the
7 symbiotic nitrogen fixation, where infection thread growth is coupled to cell cycle reactivation and
8 cell division events (Timmers *et al.*, 1999; Fournier *et al.*, 2008; Breakspear *et al.*, 2014; Downie,
9 2014).

10 During plant cell division, the massive exocytic process directed by cytoskeletal fibers in the
11 phragmoplast (Lee & Liu, 2013; Boruc & van Damme, 2015) is associated with endocytic recycling
12 of surplus membrane. In fact, next to cell plate specific markers such as the syntaxin KNOLLE, other
13 plasma membrane proteins also strongly accumulate at the forming cell plate (Van Damme *et al.*,
14 2006; Richter *et al.*, 2014), among which, the adaptin-related protein TPLATE has been shown to
15 accumulate on both the cell plate membranes and plasmalemma at the cortical division zone, where
16 the cell plate will eventually fuse (Van Damme *et al.*, 2006; Van Damme *et al.*, 2011; Gadeyne *et al.*,
17 2014). TPLATE is the founding member of an octameric protein complex, the TPLATE complex
18 (TPC), which acts largely together with Adaptor Protein complex 2 (AP-2) and other components
19 (such as Dynamin-related proteins) and the clathrin scaffolding proteins for the assembly of the
20 clathrin endocytic machinery at the plasma membrane (Gadeyne *et al.*, 2014). The accumulation of
21 TPC members at the cortical division zone has been proposed to facilitate membrane reorganization
22 during cell plate fusion with the peripheral plasmalemma (Boutté *et al.*, 2010; Van Damme *et al.*,
23 2011). In short, TPLATE appears to be a promising marker to investigate the occurrence of both cell
24 division- and endocytosis-related processes during the accommodation of AM fungi.

25 Here, we report that *TPLATE* expression is activated in cortical root cells colonized by arbuscular
26 mycorrhizal fungi, where we also observed the appearance of arbusculated cells with half the size of
27 the surrounding cortical cells. Combined with TPLATE-GFP recruitment to forming cell plates, this
28 allowed us to identify ectopic cell division events that are induced in the inner root cortex during AM
29 colonization of the outer root tissues. Lastly, *TPLATE* transcripts, alongside other cell division and
30 endocytic markers, are upregulated in the colonized area of wild-type roots but not in a CSSP mutant
31 where fungal colonization is blocked at the epidermal level. Taken together, these results suggest that
32 extant plants activate cell division-related mechanisms during AM fungal accommodation.

1 **Materials and Methods**

2

3 **Plant materials**

4 The generation of composite *L. japonicus* plants was performed using *Agrobacterium rhizogenes*
5 strain Ar1193 carrying the *pLjTPLATE:GUS* vector, according to the protocol by Stougaard (1995)
6 on wild-type seedlings of MG20 genotype (see below).

7 *Medicago truncatula* Jemalong A17 and *Lotus japonicus* MG20 plants were grown from surface
8 sterilized seeds in pots containing sterile quartz sand and respectively inoculated with *Gigaspora*
9 *margarita* or *Funnelliformis mosseae* and watered with Long-Ashton solution (Hewitt, 1966).

10 *Agrobacterium rhizogenes*-transformed root organ cultures (ROCs) expressing the *35S::AtTPLATE-*
11 *GFP* vector (Van Damme *et al.*, 2004) were obtained from *Medicago truncatula* Jemalong A17 wild-
12 type and *dmi3-1* plants (Sagan *et al.*, 1995; kindly provided by M. Chabaud, LIPM, INRA, Toulouse,
13 France), according to Boisson-Dernier *et al.* (2001). ROC generation was repeated in two independent
14 experiments for both lines, with overlapping results in terms of GFP fluorescence pattern. ROCs from
15 *Daucus carota* var *Sativus* expressing the same vector were obtained according to Bécard & Fortin
16 (1988). For both species, transformed roots with a high level of fluorescence were selected 21 days
17 after transformation, decontaminated and subcultured on M medium (Bécard & Fortin, 1988) at 25°C
18 in the dark for subsequent use as hairy roots. In each case, a single representative clone was chosen
19 for further studies. Transformation efficiency and expression of *35S::AtTPLATE-GFP* was checked
20 with GFP specific primers (Supporting Information Table S1) on both genomic DNA and cDNA
21 obtained from all the selected lines. Wild type and *dmi3-1* mutant *M. truncatula* ROCs expressing a
22 nuclearameleon YC2.1 probe were already available in the lab (Chabaud *et al.*, 2011; Genre *et al.*,
23 2013) and were used for all gene expression analyses.

24

25 **Fungal materials**

26 Different AM fungi were chosen depending on how their characteristics suited the experimental
27 setup. The AM fungus *Funnelliformis mosseae* (strain BEG 12), supplied by MycAgroLab
28 (<http://www.mycagrolab.com/>), was used to inoculate *L. japonicus* composite plants with a 10% ratio
29 of inoculum to total substrate volume; *Gigaspora margarita* (BEG34, International Bank for
30 Glomeromycota, University of Kent, Canterbury, UK) was used to inoculate *M. truncatula* whole
31 plants (50 spores per pot) and ROCs (Chabaud *et al.*, 2002) for gene expression analyses. *Gigaspora*
32 *gigantea* (isolate HC/FE30, Herbarium Cryptogamicum Fungi, University of Torino, Italy), which is
33 characterized by strong cytoplasmic autofluorescence (Genre *et al.*, 2005), was used to inoculate
34 ROCs for live confocal imaging. Spores of both *G. gigantea* and *G. margarita* were collected from

1 pot cultures in sand (with leek and clover, respectively), surface-sterilized and stored at 4°C according
2 to Bécard & Fortin (1988).

4 **GUS assay**

5 A *LjTPLATE* promoter fragment of 2073 bp (Supporting Information Fig. S1) was PCR-amplified
6 from genomic DNA by using specific primers (Supporting Information Table S1). The resulting
7 promoter fragment was cloned into pDONR221 (Invitrogen) and recombined upstream of the GUS
8 gene into pKGWFS7.0 (Gateway vector), modified to contain the red fluorescent marker DsRed
9 under the control of the constitutive *Arabidopsis* UBIQUITIN10 promoter (pUBQ10; Limpens *et al.*,
10 2005).

11 Half of the 30 *L. japonicus* composite plants selected for DsRed fluorescence were inoculated with
12 *F. mosseae* and the remaining were used as controls. Root sampling was repeated at 14, 21 and 42
13 days post inoculation.

14 Hairy roots expressing DsRed fluorescence were selected using a stereomicroscope (Leica M205 FA)
15 from both inoculated and control plants; excised roots were immediately submerged in freshly
16 prepared GUS buffer (0.1 M potassium phosphate buffer pH 7, 5 mM K₄Fe(CN)₆, 5 mM K₃Fe(CN)₆,
17 0.3% Triton X-100, 0.3% X-GlcA; Duchefa Biochemie), incubated overnight at 37°C in the dark,
18 destained with 70% ethanol and washed twice in distilled water. Part of the roots were then counter-
19 stained with 0.01% (w/v) acid fuchsin in lactoglycerol (lactic acid:glycerol: water, 14:1:1; Kormanik
20 & McGraw, 1982) to highlight intraradical fungal structures. Samples were then either directly
21 mounted on a microscope slide for observation, or previously embedded in agarose (5%) and cut into
22 100 µm thick vibratome slices. Observations were done using a bright field microscope (Nikon
23 Eclipse E400).

25 **Confocal microscopy imaging**

26 The targeted AM inoculation technique for studying early stages of the symbiotic association between
27 *Gigaspora* species and transformed root cultures, developed by Chabaud *et al.* (2002) and adapted
28 for confocal observation by Genre *et al.* (2005), was applied to both *Medicago truncatula* and *Daucus*
29 *carota* ROCs expressing *35S::AtTPLATE-GFP*. An upright Leica TCS SP2 confocal microscope
30 fitted with a long distance 40X water-immersion objective (HCX Apo 0.80) was used for imaging
31 living ROCs directly in the Petri dishes. The argon laser band of 488 nm was used to excite both GFP
32 and *G. gigantea* autofluorescence. The two signals were distinguished using specific emission
33 windows: 500 to 525 nm for GFP and 590 to 630 nm for fungal autofluorescence. The latter channel
34 was then false-colored in red to maximize the contrast in overlapping images.

1 **Morphometric analyses**

2 In order to visualize intraradical fungal structures and perform a quantitative analysis of cortical cell
3 morphology, root samples from 8 weeks old pot-grown plants were harvested and fixed in phosphate-
4 buffered saline (PBS), pH 7.2, containing 1% formaldehyde and 10% dimethyl sulfoxide (DMSO)
5 for 24 hours at 4°C. Fixed roots were sliced using a Vibratome (Oxford Vibratome® sectioning
6 system) and stained with wheat germ agglutinin and propidium iodide (Kojima *et al.*, 2014) or acid
7 fuchsin (Kormanik & McGraw, 1982) to highlight intraradical fungal structures and plant cell walls,
8 respectively. The argon laser line at 488nm was used to excite all fluorochromes and emission
9 windows were set at 500-530nm to record FITC, 600-700nm for propidium iodide and fuchsin
10 fluorescence. A total of 42 optical sections (375 x 375 µm) were recorded from 13 independent root
11 segments sampled from 3 different uninoculated plants of *M. truncatula*; 47 optical sections from 8
12 independent root segments (3 plants) were used for colonized *M. truncatula*; 22 optical sections (6
13 root segments from 3 plants) were used for uninoculated *L. japonicus* and 23 optical sections (6 root
14 segments from 3 plants) for colonized *L. japonicus*. For all our analyses, 0.5cm-long segments from
15 lateral roots of comparable size, length and position in the root system were sampled from both
16 inoculated and uninoculated plants.

17 In addition, the number of split cells (Supporting Information Table S2) was counted in the inner
18 cortical layer of live uninoculated or inoculated ROCs of *D. carota* and wild type or *dmi3-1 M.*
19 *truncatula* expressing *35S::AtTPLATE-GFP*. A minimum of 22 optical sections (375 x 375µm) from
20 at least 5 different roots of comparable size, length and position, belonging to 3 independent ROCs
21 was used for each condition.

22 **Quantitative gene expression analyses**

24 Wild type and *dmi3-1 Medicago truncatula* ROCs were inoculated with *Gigaspora margarita*; fungal
25 development and root colonization were followed under a stereomicroscope to identify hyphopodium
26 development. A few of these samples were further observed in fluorescence microscopy (Leica TCS-
27 SP2) to verify the repositioning of YC2.1 fluorescent nuclei under the hyphopodia - a hallmark of
28 epidermal cell prepenetration responses. After 48/60 hours - required for the development of young
29 colonization units – the corresponding root segments were excised and immediately frozen in liquid
30 nitrogen. All root tips were removed before sampling. Four biological replicates - each consisting of
31 a pool of ten 1cm long root segments - were collected from both uncolonized and colonized wild type
32 ROC lines; and three replicates were collected from the corresponding *dmi3-1* lines. Uncolonized
33 root segments were selected from lateral roots with a similar development and position in the root
34 system architecture as the colonized segments from inoculated ROCs.

1 Total RNA was isolated from colonized and uncolonized samples and prepared using the RNeasy™
2 Plant Mini kit (Qiagen, Hilden, Germany). Plant material was processed with a mechanical disruption
3 procedure using a TissueLyser system (Qiagen Retsch GmbH, Hannover, Germany) and further
4 treated following the manufacturer's instructions. RNA quality and quantity were evaluated by
5 photometric measurements and on agarose gels.

6 To remove traces of genomic DNA before cDNA synthesis, samples were treated with TURBO™
7 DNase (Ambion) according to the manufacturer's instructions. The RNA samples were routinely
8 checked for DNA contamination by RT-PCR analysis, using primers for *MtTEF* (Supporting
9 Information Table S1). Conventional PCR assays on fungal genomic DNA excluded any cross-
10 hybridization of *M. trunctatula* specific primers.

11 For single-strand cDNA synthesis, about 700 ng of total RNA was denatured at 65 °C for 5 min and
12 then reverse-transcribed at 25 °C for 10 min, 42 °C for 50 min and 70 °C for 15 min. The final volume
13 was 20µl and contained 10µM of random primers, 0.5mM dNTPs, 4µl 5Xbuffer, 2 µl 0.1 M DTT and
14 1µl Super-Script II (Invitrogen, Carlsbad,CA, USA).

15 Specific primers for *MtTPLATE*, *MtAP2A1*, *MtCHC1* and *MtCHC2* were designed on putative *M.*
16 *trunctatula* CDS sequences from NCBI database (<http://www.ncbi.nlm.nih.gov/>) and designed using
17 SerialCloner software specifically for qRT-PCR analyses (Supporting Information Table S1). Primer
18 validation was obtained with conventional PCR assay on *M. trunctatula* cDNA to avoid overlapping
19 with unspecified sequences, and tested on fungal genomic DNA.

20 qRT-PCR experiments were carried out in a final volume of 15µl containing 7.5µl of iTaq™
21 Universal SYBR® Green Supermix (Bio-Rad), 1µl of 3µM specific primers, and 10 ng of cDNA.
22 Samples were run in the Rotor Gene apparatus (Qiagen) using the following program: 10min pre-
23 incubation at 95°C, followed by 40 cycles of 15s at 95°C, and 30s at 60°C. Each amplification was
24 followed by melting curve analysis (60–94°C) with a heating rate of 0.5°C every 15s. All reactions
25 were performed with three technical replicates and only Ct values with a standard deviation that did
26 not exceed 0.3 were considered. The comparative threshold cycle method (Rasmussen, 2001) was
27 used to calculate relative expression levels using the plant *MtTEF* as reference genes.

28

29 **Accession numbers**

30 Sequence data from this article can be found in the NCBI data libraries under the following accession
31 numbers: LjT06B17 (*LjTPLATE* promoter from 47759 to 49830); MTR_5g012010 (*MtKNOLLE*);
32 MTR_7g089730 (*MtCYC1*); MTR_5g082900 (*MtCHCL1*); MTR_3g070940 (*MtCHCL2*);
33 MTR_7g031450 (*MtTPLATE*); MTR_2g084610 (*MtAP2A1*); MTR_1g105120 (*MtBCP*);
34 MTR_6g021805 (*MtTEF*).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

Results

***TPLATE* expression is activated in AM colonized areas of the root**

In order to highlight the involvement of the endocytosis and cell division marker *TPLATE* in the process of AM fungal accommodation, we studied the pattern of *TPLATE* promoter activity by expressing a *pLjTPLATE:GUS* transcriptional reporter (2kb promoter) in *Agrobacterium rhizogenes*-transformed roots from wild type *Lotus japonicus* and applying the GUS assay to excised roots. More than 50 roots were screened in brightfield microscopy and intense GUS-positive staining was observed in most root tips (Supporting Information Fig. S2a,b), consistent with the role of *TPLATE* in cell division. In roughly 1 out of 5 root tips, the expression pattern was less uniform and limited to small groups of meristematic cells (Supporting Information Fig. S2c,d) or isolated cell pairs (Supporting Information Fig. S2e). Taken together, the diverse meristematic patterns of *pLjTPLATE:GUS* staining appear to highlight a transient upregulation of the *TPLATE* promoter above a low basal level as previously described in Arabidopsis (Van Damme *et al.*, 2006; Brady *et al.*, 2007), even if literature data and public databases indicate that *TPLATE* mRNA is expressed in virtually all cells due to its constitutive role in endocytic processes, as confirmed by both protein localization specifically in root and hypocotyl epidermal cells as well as functional studies on *A. thaliana* seedlings (Van Damme *et al.*, 2006, Van Damme *et al.*, 2011; Gadeyne *et al.*, 2014, Sanchez-Rodriguez *et al.*, 2018). Moreover, GUS staining was often observed in the root vascular cylinder in both young (Supporting Information Fig. S2a) and differentiated areas of the root (Supporting Information Fig. S2f, h), possibly related to endocytic processes linked to phloem activity (Liesche *et al.*, 2008).

AM colonization induced an evident change in this *TPLATE* expression pattern, with the appearance of intense GUS-positive areas associated with intraradical fungal structures (Supporting Information Fig. S2g), independent of the sampling time (14, 21 or 42 days post inoculation). The observation of over 60 vibratome sections from 10 independent root samples consistently indicated that GUS activity was particularly intense in inner cortical cells containing arbuscules (Supporting Information Fig. S2i). This was further confirmed by the analysis of acid fuchsin counter-stained samples (20 root segments overall), where the precise location of intraradical fungal structures could be detected more easily in longitudinal sections (Fig. 1a-c).

In conclusion, the GUS assay provided strong indications that AM colonization boosts *TPLATE* promoter activity in arbusculated root cortical cells.

1 **Split cells are observed in the cortex of colonized roots**

2 In addition to the correlation between arbuscule presence and *TPLATE* promoter activation, a
3 consistent feature was observed in dozens of longitudinal root sections: the presence of one or more
4 couples of arbuscule-containing cells, half the size of the surrounding (either colonized or
5 uncolonized) cortical cells (Fig. 1d-f). In fact, the roughly square shape of such cells made them stand
6 out very clearly among the more elongated cortical cells.

7 To better investigate this intriguing observation and exclude that ectopic cell divisions could be
8 caused by our use of *A. rhizogenes*-transformed composite plants, we performed an extensive
9 screening of longitudinal root sections from both *L. japonicus* and *M. truncatula* wild type plants,
10 grown in pots for 8 weeks with or without AM inoculation. Fluorescent staining of fungal and plant
11 walls made split cells easy to recognize by confocal imaging (see materials and methods for details).
12 Representative images of *L. japonicus* samples (from a total of 6 colonized and 6 uninoculated root
13 segments) are shown in Fig. 2. Couples of short cortical cells were only occasionally observed in
14 uninoculated plants (Fig. 2a, b). By contrast, split cells were regularly found in AM colonized roots
15 (Fig. 2c-e), alongside arbusculated and non-arbusculated cortical cells of regular size. In more detail,
16 both unpenetrated (Fig. 2c, d) and arbusculated split cells (Fig. 2c-e) were observed in the colonized
17 area, including couples where one cell contained an arbuscule and the second one did not (Fig. 2c),
18 and couples where the two arbuscules had different (Fig. 2c,d) or comparable stages of development
19 (Fig. 2e).

20 The table in Fig. 2, resumes our quantitative analysis of split cell occurrence in control and
21 mycorrhizal plants of both *L. japonicus* and *M. truncatula*. We quantified the average number of split
22 cell couples per image and the number of images containing at least one couple of split cells. In both
23 plants, AM colonization significantly increased (t test <0.01) the average number of split cells: over
24 6 times in *L. japonicus* (from 0.71 to 4.80) and over 4 times in *M. truncatula* (from 0.58 to 2.53). The
25 percent of images containing split cells was also increased in both AM hosts, from 36% to 96% in *L.*
26 *japonicus* and from 36% to 66% in *M. truncatula*.

27 In short, the induction of anticlinal cell divisions was confirmed in the inner cortex of colonized areas
28 in untransformed *L. japonicus* and *M. truncatula* plants, excluding a causal connection with *A.*
29 *rhizogenes* infection. Secondly, our detailed quantification of split cell occurrence demonstrated a
30 statistically supported relation between AM colonization and inner cortical cell divisions. Lastly, our
31 observations of split cells where each contained differently developed arbuscules, alongside
32 uncolonized split cells next to the arbusculated area, were suggestive of cell divisions occurrence in
33 advance of arbuscule development.

34

1 **Live imaging of ectopic cortical cell divisions in mycorrhizal roots using TPLATE-GFP**

2 To clarify whether the AM-dependent induction of cortical cell division was indeed triggered before
3 fungal entry, we expressed a *35S::AtTPLATE-GFP* fusion (originally developed in *Arabidopsis*
4 *thaliana* and tobacco BY-2 cells by Van Damme *et al.*, 2004; 2006), in *A. rhizogenes*-transformed
5 root organ cultures (ROCs). Since *L. japonicus* ROCs can only very rarely be colonized in our
6 experimental conditions, we used *M. truncatula* (legume) and *D. carota* (non-legume) ROCs instead,
7 two more amenable models for live cell imaging of symbiotic interactions (Chabaud *et al.*, 2002;
8 Genre *et al.*, 2008; Fournier *et al.*, 2015).

9 *In vivo* observations of at least 10 uninoculated *M. truncatula* ROCs confirmed the predicted
10 localization of TPLATE-GFP (Van Damme *et al.*, 2004; 2006) with a major accumulation of the
11 construct along newly formed cell walls in the apical meristem (Supporting Information Fig. S3a,b).
12 By increasing microscope sensitivity, a weaker background fluorescence could also be recorded in
13 the cytoplasm (likely related to 35S-driven overexpression) and at the plasma membrane of both
14 meristematic and differentiated cells (Supporting Information Fig. S3c), in agreement with the
15 reported plasma membrane targeting of TPLATE (Van Damme *et al.*, 2011, Gadeyne *et al.*, 2014) in
16 non-dividing cells.

17 Our observation of over 50 *G. gigantea* infection units, intended as root areas underlying the sites of
18 hyphopodium contact, revealed an intense TPLATE-GFP labeling along a few anticlinal cell walls in
19 the inner root cortex of both *M. truncatula* (Fig. 3a,b) and *D. carota* ROCs (Fig. 3c; Supporting
20 Information Fig. S4a,b).

21 In particular, GFP-marked anticlinal cell walls were observed in the inner cortex of both plant species
22 as soon as hyphopodia developed on the root surface (Fig. 3a,c; Supporting Information Fig. S4a)
23 and until the first hyphae reached the inner cortex, about 2 days later (Fig. 3b; Supporting Information
24 Fig. S4b). Such GFP-labeled walls appeared to separate two square-shaped cells that corresponded in
25 shape to the previously described split cells (Fig. 1; Fig. 2). In fact, couples of split cells were
26 constantly observed in the colonized areas of the inner cortex of TPLATE-GFP-expressing ROCs
27 (Fig. 3d; Supporting Information Fig. S4b,c), although their walls were not as strongly labeled by
28 GFP in more advanced stages of root colonization (Supporting Information Fig. S4c) nor after
29 arbuscule development (Fig. 3d).

30 Comparable couples of square-shaped cells were only occasionally observed in the inner cortex of
31 corresponding control segments from uninoculated roots of the same age and size as the colonized
32 segments, (Supporting Information Fig. S4d), and their transverse wall was anyway never highlighted
33 by intense TPLATE-GFP accumulation. These observations were strongly supported by consecutive
34 imaging of a single site in an inoculated carrot root with a gap of 24 hours between each observation

1 (Fig. 3e-h). At the first time point (Fig 3e-g), hyphae developing from a single hyphopodium had
2 colonized the epidermis (Fig. 3e) and outer cortex (Fig. 3f). A deeper focal plane reaching the inner
3 cortex (Fig. 3g) revealed an intensely labeled cell plate being formed. When the same focal plane was
4 imaged 24 hours later (Fig 3h), the TPLATE-GFP signal had significantly faded from the mature
5 cross wall and - in the meantime - four additional inner cortical cells had divided in the same area.
6 On the same line, quantitative morphometric analyses performed on both *D. carota* and *M. truncatula*
7 confirmed a significant increase in the number of split cells in mycorrhizal compared to uninoculated
8 ROCs (Supporting Information Table S2).
9 Based on cell morphology and the analogous TPLATE-GFP labeling of recent cell walls in the root
10 meristem and in the colonized area of the mature cortex, we interpreted our observation as evidence
11 of an early, AM-dependent induction of cell division in the root inner cortex.

12

13 **Ectopic cortical cell division is not induced in *Mtdmi3-1* mutants**

14 Because cortical cell divisions were observed since the early steps of root colonization and often
15 occurred in cells that were not in direct contact with the fungus, we decided to investigate whether
16 this process was induced in the CSSP mutant *Mtdmi3-1*, where fungal colonization is arrested at the
17 epidermis level, with the development of extensive and branched hyphopodia (Catoira *et al.*, 2000;
18 Morandi *et al.*, 2005). By expressing the same TPLATE-GFP construct in this genotype, we first
19 confirmed construct functionality observing the expected GFP localization at newly formed cell walls
20 in the root tip meristem (Fig. 4a). Our careful observation of 8 independent hyphopodium-associated
21 root segments (Fig. 4b-e) did not reveal any fungal penetration of the root inner tissues and no
22 statistically significant change was observed in the number of split cells between uninoculated and
23 inoculated *dmi3-1* roots (Supporting Information Table S2). On this basis we concluded that the
24 activation of cortical cell divisions in *M. truncatula* is dependent on fungal presence inside the root -
25 which in turn requires DMI3-mediated signaling within the CSSP.

26

27 **Cell division and TPC-associated endocytic markers are upregulated in mycorrhizal roots**

28 Based on our observations of cell division induction and TPLATE involvement in AM fungal
29 accommodation processes, we used qRT-PCR to analyse the expression of cell division markers and
30 TPC-interacting proteins that act in endocytic processes associated with cell plate assembly.

31 Gene expression was analyzed in *M. truncatula* ROCs (Chabaud *et al.*, 2011; Genre *et al.*, 2013)
32 obtained from both wild type and *dmi3-1* plants (Genre *et al.*, 2005).

33 Two acknowledged cell division markers were chosen. *MtKNOLLE* encodes a syntaxin (homolog to
34 *AtSYP111*) exclusively localizing to cell-plate directed vesicles and essential for cytokinesis (*knolle*

1 mutants develop incomplete cell walls and multinucleated cells; Lukowitz *et al.*, 1996). Our second
2 cell division marker was *MtCYC1*, encoding a putative Cyclin-like F-box protein, previously reported
3 to be upregulated by 3,01 log₂ FC in arbusculated vs non-colonized cells (data from the *M. truncatula*
4 genome array performed on a population of root cortical cells colonized with *Rhizophagus irregularis*
5 and isolated through laser capture microdissection by Gaude *et al.*, 2012). Both transcripts were
6 significantly upregulated in mycorrhizal root samples compared to uninoculated roots in WT, while
7 no upregulation was recorded in *dmi3-1* mutants (Fig. 5). This result was consistent with our
8 hypothesis that cell division-related mechanisms are involved in AM colonization.

9 We also analysed the expression of four TPC-related endocytic markers. *MtTPLATE* and the alpha
10 subunit of the *Adaptor Protein complex 2 (MtAP2A1)* - both involved in clathrin-mediated
11 endocytosis (Gadeyne *et al.*, 2014; Di Rubbo *et al.*, 2013) - were constantly and significantly more
12 abundant in mycorrhizal root segments compared to controls in wild type *M. truncatula*, while their
13 expression did not change significantly in *dmi3-1* (Fig. 5). A weak but statistically significant
14 upregulation was observed for *Clathrin Heavy Chain 2 (MtCHC2)*, in mycorrhizal WT samples, while
15 this was not the case for *Clathrin Heavy Chain 1* (Fig. 5).

16 Lastly, as a marker of active AM colonization, we analyzed the expression of the AM-specific *Blue*
17 *Copper-binding Protein MtBCBP* (Parádi *et al.*, 2010) in the same samples. This confirmed that the
18 upregulation of our selected markers was strictly associated with *MtBCBP* expression (Supporting
19 Information Fig. S5). On this basis, we propose that cell division markers and actors of clathrin-
20 mediated endocytosis act during AM fungal accommodation, in line with our microscopic
21 observations.

22

23

1 **Discussion**

3 **The recruitment of cell division mechanisms in AM**

4 During symbiotic interface biogenesis, a novel membrane-bound compartment is assembled in the
5 middle of the cell, an area that is normally occupied by the vacuole in fully differentiated root cells.
6 Displacing the tonoplast is indeed among the first consequences of PPA-associated nuclear
7 movements in both epidermal and cortical cells (Genre *et al.*, 2005; 2008). As the PPA assembly
8 proceeds, the vacuole is further deformed (and reduced in volume), until a mass of cytoplasm (the
9 PPA aggregate) occupies the portion of the cell volume where the symbiotic interface and
10 intracellular hyphae are about to develop (Genre *et al.*, 2012). An analogous mechanism is
11 orchestrated when differentiated plant cells undergo mitosis: the vacuole has to be reduced - and
12 eventually split - so that the correct positioning of nucleus, mitotic spindle, phragmoplast and cell
13 plate can generate the cell division plane (Van Damme, 2009; Smertenko *et al.*, 2017). Also in this
14 case, initial vacuole remodeling is obtained by the coordinated movement of the nucleus and the
15 assembly of phragmosome cytoplasmic bridges that cross the vacuole along the equatorial plane and
16 eventually fuse into one central mass of cytoplasm encompassing the nucleus (Lloyd & Buschmann,
17 2007).

18 In addition to such similarities in cell dynamics, our results suggest a more direct relationship between
19 the two processes. The upregulation of several cell division markers in colonized root areas and
20 *TPLATE* expression pattern, that indicate an activation of gene expression in colonized cortical cells,
21 are both consistent with a reactivation of cell cycle-related processes. This is reinforced by the
22 recurring presence of split cells in the colonized region of roots from phylogenetically distant hosts
23 and the direct observation of dividing cortical cells in the vicinity of intraradical hyphae and
24 hyphopodia.

25 Altogether, we interpret our results as indicative of the reactivation of cell cycle-related processes in
26 the inner cortex since early root colonization. The biological meaning of this mechanism remains to
27 be fully clarified; nevertheless, literature on AM fungal accommodation provides intriguing clues:
28 the symbiotic interface materials have been described as very similar to the primordial wall found in
29 cell plates (Balestrini & Bonfante, 2014); a temporary, global re-routing of exocytic processes has
30 been proposed to take place in both cell plate (Lam *et al.*, 2008) and periarbuscular interface
31 deposition (Pumplin *et al.*, 2012); lastly, evidence of ploidy increases in mycorrhizal root systems
32 (Berta *et al.*, 2000; Lingua *et al.*, 2002; Fusconi *et al.*, 2005; Bainard *et al.*, 2011) suggests the onset
33 of endoreduplication (deriving from the arrest of the mitotic cycle) during AM colonization, even if
34 the precise location and timing of such endoreduplication events remains to be defined (Genre *et al.*,

1 2008). In the light of our current results, it is tempting to speculate that a mechanistic analogy links
2 the exocytic processes that lead to cell plate deposition and symbiotic interface biogenesis, even if
3 further investigation is needed to test this hypothesis.

4 5 **Membrane dynamics during fungal accommodation: a role for endocytosis?**

6 Our results on the upregulation of endocytic effectors such as TPLATE, CHC2 and AP2A1 during
7 AM fungal accommodation are suggestive of a role for clathrin-mediated endocytosis (CME)-related
8 processes in this symbiotic interaction.

9 Even if further investigation is needed to clarify this hypothesis, the involvement of CME
10 complements previous descriptions of the exocytic origin of the symbiotic interface in both AM and
11 nitrogen fixing symbiosis (Ivanov *et al.*, 2012; Genre *et al.*, 2012; Fournier *et al.*, 2015; Zhang *et al.*,
12 2015; Barker *et al.*, 2017): all major exocytic events in the plant cell are associated with endocytic
13 membrane recycling (Samaj *et al.*, 2004; Ketelaar *et al.*, 2008). Significantly, this also applies to cell
14 plate formation (Dhonukshe *et al.*, 2006; McMichael & Bednarek, 2013).

15 Indeed, CME has recently been shown to play a fundamental role in legume-rhizobium interactions:
16 a 60% reduction in the abundance of *L. japonicus Clathrin Heavy Chain 1 (LjCHC1)* transcripts (but
17 not *LjCHC2*) has been shown to strongly reduce rhizobial infection and nodule development (Wang
18 *et al.*, 2015). In the light of this finding, the consistent 30% upregulation of *M. truncatula CHC2* (but
19 not *MtCHC1*) during AM colonization opens the intriguing possibility that the two genes act in
20 alternative pathways controlling legume mycorrhization or nodulation, although more detailed
21 analyses going beyond the scope of the present research will be required to clarify this point.

22 23 **Evolutionary context**

24 The reactivation of the cell cycle is not unique to AM interactions, and has indeed been described
25 much earlier in legume-rhizobium interactions. The study by Xiao and colleagues (2014) provides
26 one of the most thorough descriptions of cell proliferation during nodule development, using
27 *Medicago truncatula* as a model system. In this legume, rhizobium inoculation triggers anticlinal
28 divisions that start in the pericycle and then extend to the endodermis, inner and middle cortical cell
29 layers, with the latter eventually generating the nodule meristem. The authors describe this process
30 as reminiscent of lateral root development, in line with the hypothesis of nodule evolutionary origins
31 from modified lateral roots. In the case of AM, however, cell cycle-related processes (including
32 marker gene regulation and the appearance of split cells) were restricted to a few, often sparse cells
33 in the inner cortical layer. This important difference indicates that the AM-related response is more
34 limited and also lacks the coordination observed in neo-organogenesis of both nodules (Xiao *et al.*,

1 2014) and lateral roots (Downie, 2014). On this basis, we cannot define a direct relationship between
2 the AM-dependent induction of cortical cell divisions and nodule initiation. Nevertheless, isolated
3 split or endoreduplicated cells are often observed in the vicinity of the infection thread, as it develops
4 across the root cortex. Such cell divisions are clearly unrelated to those that originate the nodule
5 primordium (Xiao *et al.*, 2014), and their observation led to the conclusion that pre-infection thread
6 assembly (a transvacuolar cytoplasmic aggregation that is remarkably similar to the PPA) could
7 involve reactivation of cell cycle (Yang *et al.*, 1994) and cell division-related mechanisms (Downie,
8 2014). Based on our current results we interpret the triggering of cortical cell divisions as a conserved
9 developmental trait in both symbioses.

10 The induction of ectopic cell divisions in roots colonized by the evolutionarily more ancient AM
11 symbiosis is particularly intriguing - also in the light of the frequent observation of couples of short
12 arbusculated cells in *Horneophyton ligneri* fossils from the Rhynie Chert (Strullu-Derrien *et al.*, 2018)
13 - and suggests that this response is not related to developmental innovations introduced by legumes
14 during their co-evolution with nitrogen-fixing bacteria, but rather a conserved strategy derived from
15 the 400 million year-old AM symbiosis.

16

17

18 **Acknowledgements**

19 We are grateful to New Phytologist reviewers and Editors for their constructive critics to the original
20 manuscript; to Mireille Chabaud, David Barker and Paola Bonfante for sharing ideas and suggestions;
21 to Beatrice Lace for starting this research with her preliminary experiments; to Paola Bonfante for
22 her support and critical revision of the text; to Christine Strullu-Derrien for highlighting
23 morphological analogies with fossil specimens; and to Dan Chamberlain for revising the English
24 language. We also thank Mara Novero and Antonella Faccio for their technical assistance during
25 sample preparation. Financial support for this research was granted by Compagnia di Sanpaolo
26 (project REPROGRAM - Progetti di Ateneo 2012, Call 01) and UNITO grant Ricerca Locale 2013-
27 2014-2017.

28

29

30 **Author contributions**

31 G.R. designed the experiments, developed the transgenic lines, performed microscopy, morphometric
32 and gene expression analyses and wrote the text. G.C. performed gene expression and morphometric
33 analyses, transgenic line production and contributed to the writing. V.F. performed gene expression
34 analyses and developed the *LjTPLATE:GUS* vector. M.C. developed the *LjTPLATE:GUS* vector.

1 V.V. performed gene expression analyses, transgenic line production and contributed to the writing.
2 D.V.D. provided the *AtTPATE-GFP* vector and contributed to the writing. A.G. designed the
3 research and experiments and wrote the text.

4

5

6 **References**

7

8 **Balestrini R, Bonfante P. 2014.** Cell wall remodeling in mycorrhizal symbiosis: a way towards
9 biotrophism. *Frontiers in Plant Science* **5**: 237.

10

11 **Bainard LD, Bainard JD, Newmaster SG, Klironomos JN. 2011.** Mycorrhizal symbiosis
12 stimulates endoreduplication in angiosperms. *Plant Cell Environment* **34**: 1577-1585.

13

14 **Barker DG, Chabaud M, Russo G, and Genre A. 2017.** Nuclear Ca²⁺ signalling in arbuscular
15 mycorrhizal and actinorhizal endosymbioses: on the trail of novel underground signals. *New*
16 *Phytologist* **214**: 533-538.

17

18 **Bécard G, Fortin JA. 1988.** Early events of vesicular-arbuscular mycorrhiza formation on Ri T-
19 DNA transformed roots. *New Phytologist* **108**: 211-218

20

21 **Berta G, Fusconi A, Sampò S, Lingua G, Perticone S, Repetto O. 2000.** Polyploidy in tomato
22 roots as affected by arbuscular mycorrhizal colonization. *Plant and Soil* **226**: 37-44.

23

24 **Boisson-Dernier A, Chabaud M, Garcia F, Bécard G, Rosenberg C, Barker DG. 2001.**
25 *Agrobacterium* rhizogenes-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing
26 and endomycorrhizal symbiotic associations. *Molecular Plant-Microbe Interactions*. **14**: 695-700.

27

28 **Bonfante P, Genre A. 2008.** Plants and arbuscular mycorrhizal fungi: an evolutionary-
29 developmental perspective. *Trends in Plant Science*. **13**: 492-498.

30

31 **Boruc J, Van Damme D. 2015.** Endomembrane trafficking overarching cell plate formation. *Current*
32 *Opinion Plant Biology*. **28**: 92-98.

33

- 1 **Boutté Y, et al. 2010.** Endocytosis restricts *Arabidopsis* KNOLLE syntaxin to the cell division plane
2 during late cytokinesis. *EMBO Journal* **29**: 546–558.
- 3
- 4 **Brady SM, Orlando DA, Lee JY, Wang JY, Koch J, Dinneny JR, Mace D, Ohler U, Benfey PN.**
5 **2007.** A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science* **318**:
6 801–806.
- 7
- 8 **Breakspear A, et al. 2014.** The Root Hair "Infectome" of *Medicago truncatula* Uncovers Changes
9 in Cell Cycle Genes and Reveals a Requirement for Auxin Signaling in Rhizobial Infection. *Plant*
10 *Cell* **25**: 4680-4701.
- 11
- 12 **Catoira R, Galera C, de Billy F, Penmetsa RV, Journet EP, Maillet F, Rosenberg C, Cook D,**
13 **Gough C, Dénarié J. 2000.** Four genes of *Medicago truncatula* controlling components of a Nod
14 factor transduction pathway. *Plant Cell* **12**: 1647–1665.
- 15
- 16 **Chabaud M, Venard C, Defaux-Petras A, Bécard G, Barker DG. 2002.** Targeted inoculation of
17 *Medicago truncatula* in vitro root cultures reveals *MtENOD11* expression during early stages of in-
18 fection by arbuscular mycorrhizal fungi. *New Phytologist*. **156**: 265–273.
- 19
- 20 **Chabaud M, Genre A, Sieberer BJ, Faccio A, Fournier J, Novero M, Barker D, Bonfante, P.**
21 **2011.** Arbuscular mycorrhizal hyphopodia and germinated spore exudates trigger Ca²⁺ spiking in the
22 legume and nonlegume root epidermis. *New Phytologist* **189**: 347-355.
- 23
- 24 **Delaux PM, et al. 2015.** Algal ancestor of land plants was preadapted for symbiosis. *Proceedings of*
25 *the National Academy of Sciences USA* **112**: 13390–13395.
- 26
- 27 **Dhonukshe, P., Baluska, F., Schlicht, M., Hlavacka, A., Samaj, J., Friml, J., Gadella, T.W. Jr.**
28 **2006.** Endocytosis of cell surface material mediates cell plate formation during plant cytokinesis.
29 *Developmental Cell* **10**: 137-150.
- 30
- 31 **Di Rubbo S, et al. 2013.** The clathrin adaptor complex AP-2 mediates endocytosis of brassinosteroid
32 insensitive1 in *Arabidopsis*. *Plant Cell* **25**: 2986–2997.
- 33
- 34 **Downie A. 2014.** Legume nodulation. *Current Biology*. **24**: R184-R190.

1
2 **Floss DS, Gomez SK, Park HJ, MacLean AM, Müller LM, Bhattarai KK, Lévesque-Tremblay**
3 **V, Maldonado-Mendoza IE, Harrison MJ. 2017.** A transcriptional program for arbuscule
4 degeneration during AM symbiosis is regulated by MYB1. *Current Biology*. **27**: 1206–1212.
5
6 **Fournier J, Timmers ACJ, Sieberer BJ, Jauneau A, Chabaud M, Barker DG. 2008.** Mechanism
7 of infection thread elongation in root hairs of *Medicago truncatula* and dynamic interplay with
8 associated rhizobial colonization. *Plant Physiol* **148**: 1985–1995.
9
10 **Fournier J, Teillet A, Chabaud M, Ivanov S, Genre A, Limpens E, de Carvalho-Niebel F,**
11 **Barker DG. 2015.** Remodeling of the infection chamber before infection thread formation reveals a
12 two-step mechanism for rhizobial entry into the host legume root hair. *Plant Physiology*. **167**: 1233-
13 1242.
14
15 **Fusconi A, Lingua G, Trotta A, Berta G. 2005** Effects of arbuscular mycorrhizal colonization and
16 phosphorus application on nuclear ploidy in *Allium porrum* plants. *Mycorrhiza* **15**: 313– 321.
17
18 **Gadeyne A, et al. 2014.** The TPLATE adaptor complex drives clathrin-mediated endocytosis in
19 plants. *Cell* **156**: 691-704.
20
21 **Gaude N, Bortfeld S, Duensing N, Lohse M, and Krajinski F. (2012).** Arbuscule-containing
22 and non-colonized cortical cells of mycorrhizal roots undergo extensive and specific reprogramming
23 during arbuscular mycorrhizal development. *Plant J.* **69**: 510–528.
24
25 **Genre A, Chabaud M, Timmers T, Bonfante P, Barker DG. 2005.** Arbuscular mycorrhizal fungi
26 elicit a novel intracellular apparatus in *Medicago truncatula* root epidermal cells before infection.
27 *Plant Cell* **17**: 3489–3499.
28
29 **Genre A, Chabaud M, Faccio A, Barker DG, Bonfante P. 2008.** Prepenetration apparatus assembly
30 precedes and predicts the colonization patterns of arbuscular mycorrhizal fungi within the root cortex
31 of both *Medicago truncatula* and *Daucus carota*. *Plant Cell* **20**: 1407–1420.
32

- 1 **Genre A, Ivanov S, Fendrych M, Faccio A, Zársky V, Bisseling T, Bonfante P. 2012.** Multiple
2 exocytotic markers accumulate at the sites of perifungal membrane biogenesis in arbuscular
3 mycorrhizas. *Plant Cell Physiology* **53**: 244–255.
- 4
- 5 **Genre A, Chabaud M, Balzergue C, Puech-Pagès V, Novero M, Rey T, Fournier J, Rochange**
6 **S, Bécard G, Bonfante P, et al. 2013.** Short-chain chitin oligomers from arbuscular mycorrhizal
7 fungi trigger nuclear Ca²⁺ spiking in *Medicago truncatula* roots and their production is enhanced by
8 strigolactone. *New Phytologist* **198**: 190–202.
- 9
- 10 **Gutjahr C, Parniske M. 2013.** Cell and developmental biology of arbuscular mycorrhiza symbiosis.
11 *Annu. Rev. Cell Developmental Biology* **29**: 593–617.
- 12
- 13 **Hewitt EJ. 1966.** Sand and Water Culture Methods Used in the Study of Plant Nutrition.
14 Commonwealth Agriculture Bureaux Farnham Royal, Bucks, U.K.
- 15
- 16 **Ivanov S, Fedorova EE, Limpens E, De Mita S, Genre A, Bonfante P, Bisseling T. 2012.**
17 Rhizobium-legume symbiosis shares an exocytotic pathway required for arbuscule formation.
18 *Proceedings of the National Academy of Sciences USA* **109**: 8316–8321.
- 19
- 20 **Ketelaar T, Galway ME, Mulder BM, Emons AMC. 2008.** Rates of exocytosis and endocytosis
21 in *Arabidopsis* root hairs and pollen tubes. *Journal of Microscopy* **231**: 265–273.
- 22
- 23 **Kistner C, Winzer T, Pitzschke A, Mulder L, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard**
24 **J, Webb KJ, et al. 2005.** Seven *Lotus japonicus* genes required for transcriptional reprogramming
25 of the root during fungal and bacterial symbiosis. *Plant Cell* **17**: 2217–2229.
- 26
- 27 **Kojima T, Saito K, Oba H, Yoshida Y, Terasawa J, Umehara Y, Suganuma N, Kawaguchi M,**
28 **Ohtomo, R. 2014.** Isolation and phenotypic characterization of *Lotus japonicus* mutants specifically
29 defective in arbuscular mycorrhizal formation. *Plant Cell Physiol* **55**: 928–941.
- 30
- 31 **Kormanik PP, McGraw, AE. 1982.** Quantification of vesicular-arbuscular mycorrhizae in plant
32 roots. In: Schenck, NC, eds. *Methods and principles of mycorrhizal research*. St. Paul: The American
33 Phytopathological Society, 37–45.
- 34

- 1 **Lam SK, Cai Y, Hillmer S, Robinson DG, Jiang L. 2008.** SCAMPs highlight the developing cell
2 plate during cytokinesis in tobacco BY-2 cells. *Plant Physiology* **147**: 1637-1645.
- 3
- 4 **Lee YR, Liu B. 2013.** The rise and fall of the phragmoplast microtubule array. *Current Opinion in*
5 *Plant Biology* **16**: 757–763.
- 6
- 7 **Lévy J et al. 2004.** A putative Ca²⁺ and calmodulin-dependent protein kinase required for bacterial
8 and fungal symbioses. *Science* **303**: 1361–1364.
- 9
- 10 **Liesche J, Schulz A, Krügel U, Grimm B, Kühn C. 2008.** Dimerization and endocytosis of the
11 sucrose transporter StSUT1 in mature sieve elements. *Plant Signaling and Behavior* **3**: 1136–1137.
- 12
- 13 **Limpens E, Mirabella R, Fedorova E, Franken C, Franssen H, Bisseling T, Geurts R. 2005.**
14 Formation of organelle-like N₂-fixing symbiosomes in legume root nodules is controlled by
15 DMI2. *Proceedings of the National Academy of Sciences USA* **102**:10375–10380.
- 16
- 17 **Lin K, et al. 2014.** Single nucleus genome sequencing reveals high similarity among nuclei of an
18 endomycorrhizal fungus. *PLoS Genetic* **10**: e1004078
- 19
- 20 **Lingua G, D’Agostino G, Massa N, Antosiano M, Berta G. 2002.** Mycorrhiza-induced differential
21 response to a yellows disease in tomato. *Mycorrhiza* **12**: 191–198.
- 22
- 23 **Lloyd C, Buschmann H. 2007.** Plant division: remembering where to build the wall. *Current Biology*
24 **17**: 1053–1055.
- 25
- 26 **Lukowitz W, Mayer U, Jurgens G. 1996.** Cytokinesis in the *Arabidopsis* embryo involves the
27 syntaxin-related *KNOLLE* gene product. *Cell* **84**: 61–71.
- 28
- 29 **McMichael CM, Bednarek SY. 2013.** Cytoskeletal and membrane dynamics during higher plant
30 cytokinesis. *New Phytologist* **197**: 1039–1057.
- 31
- 32 **Morandi D, Prado E, Sagan M, Duc G. 2005.** Characterisation of new symbiotic *Medicago*
33 *truncatula* (Gaertn.) mutants, and phenotypic or genotypic complementary information on previously
34 described mutants. *Mycorrhiza* **15**: 283–289.

1
2 **Novero M, Faccio A, Genre A, Stougaard J, Webb KJ, Mulder L, Parniske M, Bonfante P.**
3 **2002.** Dual requirement of the *LjSym4* gene for mycorrhizal development in epidermal and cortical
4 cells of *Lotus japonicus* roots. *New Phytologist* **154**: 741–749.
5
6 **Oldroyd GED. 2013.** Speak, friend, and enter: signalling systems that promote beneficial symbiotic
7 associations in plants. *Nature reviews Microbiology* **11**: 252–263.
8
9 **Parádi I, van Tuinen D, Morandi D, Ochatt S, Robert F, Jacas L, Dumas-Gaudot E. 2010.**
10 Transcription of two blue copper-binding protein isogenes is highly correlated with arbuscular
11 mycorrhizal development in *Medicago truncatula*. *Molecular Plant-Microbe Interactions* **23**: 1175–
12 1183.
13
14 **Parniske M. 2008.** Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nature Reviews*
15 *Microbiology* **6**: 763-775.
16
17 **Pumplin N, Harrison MJ. 2009.** Live-cell imaging reveals periarbuscular membrane domains and
18 organelle location in *Medicago truncatula* roots during arbuscular mycorrhizal symbiosis. *Plant*
19 *Physiology* **151**: 809–819.
20
21 **Pumplin N, Zhang X, Noar RD, Harrison MJ. 2012.** Polar localization of a symbiosis-specific
22 transporter is mediated by a transient reorientation of secretion. *Proceedings of the National Academy*
23 *of Sciences USA* **109**: E665-E672.
24
25 **Rasmussen R. 2001.** Quantification on the LightCycler instrument. In: Meuer S, Wittwer C,
26 Nakagawara K, eds. *Rapid Cycle Real-Time PCR: Methods and Applications*. Heidelberg: Springer-
27 Verlag, 21–34.
28
29 **Remy W, Taylor TN, Hass H, Kerp H. 1994.** Four hundred-million-year-old vesicular arbuscular
30 mycorrhizae. *Proceedings of the National Academy of Sciences USA* **91**: 11841–11843.
31
32 **Richter S, Kientz M, Brumm S, Nielsen ME, Park M, Gavidia R, Krause C, Voss U, Beckmann**
33 **H, Mayer U, et al. 2014.** Delivery of endocytosed proteins to the cell–division plane requires change
34 of pathway from recycling to secretion. *eLife* **3**:e02131.

- 1
- 2 **Sagan M, Morandi D, Tarengi E, Duc G. 1995.** Selection of nodulation and mycorrhizal mutants
3 in the model plant *Medicago truncatula* (Gaertn) after gamma-ray mutagenesis. *Plant Science* **111**:
4 63–71.
- 5
- 6 **Samaj J, Baluska F, Voigt B, Schlicht M, Volkmann D, Menzel D. 2004.** Endocytosis, actin
7 cytoskeleton, and signaling. *Plant Physiology* **135**:1150–1161.
- 8
- 9 **Segui-Simarro JM, Austin JR, White EA, Staehelin LA. 2004.** Electron tomographic analysis of
10 somatic cell plate formation in meristematic cells of *Arabidopsis* preserved by high-pressure freezing.
11 *Plant Cell* **16**: 836–856.
- 12
- 13 **Siciliano V, Genre A, Balestrini R, Cappellazzo G, DeWitt P, Bonfante P. 2007.** Transcriptome
14 analysis of arbuscular mycorrhizal roots during development of the prepenetration apparatus. *Plant*
15 *Physiol* **144**: 1455-1466.
- 16
- 17 **Smertenko A, Assaad F, Baluška F, Bezanilla M, Buschmann H, Drakakaki G, Hauser MT,**
18 **Janson M, Mineyuki Y, Moore I, et al. 2017.** Plant Cytokinesis: Terminology for Structures and
19 Processes. *Trends in Cell Biology* **27**: 885-894.
- 20
- 21 **Smith LG. 2001.** Plant cell division: building walls in the right places. *Nature Reviews Molecular*
22 *Cell Biology* **2**: 33–39.
- 23
- 24 **Stougaard J. 1995.** *Agrobacterium rhizogenes* as a vector for transforming higher plants: application
25 in *Lotus corniculatus* transformation. *Methods in Molecular Biology* **49**: 49–61.
- 26
- 27 **Strullu-Derrien C, Selosse MA, Kenrick P, Martin F. 2018.** The origin and evolution of
28 mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytologist*. doi:
29 10.1111/nph.15076
- 30
- 31 **Timmers, AC, Auriac MC, Truchet G. 1999.** Refined analysis of early symbiotic steps of the
32 *Rhizobium-Medicago* interaction in relationship with microtubular cytoskeleton rearrangements.
33 *Development* **126**: 3617-3628.
- 34

- 1 **Van Damme D, Bouget F-Y, Van Poucke K, Inzé D, Geelen D. 2004.** Molecular dissection of plant
2 cytokinesis and phragmoplast structure: A survey of GFP-tagged proteins. *Plant Journal* **40**: 386–
3 398.
- 4
- 5 **Van Damme D, Coutuer S, De Rycke R, Bouget FY, Inze D, Geelen D. 2006.** Somatic cytokinesis
6 and pollen maturation in *Arabidopsis* depend on TPLATE which has domains similar to coat proteins.
7 *Plant cell* **18**: 3502–3518.
- 8
- 9 **Van Damme D. 2009.** Division plane determination during plant somatic cytokinesis. *Current*
10 *Opinion in Plant Biology* **12**: 745–751.
- 11
- 12 **Van Damme D, Gadeyne A, Vanstraelen M, Inze D, Van Montagu MC, De Jaeger G, Russinova**
13 **E, Geelen D. 2011.** Adaptin-like protein TPLATE and clathrin recruitment during plant somatic
14 cytokinesis occurs via two distinct pathways. *Proceedings of the National Academy of Sciences USA*
15 **108**: 615–620.
- 16
- 17 **Wang C et al. 2015.** *Lotus japonicus* Clathrin Heavy Chain1 is associated with Rho-like GTPase
18 ROP6 and involved in nodule formation. *Plant Physiology* **167**: 1497–1510.
- 19
- 20 **Xiao TT, Schilderink S, Moling S, Deinum EE, Kondorosi E, Franssen H, Kulikova O, Niebel**
21 **A, Bisseling T. 2014.** Fate map of *Medicago truncatula* root nodules. *Development* **141**: 3517–3528.
- 22
- 23 **Yang WC, de Blank C, Meskiene I, Hirt H, Bakker J, van Kammen A, Franssen H, Bisseling**
24 **T. 1994.** *Rhizobium* nod factors reactivate the cell cycle during infection and nodule primordium
25 formation, but the cycle is only completed in primordium formation. *Plant Cell* **6**: 1415-1426.
- 26
- 27 **Zhang X, Pumplin N, Ivanov S, Harrison MJ. 2015.** EXO70I Is Required for Development of a
28 Sub-domain of the Periarbuscular Membrane during Arbuscular Mycorrhizal Symbiosis. *Curr Biol*
29 **17**: 2189-2195.
- 30
- 31 **Zipfel C, Oldroyd GED. 2017.** Plant signalling in symbiosis and immunity. *Nature* **543**: 328–336.
- 32

1 **Supporting Information**

2

3 **Table S1. Primers used in this work.**

4

5 **Table S2. AM colonization causes the appearance of split cells in the inner root cortex.**

6

7 **Fig. S1. *Lotus japonicus* TPLATE promoter sequence used for GUS analysis.**

8

9 **Fig. S2. Details of TPLATE expression pattern in uninfected roots of *L. japonicus* expressing the**
10 ***pLjTPLATE:GUS* construct.**

11

12 **Fig. S3. *At*TPLATE-GFP localization in uninfected *M. truncatula* ROCs.**

13

14 **Fig. S4. *At*TPLATE-GFP-labeled cell divisions in colonized carrot ROCs.**

15

16 **Fig. S5. Validation of sample mycorrhizal status by BCP expression analysis.**

17

18

19 **Figure legends**

20

21 **Fig. 1. TPLATE expression pattern in mycorrhizal roots of *L. japonicus*.** Following the GUS assay
22 reaction (blue), root samples were counterstained with acid fuchsin (purple) to outline *F. mosseae*
23 hyphae (hy) and arbuscules (ar). A young colonization unit is shown in (a) with limited intraradical
24 development from a single hyphopodium (hp) at 14 days post inoculation (dpi); (b) intense TPLATE
25 promoter activity is associated with arbuscules in the inner cortex. (c) Details of GUS-positive
26 staining in arbusculated cells. The largest hyphal branches are indicated by arrowheads. (d-f) GUS
27 staining of arbuscule-containing cells in the inner cortex at 14 dpi (c,d) and 21 dpi (e,f): a few
28 arbuscules have developed inside split cells (white dashed line). Bars = 100 μm in a; 20 μm in (b,d);
29 40 μm in (c,e,f).

30

31 **Fig. 2. Qualitative and quantitative analysis of cortical cell divisions in colonized roots of pot-**
32 **grown *L. japonicus* and *M. truncatula*.** (a-d) Representative images of untransformed *L. japonicus*
33 samples where the double staining of *F. mosseae* walls with WGA-FITC (green) and plant walls with
34 propidium iodide (red) indicated an increase in the number of split cells in AM colonized roots. (a,b)

1 Control uninoculated plants only occasionally displayed couples of cortical cells (white dashed line)
2 that were roughly half as long as the neighboring cells (asterisk). By contrast, split cells were frequent
3 in AM colonized roots (c-e), alongside cortical cells of regular size (asterisks). Four main conditions
4 were found: uncolonized split cells in the vicinity of arbuscules (c, left; d, right); split cells where one
5 cell contained an arbuscule and the second one did not (c, center); split cells containing arbuscules at
6 different stages of development (c, right; d, left); split cells containing arbuscules with comparable
7 morphology (e). Bars = 40 μ m (a, c); 30 μ m (b, d, e).

8 The table reports the results of our quantitative analysis of split cell occurrence in 375x375 μ m optical
9 sections (as those shown in a and c) from control and mycorrhizal plants. Two parameters were
10 considered: the average number of split cells per image (reported +/- the standard error) and the
11 number of images containing at least one couple of split cells. For *L. japonicus*, AM colonization
12 significantly increased the average number of split cells over 6 times (t test <0.01) as well as the
13 percent of images containing split cells (from 36% to 96%). An analogous situation was observed in
14 *M. truncatula*, where the average number of split cells per images increased over 4 times (T-test;
15 pvalue <0.01) and the percent of split cell-containing images passed from 36% in controls to 66% in
16 mycorrhizal roots. Overall these analyses correlate the presence of AM colonization with a consistent
17 increase in the number of split cortical cells.

18
19 **Fig. 3. *At*TPLATE-GFP highlights AM-induced cell divisions in the inner cortex.** All panels
20 show longitudinally oriented optical sections cutting through the inner cortex from living *M.*
21 *truncatula* (a,b) or *D. carota* (c-h) ROCs colonized by *G. gigantea*. In the presence of a hyphopodium
22 (hp) and penetrating hyphae (hy) in the epidermis (a,b), several cell divisions are visible in the *M.*
23 *truncatula* inner cortex, marked by the intense TPLATE-GFP fluorescence at anticlinal walls
24 (arrowheads); the faint and diffuse GFP fluorescence in the cytoplasm and on the plasma membrane
25 was obtained by increasing the microscope sensitivity to provide a general outline of root cell shapes;
26 asterisks mark neighboring undivided cortical cells. (c) shows an analogous situation in carrot, where
27 bright TPLATE-GFP labeling (arrowheads) highlights recent cell divisions in the inner cortex
28 underlying a hyphopodium; also in this case, the asterisk marks an undivided cortical cell. (d)
29 Following arbuscule development, the compact shape of several divided cells is recognizable
30 (arrowheads), but the lack of intense anticlinal wall labeling by TPLATE-GFP suggests that cell
31 divisions have not occurred in the preceding hours; one cell (arrow) appears to have divided twice,
32 as suggested by the extremely reduced size of two neighboring cells. Serial optical sections recorded
33 at increasing depth from the same carrot root are displayed in (e,f,g). A large hyphopodium is visible
34 on the root epidermis in (e). Several intracellular hyphae extend from the hyphopodium to the outer

1 cortex (f), where particularly intense accumulation of GFP signal is associated to a cortical PPA
2 (arrowheads). (g) Fluorescence of comparable intensity marks a forming cell wall (arrowhead) in the
3 inner cortex; white dashed lines mark long, undivided cortical cells. (h) displays the same focal plane
4 as (g) 24 hours later; the previously partial cell wall is now completed (arrowhead) and all the cells
5 marked by the dashed line have divided. Bars = 50µm.

6

7 **Fig. 4. *At*TPLATE-GFP localization in *M. truncatula dmi3-1* mutant ROCs.** Recent cell divisions
8 in the root tip meristem were marked by intense TPLATE-GFP accumulation (arrowheads), as shown
9 in (a). In contrast to our observations on WT ROCs, the presence of a *G. gigantea* hyphopodium (hp)
10 on the surface of a *dmi3-1* root (b) did not induce ectopic cell divisions in the underlying inner cortex
11 ((c), asterisk). An analogous example is shown in (d,e) where two optical sections from the same root
12 area are presented, one focused on the root epidermis (d) and one on the inner root cortex (e). Bars =
13 20 µm in (a); 40µm in (b-e).

14

15 **Fig. 5. Regulation of cell division and endocytic markers in wild type and *dmi3-1 M. truncatula***
16 **upon AM colonization.** The expression of the cell divisions markers *KNOLLE* and *CYC1*, alongside
17 *TPLATE* and TPLATE interactors in clathrin mediated endocytosis *AP2A1*, *CHC1* and *CHC2*, were
18 analyzed by quantitative RT-PCR 48 hours post hyphopodium development on ROC segments.
19 *TPLATE* and *AP2A1* transcripts were significantly upregulated in wild type inoculated root segments
20 (striped histograms) compared to controls (white histograms); no significant upregulation was
21 observed in *dmi3-1* mutants, where symbiotic signaling and intraradical colonization are blocked. A
22 differential regulation was observed for the two *Clathrin heavy chain* genes: while no change in gene
23 expression was observed for *CHC1*, a weak but statistically significant upregulation of *CHC2* was
24 recorded in inoculated wild type roots. Relative gene expression levels (fold change) are normalized
25 to the corresponding control; bars represent standard errors. Different letters indicate statistically
26 significant differences; T-test p value < 0.04; at least three biological replicates were used for each
27 condition.

28

29 **Table S1. Primers used in this work**

30

31 **Table S2. AM colonization causes the appearance of split cells in the inner root cortex.** Two
32 parameters were considered for quantitative analysis of split cell occurrence in 375x375 µm optical
33 sections: the average number of split cells per image (reported +/- the standard error) and the number
34 of images containing at least one couple of split cells. A marked increase in the number of split cells
35 was observed upon fungal colonization in both *D. carota* and wild type *M. truncatula* (T-test; p value

1 <0.01). By contrast, when the same analysis was done on *M. truncatula dmi3-1* mutants, no
2 significant change was observed.

3
4 **Figure S1. *Lotus japonicus* TPLATE promoter sequence used for GUS analysis.**

5
6 **Figure S2. TPLATE expression pattern in roots of *L. japonicus* expressing the *pLjTPLATE:GUS***
7 **construct.** (a) Intense blue GUS staining in a root tip meristem (asterisk). While GUS activity often
8 extended to the whole meristem, as shown in (b), alternative expression patterns were occasionally
9 observed (c, d), likely related to differences in meristematic activity: as shown here, *TPLATE*
10 promoter activity was limited to small groups of meristematic cells (arrowheads). (e) Higher
11 magnification of a root tip meristem, showing two pairs of GUS-positive meristematic cells; these
12 observations suggest that *pTPLATE* is upregulated in actively dividing cells. (f) GUS staining in the
13 central cylinder (cc) was much more prominent than in the other tissues of the differentiated zone,
14 where hardly any GUS staining could be detected during the reaction period, indicating much lower
15 expression levels than in the central cylinder. (g) 42 days post-inoculation with the AM fungus
16 *Funnelliformis mosseae*, GUS activity was evident in the colonized areas of the root cortex, where
17 arbuscules (ar) develop. (h, i) Transverse sections from control (h) and mycorrhizal (i) roots showing
18 GUS-staining in the central cylinder and arbuscule-containing cells. Bars = 200 μm in (a, f, g); 100
19 μm in (b, c, d); 30 μm in (e); 50 μm in (h, i).

20
21 **Figure S3. *AtTPLATE*-GFP localization in uninfected *M. truncatula* ROCs.** New-laid cell walls
22 are highlighted in the root apical meristem (a,b) by an intense GFP signal (arrowheads). The nuclei
23 (n) of four recently-divided meristematic cells in (a) are recognizable as dark areas in the faintly
24 fluorescent cytoplasm and allow an easy identification of relative cell positions. A deep longitudinal
25 section from an uninfected root is shown in (c); root cortex layers are recognizable between the
26 epidermis (ep) and central cylinder (cc): the lack of intense labeling in any anticlinal wall and the
27 regularity of cell sizes indicate that no recent cell division has occurred. Bars = 20 μm in (a) and (b);
28 50 μm in (c).

29
30 **Figure S4. *AtTPLATE*-GFP-labeled cell divisions in colonized carrot ROCs.** Longitudinal optical
31 sections cutting through the inner root cortex. (a) Intense GFP fluorescence (arrowheads) labels a few
32 anticlinal walls in the inner cortical layers underlying a hyphopodium (hp); such cell walls divide
33 regular cells in two smaller daughter cells. (b) (c) Couples of split cells (dashed line) and *AtTPLATE*-

1 GFP marked walls (arrowheads) are visible in the vicinity of an intraradical hypha (h) reaching the
2 inner cortex. (d) No divided cells are recognizable in uninoculated roots. Bars = 50µm.

3

4 **Figure S5. Validation of sample mycorrhizal status by *BCBP* expression analysis.** The
5 statistically significant upregulation of *Blue copper-binding protein (BCBP)* gene - an acknowledged
6 plant marker for AM colonization - confirmed AM establishment in the samples used for gene
7 expression analyses. Relative gene expression levels (fold change) are normalized to the control.
8 Different letters indicate statistically significant differences; T-test p value < 0.04.