

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Counting Roman chickens: Multidisciplinary approaches to human-chicken interactions in Roman Britain

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1689349> since 2019-02-04T10:42:40Z

Published version:

DOI:10.1016/j.jasrep.2017.09.013

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Manuscript Details

| | |
|--------------------------|--|
| Manuscript number | JASREP_2017_199 |
| Title | Counting Roman Chickens: Multidisciplinary Approaches to Human-Chicken Interactions in Roman Britain |
| Short title | Counting Chickens in Roman Britain |
| Article type | Research Paper |

Abstract

This paper discusses some of the approaches and results from two multi-disciplinary projects. The first is the AHRC-funded 'Cultural and Scientific Perceptions of Human-Chicken Interactions' Project. This is investigating the history of the exploitation of chickens in Europe. The second is the Leverhulme Trust-funded Rural Settlement of Roman Britain Project, which has collated evidence from excavation reports from thousands of sites. This paper updates the evidence for the exploitation of chickens in Roman Britain, showing that there were significant variations in the abundance of chicken bones found on different types of settlement. There was also a modest increase in their abundance during the Roman period suggesting chickens became slightly more frequent contributors to the diet, albeit still only a rare commodity. However, they continued to be frequently represented in graves, shrines and other ritual deposits. The paper also discusses evidence of egg production and avian osteopetrosis, demonstrating that when traditional zooarchaeological research is integrated with scientific analyses, a deeper understanding of past human diet can be acquired.

| | |
|---|--|
| Keywords | Zooarchaeology; Chickens; Roman Britain; Eggshell; Medullary bone; Pathology |
| Corresponding Author | Julia Best |
| Corresponding Author's Institution | Bournemouth University |
| Order of Authors | Mark Maltby, Martyn Allen, Julia Best, B. Tyr Fothergill, Beatrice Demarchi |
| Suggested reviewers | Dale Serjeantson, Tony King, Pam Crabtree, James Morris, Bea De Cupere |

Submission Files Included in this PDF

File Name [File Type]

Cover Letter Maltby et al..docx [Cover Letter]

Roman Chicken Manuscript MM MA JB TF BD.pdf [Manuscript File]

Table 1 Roman Chicken Manuscript MM MA JB TF BD.pdf [Table]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Dear Sir/Madam,

Please find attached the manuscript entitled: "Counting Roman Chickens: Multidisciplinary Approaches to Human-Chicken Interactions in Roman Britain" for consideration in the following special issue: "Interdisciplinary approaches to ancient Roman diets".

Yours Sincerely,

Mark Maltby and Julia Best

Counting Roman Chickens: Multidisciplinary Approaches to Human-Chicken Interactions in Roman Britain

Mark Maltby^a, Martyn Allen^b, Julia Best^a, B. Tyr Fothergill^c, Beatrice Demarchi^{d,e}

^aDepartment of Archaeology, Anthropology and Forensic Science, Bournemouth University, Talbot Campus, Poole

^b Department of Archaeology, School of Archaeology, Geography and Environmental Science, University of Reading

^c Department of Archaeology and Ancient History, University of Leicester

^d Department of Archaeology, University of York

^e Department of Life Sciences and Systems Biology, University of Turin, Italy

Keywords: Zooarchaeology; Chickens; Roman Britain; Eggshell; Medullary bone; Pathology

Abstract: This paper discusses some of the approaches and results from two multidisciplinary projects. The first is the AHRC-funded ‘Cultural and Scientific Perceptions of Human-Chicken Interactions’ Project. This is investigating the history of the exploitation of chickens in Europe. The second is the Leverhulme Trust-funded Rural Settlement of Roman Britain Project, which has collated evidence from excavation reports from thousands of sites. This paper updates the evidence for the exploitation of chickens in Roman Britain, showing that there were significant variations in the abundance of chicken bones found on different types of settlement. There was also a modest increase in their abundance during the Roman period suggesting chickens became slightly more frequent contributors to the diet, albeit still only a rare commodity. However, they continued to be frequently represented in graves, shrines and other ritual deposits. The paper also discusses evidence of egg production and avian osteopetrosis, demonstrating that when traditional zooarchaeological research is integrated with scientific analyses, a deeper understanding of past human diet can be acquired.

1. Introduction

The history of the domestication and westward spread of the chicken or domestic fowl (*Gallus gallus domesticus*) out of Asia is currently the focus of much debate (Xiang et al. 2014, 2015; Perry-Gal et al. 2015; Peters et al. 2015; Eda et al. 2016; Pitt et al. 2016). However, the species does not appear to have spread across Europe prior to the late prehistoric period. The earliest confirmed record for the presence of chickens in Britain is

currently from the site of White Horse Stone in Kent where a femur provided a radiocarbon date of 770-390 cal BC with modelled dates of 560-390 cal BC (Kitch 2006). However, chicken bones are rare finds in the pre-Roman period in Britain, being recorded in only around 30% of the Iron Age faunal assemblages from southern England, nearly always in very small numbers (Hambleton 2008). Only on a few Late Iron Age (100BC-AD43) sites in the south-east of England, where continental contact was more evident, did chickens appear in larger numbers (Maltby 1997; Hambleton 2008), despite the fact that images of chickens were depicted on coins minted in two areas of southern England during that period (Best et al 2016). Indeed, the regular occurrence of partial or complete skeletons of chickens along with Julius Caesar's frequently quoted, albeit enigmatic, observation from *De Bello Gallico* that the Britons kept chickens but did not eat them, has led to the very plausible contention that chickens were initially valued for some of their other qualities rather than for food (Sykes 2012).

Despite its recent introduction and continued presence in contexts associated with human burials and other ritual sites (King 2005) chickens are often summarily dismissed in zooarchaeological reports of Romano-British assemblages merely as an unremarkable addition to the diet. A previous survey (Maltby 1997) indicated that there is some evidence to suggest that chickens became more abundant during the Romano-British period but the potential complexity of production, distribution and consumption of chickens and their products in the diet was not fully explored. This potentially undervalues their impact, and their dismissal limits our understanding of their multiple roles. Two recent large multi-disciplinary research projects have provided opportunities to review the evidence for human-chicken relationships in more depth. The Arts and Humanities Research Council-funded 'Cultural and Scientific Perceptions of Human-Chicken Interactions' Project has brought together over 20 researchers from six universities to examine the social, cultural and environmental impact of chickens in Europe. This research has included the collation of zooarchaeological data from both published works and unpublished archives from all periods including the Roman era. In addition, innovative research has been carried out (inter alia) in analyses of metrical data, pathology, ancient DNA, stable isotopes, pottery residues, eggshells, ecology, material culture and anthropology associated with chickens. Meanwhile, the Leverhulme Trust-funded 'Rural Settlement of Roman Britain Project' has collated evidence from over 2,500 excavated rural settlements from England and Wales, enabling a comprehensive reassessment of the countryside of Roman Britain to be carried out (Smith et

al 2016). Over 1,600 sites have produced animal bones and counts of the bones of chickens and other species can be accessed via the wide-ranging online resource created by the project (Allen et al 2015; Allen 2017 in press).

This paper will examine whether there is evidence for an increase in importance of chickens as a source of food in Roman Britain, and whether there are variations in its abundance in different types of site and periods. It will also consider some other analyses that can be used in considering the evolution of the relationships between humans and chickens in the western provinces of the Roman Empire.

2. Chicken abundance in Romano-British zooarchaeological samples:

An initial survey into variability in the abundance of chickens from Romano-British archaeological sites was carried out by Maltby (1997). The sample consisted of 123 assemblages from 68 sites and compared data from military sites, major towns, nucleated settlements, villas and other rural settlements. Results suggested that chickens tended to be more common in assemblages from military sites and major towns, but the numbers of assemblages from some types of site rendered these conclusions tentative and precluded investigation of possible chronological variations. During the last 20 years, the number of assemblages has increased enormously principally due to the large number of developer-funded excavations that have been carried out since 1990 both on rural (Allen 2017 in press) and urban sites (Maltby 2015), thus enabling a much more comprehensive survey to be undertaken.

2.1 Materials and methods

This survey will focus on comparing the abundance of chicken bones with those of sheep/goat. Some comparisons with the abundance of pigs will also be made. Inter-site comparisons of species abundance are faced with a series of well-known challenges concerning differential identification, retrieval, preservation, quantification and deposition. With particular regard to chickens, it is not possible to distinguish all chicken bones from those of other galliforms such as pheasant (*Phasianus colchicus*) and guineafowl (*Numida melagris*) via morphological and metrical analysis, but in Roman assemblages where such distinctions have been made, nearly all the diagnostic bones have been positively identified as chicken. It is therefore assumed that the vast majority, if not all, of the galliform bones

recorded on these sites belonged to chickens. Retrieval and preservation biases have long been recognised, and bones from small birds have a greater likelihood of being destroyed or overlooked than the generally larger and more robust bones of mammal species during hand-excavation. Unfortunately, many reports do not separate or list the bones recovered by sieving. However, the great majority of the assemblages are derived mainly or totally from hand-collection and, with caution, can be compared. It is impossible, however, to fully assess whether all assemblages were recovered with the same level of efficiency. Where known, assemblages derived mainly from sieving have been excluded. Obviously sheep and pigs are larger than chickens and there will still inevitably be some bias in recovery standards, but these will not be as marked as they would be in comparisons with larger mammals such as cattle and horse. Quantification methods used by zooarchaeologists also vary. Most counts are derived from the total number of identified specimens (NISP). However, what constitutes a NISP count varies significantly. Some counts include vertebrae and ribs, whilst others do not; some zooarchaeologists count all identifiable limb bone fragments; others count only a selected suite of diagnostic elements. Another issue concerns the inclusion or exclusion of bones from partial or complete skeletons in the counts. Where known in this survey, counts exclude associated groups of bones but this was not feasible in every case. It is also quite common for urban sites, in particular, to include assemblages dominated by waste accumulated by the large-scale butchery of cattle, (Maltby 2015), which is another reason why cattle have been excluded from this survey. To minimise problems created by small samples, a minimum NISP count of 50 sheep/goat and chicken elements was set.

Data for the rural settlements, including nucleated sites, were obtained from the Roman Rural project website (Allen et al 2015). Data for the assemblages from the major urban sites were obtained from Maltby (2010a, 276) and supplemented by data obtained from more recently reported assemblages. Data from military sites were gathered from unpublished and published reports.

2.2 Farmsteads and Villages

Rural settlements were split into categories of farmsteads, villages, villas and roadside settlements based on the definitions set out by the Roman Rural Project. Many of the farmsteads could be further subdivided into unenclosed, enclosed or complex categories (Smith et al 2016). Over 67% of the 436 assemblages from farmsteads produced either no chicken bones at all or <1% of the total number of sheep/goat and chicken elements (Figure

1). A further 26% had <5% chicken. Of the few assemblages with unusually high percentages of chicken (>15%) most had extenuating reasons to explain why they were unusually well represented (Table 1). In several cases, most or all of the chicken bones accompanied human burials; in others, they were derived from single contexts and were probably part of associated bone groups (ABGs, Morris 2010). In one case they came from a site with evidence of industrial processing and specialist butchery, more commonly encountered on larger nucleated sites where chicken bones have often more commonly recovered.




Figure 1: Percentage of chicken of total sheep/goat and chicken NISP counts from farmsteads (n=436)

Table 1: Rural assemblages with high percentages of chicken bones. Data derived from Allen et al. (2015)

Thirty-two assemblages came from sites categorised by the Roman Rural Project as villages (Smith et al. 2016). Of these, 18 (56%) contained <1% chicken and 10 (31%) 1%-5% chicken of the total sheep/goat and chicken NISP counts. Three contained between 6% and 10% chicken and only one, a very small assemblage from Abingdon, Oxfordshire, produced an

assemblage with over 15% chicken (Table 1). Generally, however, chicken bones were very uncommon components of faunal assemblages from all types of farmsteads and villages.

2.3 Villas

Most assemblages from villas also produced few chicken bones. In 33% of the 79 assemblages, chickens contributed <1% of the total number of sheep/goat and chicken elements (Figure 2). However, chicken bones did quite commonly form higher percentages in villa assemblages, providing 1%-5% of sheep/goat and chicken elements in 34% of the assemblages and between 6%-10% in a further 18%. However, in only six cases did chickens provide over 20% of the sheep/goat and chicken elements (Table 1). Unsurprisingly, these included an assemblage from the spectacular Fishbourne Palace in West Sussex, a site which also produced exceptionally high percentages in the earlier Late Iron Age and Flavian deposits and continued to produce quite large quantities in the later Roman period (Allen 2011). In two cases (Bancroft and Yarford), percentages of chicken bones increased significantly from assemblages that accumulated prior to the construction of the villas. The Castle Copse (Wiltshire) assemblage was the only one to produce more chicken than sheep/goat bones. This was partly due to their increased abundance in sieved deposits, but the assemblage was also remarkable for the dominance of pig bones (Payne 1997). None of these six assemblages had evidence for biases created by the presence of associated bone groups. There is therefore some evidence that chickens made a significantly greater contribution to the diet at some high status villa sites.




Figure 2: Percentage of chicken of total sheep/goat and chicken NISP counts from villas (n=79)

2.4 Roadside settlements

These produced results similar to those obtained from villas (Figure 3). In 40% of the 115 assemblages, chickens provided <1% of the total number of sheep/goat and chicken elements, and in a further 37% of the assemblages this figure lay between 1% and 5%. Chicken bones contributed 6%-10% in a further 11% of the assemblages. In only six assemblages did chickens provide over 15% of the sheep/goat and chicken elements (Table 1). Of these, the assemblage from Skeleton Green (Ashdown and Evans 1981) is better characterised as a late Iron Age oppidum displaying significant evidence of continental influence. It also produced unusually large percentages of pig bones (Maltby 1997; Hambleton 2008). The two assemblages from Staines, Surrey, are from a settlement where several excavations have revealed evidence that indicates that the settlement had many urban characteristics, including dumps of specialist butchery waste (Chapman 1984; 2010). The same case could be argued for the settlements of Elms Farm, Heybridge, Essex (Johnstone and Albarella 2002) and Shadwell, Greater London (Douglas et al. 2011).




Figure 3: Percentage of chicken of total sheep/goat and chicken NISP counts from roadside settlements (n=115)

2.5. Chronological Variations.

Rural assemblages (n=587) were sub-divided where possible into five broad periods ranging from the late Iron Age through to the late Roman period (Figure 4). These confirmed that the great majority had <1% chicken in the total sheep/goat NISP counts. However, the percentage of assemblages in this category fell in each period from >90% in the Late Iron Age down to 43% in the Late Roman period. Assemblages with 1%-5% chicken increased from 7% in the late Iron Age sample to over 30% in the early Roman and later periods. Assemblages with 6%-10% chicken bones formed over 8% of the early Roman sample, rising to over 13% in the assemblages from the late Roman period. Chickens gradually became a more consistent, albeit still minor component, of rural assemblages.




Figure 4: Percentage of chicken of total sheep/goat and chicken NISP counts from rural settlements by period (n=587)

2.6. Urban assemblages

A total of 91 assemblages were obtained from 16 *civitas* capitals and *colonia* from Britain. These showed a marked contrast with those from rural settlements (Figure 5). None of the assemblages produced <1% chicken of the total sheep/goat and chicken NISP counts and only 13% fell into the second lowest category (1%-5%). In contrast, 58% of the assemblages included >15% chicken and the mode (21%) lay between 16%-20% chicken. Most of these counts excluded bones in associated bone groups and bones from sieved assemblages were not included. Although urban sites tend to produce better preserved assemblages than those from rural settlements, it is very unlikely that this could account for all of the urban-rural contrasts. Put simply, people living in towns were much more likely to eat chickens than those living in the countryside. There is abundant butchery evidence (Figure 6) that supports the increased use of chickens for meat in both the urban context, such as Exeter (e.g. Coles in press). Similar evidence has been found on some rural sites including Fishbourne (Allen 2011, 223) and Shefford, Bedfordshire (Maltby 2010b).




Figure 5: Percentage of chicken of total sheep/goat and chicken NISP counts from urban settlements (n=91)



Figure 6: Chicken tibiotarsus from Exeter Princesshay showing diagonal knife-cuts on the distal condyles characteristic of disarticulating the lower leg (Photo J. Best).

The contrast between urban and rural chicken abundance can be seen at a regional level, as demonstrated by comparing sites from within the *civitas* capital of Cirencester and rural sites in the local hinterland (Figure 7). This is not to say that the pattern is totally consistent. Sites

from Winchester have consistently produced assemblages in the 1%-5% chicken category, whereas those from Dorchester, Exeter and Caerwent have nearly all produced over 15% chickens (Maltby 2010a). The fact that most of the Winchester assemblages are from extra-mural sites, whereas most of the assemblages from the other towns are from sites from central areas of the towns may be significant, perhaps reflecting socio-cultural variations of diet in different areas of the towns.




Figure 7: Percentage of chicken of total sheep/goat and chicken NISP counts from sites in Cirencester (checked pattern) and its hinterland (grey)

King (1984) observed that pigs often are more prominent in more Romanised settlements in Britain. This updated review generally supports this interpretation, with assemblages from both villas and towns that had higher percentages of chickens to sheep/goat also having higher percentages of pig in relation to sheep/goat, although there is substantial variation (Figure 8).




Figure 8: Comparisons of chicken/sheep and pig/sheep ratios in (a) urban (n=91) and (b) villa (n=63) assemblages in Britain

2.7 Military Sites

Excluding vici, 30 assemblages from military sites were considered (Figure 9). Nine (30%) of these fell within the 1%-5% chicken bracket but a similar number produced >15% chicken. A wide range of variability is to be expected as this category covers a wide range of sites ranging from large fortresses to small auxiliary forts in different parts and periods in Roman Britain. However, the tendency was for chickens to be better represented than on rural settlements, but not as consistently as well represented as in towns. There are also indications that chicken meat may have been more available to high-ranking officers at the supply fortress at South Shields (Stokes 2000) and the legionary fort in Caerleon (Hamilton-Dyer 1993). At the latter, chicken bones were particularly prominent in the drains of the baths, (O'Connor 1986), indicating that chickens were commonly eaten by the bathers.




Figure 9: Percentage of chicken of total sheep/goat and chicken NISP counts from military sites (n=30)

2.8 Religious and Burial Sites and other Depositions

King (2005) demonstrated that chickens were sometimes very well represented at temples and shrines in Roman Britain. The best known example comes from Uley, Gloucestershire, where goat and chickens were sacrificed in large numbers at a temple dedicated to Mercury (Levitan 1993; Brothwell 1997). Substantial amounts of chicken bones have also been

reported from other temple sites at Brigstock, Northamptonshire, and Folly Lane, St Albans, Hertfordshire (King 2005). The highest percentage of chickens (87%) from the 91 urban assemblages discussed above came from near the Temple of Mithras in London (Macready and Sidell 1998). Continental examples are also well known, including amongst many others, the temple associated with Mithras at Tienen, Belgium (Lentacker et al. 2003a; 2003b) and the temple at Carnuntum–Mühläcker, Austria dedicated to Jupiter (Gál and Kunst 2010). It should be noted, however that by no means every temple and shrine has evidence of votive offerings of chickens, even where the sacrifice of other animals is prominent (King 2005). On the other hand, in Roman Britain chicken bones have quite commonly been found in association with inhumations and cremations in both urban and rural cemeteries, showing that they had multiple roles, including food for the dead and votive offerings (Morris 2011). As discussed above, they quite often feature much more prominently in grave deposits than in other contexts.

3. The exploitation of chicken eggs

When considering chickens in Roman diet, it is also important to recognise the secondary products that they can provide, particularly eggs. Chicken eggs become increasingly prominent as food items in Roman and Roman-influenced contexts. Their production and use can be traced by integrating multiple lines of evidence and analytical techniques including historical sources, archaeological eggshell and medullary bone.

3.1 Documentary evidence

Documentary sources can provide insights into productivity, use and trade. On Hadrian's Wall, tablets from the fort of Vindolanda written in the 1st and 2nd centuries AD indicate that eggs as well as chickens were valuable resources to be acquired:

"... bruised beans, two modii, **chickens, twenty**, a hundred apples, if you can find nice ones, **a hundred or two hundred eggs, if they are for sale there at a fair price**. ... 8 sextarii of fish-sauce ... a modius of olives ... (Back) To ... slave (?) of Verecundus" (Tablet 302, Translation: Bowman and Thomas 1983).

This particular statement does not indicate specifically that these were chicken eggs, but given the reference to chickens earlier in the tablet, it is a fair assumption to make, particularly given the quantity requested. No eggshell has yet been recovered from

excavations at Vindolanda, and whilst this may result from preservation conditions, it could be that eggs were not locally available. The desire to obtain them as a special order probably reflects their high value.

Columella's *Res Rustica* is one of several agricultural works that provide instructions for the care of egg-laying chickens, including housing requirements and modifying feed to make hens lay sooner, more often and with larger eggs. He also describes aspects of productivity and preservation, such as transferring eggs for hatching to capable broody hens, and using chaff, bran and salt for egg storage. Columella and other ancient sources such as Varro even suggest that certain types of chicken, including those with five toes, were the best for egg laying and brooding.

Although rare, recipes can demonstrate how eggs could contribute to diet. Apicius' *De Re Coquinaria* a collection of recipes, compiled in the late 4th or early 5th AD, shows that they had a wide range of culinary uses, including clarifying muddy wine, and an ingredient in brain sausages and many sauces. Of course, it is unknown how widespread these recipes and agricultural guides were practised beyond Italy, as documentary sources are often limited in applicability by being restricted in period and place.

3.2 Eggshell

Eggshell is found fairly regularly on archaeological sites, although thorough soil processing is generally needed for its recovery. Eggshells were recorded on 38 sites collated by the Romano-British rural settlement project (Allen et al. 2015), although rarely were the eggshells further identified. Eggshell can be identified to species via microscopy (Sidell 1993), although this has significant limitations (Best et al. in prep), and more recently by ZooMS (Zooarchaeology by Mass Spectrometry) which identifies taxa-specific peptide mass markers (Demarchi et al. 2016; Stewart et al. 2013; Presslee et al. in prep.). These two methods can be combined: using ZooMS for species identification and microscopy to identify the stage of chick development within the egg (since the developing chick takes calcium from the eggshell to aid bone formation, causing changes to the interior surface of the eggshell) (Beacham and Durand 2007; Best et al. in prep).

One of the first archaeological eggshell assemblages to be analysed using both techniques came from the military amphitheatre at Chester, Cheshire, where substantial amounts of eggshell were found. The bulk of this material came from two deposits: a well-stratified early assemblage from AD70-80, which correlates with the first phase of amphitheatre use, and a

second dating to AD100 from substantial deposits underneath the seating banks (Wilmott pers. comm.). The ZooMS results indicate that all analysed fragments were from chicken eggs (a representative ZooMS spectrum is shown in Figure 10). Microscopy revealed that *c.*90% of the analysed fragments from the AD100 deposits showed no signs of reabsorption associated with chick development. Therefore almost all of the eggs were freshly laid, halted early in their incubation sequence, or infertile. In this instance the assemblage appears to represent food consumed by spectators watching events at the amphitheatre. Such snacks foods may have been on sale outside the amphitheatre as appears to be depicted in a fresco of the Pompeii amphitheatre (Ellis 2004). This evidence suggests that chicken eggs were traded from a relatively early period of Roman occupation in Britain, at least on military and associated sites.

The eggshells from the AD70-80 phase at the Chester amphitheatre, whilst all identified as chickens, had more varied stages of development, potentially indicating that not all of the eggs were consumed fresh.




Figure 10: Annotated mass spectrum (ZooMS) of Chester Amphitheatre AD70-80 chicken eggshell from context 625. Species and family specific markers are represented by star symbols (following Presslee et al. in prep.).

3.3 Medullary Bone

The analysis of medullary bone, a calcium deposit for egg production laid down on the endosteal surface of the medullary cavity, is a useful method for identifying the presence of laying hens in the archaeological record (van Neer et al. 2002, 129-132). It can be used to give an indirect insight into breeding and egg production on sites where eggshell is not recovered. It can be identified by macroscopic assessment of fragmented bones. However, by employing non-destructive x-ray analysis its presence or absence can also be determined for complete bones. This combined approach allows broad sex profiles to be identified for whole assemblages (Best in prep.). For example, no eggshell was available for identification at Fishbourne Palace, but observations of medullary bone in the fragmented bone assemblage (Allen 2011) indicated that laying hens were present at the site, either as live birds or dead meat resources. The femur is the best element for examining medullary bone in chickens since the fill is most substantial and enduring in this bone. X-ray analysis of the Fishbourne assemblage increased the overall recorded occurrence of medullary bone from 8% to 20% of the femora. The majority of the deposits only occupied a small proportion of the bone cavity, perhaps indicating that these birds were killed for meat at the end of lay, either permanently or temporarily as a result of moulting, illness or dietary deficiencies. This suggests that these birds were kept for egg production, with meat being a secondary consideration. The hens at Fishbourne may have been kept on site, but the possibility that some were traded from elsewhere, such as the nearby town of Chichester, should not be ruled out.

Absence of medullary bone can also be valuable for profiling the birds that were contributing to Roman diet and life. Bones without medullary deposits can belong to males, but also to females not in lay, or with no deposit in that specific skeletal element. At the temple site of Uley, medullary bone was scarce. When combined with spur evidence and metrics, these data support the interpretation that a large proportion of the birds sacrificed were male (Brothwell 1997; Fothergill and Best in prep.). These birds would probably have been consumed in multiple ways: as meat, but also psychologically and metaphorically as spiritual offerings. A similar pattern can also be seen on the continent at sites such as Tienen in Belgium where over 7,600 chicken bones were found, representing at least 238 individuals (155 adults; 83 subadults) which were deposited in a pit after what appears to have been a single very large ritual feasting event (Lentacker *et al.* 2004a, 77-81; 2004b). This site was associated with the god Mithras, who in turn was often connected with the cockerel. Again, several lines of evidence indicate that these birds were primarily males and no medullary bone was identified

in the fragmented material or in x-rayed whole bones. This demonstrates that several elements of ritual consumption in the Roman world are found in multiple geographic locations.

4. Pathology

One palaeopathological hallmark of Roman-era avian bone assemblages is the presence of avian osteopetrosis, a pathology which is routinely identified in material from sites across Europe. These lesions are caused by a range of avian leucosis viruses, spread through contact as well as from hen to chick and through genomic transmission (Pruková et al., 2007). Avian osteopetrosis lesions are distinctive in appearance, consisting of hypermineralised endosteal and periosteal new bone formation in the diaphyses of affected elements (Figure 11), which can be differentially diagnosed through radiography (O'Connor and O'Connor 2005). Avian leucosis viruses affect various species of domestic poultry and cause a number of detrimental physical and behavioural symptoms which negatively impact vivacity, egg-laying, and weight gain (Holmes, 1961; Payne 1992; Uzunova et al. 2014; Vogt, 1977).



Figure 11: Tibiotarsus with avian osteopetrosis lesions from Uley, shown with a modern comparative element

Although it is possible that avian leucosis viruses affected poultry flocks in earlier periods (particularly as infection does not always result in bony lesion formation), the earliest archaeological evidence of avian osteopetrosis originates from Tiberian contexts at Roman military sites: the fort and naval base at Velsen in the Netherlands and the fort at Aulnay in France (Prummel, 1987; Lignereux and Peters, 1997). The Roman assemblage from Carlisle (Old Grapes Lane), also dating to the 1st century AD, contained two elements described as osteopetrotic (Allison 2010). The proportional frequency of avian osteopetrosis lesions identified in archaeological assemblages increases in the first and second centuries AD, and the initial geographic spread of avian leucosis viruses is likely to be linked to the movement of people and their animals around the Empire (Fothergill in press). Since animal husbandry plays a key role in pathogenesis, it is possible that Roman chicken-keeping methods and the environments in which these birds were kept fostered the transmission of avian leucosis viruses. These husbandry techniques have a direct link to human diet in terms of the quantity and quality of chicken resources available. These data also provide insights into how the diet-related cycle of production, distribution and consumption affected many aspects of animal health and avian-human interactions.

5. Discussion

Although there is evidence that the consumption of chicken meat and eggs increased during the Romano-British period, they were still nevertheless a rare commodity. The zooarchaeological data has shown that meat supply was heavily dependent upon the provision of beef, particularly in towns (King 1999; Hesse 2011; Maltby 2015). This is supported by lipid residue analysis. In Silchester, for example, most residues were composed of ruminant fats (Marshall et al 2008; Colonese et al. in press). In Britain, chicken meat and eggs would have been regarded as luxury foods obtained from an exotic, recently introduced, species. It is no surprise that they were consumed more readily on settlements where Roman and other continental influences were more prominent, reflecting the greater cultural and culinary diversity of the inhabitants. The greater dominance of chicken in Romano-British urban deposits is mirrored in other parts of the western Roman Empire, including northern France (Lepetz 1996) and Switzerland (Groot and Deschler-Erb 2015), as well as across much of North Africa (Fothergill and Sterry in press; Fothergill et al. in press). Given their special status combined with their convenient small size, it is understandable that chickens continued

to be sacrificed as votive offerings, linked with a number of deities and buried with humans even on settlements where they were probably rarely eaten. The supply of chickens may sometimes have been challenging, as indicated by the Vindolanda tablet and this challenge would have been heightened by the need to supply birds for sacrifice some of the temple sites. It is also likely that many chickens were raised in towns, where there was, at least initially, a greater demand for their products. Bones of very young chicks have been found in Winchester, Hampshire indicating at least some of the birds were being bred in the town (Maltby 2010a). The appearance of avian osteopetrosis lesions may also be linked to keeping chickens in more confined environments (Fothergill in press).

However, whilst all strands of evidence examined here indicate that the Roman period in Britain sees an increased use of chicken meat and eggs for food, these animals continued to hold multiple other roles within society and culture; from deity companions to luxury goods. Therefore, whilst frequently the archaeology of chickens, and particularly their zooarchaeological record, is seen primarily in terms of diet, this is not the only avian-human interaction that needs to be considered.

As such, this integrated approach incorporating traditional zooarchaeological methods alongside historical sources and a suite of scientific analyses shows that the investigation of avian demography can provide insights into their complex relationships with humans and resultantly inform upon and beyond human diet.

Acknowledgements:

This work was supported by the Arts and Humanities Research Council as part of the project “Cultural and Scientific Perspectives of Human-Chicken Interactions” (Grant No AH/L006979/1) and by the Leverhulme Trust as part of the project “The Rural Settlement of Roman Britain”. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors also wish to thank the following people for their help and collaboration: Tony Wilmott, Alice Forward, Robert Symmons, the Intergemeentelijke Archeologische Dienst PORTIVA, Anton Ervynck, An Lentacker, Wim van Neer and Charlotte Coles. We are very grateful to Samantha Presslee and Matthew Collins for providing information and discussion in relation to ZooMS.

References:

- Allen M., 2006 The animal bones from Yarford Villa, Somerset. University of Winchester: Unpublished report to the Southern Quantocks Archaeological Survey.
- Allen, M., 2011. *Animalscapes and Empire: New Perspectives on the Iron Age/Romano-British Transition*. University of Nottingham PhD Thesis.
- Allen, M., 2017 (in press). Pastoral farming. In M. Allen, L. Lodwick, T. Brindle, M. Fulford and A. Smith, *New Visions of the Countryside of Roman Britain, Vol. 2: The Rural Economy of Roman Britain*. London: Britannia Monograph Series.
- Allen, M., Brick, N., Brindle, T., Fulford, M., Holbrook, N., Richards, J.D., and Smith, A., 2015. *The Rural Settlement of Roman Britain: an Online Resource*. data-set]. York: Archaeology Data Service [distributor] <https://doi.org/10.5284/1030449>
- Allison, E.P., 2010. The bird bones. In M. McCarthy (ed.), *The South Lanes, Carlisle: Specialist Fascicules*. (Oxford Archaeology North/English Heritage) York: Archaeology Data Service [distributor], Fascicule 1, 161-162. <https://doi.org/10.5284/1000182>
- Hamilton-Dyer, S., 2008. The animal bones. In G. Anelay, *Liss Roman Villa, Liss, Hampshire: Report on the Archaeological Excavations*. Liss Archaeological Group.
- Ashdown, R., and Evans, C., 1981. The animal bones. In C. Partridge, *Skeleton Green: a Late Iron Age and Romano-British Site*. London: Britannia Monograph 2, 205-237.
- Atkins, A., Popescu, E., Rees, G., and Stansbie, D. 2014. *Broughton, Milton Keynes, Buckinghamshire: The Evolution of a South Midlands Landscape*. Oxford: Archaeology Monograph 22.
- Barker, B., Luke M and Wells, J., 2006. *A421 Improvements: M1 Junction 13 to Bedford. Archaeological Field Evaluation*. Bedford: Albion Archaeology <https://doi.org/10.5284/1010377>
- Beacham, E.B. and Durand, S.R., 2007 Eggshell and the archaeological record: new insights into turkey husbandry in the American Southwest. *Journal of Archaeological Science*, 34: 1610–1621. <https://dx.doi.org/10.1016/j.jas.2006.11.015>
- Best, J., In prep. Non-destructive medullary bone analysis: exploring advantages and limitations of x-ray analysis.

520 Best, J., Demarchi, B., Presslee, S., Collins, M., and Maltby, M. In prep. Integrated
521 application of ZooMS and microscopy in the study of archaeological eggshell.

522 Best, J., Fieder, M. and Pitts, J., 2016 Introducing chickens - arrival, uptake and use in
523 prehistoric Britain. *Past: the Newsletter of the Prehistoric Society* 84, 1-3.

524 Bowman, A.K. and Thomas J.D. 1983. *Vindolanda: the Latin writing-tablets*. London:
525 Britannia Monograph 4.

526 Branigan, K., 1971. *The Roman Villa at Latimer*. Chess Valley Archaeology and History
527 Society.

528 Brothwell, D., 1997, Interpreting the immature chicken bones from the Romano-British ritual
529 complex on West Hill, Uley. *International Journal of Osteoarchaeology* 4, 330-332.

530 Bullock, A., and Allen, M., 1997. Animal bones (Flagstones). In R. Smith, F. Healy, M.
531 Allen, E. Morris, I. Barnes and P. Woodward, *Excavations along the Route of the Dorchester*
532 *By-pass, Dorset, 1986-8*. Salisbury: Wessex Archaeology Monograph 11, 191-193.

533 Chapman, J., 1984. Animal and human bone. In K. Crouch and S. Shanks, *Excavations at*
534 *Staines 1975-76: The Friends' Burial Ground Site*. London: London and Middlesex
535 Archaeological Society /Sussex Archaeological Society Joint Publication No 2, 115-123.

536 Chapman, J., 2010. Animal bone, In P. Jones, *Roman and Medieval Staines. The*
537 *Development of the Town*. Woking: SpoilHeap Monograph 2, 169–171.

538 Colonese, A.C., Lucquin, A., Timby, J. and Craig, O., In press. Organic residue analysis of
539 ceramic vessels. In M. Fulford, A Clarke, E. Durham and N. Pankhurst, *Late Iron Age*
540 *Calleva: the Pre-conquest Occupation at Silchester Insula IX*. London: Society for the
541 Promotion of Roman Studies Britannia Monograph.

542 Coles, C. In press. The Animal Bones of Roman Princesshay. In J. Allan Roman archaeology
543 of Princesshay, Exeter. *Devon Archaeological Society proceedings*.

544 Demarchi, B., Hall, S., Roncal-Herrero, T., Freeman, C.L., Woolley, J., Crisp, M.K., Wilson,
545 J., Fotakis, A., Fischer, R., Kessler, B.M., Rakownikow Jersie-Christensen, R., Olsen, J.V.,
546 Haile, J., Thomas, J., Marean, C.W., Parkington, J., Presslee, S., Lee-Thorp, J., Ditchfield, P.,
547 Hamilton, J.F., Ward, M.W., Wang, C.M., Shaw, M.D., Harrison, T., Domínguez-Rodrigo,
548 M., MacPhee, R.D.E., Kwekason, A., Ecker, M., Kolska Horwitz, L., Chazan, M., Kröger,
549 R., Thomas-Oates, J., Harding, J.H., Cappellini, E., Penkman, K., and Collins, M.J. 2016.

550 Protein sequences bound to mineral surfaces persist into deep time. *eLife* [online] 5, e17092.
551 <https://dx.doi.org/10.7554/eLife.17092>

552 Dobney, K. and Jaques, S.D., 1996. Animal bones. In R. J. Williams, P.J. Hart and A.T.L.
553 Williams, *Wavendon Gate. A late Iron Age and Roman Settlement in Milton Keynes*
554 *Buckinghamshire Archaeological Society Monograph Series* 10, 203-230.

555 Douglas, A., Gerrard, J., and Sudds, B., 2011. *A Roman settlement and Bath House at*
556 *Shadwell: excavations at Tobacco Dock and Babe Ruth Restaurant, The Highway, London.*
557 London: Pre-Construct Archaeology Monograph 12.

558 Eda, M., Lu, P., Kikuchi, H., Li, Z., Li, F. and Yuan, J., 2016. Reevaluation of early
559 Holocene chicken domestication in northern China. *Journal of Archaeological Science* 67,
560 25-31.

561 Ellis, S., 2004. The distribution of bars at Pompeii: archaeological, spatial and viewshed
562 analyses. *Journal of Roman Studies* 17, 371-384.

563 Fothergill, B.T., In Press. Human-aided movement of viral disease and the archaeology of
564 avian osteopetrosis. *International Journal of Osteoarchaeology*.

565 Fothergill, B.T., Linseele, V., and Valenzuela-Lamas, S., In Press. Movement and
566 Management of Animals in the North and West of Africa from 1000 BC to AD 1000. In D.J.
567 Mattingly, A. Cuenod and C. Duckworth, eds. *Trans-Saharan: Mobile Technologies*.
568 Cambridge University Press.

569 Fothergill, B.T., and Sterry, M., In Press. Pouligraphy and “Poultrymen” in North Africa. In
570 Proceedings of XIe Colloque international Histoire et Archéologie de l’Afrique du Nord,
571 *Hommes et animaux au Maghreb, de la Préhistoire au Moyen Age: explorations d’une*
572 *relation complexe*. Marseille et Aix-en-Provence.

573 Fothergill, B.T., and Best, J. In Prep. Hens, health and husbandry: new approaches to past
574 poultry-keeping in England.

575 Gál, E., and Kunst, G., 2010. Offered to gods, eaten by people: bird bones from the Sanctuary
576 of Jupiter Heliopolitanus in Carnuntum–Mühlacker (Austria). *International Journal of*
577 *Osteoarchaeology* 24, 336-346.

578 Groot, M., and Deschler-Erb, S., 2015. Market strategies in the Roman provinces: different
579 animal husbandry systems explored by a comparative regional approach. *Journal of*
580 *Archaeological Science Reports* 4, 447-460.

581 Hambleton E., 2008. *Review of middle Bronze Age - late Iron Age Faunal Assemblages from*
582 *Southern Britain*. Portsmouth: English Heritage Research Department Report Series number
583 71-2008.

584 Hamilton, J., 2000. Animal bones. In: B. Cunliffe and C. Poole (eds.), *The Danebury*
585 *Environs Programme: the Prehistory of a Wessex Landscape. Vol. 2, Part 6, Houghton*
586 *Down, Stockbridge, Hants, 1994*. Oxford: University of Oxford, Committee for
587 Archaeology, 67-73.

588 Hamilton-Dyer, S., 1993. The animal bones, in V. Zienkiewicz, *Excavations in the Scamnum*
589 *Tribunorum* at Caerleon. *Britannia* 24, 132-136.

590 Harman M., 1994, Mammal and bird bones. In R.S. Leary (ed.), *Excavations at the Romano-*
591 *British Settlement at Pasture Lodge Farm, Long Bennington, Lincolnshire, 1975-77 by H.M.*
592 *Wheeler*. Nottingham: Occasional Papers in Lincolnshire History and Archaeology 10
593 (Nottingham, Trent and Peak Archaeological Trust), 49-53.

594 Halkon, P., Millett, M., and Woodhouse, H., 2015. *Hayton, East Yorkshire: Archaeological*
595 *Studies of the Iron Age and Roman Landscapes Volumes 1-2* (Yorkshire Archaeological
596 Report 7). Leeds: Yorkshire Archaeological Society.

597 Hesse, R., 2011. Reconsidering animal husbandry and diet in the northwest provinces”,
598 *Journal of Roman Archaeology* 24, 215-248.

599 Higbee, L. 2004. Mammal, bird and fish bone. In R. Regan, C. Enans and G. Webley, *The*
600 *camp Ground Excavations: Colne Fen, Earith Assessment Report*. Cambridge: Cambridge
601 Archaeological Unit Report 164, 160-200. <https://doi.org/10.5284/1021819>

602 Holmes, J.R., 1961. Postmortem findings in avian osteopetrosis. *Journal of Comparative*
603 *Pathology and Therapeutics* 71, 20-27.

604 Horton, W., Lucas, G., and Wait, G.A., 1994, Excavation of a Roman site near Wimpole,
605 Cambs., 1989. *The Proceedings of the Cambridge Antiquarian Society* 83, 31-74

606 Johnstone, C., and Albarella, U., 2002, *The late Iron Age and Romano-British Mammal and*
607 *Bird Bones Assemblage from Elms Farm, Heybridge, Essex*. Portsmouth: English Heritage
608 Ancient Monuments Laboratory Report 45/02.

609 King, A., 1984. Animal bones and the dietary identity of military and civilian groups in
610 Roman Britain, Germany and Gaul. In T. Blagg and A. King (eds.), *Military and Civilian in*
611 *Roman Britain: Cultural Relationships in a Frontier Province*. Oxford: British
612 Archaeological Reports (British Series) 136, 187-218.

613 King, A., 1999. Diet in the Roman world: a regional inter-site comparison of the animal
614 bones. *Journal of Roman Studies* 12, 168-202.

615 King, A., 2005. Animal remains from temples in Roman Britain. *Britannia* 36, 329-369.

616 Kitch, J., 2006. *Animal Bone from White Horse Stone, Aylesford, Kent*. CTRL Specialist
617 Report Series.

618 Lentacker A., Ervynck, A., and Van Neer, W., 2004a. Gastronomy or religion? The animal
619 remains from the Mithraeum at Tienen (Belgium). In S. O'Day, W. Van Neer and A. Ervynck
620 (eds.), *Behaviour Behind Bones: The zooarchaeology of ritual, religion, status and identity*.
621 Oxford: Oxbow, 77-94.

622 Lentacker, A., Ervynck, A., and Van Neer, W., 2004b. The symbolic meaning of the cock.
623 The animal remains from the Mithraeum at Tienen (Belgium). In M. Martens and G. De Boe
624 (eds.), *Roman Mithraism: the Evidence of the Small Finds*. Brussels: Archeologie in
625 Vlaanderen Monografie 4, 57-80.

626 Lepetz, S., 1996. L'animal dans la société gallo-romaine de la France du nord. *Revue*
627 *archéologique de Picardie* Numéro spécial 12.

628 Levitan, B., 1993. The vertebrate remains. In A. Woodward and P. Leach, *The Uley Shrines:*
629 *Excavation of a Ritual Complex on West Hill, Uley, Gloucestershire*. London: English
630 Heritage Archaeological Report 17, 257-345.

631 Levitan B., 1994. Vertebrate remains from the villa. In R.J. Williams and R. Zeepvat (eds.),
632 *Bancroft: A late Bronze Age/Iron Age Settlement, Roman Villa and Temple-Mausoleum:*
633 *Volume II, Finds and Environmental Evidence*. Aylesbury: Buckinghamshire Archaeology
634 Monograph Series 7, 515-536.

635 Lignereux, Y., Peters, J., Tassaux, F., and Tronche, P., 1997. Viandes, volailles et fruits de
636 mer a la table des légions Romaines d'Aunedomnacum, 20-30 après Jésus-Christ (Aulnay-de-
637 Saintonge, Charente-Maritime). *Revue de Médecine Vétérinaire* 148 (5), 399-412.

638 Macready, S., and Sidell, J., 1998: *The animal bones*. In J. Shepherd, *The Temple of Mithras*,
639 *London: Excavations by W.F. Grimes and A. Williams at the Walbrook*, London: English
640 Heritage Archaeological Report 12, 208-215.

641 Magilton, J., 2006. A Romano-Celtic temple and settlement at Grimstock Hill, Coleshill,
642 Warwickshire. *Birmingham and Warwickshire Archaeological Society Transactions* 110, 1-
643 236.

644 Maltby, M., 1997. Domestic fowl on Romano-British sites: inter-site comparisons of
645 abundance. *International Journal of Osteoarchaeology* 7, 402-414.

646 Maltby, M., 2010a. *Feeding a Roman Town: Environmental Evidence from Excavations in*
647 *Winchester, 1972-1985*. Winchester: Winchester Museums Service.

648 Maltby, M., 2010b. Animal bone. In M. Luke, T. Preece and J. Wells, A Romano-British
649 aisled building and associated settlement south of Ampthill Road, Shefford. *Bedfordshire*
650 *Archaeology* 26, 317-320.

651 Maltby, M., 2015. Commercial archaeology, zooarchaeology and the study of Romano-
652 British towns. In M. Fulford and N. Holbrook (eds.), *The Towns of Roman Britain: the*
653 *Contribution of Commercial Archaeology since 1990*. London: Britannia Monograph, 175-
654 193.

655 Marshall, L.-J., Cook, S. R., Almond, M. J., and Fulford, M. G., 2008. Roman diet and trade:
656 evidence from organic residues on pottery sherds recovered at the Roman town of Calleva
657 Atrebatum (Silchester Hants.). *Britannia* 39, 245-254.

658 Maynard, D., Cleary, R., Moore, R., Brooks, I., and Price. J., 1997. Excavations at Foxton,
659 Cambridgeshire 1994. In J. Price and D. Maynard (eds.), *The Archaeology of the St Neots to*
660 *Duxford Gas Pipeline 1995* (British Archaeological Reports (British Series) 255). Oxford:
661 Archaeopress, 21-39.

662 Morris, J., 2010. Associated bone groups; beyond the Iron Age. In J. Morris and M. Maltby
663 (eds.), *Integrating Social and Environmental Archaeologies; Reconsidering Deposition*.
664 (British Archaeological Reports (International Series) 2077) Oxford: Archaeopress, 12-23.

665 Morris, J., 2011. *Investigating Animal Burials: Ritual, Mundane and Beyond*. British
666 Archaeological Reports 535) Oxford: Archaeopress.

667 Morrison, A., 2000. The animal bone. In N. Cooper (ed.), *The Archaeology of Rutland Water:
668 Excavations at Empingham in the Gwash Valley, Rutland, 1967-73 and 1990*. Leicester:
669 Leicester Archaeology Monographs 6, 132-136.

670 O'Connor, T. 1986. The animal bones. In J. D. Zienkiewicz, *The Legionary Fortress Baths at
671 Caerleon: Volume II. The Finds*. Cardiff: National Museum of Wales/CADW, 225-248.

672 O'Connor, T., and O'Connor, S., 2005. Digitising and image-processing radiographs to
673 enhance interpretation in avian palaeopathology. In G. Grupe and J. Peters (eds.),
674 *Documenta Archaeobiologiae 3: Feathers, Grit and Symbolism. Birds and Humans in the
675 Ancient Old and New Worlds*. Rahden: Verlag Marie Leidorf GmbH, 69-82.

676 Payne, L.N., 1992. Biology of avian retroviruses. In J.A. Levy (ed.), *The Retroviridae, Vol. 1*.
677 New York: Plenum Press, 299-404.

678 Payne, S., 1997. Animal remains. In E. Hostetter and T. Howe (eds.), *The Romano-British
679 Villa at Castle Copse, Great Bedwyn*. Indianapolis: Indiana University Press, 322-330.

680 Perry-Gal, L., Erlich, A., Gilboa, A., and Bar-Oz, G., 2015. Earliest economic exploitation of
681 chicken outside East Asia: Evidence from the Hellenistic Southern Levant. *Proceedings of
682 the National Academy of Sciences* 112 (32), 9849-9854.

683 Peters, J., Lebrasseur, O., Best, J., Miller, H., Fothergill, T., Dobney, K., Thomas, R., Maltby,
684 M., Sykes, N., Hanotte, O., O'Connor, T., Collins, M., and Larson, G., 2015. Questioning
685 new answers regarding Holocene chicken domestication in China. *Proceedings of the
686 National Academy of Sciences* 112 (19), E2415-E2415.

687 Pitt, J., Gillingham, P., Maltby, M., and Stewart, J., 2016. New perspectives on the ecology of
688 early domestic fowl: an interdisciplinary approach. *Journal of Archaeological Science* 74, 1-
689 10.

690 Presslee, S., Wilson J., Russell D., Fischer, R., Kessler, B., Best, J., Radini, A., Collins, M.,
691 and Demarchi, B., In Review. Identifying taxon-specific peptide markers for archaeological
692 eggshell: a new proteomics approach. *Science and Technology of Archaeological Research*.

693 Pruková, D., Vernerová, Z., Pilčík, T., Stepanets, V., Indrová, M., Geryk, J., Plachý, J.,
 694 Hejnar, J., and Svoboda, J., 2007. Differences in pathogenicity among strains of the same or
 695 different avian leukosis virus subgroups. *Avian Pathology* 36 (1), 15-27.

696 Prummel, W., 1987. Poultry and fowling at the roman castellum Velsen I. *Palaeohistoria* 29,
 697 183-201.

698 Sidell, J., 1993. *A Methodology for the Identification of Archaeological Eggshell*.
 699 Philadelphia, PA: The University Museum of Archaeology and Anthropology.

700 Smith, A., Allen, A., Brindle, T., and Fulford, M., 2016. *The Rural Settlement of Roman*
 701 *Britain*. London: Britannia Monograph 29.

702 Stokes, P., 2000. A cut above the rest? Officers and men at South Shields Roman fort. in P.
 703 Rowley-Conwy (ed.), *Animal Bones, Human Societies*. Oxford: Oxbow, 146-151.

704 Stewart, J.R.M., Allen, R.B., Jones, A.K.G., Penkman, K.E.H., Collins, M.J., 2013. ZooMS:
 705 making eggshell visible in the archaeological record. *Journal of Archaeological Science* 40,
 706 1797–1804.

707 Sykes, N., 2012. A social perspective on the introduction of exotic animals: the case of the
 708 chicken. *World Archaeology* 44 (1), 158-169.

709 Uzunova, K., Stamatova-Yovcheva, K., Dimova, V., Yovchev, D., and Halil, M., 2014.
 710 Anatomical and ethological changes in poultry affected by osteopetrosis. *Scientific Papers on*
 711 *Animal Science and Biotechnologies* 47 (1), 188-191.

712 Vogt, P.K., 1977. Genetics of RNA tumor viruses. In H. Fraenkel-Conrat and R.R. Wagner
 713 (eds.), *Comprehensive Virology, Vol. 9: Regulation and Genetics, Genetics of Animal*
 714 *Viruses*. New York and London: Plenum Press, 341-455.

715 Wilson, B., 1993. The animal bones. In Anon, *Abingdon Vineyard Area 3. Summary Report*
 716 *of Excavations and Statement of Potential*. Oxford: Oxford Archaeological Unit, 12-13.
 717 <https://doi.org/10.5284/1028217>

718 Wood, M., 2006. *Archaeological Assessment Report on Land at Wygate Park, Spalding,*
 719 *Lincolnshire. Volume 2*. Sleaford: Archaeological Project Services.
 720 <https://doi.org/10.5284/1013596>

721 Xiang, H., Gao, J., Yu, B., Hofreiter, M. and Zhao, X., 2015. Reply to Peters et al.: further
722 discussions confirm early Holocene chicken domestication in northern China. *Proceedings of*
723 *the National Academy of Sciences* 112 (19), E2416-E2416.

724 Xiang, H., Gao, J., Yu, B., Zhou, H., Cai, D., Zhang, Y., Chen, X., Wang, X., Hofreiter, M.
725 and Zhao, X., 2014. Early Holocene chicken domestication in northern China. *Proceedings of*
726 *the National Academy of Sciences* 111 (49), 17564-17569.

727 van Neer, W, Noyen, K and de Cupere, B 2002. On the use of endosteal layers and medullary
728 bone from domestic fowl in archaeozoological studies. *Journal of Archaeological Science*
729 29(2), 123–134. <https://doi.org/10.1006/jasc.2001.0696>

| Region | Site | Type | Assemblage Date | NISP | S/G | Chicken | %Chicken | Comments and original source |
|---------------|--------------------------------------|--------------------------|----------------------------|------|-----|---------|----------|---|
| Central Belt | Broughton Manor Farm | unenclosed farmstead | 1st C BC-mid 1st C AD | 97 | 78 | 19 | 19.59 | Chicken bones from cremations (Atkins et al 2014) |
| Central Belt | Wavendon Gate, Milton Keynes | enclosed farmstead | 1st C BC/AD | 209 | 171 | 38 | 18.18 | Chicken bones from cremations (Dobney and Jaques 1996) |
| Central Belt | Pasture Lodge Farm, Long Bennington | farmstead (unclassified) | 3rd-4th C AD | 412 | 342 | 70 | 16.99 | Includes chicken ABG (Harman 1994) |
| Central Belt | Woolram Wygate, Spalding | farmstead (unclassified) | 3rd-4th C AD | 72 | 55 | 17 | 23.61 | Includes chicken ABG (Wood 2006) |
| South | Maiden Castle Road | farmstead (unclassified) | 1st-4th C AD | 224 | 186 | 38 | 16.96 | Chicken bones from inhumation (Bullock and Allen 1997) |
| East Anglia | Foxton | complex farmstead | 1st-4th C AD | 366 | 297 | 69 | 18.85 | Chicken bones from inhumation (Maynard et al 1997) |
| Central Belt | Empingham | enclosed farmstead | 3rd-4th C AD | 273 | 221 | 52 | 19.05 | Most chicken bones from a well (Morrison 2000) |
| South | St Georges Road, Dorchester By-pass | field system | 3rd-4th C AD | 135 | 106 | 29 | 21.48 | Chicken bones all from one pit (Bullock and Allen 1997) |
| Central Belt | Brogborough Hill (A421 Site 2) | complex farmstead | 2nd-3rd C AD | 60 | 34 | 26 | 43.33 | All chicken bones from one oven (Barker et al 2006) |
| Central Belt | Langdale Hale, Earith, Colne Fen | complex farmstead | 2nd-3rd C AD | 250 | 182 | 68 | 27.20 | Site includes specialist butchery deposits (Higbee 2004) |
| West Midlands | Grimstock Hill, Coleshill | enclosed farmstead | 1st-2nd C AD | 84 | 64 | 20 | 23.81 | All chicken bones from one context (Magilton 2006) |
| North-East | Burnby Lane, Hayton | farmstead (unclassified) | 3rd-4th C AD | 185 | 131 | 54 | 29.19 | Many bones from well (Halkon et al. 2017) |
| Central Belt | Abingdon, The Vineyard | village | 1st-4th C AD | 50 | 42 | 8 | 16.00 | (Wilson 1993) |
| South | Fishbourne | palace | 1st-2nd C AD | 1035 | 797 | 238 | 23.00 | 34% in 1st C BC/AD deposits; 15% n 3rd-4th C AD (Allen 2011) |
| Central Belt | Latimer | villa | 2nd-3rd C AD | 61 | 43 | 18 | 29.51 | (Branigan 1971) |
| Central Belt | Bancroft | villa | 2nd-3rd C AD | 111 | 74 | 37 | 33.33 | 1% in 1st-2nd C AD; 5% in 3rd-4th C AD (Levitan 1994) |
| South | Liss | villa | 3rd-4th C AD | 115 | 75 | 40 | 34.78 | (Hamilton-Dyer 2008) |
| Central Belt | Yarford, Kingston St Mary | villa | 3rd-4th C AD | 291 | 220 | 71 | 24.40 | 7% in 1st C BC/AD farmstead (Allen 2006) |
| Central Belt | Castle Copse, Great Bedwyn | villa | 3rd-4th C AD | 1251 | 367 | 884 | 70.66 | Very high % of pig; sieved (Payne 1997) |
| East Anglia | Braughing, Skeleton Green | roadside settlement | Late 1st C BC-mid 1st C AD | 586 | 449 | 137 | 23.38 | LIA oppidum (Ashdown and Evans 1981) |
| South | Heybridge, Elms Farm | roadside settlement | 2nd-3rd C AD | 302 | 247 | 55 | 18.21 | 2% in 1st-2nd C AD; 7% in 3rd-4th C AD (Johnstone & Alberella 2002) |
| South | Staines, Friends' Burial Ground site | roadside settlement | 2nd-3rd C AD | 432 | 342 | 90 | 20.83 | 9% in 1st-2nd C AD; 0% in 3rd-4th C AD (Chapman 1984) |
| South | Staines, Elmsleigh Centre 1975-78 | roadside settlement | 3rd-4th C AD | 318 | 260 | 58 | 18.24 | 5% in 1st-2nd C AD (Chapman 2010) |
| Central Belt | Wimpole | roadside settlement | 3rd-4th C AD | 92 | 70 | 22 | 23.91 | (Horton et al 1994) |
| South | Shadwell, Tobacco Dock | roadside settlement | 3rd-4th C AD | 292 | 211 | 81 | 27.74 | Sieved; dominated by cattle (Douglas et al. 2011) |