

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Probiotic potential of a *Lactobacillus rhamnosus* cheese isolate and its effect on the fecal microbiota of healthy volunteers

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1693026> since 2019-02-18T14:44:08Z

Published version:

DOI:10.1016/j.foodres.2019.02.004

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Probiotic potential of a *Lactobacillus rhamnosus* cheese isolate and its effect on the**
2 **fecal microbiota of healthy volunteers**

3

4 Joaquin Bautista-Gallego^{1§}, Ilario Ferrocino^{2§}, Cristian Botta², Danilo Ercolini^{3,4}, Luca
5 Cocolin², Kalliopi Rantsiou^{2*}

6

7 ¹ Food Biotechnology Department, Instituto de la Grasa, Consejo Superior de
8 Investigaciones Científicas, Universidad Pablo de Olavide, Seville, Spain

9 ² Department of Agricultural, Forest and Food Sciences, University of Torino, Italy

10 ³ Department of Agricultural Sciences, University of Naples “Federico II”, Italy

11 ⁴ Task Force on Microbiome Studies, University of Naples “Federico II”, Italy

12

13 §authors contributed equally to the work

14

15

16

17

18

19

20

21

22

23

24 *Corresponding author: kalliopi.rantsiou@unito.it, Largo Paolo Braccini 2, 10095

25 Grugliasco, Italy, 0039-011-6708870

26 **Abstract**

27 The present study describes an *in vitro* characterization of strains of lactic acid bacteria,
28 focusing on physiological characters of probiotic interest, and a subsequent placebo-
29 controlled, crossover administration trial, with a cohort of healthy volunteers. The
30 strains of lactic acid bacteria were previously isolated from a fermented food (long
31 ripened cheese) and several ones resulted to have promising probiotic characteristics.
32 Based on comprehensive evaluation of the data obtained, one strain was chosen and
33 supplemented in a fermented milk. The fermented milk was then used in the
34 administration trial with the goal of assessing its effect on the composition of the
35 intestinal microbiota, as reflected in the feces. The fermented milk, with or without
36 probiotic, had an effect on the intestinal microbiota and significant inter-individual
37 differences were observed in response to the intervention. A common trend was
38 observed related to two important populations of the human gut microbiota; a reduction
39 in the relative abundance of *Bacteroides* and increase in the abundance of *Prevotella* in
40 subjects during treatment compared to baseline were registered.

41

42 **Keywords:** probiotic, fermented milk, intestinal microbiota, *Lactobacillus*

43

44

45

46

47

48

49

50

51 **1. Introduction**

52 Fermented foods are colonized by Lactic Acid Bacteria (LAB), not only during
53 production but also at the moment of consumption. Increasing evidence suggests that
54 consumption of some fermented foods can promote human health and the live microbes
55 within them contribute to such effect (Marco et al, 2017). Delivery of high numbers of
56 commensal microorganisms that transiently interact with the intestinal microbiota and
57 the host may play a role in gut physiology (Plé et al., 2013). The diversity of microbes
58 associated with fermented foods represents also a rich source of potential probiotic
59 microorganisms. Several studies have focused on isolation and probiotic
60 characterization of LAB from fermented foods (olives, sausages, dairy). Consequently,
61 fermented foods can serve as vehicles for probiotic delivery (van Hylckama Vlieg et al.,
62 2011).

63 Probiotics are “live microorganisms that, when administered in adequate amounts,
64 confer a health benefit on the host” (Hill et al., 2014, FAO 2001, FAO 2002). Among
65 other beneficial health effects, modulation of the host immunity, pathogen exclusion
66 and reinforcement of the gut barrier are most cited. Such effects may be the result of
67 direct probiotic-host interaction or indirect, through modulation of the intestinal
68 microbiota. Several mechanisms have been described that lead to the above described
69 effects. Such mechanisms may be widely spread among different genera of LAB and
70 therefore are considered core benefits, or less commonly encountered and can be
71 considered strain-specific (Sanders et al, 2018, Hill et al, 2014).

72 Among probiotic intervention outcomes, the modulation of intestinal microbiota is
73 commonly sought after as it is considered an indirect way of health maintenance or
74 promotion (Walter et al, 2018). Advancements achieved in our understanding of the role
75 of intestinal microbiota in human health corroborate the need to explore how

76 consumption of foods supplemented with probiotics influences microbiota composition
77 and human health (Marco et al., 2017).

78 In a previous work, we studied a large collection of isolates, originating from a ripened
79 cheese, focusing on certain functional properties. Most of the isolates showed a high
80 auto-aggregation property, low hydrophobicity values and a general low survival to
81 simulated digestion process. However, sixteen strains showed promising functional
82 characteristics (Bautista-Gallego et al., 2014). The purpose of this study was to (i) refine
83 the characterization of the sixteen strains, focusing on probiotic properties (adhesion to
84 cell lines, inhibition of *Listeria monocytogenes* adhesion and invasion); (ii) to
85 investigate the consequences of the consumption of a fermented milk supplemented
86 with a probiotic, selected based on the characterization performed, on intestinal
87 microbiota composition of 14 healthy volunteers.

88
89

90 **2. Materials and methods**

91 *2.1. Bacterial isolates*

92 Sixteen different strains, previously isolated during ripening of an Italian hard
93 cheese, were used in this work. The strains were selected from a larger collection, based
94 on a functional characterization that included testing for autoaggregation and
95 hydrophobicity properties as well as survival in simulated gastrointestinal passage
96 (Bautista-Gallego et al., 2014). They belong to *Lb. helveticus* (6), *Lb. rhamnosus* (9)
97 and *Lb. fermentum* (1) species (Table 1). *Lb. rhamnosus* GG, a known probiotic
98 microorganism, was used as a reference strain (Caggia et al, 2015). *Lactococcus lactis*
99 40FEL3 was also used in this study since it is a potent bacteriocin-producing
100 microorganism (Dal Bello et al., 2012).

101

102 *2.2. Antibiotic susceptibility*

103 Antibiotic resistance or susceptibility were tested following the method
104 described by Argyri et al. (2013) and Botta et al. (2014). Minimal inhibitory
105 concentrations (MICs) of 9 antibiotics (ampicillin, AMP; gentamicin, GEN; kanamycin,
106 KAN; streptomycin, STR; erythromycin, ERY; vancomycin, VAN; chloramphenicol,
107 CHL; tetracycline: TET; and clindamycin, CLY) (Sigma, Milan, Italy) were determined
108 in MRS broth. All antibiotics were added according to the breakpoint concentrations
109 proposed by the European Food Safety Authority (EFSA, 2008). If an isolate resulted
110 resistant to a breakpoint concentration, it was progressively tested with higher
111 concentrations until the MIC was determined. All the assays were done in triplicate.

112

113 *2.3. Cell assays*

114 The experiments were performed using two intestinal epithelial cell lines of human
115 origin, namely Caco-2 (ATCC HTB-37) and HT-29 (ATCC HTB-38). Caco-2 cell line
116 was grown in an advanced Dulbecco Modified Eagle Medium (DMEM) (Thermo Fisher
117 Scientific, Milan, Italy) whereas HT-29 was grown in McCoy's Medium Modified
118 (Thermo Fisher Scientific). Both media were supplemented with 5% foetal calf serum
119 (Lonza, Basel, Switzerland), L-glutamine (2 mM, Sigma), penicillin (100 U mL⁻¹,
120 Sigma) and streptomycin (1 mg mL⁻¹, Sigma). The cell lines were routinely grown in 25
121 cm² culture flasks (Corning, New York, USA) at 37 °C in a humidified atmosphere
122 containing 5% CO₂ and 95% air, until confluent monolayers were obtained. The culture
123 medium was changed routinely and once the cells reached confluence they were
124 subpassaged. All tests were carried out in triplicate in an undifferentiated monolayer,
125 with a suspension of 30000 cells/well (c/w) for Caco-2 and 50000 c/w in the case of
126 HT-29.

127 2.3.1. Adhesion assay

128 The ability of the 16 *Lactobacillus* strains to adhere to an undifferentiated cell
129 monolayer (Caco-2 and HT-29) was studied using the protocol described by Botta et al.
130 (2014).

131 2.3.2. Inhibition of *Listeria monocytogenes* adhesion and invasion

132 Determination of the inhibition of *Listeria monocytogenes* adhesion and invasion
133 of a cell monolayer, by the 16 *Lactobacillus* strains, was carried out following the
134 method described by Nakamura et al. (2012) with some modifications. Briefly, 0.1 ml
135 of fresh medium (DMEM or McCoy's, depending of the cell line used) containing 7 log
136 CFU/ml of each of the *Lactobacillus* strains was added to the monolayer. After 90 min
137 of incubation at 37 °C, the medium was removed and the monolayer washed twice with
138 Phosphate Buffered Saline (PBS, pH 7.3). Then, 0.1 ml of fresh medium containing 7
139 log CFU/ml of *L. monocytogenes* (without antibiotics) was placed in each well of a 96
140 Well TC-Treated Microplate (Corning). After 2h of incubation at 37 °C, bacterial cells
141 that did not adhere to the monolayer were washed away with PBS. Then to quantify the
142 *L. monocytogenes* adhesion, 0.1 ml of Triton-X100 (0.25% in PBS) was added for 30
143 minutes at 37 °C. Subsequently, *L. monocytogenes* adhered to the monolayer was
144 quantified by plating on BHI agar (Oxoid, Milan, Italy). To measure the intracellular *L.*
145 *monocytogenes*, after the 2 hours incubation and the removal of non-adhered *L.*
146 *monocytogenes*, 200 µl of fresh medium containing 50 mg/ml gentamicin (Sigma) was
147 added for 1 hour to kill extracellularly adhered bacteria (Botta et al., 2014, Corr et al.,
148 2007). The cells were then washed 3 times with PBS and lysed by the addition of 0.1 ml
149 of Triton-X100 (0.25% in PBS). Viable *L. monocytogenes* were determined by plating
150 on BHI agar (Oxoid). Inhibition of *L. monocytogenes* adhesion and invasion for each
151 *Lactobacillus* strain were referred to the control obtained only with the addition of *L.*

152 *monocytogenes* (without prior treatment with a *Lactobacillus* strain). Results were
153 subjected to one-way ANOVA and Fisher Test was used to determine differences at $P <$
154 0.05, using the statistical software, Statistica 7.0 for Windows (Statsoft, Tulsa, USA).

155

156 2.4. Technological characterization

157 The growth/survival in skim milk (Biogenetics, Padova, Italy) of the *Lactobacillus*
158 strains was evaluated during 7 days. Briefly, Skim Milk was inoculated firstly with 6
159 log CFU/ml of *Streptococcus thermophilus* (Sacco, Milan, Italy). After 6 hours at 40 °C,
160 the *Lactobacillus* strains were added at ~8 log CFU/ml. Cell counts were determined, at
161 time of inoculation and after 7 days at 4 °C, on MRS agar and the survival rate of the
162 *Lactobacillus* strains was calculated $[(CFU/mL_{t=0}/CFU/mL_{t=7}) \times 100]$. The pH was
163 monitored daily throughout the period of 7 days using a pH meter (Crison, Modena,
164 Italy).

165

166 2.5. Fermented milk production and administration

167 Based on the previously obtained results, *Lb. rhamnosus* 14E4 was chosen to be used in
168 the production of a probiotic fermented milk. UHT milk was heated at 42 °C and then a
169 *S. thermophilus* inoculum (Sacco, Milan, Italy) (≈ 6 log cfu/mL) was added. After 15
170 hours, milk was divided in two different batches. One was stored directly at 4 °C
171 (placebo fermented milk). The other one was inoculated with *Lb. rhamnosus* 14E4 at a
172 final concentration of ~10 log CFU/mL and the fermented milk was incubated at 37°C
173 for 2 hours (probiotic fermented milk). Subsequently, the probiotic fermented milk was
174 stored at 4 °C until its consumption.

175 The fermented milk was administered to fourteen healthy volunteers ($n = 14$) with about
176 equal portions of men and women aged 25–40 (30.7 ± 4.0), with BMI > 18 ($25.18 \pm$

177 5.21) who habitually followed an omnivore diet. The exclusion criteria were: acute or
178 chronic gastrointestinal diseases; eating disorders, such as anorexia, bulimia or other
179 specified feeding or eating disorder; prevalent chronic diseases, such as diabetes
180 mellitus and cancer; antibiotic treatment or surgical operations during the previous 3
181 months; pregnancy and breastfeeding.

182 Volunteers were divided in two groups; group 1 consumed the probiotic fermented
183 milk and the group 2 consumed the placebo fermented milk (Fig. 1). The probiotic or
184 placebo product was administered daily (100 mL) for 15 days. After a week of wash
185 out, intervention was inverted and the fermented milk (placebo for group 1 and
186 probiotic for group 2) was administered for another 15 days.

187 Volunteers were provided with a series of containers to collect the faeces (VWR, Milan,
188 Italy) containing 10 mL of *RNAlater* (Thermo Fisher Scientific). The faecal samples
189 were collected at home and transferred to the sterile sampling containers using a
190 polypropylene spoon (3 spoons of about 15g of faecal samples) and immediately stored
191 at 4 °C. The specimens were transported to the laboratory within 12 hours of collection
192 at a refrigerated temperature and stored at -80 °C for further analysis. During the whole
193 study faeces were sampled from both groups collecting a total of 9 samples from each
194 volunteer (Fig. 1).

195

196 *2.6. RNA extraction and analysis by pyrosequencing.*

197 At each sampling point, RNA was extracted directly from faecal samples using the
198 Electrofor-Norgen RNA purification kit (Norgen Biotek Corp. Ontario, Canada),
199 according to the manufacturer's instructions. Seven µl of TURBO-DNase (Life
200 Technologies) were added to digest the DNA in the RNA samples, with an incubation
201 of 3 h at 37°C. RNA was quantified using a NanoDrop 1000 spectrophotometer
202 (Thermo Scientific, Milano, Italy) and standardized at 500 ng/µl. Each RNA solution

203 was checked for the presence of residual DNA by performing PCR amplification
204 (Ferrocino et al., 2015). When positive signals were detected, the DNase treatment was
205 repeated.

206 Reverse transcription (RT) reactions were performed using an M-MLV reverse
207 transcriptase (Promega, Milan, Italy). Five hundred ng of RNA were mixed with 1 µl of
208 10 µM Random Primers (Promega, Milan, Italy) and DNase- and RNase-free sterile
209 water (Sigma) to a final volume of 10 µl and then incubated at 75 °C for 5 min. The mix
210 was placed on ice and a mixture containing 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3
211 mM MgCl₂, 10 mM DTT, 2 mM of each dNTP, 1 µl of 200 U/µl M-MLV and 0.96 U of
212 RNasin ribonuclease inhibitor (Promega) was transferred to the reaction tube. Reverse
213 transcription was carried out at 42 °C for 1 hour.

214 Complementary DNA (cDNA) was used to study the bacterial diversity in the faecal
215 samples. Primers Gray28F (5'-TTTGATCNTGGCTCAG) and Gray519r (5'-
216 GTNTTACNGCGGCKGCTG), which amplify a fragment of 520 bp (Ercolini et al
217 2012) were used to amplify the V1-V3 region of the 16S rRNA and 454 adaptors were
218 included in the forward primer, followed by a 10-bp sample-specific multiplex identifier
219 (MID).

220 The PCR mixture contained 50 ng of template DNA, 0.4 µM of each primer, 0.50 mmol
221 L⁻¹ of each deoxynucleoside triphosphate, 2.5 mM MgCl₂, 5 µl of 10 X PCR buffer, and
222 2.5 U of native Taq polymerase (Thermo Fisher Scientific). The following PCR
223 conditions were used: 94 °C for 2 min, 35 cycles of 95 °C for 20 s, 56 °C for 45 s, and
224 72 °C for 5 min, and a final extension at 72 °C for 7 min. PCR products were purified
225 twice with Agencourt AMPure kit (Beckman Coulter, Milano, Italy) prior to further
226 processing. An equimolar pool of the PCR templates was obtained prior to further
227 processing. The amplicon pool was processed by using Titanium chemistry on a GS

228 Junior platform (454 Life Sciences, Roche, Monza, Italy) according to the
229 manufacturer's instructions.

230

231 *2.7. Bioinformatics*

232 The sequencing data obtained were analyzed by using QIIME 1.9.0 software (Caporaso
233 et al., 2010), and a pipeline previously described (Ferrocino et al., 2016). OTUs were
234 picked at 99% of similarity and representative sequences of each cluster were used to
235 assign taxonomy using the Greengenes 16S rRNA gene database (McDonald et al.,
236 2012). Alpha diversity indices were calculated by using the *diversity* function of the
237 vegan package. The Shannon-Wiener diversity index (H') was further analyzed using t-
238 test, with intervention (fermented milk consumption) being the main factor. Weighted
239 UniFrac distance matrices obtained through QIIME were imported in R to obtain
240 Principal Coordinates Analysis (PCoA) plots. The OTU table obtained through QIIME
241 displays the higher taxonomy resolution that was reached by the 16S data; when the
242 taxonomy assignment was not able to reach the species level, the genus or the family
243 was displayed. OTU tables and Weighted UniFrac distance matrices were used for
244 Adonis and Anosim statistical tests in R environment (www.r-project.org) in order to
245 verify differences across subjects (volunteers) and between the samples as a function of
246 intervention (placebo and probiotic) and groups (1 and 2). OTU tables filtered for OTUs
247 presence > 3% relative abundance in at least 20 samples, were then used to build a
248 principal component analysis (PCA) as a function of the groups by using the “*made4*”
249 package of R. Kruskal–Wallis tests were used to find significant differences in
250 microbial taxa abundance according to intervention or group. A P value of 0.05 or lower
251 was considered as statistically significant. PICRUSt (Langille et al., 2013) was used to
252 predict abundances of gene families based on 16S rRNA sequences data as recently

253 described (Ferrocino et al., 2015). KEGG orthologs were then collapsed at level 3 of
254 hierarchy, and the table was imported in the “GAGE” Bioconductor package (Luo et al.,
255 2009) to identify inferred metabolic pathways overrepresented or underrepresented
256 between placebo and probiotic consumption.

257

258 2.8. Nucleotide sequence accession number

259 All the sequencing data were deposited at the Sequence Read Archive of the National
260 Center for Biotechnology Information (Sequence accession number SRP067522).

261

262 3. Results

263 3.1. Probiotic characterization of *Lactobacillus* strains

264 3.1.1. Antibiotic resistance

265 The results of the antibiotic resistance trials are shown in Table 1. For the antibiotics
266 ampicillin, vancomycin, gentamycin, kanamycin and streptomycin, the strains
267 belonging to 3 different species of *Lactobacillus* showed comparable levels of
268 resistance. For the antibiotics erythromycin, clindamycin, tetracycline and
269 chloramphenicol, the resistance varied among strains belonging to the same species as
270 well as between different species. *Lb. rhamnosus* 10D9 showed high resistance to all
271 antibiotics, even up to the breakpoints proposed by EFSA (2008).

272 3.1.2. Adhesion to human cell lines and inhibition of *L. monocytogenes*

273 The sixteen *Lactobacillus* strains were tested for their ability to adhere to Caco-2 and
274 HT-29 human-derived cell lines. In the case of the Caco-2 undifferentiated monolayer
275 (Fig. 2a), the adhesion capability was variable (from 0.51% to 13.65%), but ten strains
276 showed higher values than the reference probiotic *Lb. rhamnosus* GG (with five of them
277 over 10% of adhesion). The adhesion of *Lb. rhamnosus* GG was higher on the HT-29
278 cell line, reaching a value above 12 % (Fig. 2b). Also in this case, strains 2D5, 4D5,

279 5D10, 10D9, 13E7, 14E4 and 15F9 showed similar or higher percentages of adherence.
280 Globally, a better capability to adhere to the HT-29 monolayer than to Caco-2 was
281 observed.

282 Further to the adhesion to a monolayer, the strains were also tested for their ability to
283 inhibit adhesion or invasion of *L. monocytogenes* to the same two monolayers. As
284 shown in Figure 3, most strains tested, including the *Lb. rhamnosus* GG, showed limited
285 effect on adhesion of *L. monocytogenes* to Caco-2 cells. Only strains 5D10, 6E8, 13E7,
286 21E1, 10F3 and 11F9 reduced significantly the adhesion. On the other hand, 10D9
287 increased significantly the adhesion (114.81%). Twelve of the sixteen strains decreased
288 the *L. monocytogenes* invasion into Caco-2 cells. The degree of inhibition varied
289 significantly among the strains. Strain 10F3 demonstrated the highest reduction of *L.*
290 *monocytogenes* invasion, similar to the level of inhibition of the bacteriocinogenic *L.*
291 *lactis* 40FEL3 (26.08% of *L. monocytogenes* invasion in the presence of *L. lactis*).
292 Interestingly, certain *Lactobacillus* strains, including the reference *Lb. rhamnosus*, had a
293 positive effect and promoted the invasion of *L. monocytogenes* into the Caco-2 cells. In
294 the case of the inhibition of *L. monocytogenes* adhesion to HT-29 cells (Fig. 3b), some
295 strains showed a similar behaviour with respect to *Lb. rhamnosus* GG while 12D5, 6E8,
296 13E7, 14E4, 21E1, 1F4, 10F3 and 15F9 promoted a significant adhesion reduction. In
297 the case of the inhibition of the invasion, only three strains (13E7, 14E4 and 10F3) were
298 able to reduce the entry of *L. monocytogenes* into HT-29 cells better than *Lb.*
299 *rhamnosus* GG. As observed for the Caco-2 cell line, also in the case of the HT-29,
300 certain strains improved significantly the invasion of *L. monocytogenes* (21E1 and 1F8).
301 *3.2. Technological characterization of Lactobacillus strains*

302 In order to complement the probiotic characterization of the strains, some technological
303 parameters were also taken into consideration. More specifically strains were tested for

304 their ability to survive during storage of fermented milk and for their impact on the pH
305 during storage. As shown in Table 1, the highest survival rate after 7 days was detected
306 for strains 2D5, 4D5, 5D10, 10D9, 14E4, 21E1, 1F4, 11F9 and 15F10 (values > 100%
307 represent growth of the strain during storage). Conversely, strains 12D5, 17D10, 6E8,
308 13E7, 1F8, 10F3 and 15F9 displayed the lowest survival (values < 100% represent
309 reduced cell concentration after 7 days). Some strains showed limited acidification
310 ability and at the end of the period reduced the pH by 0.2 units while others (21E1,
311 10D9, 12D5, 1F8, 1F4) sharply decreased the pH within the first day of storage,
312 reaching final values of pH around 4.4 (initial pH was 4.88) (data not shown).

313 3.2. Impact of fermented milk on intestinal microbiota

314 RNA extracted from feces of volunteers that consumed a probiotic (fermented milk
315 produced with the use of *S. thermophilus* and supplemented with *Lb. rhamnosus* 14E4)
316 or placebo fermented milk (fermented milk without addition of *Lb. rhamnosus* 14E4)
317 was used as template for the V1-V3 16S rRNA gene pyrosequencing. A total of
318 1,762,374 raw sequences was obtained and analyzed; 938,961 reads passed the filters
319 applied through the QIIME split_library.py script, with an average value of 7,890
320 reads/sample and an average length of 452 bp. Good's estimated sample coverage
321 (median value of 99%; Range 97-99%) indicated that satisfactory coverage was reached
322 for all samples analyzed (Tab. S1). A high level of microbial diversity was found with
323 an average of OTUs of $247,80 \pm 84,91$. A core microbiota was identified and included
324 9 OTUs across 90% of the samples (Fig. 4). As can be deduced from the figure, among
325 the 9 OTUs of the core microbiota, *Bacteroides*, *Roseburia* and *Faecalibacterium* that
326 had a median relative abundance higher than 10% also showed important variability
327 between samples. On the other hand, OTUs with lower relative abundance (< 10%)
328 presented limited variability between samples. Pairwise comparison of the Shannon-

329 Wiener diversity index (H') revealed a biological diversity ($P < 0.05$) comparing
330 samples before and after the intervention or comparing group 1 vs group 2 (data not
331 shown).

332 The fecal microbiota of each individual was clearly differentiated based on principal-
333 coordinate analysis with a Weighted UniFrac distance matrix. In addition, both Adonis
334 and Anosim statistical tests confirmed that there was a significant difference between
335 the individuals based on their microbiota ($P < 0.001$) (data not shown). Plotting the
336 Weighted UniFrac distance matrix of each subject's dataset it was possible to observe a
337 shift in the gut microbiota composition before and after intervention in 7 out of the 14
338 volunteers. Fig. S1 concerns the 7 individuals for which a shift was observed with the
339 intervention while fig. S2 concerns the 7 individuals for which such shift was not
340 evident. However this separation (of data before/after intervention) was not statistically
341 significant (ANOSIM statistical test $P > 0.05$). Further, the most separated samples were
342 the ones corresponding to the subjects at time zero (Fig. S1). By comparing the initial
343 microbiota of all the subject (time 0) we did not observe a separation in terms of
344 composition across samples (ANOSIM statistical test $P > 0.05$).

345 The discrimination (ANOSIM, $P < 0.01$) on the relative abundances of OTUs associated
346 with the two groups was also taken into account (Fig. 6). The Principal Component
347 Analysis (PCA) on the OTUs table at genus level, showed a certain degree of separation
348 between subjects that used first the probiotic and then the placebo (G1) and subjects that
349 *vice versa* consumed first the placebo and then the probiotic (G2). The plot shows that
350 G1 were characterized by significantly elevated abundance of *Prevotella*, *Collinsella*,
351 *Faecalibacterium* and *Oscillospira* while G2 were characterized by *Sutterella*,
352 *Eubacterium* and *Odoribacter* (Fig. 6).

353 Comparing sampling during probiotic treatment against sampling at time 0 (Fig. 6A), it
354 was possible to observe an increment ($P < 0.001$) of the minor OTUs fraction (0.5-4%
355 of the total OTUs) comprising *Prevotella*, *Clostridium*, *Roseburia* and *Succinivibrio*.
356 The most remarkable differences appeared on the decrement of the abundance of
357 *Bacteroides* and *B. dorei* in samples during probiotic consumption compared to
358 baseline. Moreover, samples during placebo intervention (Fig. 6B) showed an increase
359 of *Prevotella*, *Ruminococcaceae* and *Sutterella* and a decrease of *Bacteroides* and
360 *Lachnospiraceae* compared to samples at baseline. Regarding the predicted
361 metagenomes, NSTI index for the samples was 0.079 ± 0.040 , indicating 92% accuracy
362 for all the samples. The pathway enrichment analysis performed by GAGE of the
363 predicted metagenomes showed an enrichment of Phosphonate and phosphinate
364 metabolism (ko00440), Pentose and glucuronate interconversions (ko00040) and ABC
365 transporters (ko02010) in samples under intervention (placebo and probiotic) compared
366 to baseline (data not shown).

367

368 **Discussion**

369 In the last decades several studies report how the diet can influence the composition of
370 the human gut microbiota. Food containing LAB, such as fermented milks or cheeses
371 can modify the gut microbiota since these bacteria can easily survive and colonize the
372 gastrointestinal tract (Putignani et al. 2014). The consumption of probiotics can have
373 significant effects on human health (Larsen et al., 2010) and may change the metabolic
374 gene content in the gut (Claus et al., 2011). Short-term dietary interventions on the
375 contrary have shown to only have a minor influence on microbiota composition (Wu et
376 al. 2011; Lappi et al., 2013).

377 In this work we expanded the probiotic characterization of selected LAB strains with
378 the purpose of identifying a probiotic candidate to be used in an intervention study with
379 healthy volunteers. Antibiotic resistance was evaluated and β -Lactam antibiotic
380 ampicillin and the broad-spectrum clindamycin were the most effective antibiotics, as
381 observed by Botta et al. (2014). Conversely, vancomycin, kanamycin and streptomycin
382 resulted to be less effective and higher concentrations (256 μ g/ml) were required to
383 reach inhibition. Biogenic amine production was also evaluated and only tyramine was
384 produced by three strains at levels below 1 mg/mL (data not shown). Adhesion to Caco-
385 2 and HT-29 was variable among the strains tested, but for several ones it was higher
386 than the adhesion observed for *Lb. rhamnosus* GG, used as a reference probiotic strain.
387 Variability was also observed in the capacity of the strains to limit adhesion or invasion
388 of *L. monocytogenes* in the same two cell lines. Overall the strains did not significantly
389 inhibit adhesion, but had an effect (mostly negative) in the invasion. Based on the
390 results obtained, a *Lb. rhamnosus* strain was chosen and inoculated in milk fermented
391 by *S. thermophilus* to produce a probiotic fermented milk.

392 The influence of the fermented milk on gut microbiota was assessed using RNA as
393 target molecule to investigate whether the consumption could result in alterations in the
394 microbial community structure. It was not possible to trace the *Lb. rhamnosus* or *S.*
395 *thermophilus*, that were ingested with the fermented milk, in the sequencing data or by
396 specifically targeting them by real time PCR (data not show).

397 The results indicate a significant inter-individual difference in response to the
398 intervention; possibly due to the relatively small size of the cohort, such inter-individual
399 differences may not allow for a clear effect to emerge. In fact, no clear effect on the
400 major components of gut microbiota, similar among subjects, was found when
401 comparing intervention and baseline samples. This is in agreement with recent DNA-

402 based studies evaluating the fecal microbiota before and after probiotic intervention
403 from 12 elderly (Eloe-Fadrosh et al. 2015), 25 healthy Finnish adults (Lahti et al. 2013),
404 seven adult female monozygotic twin pairs (Unno et al., 2015), 18 healthy Japanese
405 adults (Kim et al., 2013) and 10 New Zealand adult subjects (Tannock et al., 2010). We
406 found however a shift in the minor OTUs during probiotic (or placebo) intervention
407 compared to baseline. In particular, such shift concerned an overall increase in the
408 *Firmicutes* and a decrease in *Bacteroidetes* population as affected by the fermented
409 milk. These results are in contrast with Unno et al. (2015) where the relative abundance
410 of *Bacteroidetes* species increased during the intervention period and decreased during
411 the non-ingestion period. Notwithstanding the obvious heterogeneity of the individuals,
412 the discrepancy observed with this study may also be due to the different target
413 molecules used, RNA instead of DNA and to the different number of subjects in the
414 cohort (14 instead of 6). This is clearly shown from the core microbiota of this cohort,
415 mainly represented by members belonging to *Firmicutes*. Regarding the OTUs that
416 changed as affected by the intervention, it was possible to observe a reduction in the
417 relative abundance of *Bacteroides* in subjects during intervention (placebo and
418 probiotic) compared to baseline and an increase in the abundance of *Prevotella*. It has
419 been recently reported that high *Bacteroides* levels are associated to an increased risk of
420 weight gain and insulin resistance (Le Chatelier et al., 2013), while *Prevotella* is mostly
421 associated with the production of short chain fatty acid (SCFA) and in details is mainly
422 associated with the production of acetate from pyruvate via acetyl-CoA (Rey et al.,
423 2010; Louis et al., 2014). The protective role of SCFAs against different types of
424 disease is well documented (De Filippis et al., 2015). A recent metatranscriptomic study
425 in a mouse model showed that the genes that exhibited the highest fold-change in
426 expression were associated with the KEGG categories ‘carbohydrate metabolism’ and

427 'membrane transport' (McNulty et al. 2015). Consistent with this, the predicted
428 metagenomes in this study showed a significant increase in abundance of genes
429 associated with sugar interconversions and genes involved in membrane transport in
430 subjects during intervention compared to baseline (data not shown).

431 The main finding from the investigation of the intestinal microbiota of subjects that
432 consumed a fermented milk, with or without a probiotic, is that changes occur for minor
433 bacterial populations. This finding is in line with previous reports that interested
434 subjects with compromised health (Ceapa et al, 2013). In probiotic interventions
435 performed on healthy adults, little or no effect on the composition of the microbiota was
436 observed (Kristensen et al, 2016). Low abundance populations, may be metabolically
437 active and interfere with metabolic networks in the large intestine, modulating intestinal
438 microbiota (Ceapa et al, 2013).

439 The strains that were explored in this study originate from a fermented food. It has been
440 proven here that a ripened cheese harbors microbes that, when ingested in large
441 quantities, have the potential to interfere and modulate the composition of the intestinal
442 microbiota. It remains to be seen what is the effect of these microbes when ingested in
443 lower concentrations and concomitantly with other, closely related and competitive
444 microbes, within a fermented food.

445

446

447 **Acknowledgements**

448 This research program has received funding from the Compagnia di San Paolo, through
449 the University of Torino (ORTO11HEXP).

450 **References**

451 Argyri, A.A., Zoumpopoulou, G., Karatzas, K.-A.G., Tsakalidou, E., Nychas, G.-J.E.,
452 Panagou, E.Z., Tassou, C. (2013). Selection of potential probiotic lactic acid bacteria
453 from fermented olives by in vitro tests. *Food Microbiology*, 33, 282–291.

454 Bautista-Gallego, J., Alessandria, V., Fontana, M., Bisotti, S., Taricco, S., Dolci, P.,
455 Cocolin, L., Rantsiou, K. (2014). Diversity and functional characterization of
456 *Lactobacillus* spp. isolated throughout the ripening of a hard cheese. *International*
457 *Journal of Food Microbiology*, 181, 60-66.

458 Botta, C., Langerholc, T., Cencič, A., Cocolin, L. (2014). In vitro selection and
459 characterization of new probiotic candidates from table olive microbiota. *Plos one*,
460 9, e94457-e94457.

461 Caggia, C., De Angelis, M., Pitino, I., Pino, A., Randazzo, C.L. (2015). Probiotic
462 features of *Lactobacillus* strains isolated from Ragusano and Pecorino Siciliano
463 cheeses. *Food Microbiology*, 50: 109-117.

464 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello,
465 E.K., Fierer, N., Peña, A.G., Goodrich, K., Gordon, J.I., Huttley, G., Kelley, S.T.,
466 Knights, D., Jeremy, E., Ley, R.E., Lozupone, C., Mcdonald, D., Muegge, B.D.,
467 Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W. (2010). QIIME allows
468 analysis of high-throughput community sequencing data. *Nature Methods*, 7, 335–
469 336. doi:10.1038/nmeth.f.303.

470 Ceapa, C., Wopereis, H., Rezaïki, L., Kleerebezem, M., Knol, J., Oozeer, R. (2013).
471 Influence of fermented milk products, prebiotic and probiotics on microbiota
472 composition and health. *Best Practice & Research Clinical Gastroenterology*, 27,
473 139–155.

474 Claus, S. P., S. L. Ellero, B. Berger, L. Krause, A. Bruttin, J. Molina, A. Paris, E. J.
475 Want, I. de Waziers, O. Cloarec, S. E. Richards, Y. Wang, M. E. Dumas, A. Ross, S.
476 Rezzi, S. Kochhar, P. Van Bladeren, J. C. Lindon, E. Holmes, J. K. Nicholson.
477 (2011). Colonization-induced host-gut microbial metabolic interaction. *MBio*,
478 2:e00271-10.

479 Corr, S.C., Grahan, C.G.M., Hill, C. (2007). Impact of selected *Lactobacillus* and
480 *Bifidobacterium* species on *Listeria monocytogenes* infection and the mucosal
481 immune response. *FEMS Immunology and medical microbiology*, 50, 380–388

482 Dal Bello, B., Cocolin, L., Zeppa, G., Field, D., Cotter, P.D., Hill, C. (2012).
483 Technological characterization of bacteriocin producing *Lactococcus lactis* strains
484 employed to control *Listeria monocytogenes* in Cottage cheese. *International*
485 *Journal of Food Microbiology*, 153, 58-65.

486 De Filippis, F., Pellegrini, N., Vannini, L., Jeffery, I., La Stora, A., Laghi, L.,
487 Serrazanetti, D.I., Di Cagno, R., Ferrocino, I., Lazzi, C., Turrone, S., Cocolin, L.,
488 Brigidi, P., Neviani, E., Gobbetti, M., O’Toole, P.W., Ercolini, D. (2015). High-
489 level adherence to a Mediterranean diet beneficially impacts the gut microbiota and
490 associated metabolome. *Gut*, 0,1–10

491 EFSA (2008). Update of the criteria used in the assessment of bacterial resistance to
492 antibiotics of human or veterinary importance (Technical guidance). *The EFSA*
493 *Journal*, 732, 1–15.

494 Eloje-Fadros, E., Brady, A., Crabtree, J., Drabek, E.F., Ma, B., Mahurkar, A., Ravel, J.,
495 Haverkamp, M., Fiorino, A., Botelho, C., Andreyeva, I., Hibberd, P.L., Fraser, M.
496 (2015). Functional dynamics of the gut microbiome in elderly people during

497 probiotic consumption. *mBio*, 6, 1–12. doi:10.1128/mBio.00231-15.

498 Ercolini, D., De Filippis, F., La Stora, A., Iacono, M. (2012). "Remake" by high-
499 throughput sequencing of the microbiota involved in the production of water
500 buffalo mozzarella cheese. *Applied and Environmental Microbiology*, 78, 8142-
501 8145.

502 Food and Agricultural Organization of the United Nations and World Health
503 Organization. 2001. Health and nutritional properties of probiotics in food including
504 powder milk with live lactic acid bacteria. *World Health Organization*,
505 [http://www.fao.org/3/a-](http://www.fao.org/3/a-a0512e.pdf)
506 [a0512e.pdf](http://www.who.int/foodsafety/publications/fs_management/en/probiotics.pdf)[http://www.who.int/foodsafety/publications/fs_management/en/p](http://www.who.int/foodsafety/publications/fs_management/en/probiotics.pdf)
507 [robiotics.pdf](http://www.who.int/foodsafety/publications/fs_management/en/probiotics.pdf).

508
509 Food and Agricultural Organization of the United Nations and World Health
510 Organization. 2002. Joint FAO/WHO working group report on drafting guidelines
511 for the evaluation of probiotics in food. *Food and Agricultural Organization of the*
512 *United Nations*.
513 https://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.pdf.
514

515 Ferrocino, I., Di Cagno, R., De Angelis, M., Turrone, S., Vannini, L., Bancalari, E.,
516 Rantsiou, K., Cardinali, G., Neviani, E., Cocolin, L. (2015). Fecal microbiota in
517 healthy subjects following omnivore, vegetarian and vegan diets: culturable
518 Populations and rRNA DGGE profiling. *PLoS One* 10, e0128669.
519 doi:10.1371/journal.pone.0128669

520 Ferrocino, I., Greppi, A., La Stora, A., Rantsiou, K., Ercolini, D., Cocolin, L. (2016).
521 Impact of nisin-activated packaging on microbiota of beef burgers during storage.
522 *Applied and Environmental Microbiology*, 82, 549–559. doi:10.1128/AEM.03093-
523 15.

524 Hill, C., Guarner, F., Reid, G., Gibson, G.R., Merenstein, D. J., Pot, B., Morelli, L.,
525 Canani, R. B., Flint, H.J., Salminen, S., Calder, P.C., Ellen Sanders, M.E. (2014).
526 The International Scientific Association for Probiotics and Prebiotics consensus
527 statement on the scope and appropriate use of the term prebiotic. *Nature Reviews*
528 *Gastroenterology & Hepatology*, *11*, 506–514.

529 Kim, S.W., Suda, W., Kim, S., Oshima, K., Fukuda, S., Ohno, H., Morita, H., Hattori,
530 M. (2013). Robustness of gut microbiota of healthy adults in response to probiotic
531 intervention revealed by high-throughput pyrosequencing. *DNA Research*, *20*,
532 241–253. doi:10.1093/dnares/dst006

533 Kristensen, N.B., Bryrup, T., Allin, K.H., Nielsen, T., Hansen, T.H., Pedersen, O.
534 (2016). Alterations in fecal microbiota composition by probiotic supplementation
535 in healthy adults: a systematic review of randomized controlled trials. *Genome*
536 *Medicine*, *8*, 52.

537 Lahti, L., Salonen, A., Kekkonen, R.A., Salojärvi, J., Jalanka-Tuovinen, J., Palva, A.,
538 Orešič, M., de Vos, W.M. (2013). Associations between the human intestinal
539 microbiota, *Lactobacillus rhamnosus* GG and serum lipids indicated by integrated
540 analysis of high-throughput profiling data. *PeerJ*, *1*, e32. doi:10.7717/peerj.32

541 Lappi, J., Salojärvi, J., Kolehmainen, M., Mykkänen, H., Poutanen, K., de Vos, W.M.,
542 Salonen, A. (2013). Intake of whole-grain and fiber-rich rye bread versus refined
543 wheat bread does not differentiate intestinal microbiota composition in Finnish
544 adults with metabolic syndrome. *Journal of Nutrition*, *143*, 648–55.
545 doi:10.3945/jn.112.172668

546 Larsen, N., Vogensen, F.K., Van Den Berg, F.W.J., Nielsen, D.S., Andreasen, A.S.,

547 Pedersen, B.K., Al-Soud, W.A., Sørensen, S.J., Hansen, L.H., Jakobsen, M.
548 (2010). Gut microbiota in human adults with type 2 diabetes differs from non-
549 diabetic adults. *PLoS One* 5, doi:10.1371/journal.pone.0009085

550 Le Chatelier, E., Nielsen, T., Qin, J., Prifti, E., Hildebrand, F., Falony, G., Almeida, M.,
551 Arumugam, M., Batto, J.-M., Kennedy, S., Leonard, P., Li, J., Burgdorf, K.,
552 Grarup, N., Jørgensen, T., Brandslund, I., Nielsen, H.B., Juncker, A.S., Bertalan,
553 M., Levenez, F., Pons, N., Rasmussen, S., Sunagawa, S., Tap, J., Tims, S.,
554 Zoetendal, E.G., Brunak, S., Clément, K., Doré, J., Kleerebezem, M., Kristiansen,
555 K., Renault, P., Sicheritz-Ponten, T., de Vos, W.M., Zucker, J.-D., Raes, J.,
556 Hansen, T., Bork, P., Wang, J., Ehrlich, S.D., Pedersen, O., Guedon, E., Delorme,
557 C., Layec, S., Khaci, G., van de Guchte, M., Vandemeulebrouck, G., Jamet, A.,
558 Dervyn, R., Sanchez, N., Maguin, E., Haimet, F., Winogradski, Y., Cultrone, A.,
559 Leclerc, M., Juste, C., Blotière, H., Pelletier, E., LePaslier, D., Artiguenave, F.,
560 Bruls, T., Weissenbach, J., Turner, K., Parkhill, J., Antolin, M., Manichanh, C.,
561 Casellas, F., Boruel, N., Varela, E., Torrejon, A., Guarner, F., Denariáz, G.,
562 Derrien, M., van Hylckama Vlieg, J.E.T., Veiga, P., Oozeer, R., Knol, J.,
563 Rescigno, M., Brechot, C., M'Rini, C., Mérieux, A., Yamada, T. (2013). Richness
564 of human gut microbiome correlates with metabolic markers. *Nature*, 500, 541–
565 546. doi:10.1038/nature12506

566 Louis, P., Hold, G.L., Flint, H.J. (2014). The gut microbiota, bacterial metabolites and
567 colorectal cancer. *Nature Reviews Microbiology*, 12, 661–672.

568 Marco, M.L., Heeney, D., Binda, S., Cifelli, C.J., Cotter, P.D., Foligné, B., Gänzle M.,
569 Remco Kort, R., Pasin, G., Pihlanto, A., Smid, E.J., Robert Hutkins, R. (2017).
570 Health benefits of fermented foods: microbiota and beyond. *Current Opinion in*

571 *Biotechnology*, 44, 94–102.

572 McDonald, D., Price, M.N., Goodrich, J., Nawrocki, E.P., DeSantis, T.Z., Probst, A.,
573 Andersen, G.L., Knight, R., Hugenholtz, P. (2012). An improved Greengenes
574 taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria
575 and archaea. *ISME Journal*, 6, 610–8. doi:10.1038/ismej.2011.139

576 McNulty, N.P., Yatsunenko, T., Hsiao, A., Faith, J.J., Muegge, B.D., Goodman, A.L.,
577 Henrissat, B., Oozeer, R., Cools-Portier, S., Gobert, G., Chervaux, C., Knights, D.,
578 Lozupone, C.A., Knight, R., Duncan, A.E., Bain, J.R., Muehlbauer, M.J.,
579 Newgard, C.B., Heath, A.C., Gordon, J.I. (2011). The Impact of a consortium of
580 fermented milk strains on the gut microbiome of gnotobiotic mice and
581 monozygotic twins. *Science Translational Medicine*, 3, 106ra106-106ra106.
582 doi:10.1126/scitranslmed.3002701.

583 Nakamura, S., Kuda, T., An, C., Kanno, T., Takahashi, H., Kimura, B. (2012). Inhibitory
584 effects of *Leuconostoc mesenteroides* 1RM3 isolated from narezushi, a
585 fermented fish with rice, on *Listeria monocytogenes* infection to Caco-2 cells
586 and A/J mice. *Anaerobe*, 18, 19-24.

587 Plé, C., Breton, J., Daniel, C., Foligné, B. (2015). Maintaining gut ecosystems for
588 health: are transitory food bugs stowaways or part of the crew? *International*
589 *Journal of Food Microbiology*, 213, 139–43.

590 Putignani, L., Del Chierico, F., Petrucca, A., Vernocchi, P., Dallapiccola, B. (2014).
591 The human gut microbiota: a dynamic interplay with the host from birth to
592 senescence settled during childhood. *Pediatric Research*, 76, 2–10.
593 doi:10.1038/pr.2014.49

594 Rey, F.E., Faith, J.J., Bain, J., Muehlbauer, M.J., Stevens, R.D., Newgard, C.B.,
595 Gordon, J.I. (2010). Dissecting the in vivo metabolic potential of two human gut
596 acetogens. *Journal of Biological Chemistry*, 285, 22082–22090.

597 Sanders, M.E., Benson, A., Lebeer, S., Merenstein, D.J., Klaenhammer, T.R. (2018).
598 Shared mechanisms among probiotic taxa: implications for general probiotic
599 claims. *Current Opinion in Biotechnology*, 49, 207–216.

600 Tannock, G.W., Munro, K., Harmsen, H.J.M., Welling, G.W., Smart, J., Gopal, P.K.
601 (2000). Analysis of the fecal microflora of human subjects consuming a probiotic
602 product containing *Lactobacillus rhamnosus* DR20. *Applied and Environmental*
603 *Microbiology*, 66, 2578–2588. doi:10.1128/AEM.66.6.2578-2588.2000

604 Unno, T., Choi, J.-H., Hur, H.-G., Sadowsky, M.J., Ahn, Y.-T., Huh, C.-S., Kim, G.-B.,
605 Cha, C.-J. (2015). Changes in human gut microbiota influenced by probiotic
606 fermented milk ingestion. *Journal of Dairy Science*, 98, 3568–3576.
607 doi:10.3168/jds.2014-8943

608 van Hylckama Vlieg, J. E.T., Veiga, P., Zhang, C., Derrien, M., Zhao, L. (2011). Impact
609 of Microbial Transformation of Food on Health-from Fermented Foods to
610 Fermentation in the Gastro-Intestinal Tract. *Current Opinion in Biotechnology* 22,
611 211–19.

612 Walter, J., Maldonado-Gómez, M.X., Martínez, I. (2018). To engraft or not to engraft:
613 an ecological framework for gut microbiome modulation with live microbes.
614 *Current Opinion in Biotechnology*, 49, 129–139.

615 Wu, G.D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y.-Y., Keilbaugh, S. a, Bewtra,

616 M., Knights, D., Walters, W. a, Knight, R., Sinha, R., Gilroy, E., Gupta, K.,
617 Baldassano, R., Nessel, L., Li, H., Bushman, F.D., Lewis, J.D. (2011). Linking
618 long-term dietary patterns with gut microbial enterotypes. *Science* 334, 105–8.
619 doi:10.1126/science.1208344

620 **Figure legends**

621 **Figure 1.** Schematic representation of the experimental trial. The figure shows the
622 intervention for each group. Group 1 consumed first the probiotic for a period of 14
623 days, then there was a 1-week washout followed by placebo consumption for a period of
624 14 days. Group 2 consumed first the placebo for 14 days, then there was a 1 week
625 washout followed by probiotic consumption for 14 days. The figure also shows the days
626 of the sampling; 1 sample before the intervention, followed by three samples during the
627 first part of the trial, 1 sample during washout, 3 more samples during the second part of
628 the trial, followed by one last sample after the intervention (total 9 samples for each
629 individual).

630 **Figure 2.** Adhesion profile of the lactic acid bacteria, expressed as the ratio (%) of
631 bacteria recovered from the human cells after incubation (37 °C for 90 minutes) and the
632 initial bacterial count of the inoculum. A) Caco-2 undifferentiated monolayer; B) HT-29
633 undifferentiated monolayer.

634 **Figure 3.** Adhesion and invasion inhibition of *Listeria monocytogenes* to an A) Caco-2
635 undifferentiated monolayer; B) HT-29 undifferentiated monolayer. Results are
636 expressed relative to control (i.e. *Listeria monocytogenes* adhesion and invasion without
637 the presence of a *Lactobacillus* strain, corresponding to 100% adhesion or invasion).

638 **Figure 4.** Abundance (%) of the 9 OTUs identified in 90% of the samples. Boxes
639 represent the interquartile range (IQR) between the first and third quartiles, and the line
640 inside represents the median (2nd quartile). Whiskers denote the lowest and the highest
641 values within 1.56 IQR from the first and third quartiles, respectively. Circles represent
642 outliers beyond the whiskers.

643 **Figure 5.** PCA based on the OTU abundance. G1 (subjects that used first the probiotic
644 and then the placebo); G2 (subjects that vice versa consumed first the placebo and then
645 the fermented milk).

646
647 **Figure 6.** Relative abundance (percent) of the OTUs ($> 0.2\%$) that are differentially
648 abundant ($P < 0.01$) across baseline and probiotic treated samples (Plot A), and across
649 baseline and placebo treated samples (Plot B). The red bar corresponds to the mean
650 value at time zero (baseline) for the 14 individuals. The blue bar in plot A, shows the
651 mean value for the 14 individuals during probiotic consumption (for each individual the
652 mean value of the three sampling points during probiotic consumption was used).
653 Similarly, the yellow bar in plot B, shows the mean value of the 14 individuals during
654 placebo consumption (for each individual the mean value of the three sampling points
655 during placebo consumption were considered). Refer to figure 1 for sampling points
656 during probiotic and placebo consumption.

657

658

659

660

661 **Table 1.** Antibiotic MICs values ($\mu\text{g/mL}$) and survival rate in skim milk after 7 days for
 662 the 16 *Lactobacillus* strains.

Strain	Species	AMP.	VAN.	GEN.	KAN.	STR.	ERY.	CL.	TET.	CHL.	Survival rate (%)
2D5	<i>L. helveticus</i>	1	256	64	256	128	2	1	8	8	93.88
4D5		1	256	32	256	256	2	1	8	8	134.93
5D10		1	256	256	256	256	8	8	64	64	85.71
6E8		1	256	256	256	256	2	2	64	32	26.70
1F4		1	256	256	256	256	2	8	64	32	128.76
1F8		1	256	128	256	256	2	1	64	32	64.31
10D9		<i>L. rhamnosus</i>	1	256 ^{n.r.}	256	256	256	16	16	256	256
12D5	1		256 ^{n.r.}	128	256	256	1	1	8	8	68.86
17D10	1		256 ^{n.r.}	128	256	128	1	1	8	8	69.97
13E7	1		256 ^{n.r.}	128	256	256	2	1	16	8	32.56
14E4	1		256 ^{n.r.}	256	256	256	8	1	64	32	128.76
21E1	1		256 ^{n.r.}	128	256	256	8	1	64	32	171.67
11F9	1		256 ^{n.r.}	256	256	256	8	8	64	256	92.08
15F9	1		256 ^{n.r.}	128	256	256	2	2	64	32	36.22
15F10	1		256 ^{n.r.}	256	256	256	4	2	64	32	127.27
10F3	<i>L. fermentum</i>		1	256 ^{n.r.}	256	256	256	8	8	64	32

663 **Note:** ampicillin, AMP; gentamicin, GEN; kanamycin, KAN; streptomycin, STR; erythromycin, ERY;

664 vancomycin, VAN; chloramphenicol, CHL; tetracycline: TET; and clindamycin, CLI.

665 **n.r.:** not required (EFSA, 2008).

666

667 **Figure 1.** Schematic representation of the experimental trials.

668


669

670

671

672

673



674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704


705

706

707



708
709

Figure 2.




710
711
712
713
714
715
716
717
718
719
720

721 **Figure 3.**




722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732

733 **Figure 4.**
 734
 735




736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757 **Figure 5.**
 758

759
760
761



762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783


Figure 6.



785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796


Supplementary Figure S1. Principal-coordinate analysis based on Weighted UniFrac analysis of the 16S rRNA gene sequences for selected subjects. Samples from each plot

797 are color coded according to: baseline (red), probiotic (blue), placebo (yellow), washout (gray) and after the trial (black). Sampling day are also reported.
 798
 799
 800
 801
 802
 803



804
 805
 806
 807
 808
 809
 810
 811
 812
 813

814 **Supplementary Figure S2.** Principal-coordinate analysis based on Weighted UniFrac
 815 analysis of the 16S rRNA gene sequences for selected subjects. Samples from each plot
 816 are color coded according to: baseline (red), probiotic (blue), placebo (yellow), washout
 817 (gray) and after the trial (black). Sampling day are also reported.
 818



819