

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Polyphenolic diversity in *Vitis* sp. leaves

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1704440> since 2020-04-01T15:38:32Z

Published version:

DOI:10.1016/j.scienta.2019.108569

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Polyphenolic diversity in *Vitis* sp. leaves

2

3

4 Olga Kedrina-Okutan¹, Vittorino Novello¹, Thomas Hoffmann², Johannes Hadersdorfer³, Anna

5 Schneider⁴, Wilfried Schwab², and Alessandra Ferrandino^{1*}

6 ¹Dipartimento Scienze Agrarie, Forestali, Alimentari (DISAFA), Università degli Studi di Torino,

7 Largo P. Braccini, 2 10095, Grugliasco (TO), Italy.

8 ²Biotechnology of Natural Products, Technical University Munich, Liesel-Beckmann-Str. 1, Freising

9 85354, Germany.

10 ³Fruit Science, Technical University Munich, Dürnast 2, Freising 85354, Germany.

11 ⁴ Istituto Protezione Sostenibile delle Piante (IPSP), CNR, Strada delle Cacce, 73, 10135 Torino (TO),

12 Italy.

13 ***corresponding author: tel. +39-011-6708755; email: alessandra.ferrandino@unito.it**

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28 ABSTRACT

29 We identified and quantified the main constitutive polyphenolic compounds of the leaves of
30 seven *Vitis* species and of one interspecific cross, analysing leaf blades and veins separately, to spread
31 light on the strategic localization of polyphenols in leaf tissues. To the best to our knowledge, the main
32 leaf constitutive polyphenols of *V. candicans*, *V. coignetiae*, *V. vinifera sylvestris* and Börner were never
33 described. *V. riparia* and *V. rupestris* (belonging to the same botanical series of *Ripariae*) displayed
34 similar flavan-3-ol and dimeric proanthocyanidin concentration and similar percentage incidence of
35 caftaric acid over total phenolic acids. *V. riparia* distinguished respect to the other genotypes for its high
36 flavonol content, the highest percentage incidence of myricetin derivatives and an important
37 diversification in the type of accumulated flavonol. *V. berlandieri* (series *Cinereae*) and Börner (hybrid
38 of *V. riparia* x *V. cinerea*) accumulated low amounts of flavonol-glucosides comparing to the other
39 species, but they showed a wide profile diversification, as well. However, it was *V. v. sylvestris*, the wild
40 ancestor of *Vitis vinifera* subsp. *sativa* that displayed the widest flavonol profile diversification. The
41 differences in the flavonol profile could be related to the genus *Vitis* evolution: in fact, with
42 domestication, the flavonoid biosynthetic pathway underwent a progressive simplification; for this, the
43 highest flavonol diversity found in *Vitis v. sylvestris* is probably a demonstration of its reduced or nil
44 level of domestication. *V. amurensis*, known for its cold tolerance and resistance to downy mildew,
45 anthracnose and white rot, markedly differentiated respect to the other genotypes, for its high
46 concentration of polyphenols, particularly of vein flavonols and flavanonols. Moreover, *V. amurensis*
47 leaves generally presented a constantly high concentration of constitutive polyphenols throughout the
48 season that probably contributes to protect against adverse environmental condition. The abundance of
49 polyphenols in *V. amurensis* leaves emphasizes that this species is a source of natural bioactive
50 compounds that could find application for nutraceutical and pharmacological uses. *V. berlandieri* and
51 Börner markedly distinguished respect to the other studied species for their exclusive capability to
52 accumulate flavones (mainly orientin, isoorientin, vitexin and isovitexin) in blades and in veins, in
53 considerable amounts.

54 Knowledge about these subjects could contribute to shed light on the identification of species-
55 related molecules involved in the plant-defense mechanisms, to the chemotaxonomy of the genus *Vitis*,
56 to the possibility of identifying specific natural bioactive compounds to use in plant-based preparation
57 for nutraceutical, cosmetic, feed/food-additive purposes.

58

59 KEYWORDS: *V. amurensis*, *V. candicans*, *V. riparia*, *V. rupestris*, *V. berlandieri*, *V. coignetiae*, *V.*
60 *vinifera sylvestris*, Börner.

61

62

63 1. INTRODUCTION

64 The genus *Vitis* comprises more than 60 species, mostly inhabiting temperate regions. The most
65 known and economically important one is the western Eurasian species *Vitis vinifera* subsp. *vinifera*,
66 which ancestor is the wild *Vitis vinifera* subsp. *sylvestris*, naturally occurring in Europe, Middle East
67 and Northern Africa. Up to thirty grapevine species are native of East Asia and North America, areas
68 that host a wide diversity of *Vitis* species (Wan et al., 2013).

69 *Vitis vinifera* is susceptible to many pests. *Phylloxera* (*Daktulosphaira vitifoliae*), downy and
70 powdery mildew (caused by *Plasmopara viticola* and *Erysiphe necator*, respectively), imported almost
71 simultaneously from North America to Europe in the XIX century, are among those causing important
72 economic loss in viticulture. Viticulturists, technicians and researchers overcame the problem tied to
73 *vinifera* sensitiveness to *Phylloxera* by using American species or their hybrids as rootstocks.
74 Furthermore, the introduction of downy and powdery mildew pushed to develop breeding programs
75 based on the constitution of interspecific crosses *vinifera*-American species that, together with Asian
76 species, were disease-tolerant or resistant due to their coevolution with the pathogen. *Vitis* species such
77 as *V. riparia*, *V. rupestris*, *V. californica* and *V. amurensis* are known as highly tolerant to downy mildew
78 (Gómez-Zeledón et al., 2013; Jürges et al., 2009); *V. riparia*, *V. munsoniana*, *V. candidans*, *V.*
79 *rotundifolia* as highly resistant to powdery mildew (Staudt, 1997; Wan et al., 2007) and *V. rotundifolia*
80 (subspecies *Muscadinia*) and *V. arizonica* as resistant to Pierce's disease (Ruel and Walker, 2006).
81 Besides, some species distinguish for their tolerance to environmental stressors thanks to their origin: it
82 is the case of *V. amurensis*, native of cold areas in northeastern China and Russian Siberia, that is highly
83 frost tolerant (Zhang et al., 2012).

84 To prevent disease or damage caused by biotic or abiotic stresses, plants employ a complex
85 defense system, which involves a broad spectrum of physical and biochemical changes. Biochemical
86 resistance, such as preformed defenses, has evolved to face environmental stressors and is also related,
87 among others, to secondary metabolites, including polyphenols (reviewed in Llorens et al., 2017 and in
88 Dixon, 2001). *Vitis vinifera* is known for its abundance and richness in polyphenols that have been and

89 currently are widely studied in the berries due to their implications in the technology of winemaking and
90 in the quality of derived wines. Thousands of scientific studies were devoted to define the polyphenolic
91 composition of *Vitis vinifera* berries, and, to a much lesser extent, to vegetative organs; oppositely, less
92 is known about other *Vitis* species polyphenolic composition, particularly as to vegetative organs. Moore
93 and Giannasi (1994) described the qualitative composition of flavonols and flavones in leaves of some
94 North American *Vitis* species. Flavonol profiles were described also in *V. amurensis* (Hmamouchi et al.,
95 1996) and *V. rotundifolia* (Pastrana-Bonilla et al., 2003) leaves. Main phenolic compounds were also
96 analyzed in *V. labrusca* leaves (Dani et al., 2010; Dresch et al., 2014). Chen et al. (2018) reviewed *V.*
97 *amurensis* polyphenolic composition and related pharmacological properties. At present, the scientific
98 interest about *Vitis* sp. polyphenolic composition is at least dual: to investigate the health beneficial
99 effects of grapevine leaves as a possible source of natural bioactive compounds, that can be used in
100 nutraceutical and pharmaceutical applications, and to deepen knowledge about their involvement in
101 grapevine defense mechanisms. Different *Vitis* species might display different metabolic pathways
102 and/or different gene regulations, resulting in the production of specific classes of bioactive polyphenols,
103 or of individual molecules. Knowledge about these subjects could contribute to shed light on the
104 identification of species-related molecules involved in the plant-defense mechanisms, to the
105 chemotaxonomy of the genus *Vitis* and to the possible identification of specific molecules to use in
106 specific plant-based preparation for nutraceutical, cosmetical, nutritional purposes.

107 The aim of the present study was to identify the main constitutive polyphenolic compounds in
108 the leaves of seven *Vitis* species and of one interspecific cross (the rootstock Börner, *V. riparia* x *V.*
109 *cinerea*; Figure 1). To the best to our knowledge *V. candicans*, *V. coignetiae*, *V. v. sylvestris* and Börner
110 main leaf constitutive polyphenols were never described. Leaf blades and veins were analysed separately
111 to spread light on polyphenol strategic localization in leaf tissues, to contribute to explain the grapevine-
112 pathogen interaction in specific biological systems.

113

114

115 2. MATERIALS AND METHODS

116 **2.1 Plant Material.** Four American (*V. candicans*, *V. riparia*, *V. berlandieri* and Börner), two
117 Asian (*V. amurensis*, *V. coignetiae*) and one Eurasian (*V. v. sylvestris*) grapevine genotypes were studied
118 (Table 1; Figure 1). All the studied accessions were previously checked for their trueness to type by
119 morphology and molecular markers. The studied *Vitis* genotype leaves were collected in the collection
120 vineyard of DISAFA, University of Turin, Grugliasco (Piedmont, Italy), sampled five times during the
121 season: 1 = 28th of May (148 day of the year - DOY), 2 = 22nd of June (DOY 173), 3 = 14th of July (DOY
122 195), 4 = 3rd of August (DOY 215), 5 = 28th of August (DOY 214), in 2015. The collection vineyard
123 was managed routinely during spring and summer as already described (Kedrina-Okutan et al., 2018)
124 (Kedrina-Okutan et al., 2018). One field parcel was generally constituted by four consecutive plants
125 used to collect the leaves; leaves were detached between the fourth and seventh node of the main shoots
126 from the west side of the row. Pools of 15 adult and healthy leaves (visual evaluation) were collected at
127 each sampling time and divided into three replicates. After sampling, leaves were immediately
128 transported to the laboratory for further analyses.

129 **2.2 Reference Compounds and Reagents.** Orientin, isoorientin, astilbin and caftaric acid were
130 purchased from Sigma-Aldrich S.r.l. (Milan, Italy). Vitexin, *trans*-ferric acid and *trans*-coutaric acid
131 were purchased from Phytolab (Vestenbergsgreuth, Germany). (+)-catechin, (-)-epicatechin, (-)-
132 epicatechin gallate, (-)-epigallocatechin gallate, proanthocyanidin B₁, proanthocyanidin B₂, quercetin 3-
133 *O*-glucoside, quercetin 3-*O*-glucuronide, kaempferol 3-*O*-glucoside, kaempferol 3-*O*-glucuronide,
134 myricetin 3-*O*-glucoside and isorhamnetin 3-*O*-glucoside from Extrasynthèse (Genay, France).
135 Folin–Ciocalteu reagent and tartaric acid were purchased from Merck (Darmstadt, Germany); sodium
136 sulfate and sodium metabisulfite were purchased from BDH Laboratory Supplies (Poole, England).

137 **2.3 Measurement of Dry Matter.** Leaf tissue dry matter was measured three times during the
138 season at DOY 186, DOY 200 and DOY 241. Leaf veins and blades were separated and dried inside an
139 oven for 72 hours at constant temperature of 110 °C. The weight differences after exsiccation were
140 measured, and results were expressed as percentage of dry weight over fresh weight.

141 **2.4 Sample Extraction for Polyphenol Analyses.** The 15 collected leaves were divided into
142 sub-samples of five, and veins and blades were immediately separated. Two grams of tissue were
143 randomly chosen from the three sub-samples and extracted in a hydroalcoholic buffer (pH = 3.9, 40%
144 ethanol, 22 mL L⁻¹ of 1 N NaOH, 5 g L⁻¹ of tartaric acid, 2 g L⁻¹ of sodium metabisulfite). For sample
145 homogenization and polyphenol extraction, an Ultraturrax dispersing machine (IKA, Staufen, Germany)
146 was used for around 1 min setting the speed at 10 000 rpm followed by 10 min of centrifugation at 4000
147 rpm. The supernatant was separated, and the pellet was resuspended with the same buffer; the
148 resuspension was kept in the dark for 30 min and then centrifuged again. The two supernatants were
149 combined and brought to a final volume of 50 mL. Obtained leaf extracts were stored at -20 °C until
150 further analysis.

151 **2.5 Measurement of Total Polyphenols.** The method of Singleton et al. (1999) was used and
152 results were expressed as gram of (+)-catechin equivalents (CE) per kg of blades/veins dry weight (DW).
153 Briefly, 100 µL of leaf extract were mixed with 5 mL ultrapure water and 1 mL of Folin-Ciocalteu
154 reagent was added, mixed and incubated at room temperature for 5 min. After incubation, 4 mL of 10%
155 Na₂CO₃ and ultrapure water were added until final volume of 20 mL. The mixture was incubated in the
156 dark for 90 min and the absorbance was read at 760 nm by UV/Vis spectrophotometer (PerkinElmer,
157 Lambda 25, Beaconsfield, Bucks, UK).

158 **2.6 Analysis of individual polyphenols.**

159 *2.6.1 Sample preparation.* Prior to the chromatographic analysis, the extracts were diluted with
160 1 M phosphoric acid 90/10 v/v and passed through 0.20 µm membrane filter GHP Acrodisc (PALL
161 Italia, Buccinasco, Milan, Italy) (Di Stefano and Cravero, 1992).

162 *Individual phenolic compound analysis by HPLC-ESI-MS/MS.* Chromatographic separation was
163 performed by HPLC-DAD system (HPLC 1100, Agilent Technologies, U.S.) equipped with column
164 Luna C-18 150 x 2 mm (Phenomex Aschaffenburg, Germany). Mobile phase solvent A was water with
165 formic acid 0.1% and solvent B was methanol with formic acid 0.1%; the gradient program for solvent
166 B was the following: 0-30 min 0-50 % B; 30-35 min 50-100 % B; 35-50 min 100 % B and back to initial

167 conditions from 50 to 55 min, followed by 10 min of isocratic flow. Injection volume was 5 μL , flow
168 rate 200 $\mu\text{L min}^{-1}$ and individual phenolic compounds were detected at 280, 320, and 370 nm. For the
169 mass spectrometry analyses, a Bruker Daltonics esquire 3000^{plus} ion trap spectrometer (Bruker
170 Daltonics, HB, Germany) equipped with electrospray (ESI) was used, operating at positive and negative
171 modes. The scan mode was between 100-800 m/z, with a scan resolution of 13 000 m/z/s until the ICC
172 target reached either 20 000 or maximum accumulation time of 200 ms. The MS instrument operated
173 with nitrogen as drying gas at a temperature of 330 °C (flow rate of 9 L min^{-1}). Ionization voltage of the
174 electrospray capillary source was 4000 V and tandem MS was carried out using helium as the collision
175 gas (4.21×10^{-6} mbar) with 1 V collision voltage. MS identification of metabolites was according to
176 mass spectra, product ion spectra, retention time and confirmed with authentic standards and published
177 data.

178 *2.6.2 Individual phenolic compound quantification by HPLC-DAD.* The quantification of
179 phenolic compounds was performed by HPLC-DAD according to previously published methods (Di
180 Stefano and Cravero, 1992; Ferrandino and Guidoni, 2010). Stationary phase column was Licrosphere
181 100 RP5 μm) packed into LiChroCART 250-4 (25 \times 0.4 cm ID) HPLC-Cartridge (Merck KGaA,
182 Germany) with guard column (LiChroCART 4-4). Mobile phase solvent A was phosphoric acid 10^{-3} M
183 and solvent B was pure methanol. Run time was 50 min, temperature 25 °C and DAD peaks were
184 detected at 280 nm, 320 nm and 360 nm. Compounds were identified based on compliance with data
185 obtained from available pure standards and quantified based on standard curve constructed per each
186 individual molecule. Among flavanone, as exclusively dihydroquercetin-rhamnoside (astilbin) was
187 available as commercial standard, semi-quantification of individual compounds was carried out using
188 the astilbin standard curve. The average flavonol profile of individual genotype was calculated averaging
189 results of five sampling dates.

190 **2.7 Statistical analysis.**

191 All data are averages of three biological replicates and standard errors. The analysis of variance was
192 performed by one-way ANOVA with IBM SPSS Statistics software program version 24.0 for Windows

193 (SPSS Inc., Chicago, IL). In case of significant differences ($P \leq 0.05$), means were compared by Tukey-
 194 b post-hoc test. Results related to leaf blade and vein polyphenolic composition taken separately were
 195 used to run a series of principal component analysis (PCA); correlated variables were progressively
 196 excluded, when this correlation had a biological meaning. Species and dates of sampling were separated
 197 on the basis of specific classes of polyphenols (i.e. variables). PCA analysis was performed by SAS 9.4
 198 for Windows (SAS Institute Inc., Cary, NC, US).


199

200 Table 1. Origin and botanical sub-generic series of the studied genotypes.

Plant name	Abbreviation	Origin	Series ^a
<i>Vitis candicans</i>	CAN	American	<i>Candicansae</i>
<i>Vitis riparia</i> cv Gloire de Montpellier	RIP	American	<i>Ripariae</i>
<i>Vitis rupestris</i> cv du Lot	RUP	American	<i>Ripariae</i>
<i>Vitis berlandieri</i>	BER	American	<i>Cinereae</i>
Börner (<i>V. riparia</i> x <i>V. cinerea</i>)	BOR	American	<i>Ripariae X Cinereae</i>
<i>Vitis coignetiae</i>	COI	Asian	<i>Lambruscae</i>
<i>Vitis amurensis</i>	AMU	Asian	<i>Flexuosae</i>
<i>Vitis vinifera</i> subsp. <i>sylvestris</i>	SYL	Eurasian	<i>Viniferae</i>

201 ^aSeries were defined according to Galet (1988).

202



203
204 **Figure 1.** Leaves of eight *Vitis* genotypes.

205

206 **3. RESULTS**

207 Significant differences of water content occurred among *Vitis* species at the same picking time
 208 (Table 2); however, differences did not exceed 7% in blades and 8% in veins. Within the species, water
 209 content evolution during the vegetative season was not higher than 6%; in blades of *V. candicans*, *V. v.*
 210 *sylvestris*, Börner and *V. coignetiae* dry matter (DM) content changes during the season were never
 211 significant. The comparison between averages of all analyzed grapevine blades and veins showed that
 212 dry matter content in veins was around 10% lower than in blades. Concentrations of compounds were
 213 calculated and expressed as amounts per dry matter, even though leaf extracts were prepared from fresh
 214 leaves: this choice was done to minimize the well detailed polyphenol analysis perturbations due to
 215 losses or to chemical alterations during preparation (Abascal et al., 2005).


216

217 Table 2. Dry matter (%) in leaves of *Vitis* genotypes during the vegetative season ^a

	DOY 186		DOY 200		DOY 241		date
		#	Blades	#		#	Sampling dates
AMU	32.09 ± 0.03a	e	34.43 ± 0.58b	c	35.29 ± 0.23b	bc	*
CAN	29.71 ± 0.30	cde	29.68 ± 0.18	ab	28.94 ± 1.63	a	ns
RIP	27.78 ± 0.69a	ab	28.05 ± 0.79a	a	31.95 ± 1.21b	ab	*
RUP	27.35 ± 0.75a	a	27.16 ± 0.80a	a	33.51 ± 1.01b	b	**
BER	31.45 ± 0.18a	de	31.86 ± 1.07a	bc	37.57 ± 0.61b	c	**
COI	29.59 ± 0.43a	bc	32.09 ± 0.70b	bc	32.93 ± 0.53b	ab	*
SYL	31.39 ± 0.62	de	29.70 ± 0.83	ab	31.76 ± 0.61	ab	ns
BOR	30.57 ± 0.66	de	31.49 ± 0.74	bc	32.88 ± 0.35	ab	ns
average species	29.99 ± 0.37	**	30.56 ± 0.51	**	33.10 ± 0.56	**	
			Veins				
AMU	19.01 ± 0.42a	ab	22.61 ± 0.47b	bc	22.32 ± 0.33b	b	**
CAN	23.14 ± 0.70b	b	23.61 ± 0.99b	c	19.32 ± 0.60a	a	*
RIP	21.21 ± 0.71ab	b	20.52 ± 0.52a	ab	22.88 ± 0.21b	bc	*
RUP	19.23 ± 1.43a	ab	20.02 ± 0.52a	ab	25.13 ± 0.34b	cd	**
BER	21.37 ± 0.40a	ab	23.08 ± 1.12a	bc	26.30 ± 0.58b	d	*
COI	17.82 ± 0.53	a	18.89 ± 0.72	a	19.27 ± 0.66	a	ns
SYL	21.22 ± 0.47a	ab	23.74 ± 0.15b	c	27.28 ± 0.48c	d	**
BOR	20.59 ± 0.45a	ab	20.26 ± 0.33a	ab	23.88 ± 0.81b	bc	**
average species	20.45 ± 0.38	*	21.59 ± 0.42	**	23.30 ± 0.60	**	

218 ^a Results are expressed as means ± standard errors (SE; n=3). Different small letters in each individual row after averages ±
 219 SE represent statistical differences among sampling dates, within the same species. In columns individuated with #, different
 220 letters represent statistical differences among genotypes, within the same tissue and date. General statistical differences
 221 among sampling dates and species were assessed by a post-hoc Tukey-b test for P ≤ 0.05 (*), P ≤ 0.01 (**); ns – not
 222 significant. Day of the year (DOY) refers to 5th of July (DOY 186); 19th of July (DOY 200); 29th of August (DOY 241).

223
 224
 225 The total polyphenol (TP) content ranged from 60.2 to 165.9 g kg⁻¹ DW in blades and from 28.6 to 130.1
 226 g kg⁻¹ DW in veins. *V. amurensis* blades (Figure 2) displayed higher TP concentration during the entire
 227 vegetative season compared to the other genotypes. TP was on average about 1.5 times lower in veins
 228 than in blades. TP accumulation trend was different among **genotypes**: in *V. riparia*, the accumulation
 229 trends in veins and blades were similar, whereas in *V. candicans*, TP concentration increased in blades
 230 and decreased in veins during the season.



231

232 **Figure 2.** Total polyphenol (TP) accumulation in leaves of eight *Vitis* genotypes during the vegetative
 233 season; A = blades and B = veins. Means of three replicates \pm SE.

234

235 **3.1 Phenolic Compound Accumulation in *Vitis* sp. Leaves.** Twenty phenolic compounds were
 236 identified in the two Asian species, 27 in the four American species and in the hybrid, and 18 in *V. v.*
 237 *sylvestris*. The detected compounds belonged to the groups of flavonols, phenolic acids, flavan-3-ols,
 238 proanthocyanidins, flavanonols and flavones. Anthocyanins were exclusively detected in trace amounts
 239 or in extremely low concentrations (data not shown) and, in line with what we previously found in
 240 varieties of *V. vinifera sativa* (Kedrina-Okutan et al., 2018), in healthy leaves they were found
 241 exclusively in veins.

242 3.1.1 Flavonols.


243 The concentration of flavonol glycosides ranged from 3.6 to 20.6 g kg⁻¹ DW in blades and from
 244 0.8 to 7.7 g kg⁻¹ DW in veins (Figure 3); they were the most abundant phenolic compounds in leaves.
 245 We detected a net separation among genotypes based on their ability to accumulate flavonol glycosides
 246 during the season. A first group included *V. amurensis* and *V. riparia*, both species displaying the highest

247 total flavonol concentration, with a peak at DOY 196 in *V. riparia* and at DOY 215 in *V. amurensis*
248 (Figure 3A). A second group with medium flavonol concentration included *V. v. sylvestris* and *V.*
249 *rupestris* with a stable concentration trend during the vegetative season. A third group included the
250 remaining species, displaying relatively low total flavonol concentration and slightly decreasing trends
251 during the examined period. Similar results were found in veins: genotypes of the first two groups
252 displayed higher total flavonol concentration and genotypes of the third group lower concentration with
253 decreasing trend during the season (Figure 3B). The comparison between the two compartments within
254 each genotype showed that total flavonol concentration was always higher in blades than in veins. The
255 prevalent flavonol was quercetin 3-*O*-glucuronide, ranging between 60% and 92% of all the detected
256 flavonols (Figure 4; Supplementary Table 1A,B). The most complex flavonol profile was found in *V.*
257 *v. sylvestris* leaves accumulating seven out of the eight identified flavonol glycosides and trace amount
258 of kaempferol 3-*O*-glucoside (Figure 4). Kaempferol 3-*O*-rhamnoside was exclusively found in *V. v.*
259 *sylvestris* that also accumulated significant amounts of quercetin 3-*O*-rhamnoside. The simplest flavonol
260 profile was found in *V. amurensis* and *V. candicans* that exclusively accumulated quercetin and
261 kaempferol glucuronides and glucosides. In *V. berlandieri* and in Börner, no kaempferol glycosides
262 were detected.


263 3.1.2 Phenolic acids.

264 Total phenolic acid content in blades ranged from 3.2 to 10.2 g kg⁻¹ dry weight and showed
265 different accumulation trends among genotypes. In Asian species, *V. amurensis* and *V. coignetiae*, total
266 phenolic acid concentration fell during the vegetative season. In American species, concentration firstly
267 increased and then it started to fall between July and beginning of August (DOY 173-215), whereas in
268 *V. v. sylvestris*, concentration decreased between the first and the second sampling date and then rose
269 from the end of June (DOY 173) to the beginning of August (DOY 215) (Figure 5A). In leaf veins,
270 phenolic acid content showed a significant drop between the first and the second sampling date in all
271 genotypes, followed by a relatively steady period (Figure 5B). Total phenolic acid concentration in veins
272 was lower than in blades in each individual species, except at the first sampling date in *V. candicans*, *V.*


273 *riparia* and *V. rupestris* when concentration in veins was higher than in blades. Hydroxycinnamoyl
 274 tartaric acids were the prevalent phenolic acids in leaves, and, based on their characteristic UV maximum
 275 absorbance around 320 nm and mass spectra, they were identified as *cis*- and *trans*-forms of caftaric,
 276 coutaric and fertaric acid (Table 3). *Trans*-forms were always prevalent over *cis*-forms. The main
 277 phenolic acid in leaves was *trans*-caftaric acid, which comprised up to 90% followed by *trans*-coutaric
 278 and *trans*-fertaric acid (Figure 5C, Supplementary Table 2A, B). On the contrary, in *V. rupestris* and *V.*
 279 *riparia* *trans*-fertaric acid displayed higher concentration respect to *trans*-coutaric acid. Moreover, these
 280 two species accumulated an additional molecule, a coutaric acid isomer, that was tentatively identified
 281 through its pseudomolecular ion $[M-H]^-$ at m/z 295 and two product ions at m/z 163, 131 in MS^2 . Low
 282 amounts of protocatechuic acid-glucoside were detected in *V. rupestris*, *V. riparia*, *V. amurensis* and
 283 Börner blades, with a slight increasing trend during the season (Supplementary Table 2A).



284
 285 **Figure 3.** Total flavonol concentration in leaves of eight *Vitis* genotypes during the vegetative season;
 286 A = blades and B = veins. Means of three replicates \pm SE.



287
 288 **Figure 4.** Flavonol profile of leaf blades of eight *Vitis* genotypes calculated as average of results
 289 obtained at five sampling times performed during the vegetative season.



290

291 **Figure 5.** Accumulation of phenolic acids in *Vitis sp.* leaves during the vegetative season. Total phenolic
 292 acid concentration in blades (A) and veins (B); results are means of three replicates \pm SE. Phenolic acid
 293 profile of leaf blades (C) and veins (D) during the season. For species acronyms see Table 1; numbers
 294 after the species acronyms refer to 1 = DOY 148 (28th of May); 2 = DOY 173 (22nd of June); 3 = DOY
 295 195 (14th of July); 4 = DOY 215 (3rd of August); and 5 = DOY 240 (28th of August).

296

297 Table 3. Phenolic compound identification in leaf blades and veins of eight *Vitis* genotypes by HPLC-ESI-
 298 MS/MS

ID	Rt (±0.2 min)	[M] ⁻ (m/z)	[MS ²] ⁻ (m/z)	[M] ⁺ (m/z)	[MS ²] ⁺ (m/z)	compound identification
1	14.5	593	425, 407, 289	595	291	(epi)galocatechin-(epi)catechin
2	16.5	315	153, 123			protocatechuic acid-glucoside
3	18.3	311	179			<i>cis</i> -caftaric acid
4	18.8	577	451, 407, 289			procyanidin B ₃
5	19.9	311	179			<i>trans</i> -caftaric acid
6	21.8	289	245, 205, 179	291	273, 165, 123	catechin
7	22.5	295	163			<i>cis</i> -coutaric acid
8	23.0	577	451, 425, 407, 289	579	427, 409, 289	procyanidin B ₁
9	23.5	295	163			<i>trans</i> -coutaric acid
10	23.8	729	577, 441, 289	731	579, 409	procyanidin dimer gallate
11	24.9	595	576, 385, 355			unknown proanthocyanidin
12	25.1	325	193			<i>trans</i> -fertaric acid
13	25.7	289	245, 205, 179	291	273, 165, 123	epicatechin
14	27.7	465	303, 285, 151			dihydroquercetin-hexoside
15	27.7	295	163, 131			coutaric acid isomer
16	29.2	615	481, 407, 359			unknown
17	30.8	447	393, 357, 327	449	413, 383, 329	luteolin 8- <i>C</i> -glucoside (orientin)
18	31.1	449	287, 269, 151			dihydrokaempferol hexoside
19	31.4	447	429, 357, 327, 285	449	431, 413, 383	luteolin 6- <i>C</i> -glucoside (isoorientin)
20	31.8	479	317	481	319	myricetin 3- <i>O</i> -glucoside
21	32.3	431	341, 311, 283	433	415, 367, 283	apigenin 8- <i>C</i> -glucoside (vitexin)
22	32.7	493	317	495	319	myricetin 3- <i>O</i> -glucuronide
23	33.7	449	303, 285, 151			dihydroquercetin-rhamnoside (astilbin)
24	33.8	431	413, 341, 311, 283	433	415, 367, 337	apigenin 6- <i>C</i> -glucoside (isovitexin)
25	34.2	447	285	449	287	luteolin-7- <i>O</i> -glucoside (luteoloside)
26	35.1	463	301	465	303	quercetin 3- <i>O</i> -glucoside
27	35.6	477	301	479	303	quercetin 3- <i>O</i> -glucuronide
28	37.1	447	301	449	303	quercetin 3- <i>O</i> -rhamnoside
29	37.2	447	285	449	287	kaempferol 3- <i>O</i> -glucoside
30	37.5	461	285	463	287	kaempferol 3- <i>O</i> -glucuronide
31	38.3	431	285	433	287	kaempferol 3- <i>O</i> -rhamnoside

299 ^aID numbers correspond to peaks reported in Supplement Figure 1, 2.

300

301 *3.1.3 Flavan-3-ols.*

302 Important quantities of total flavan-3-ols were detected in grapevine leaves. The concentration

303 of flavan-3-ols was rather similar between blades and veins, oppositely to flavonols and phenolic acids

304 that prevalently accumulated in blades. Specifically, the concentration of flavan-3-ols ranged from 0.44

305 to 3.39 g kg⁻¹ in blades and from 0.04 to 1.55 g kg⁻¹ in veins (Figure 6A and B). In *V. amurensis* blades,

306 the concentration of total flavan-3-ols was significantly higher respect to the other genotypes, and it

307 displayed a decreasing trend, so that at the end of the examined period the differences respect to the
308 other genotypes were less marked. In *V. rupestris* flavan-3-ol concentration increased almost
309 continuously during the entire vegetative season, oppositely to the other analyzed genotypes. The flavan-
310 3-ol profile analysis showed that (+)-catechin was generally the prevalent flavan-3-ol, corresponding up
311 to 73 % of total flavan-3-ols in blades (Figure 5C; Supplementary Table 3A) or the exclusive one in *V.*
312 *coignetiae* veins (Figure 5D; Supplementary Table 3B). In the Asian species *V. amurensis*, (-)-
313 epicatechin was the prevalent flavan-3-ol in blades, similarly to *V.v. sylvestris* and *V. rupestris* veins
314 from the second sampling date onwards. Concentration of (-)-epigallocatechin showed an increasing
315 trend during the examined season and in blades it accounted up to 38% by the end of the season.
316 Interestingly, in veins of *V. coignetiae* and Börner, (-)-epicatechin was absent during the whole season,
317 and in *V. candicans* it was exclusively detected at the first sampling date. The relative abundance of (-
318)-epicatechin gallate ranged between 4.7% and 14.9% and it was exclusively found in blades of *V.*
319 *rupestris*, *V. berlandieri* and Börner.


320 3.1.4 Proanthocyanidins.

321 Proanthocyanidin concentration ranged from 0.21 to 3.09 g kg⁻¹ in blades. Particularly high
322 concentration was detected in *V. amurensis* at first four sampling dates and in *V. berlandieri* at first two
323 sampling dates, followed by a sharp concentration fall (Figure 7A). In veins, PA exclusively
324 accumulated in *V. amurensis* and *V. candicans* during the whole season and in some dates in four
325 species, also in *V. coignetiae* and *V. v. sylvestris*. PA content was generally higher in blades than in
326 veins, with the only exception of *V. amurensis* and *V. candicans* at DOY 148 (end of May) when vein
327 PA content was higher respect to that of blades (Figure 7; Supplementary Table 4A,B). Procyanidin B₁
328 accumulated both in blades and veins. Additionally, in *V. amurensis*, *V. berlandieri* and in Börner blades
329 and in *V. amurensis* veins, a procyanidin dimer gallate was tentatively identified by pseudomolecular
330 ion [M-H]⁻ at *m/z* 729; product ions at *m/z* 577, 441, 289 in MS² and characteristic UV maximum
331 absorbance at 280 nm, in line with other report (Flamini, 2013). In *V. berlandieri* and Börner blades,

332 procyanidin dimer gallate accounted for around 50% of total PA, whereas in *V. amurensis*, it accounted
333 for 20-25 % of total PA during the vegetative season.


334 Furthermore, by HPLC-ESI-MS/MS (epi)gallocatechin-(epi)catechin was tentatively identified
335 in blades (pseudomolecular ion $[M-H]^-$ at m/z 593; product ions at m/z 425, 407, 289 in MS^2 ; Table 3),
336 In addition, in *V. amurensis* blades, another B-type of procyanidin was identified, most likely
337 procyanidin B₃ and one further proanthocyanidin that remains unknown (Table 3).

338



339

340 **Figure 6.** Accumulation of flavan-3-ols in *Vitis sp.* leaves during the vegetative season. Total flavan-3-
 341 ol concentration in blades (A) and veins (B); results are means of three replicates \pm SE. Flavan-3-ol
 342 profile of leaf blades (C) and veins (D) during the season; see Figure 5 for genotype acronym
 343 specification.



344


345 **Figure 7.** Accumulation of proanthocyanidin in *Vitis sp.* leaves (A = blades; B = veins) during the
346 vegetative season. Means of three replicates \pm SE; see Figure 5 for genotype acronym specification.

347

348 3.1.5 Flavanonols.

349 In blades, flavanonols were detected exclusively in *V. amurensis* and *V. coignetiae*, whereas in
350 veins, they were also found in *V. v. sylvestris*. (Figure 8; Supplementary Table 4A, B). In blades,
351 concentration was much lower than in veins. Particularly high concentration was found in *V. amurensis*
352 veins with values ranging between 0.95 and 2.12 g kg⁻¹ (Figure 8B). In both Asian species, vein total
353 flavanonol concentration showed two peaks, the first at DOY 148 and the second at DOY 215 whereas
354 in *V. v. sylvestris* no differences among sampling times were noticed. Three flavanonol glycosides were
355 quantified and identified in leaves by the characteristic absorbance maximum at 290 nm and
356 correspondent mass spectra (Table 3): dihydroquercetin-hexoside was the main flavanonol found in *V.*

357 *amurensis* and *V. coignetiae* where it was the exclusive flavanone in blades and it accounted for up to
 358 90% of total flavanones in veins (Figure 8A,B). In addition, dihydrokaempferol-hexoside was detected
 359 in *V. amurensis* and in *V. coignetiae* veins with no differences among sampling dates. Astilbin
 360 (dihydroquercetin-rhamnoside) was exclusively detected in the Eurasian species *V. v. sylvestris* veins.
 361




362
 363 **Figure 8.** Accumulation of flavanone content in *Vitis* sp. leaves (A = blades; B = veins) during the
 364 vegetative season. Means of three replicates \pm SE; DHQuer - dihydroquercetin; DHKaemp -
 365 dihydrokaempferol; hex - hexoside; rham - rhamnoside; see Figure 5 for genotype acronym
 366 specification.

367
 368 *3.1.6 Flavones*
 369 Flavones have rarely been detected in leaves and other vegetative organs of grapevine. In fact,
 370 also in our analyzed genotypes, flavones accumulated exclusively in *V. berlandieri* and Börner. Total
 371 flavone concentration ranged between 2.5 and 4.9 g kg⁻¹ DW in blades (Figure 9A). In comparison with
 372 *V. berlandieri*, Börner blades accumulated significantly higher concentration of total flavones from
 373 DOY 148 (end of May) until DOY 195 (mid-July). Similar to the other classes of measured polyphenols,
 374 flavone concentration was lower in veins respect to blades, ranging from 0.7 to 1.2 g kg⁻¹ DW (Figure
 375 9B). Five flavones accumulated in both tissues, two were apigenin glycosides and three luteolin
 376 glycosides. Isomers of apigenin glucosides were differentiated by pseudomolecular ion [M - H]⁻ at *m/z*
 377 431 and absorbance maximum at 270 nm and 337 nm. Moreover, the product ions of luteolin glycoside

378 and their intensities provided insight into individual compound structure. Apigenin 8-*C*-glucoside
 379 (isovitexin) product ions were detected at m/z 283, 311, 341 and apigenin 6-*C*-glucoside (vitexin)
 380 product ions were detected at m/z 283, 311, 341, 413 (Table 3). The absence of product ion m/z 413 and
 381 the lower intensity of product ion at m/z 341 $[^{0,3}X_0 - H]^-$ helped to differentiate apigenin 8-*C*-glucoside
 382 from apigenin 6-*C*-glucoside, accordingly to Kim et al. (2018). Isomers of luteolin glucoside had
 383 pseudomolecular ion $[M - H]^-$ at m/z 447 and absorbance maximum at 269, 350 nm. Furthermore,
 384 comparison of product ion pattern of three luteolin glucoside isomers let us tentatively identify luteolin
 385 8-*C*-glucoside (isoorientin), luteolin 6-*C*-glucoside (orientin) and luteolin 7-*O*-glucoside (luteolin)
 386 (Table 3). The compound with absence of product ion at m/z 429 and high intensity of product ion at
 387 m/z 357 $[^{0,3}X_0 - H]^-$ was tentatively identified as luteolin 6-*C*-glucoside. Luteolin glucoside isomer with
 388 product ion pattern of two high abundance fragment at m/z 285 $[M - H - 162]^-$ and m/z 284 $[M - H -$
 389 $162]^-$ was tentatively identified as luteolin 7-*O*-glucoside, according to Li et al. (2016). Leaf flavone
 390 profile analysis showed that isoorientin together with isovitexin comprised up to 70% of all detected
 391 flavones, thus 8-*C* was the predominant glycosylation site of flavone aglycones in *Vitis sp.* leaves
 392 (Figure 9; Supplementary Table 5).

393



394

395 **Figure 9.** Accumulation of flavones in *Vitis sp.* leaves (A = blades; B = veins) during the vegetative
 396 season. Means of three replicates \pm SE; see Figure 5 for genotype acronym specification.

397


398 **3.2 Overall polyphenolic patterns and genotype associations**

399 In blades, the PCA model that explained the highest variance with three PRINs (72 %), allowing
400 a net separation of genotypes, based on eleven variables (Table 4) that were gradually chosen to exclude
401 redundancy. Variables associated to the first PRIN were phenolic acids (the sum of fertaric acid, coutaric
402 acid isomer and the *cis* and *trans* forms of caftaric acid) and the total concentration of dimeric
403 proanthocyanidins. Genotypes separated on the second PRIN on the basis of myricetin glycoside
404 (glucoside + glucuronide) concentration and of flavonol total amount. The third PRIN separated
405 genotypes based on total flavan-3-ol concentration, evidencing in particular, *V. amurensis*. Considering
406 that the eigenvalue on PRIN3 was almost 2.0 (1.79, Table 4), the flavone total concentration could be
407 considered as the second variable associated (negatively) to the third PRIN. On this basis, Börner and
408 *V. berlandieri*, highly associated to the third PRIN negative values, evidenced their peculiar capability
409 to accumulate high amounts of flavones in leaf blades (Figure 9; Figure 10).

410 In veins, the proposed PCA model (Table 5; Figure 11) explained the 75 % of total variance with
411 the first three PRINs. Variables associated to the first PRIN were total polyphenols, total flavan-3-ols
412 and flavonol glucuronic derivatives; these variables did not markedly identify varieties or specific time-
413 points of the leaf vegetative cycle, with the only exception of *V. amurensis*, generally associated to
414 positive values of the first PRIN. Flavanonols and glucosides of flavonols were negatively associated to
415 the second PRIN. Dimeric proanthocyanidins associated to the third PRIN, allowed to evidence that in
416 genotypes accumulating this class of compounds, concentration was higher at the first sampling
417 (particularly in *V. amurensis*, *V. candicans*, *V. berlandieri* and in Börner).

418 **3.3 Seasonal polyphenol-related traits**

419 The flavonol concentration varied during the vegetative season, but the profile, strictly related
420 to a specific genotype, did not change (Figure 4).




446
 447 **Figure 10.** Tridimensional distribution of *Vitis* species during the vegetative season (1 to 5) on the first
 448 three principal components, obtained by polyphenol quantifications and profiles of leaf blade extracts
 449 (see Table 4 for the list of used variables).

450
 451 **Table 5.** Eigenvectors of nine variables (polyphenols in leaf veins) on the first three principal
 452 components (PRIN1, PRIN2 and PRIN3).

	PRIN1	PRIN2	PRIN3
<i>Total polyphenols</i>	0.50	0.11	-0.14
<i>Total phenolic acids</i>	0.36	0.39	0.25
<i>Total flavan-3-ols</i>	0.38	0.32	0.38
<i>Dimeric proanthocyanidins</i>	0.30	-0.30	0.52
<i>Total flavanonols</i>	0.18	-0.56	0.21
<i>Total flavones</i>	-0.22	0.35	0.28
<i>Sum of glucuronide flavonols</i>	0.38	0.14	-0.33
<i>Sum of glucoside flavonols</i>	0.36	-0.38	-0.29
<i>Sum of rhamnoside flavonols</i>	0.15	0.21	-0.42
Eigenvalues	3.27	2.17	1.34
Total variance	0.36	0.24	0.15

466 Eigenvalues of the three PRINs and their contribution to total variance. In bold letters, the variables associated to the
 467 appropriate PRIN.



468
 469 **Figure 11.** Tridimensional distribution of *Vitis* species during the vegetative season (1 to 5) on the first
 470 three principal components, obtained by polyphenol quantifications and profiles of leaf vein extracts
 471 (see Table 5 for the list of variables used).
 472

473 4. DISCUSSION

474 There is increasing interest in grapevine leaf polyphenols to deepen knowledge about their
 475 biological role in plant-defense mechanisms and to answer to the growing demand of natural bioactive
 476 compounds for the feed/food, pharmaceutical and cosmetic sectors. Grapevine leaves are already
 477 employed in the production of food ingredients, dietary supplements, pharmaceutical products and
 478 medicated cosmetics (Dani et al., 2010). The evolution of the polyphenolic composition of eight *Vitis*
 479 genotypes during the vegetative season allowed pointing out significant qualitative and quantitative
 480 differences.

481 4.1 Genotype-related specific traits of leaf polyphenols

482 The two studied American species, *V. riparia* and *V. rupestris*, belonging to the same subgeneric
 483 group of *Ripariae* (Table 1), involved in crossbreed of Regent, a cultivar displaying low susceptibility
 484 to *Plasmopara viticola* and to *Erysiphe necator*, shared common phenolic composition and
 485 accumulation trends. In particular, they showed similarities in the accumulation trend of total phenolic

486 acids and total flavan-3-ols. Additionally, they displayed identical flavonol profiles (Figure 4). However,
487 *V. riparia* accumulated higher concentration of total polyphenols, total flavonols and total phenolic acids
488 respect to *V. rupestris*, which could justify its specifically high resistance to downy mildew, as both
489 flavonols and phenolic acids have been associated with the ability to limit this pathogen diffusion (Ali
490 et al., 2012). Moreover, *V. riparia* total flavonol concentration both in blades and in veins was the
491 highest comparing to all the other analyzed genotypes, almost during the entire vegetative season (Figure
492 3).

493 The American genotypes Börner and *V. berlandieri*, known as low-susceptible to many diseases,
494 were classified by Galet (1988) into the same subgeneric serie of *Cinereae* (Table 1). Their main leaf
495 polyphenolic traits showed many similarities, such as the concentration and the accumulation trend of
496 total polyphenols, total flavonols and total flavones, together with identical profiles of flavonols,
497 proanthocyanidins and flavones. Therefore, Galet's classification based on plant morphology matched
498 with phenol profile patterns of these two genotypes, suggesting a phylogenetic link between *Vitis*
499 *cinerea* (Börner parent) and *Vitis berlandieri*. Flavones, bioactive compounds often extracted from
500 medical and herbaceous plants and involved in plant signaling and defense responses (Jiang et al., 2016),
501 were exclusively detected in Börner and in *V. berlandieri*, among the studied genotypes. In grapevine,
502 they are generally considered as minor compounds mostly represented by luteolin, but in Börner and *V.*
503 *berlandieri* leaves their concentration reached 5 g kg⁻¹ DW (Figure 9) and showed a much wider profile,
504 consisting of five apigenin and luteolin derivatives: specifically, four of them were *C*-glycosides and
505 one *O*-glycoside (Table 3; Figure 9). *C*-glycoside flavonoids generally display higher antioxidant
506 potential and better therapeutic properties compared to aglycons or to *O*-glycosylated flavonoids (Xiao
507 et al., 2016). However, as most studies focused on biological activities of the more common plant *O*-
508 glycosylated-flavonoids, further research is required on grapevine *C*-glycosylated derivative health
509 beneficial effect. McNally et al. (2003) demonstrated that in cucumber *C*-flavones acted as phytoalexins
510 in response to powdery mildew fungus *Podosphaera xanthii* and concentration increases of orientin and
511 isoorientin were reported to be induced by soil salinity in buckwheat (*F. esculentum*) (Yang et al., 2018).

512 *V. berlandieri* and Börner were also similar in the fact that they did not accumulate kaempferol
513 glycosides, in line with what was previously demonstrated in the grapevine series *Cinerea* (Moore and
514 Giannasi, 1987).

515 The Asian wild species *V. coignetiae* has been cultivated during the last two decades in Japan
516 for wine, juice and jam making (Kamiya et al., 2018). *V. coignetiae* leaf extract showed a strong radical-
517 scavenging activity and potential hepato-protective effect on nonalcoholic steatohepatitis in liver
518 (Takayama et al., 2009). However, in *V. coignetiae* leaves of the present study, the concentration of leaf
519 polyphenols was generally lower comparing to the other genotypes (Figure 2). *V. coignetiae* along with
520 *V. amurensis* and *V. v. sylvestris* was one of the few species which accumulated flavanonols; two
521 flavanone compounds were detected in *V. coignetiae*, dihydrokaempferol-hexoside in veins and
522 dihydroquercetin-hexoside in both blades and veins (Figure 8A, B). Taxifolin (dihydroquercetin)
523 glycoside is known to have anti-inflammatory activity (Kim et al., 2008) and potential positive effect
524 on atopic dermatitis treatment (Ahn et al., 2010).

525 The most peculiar species among the analyzed genotypes was *V. amurensis*, which is known for
526 its cold tolerance and disease resistance to downy mildew (Jürges et al., 2009), anthracnose and white
527 rot (Li et al., 2008). The leaves of *V. amurensis* reached the highest concentration of total polyphenols
528 and flavan-3-ols in blades, important amounts of flavonols, and the highest concentration of flavanone
529 in veins. *V. amurensis* leaves are used in conventional Chinese medicine and included in Korean herbal
530 Pharmacopoeia (Chen et al., 2018). Moreover, leaf extracts showed antimicrobial activity against
531 *Streptococcus mutans* and *Streptococcus sanguis* (Yim et al., 2010) and neuroprotective effect (Jeong
532 et al., 2010). Bak et al. (2012, 2016) demonstrated that *V. amurensis* seed proanthocyanidins have
533 hepato-protective and antioxidative properties and a possible chemopreventive role in humans
534 hepatocarcinoma cells. In *V. amurensis* blades, concentration of total flavan-3-ols was significantly
535 higher during the entire vegetative season comparing to the other genotypes (Figure 6A). Moreover, also
536 the concentration of (-)-epicatechin was particularly high. This compound was previously individuated
537 as a reaction of pear leaves against *Erwinia amylovora* and of apple leaves against *Venturia inaequalis*

538 (as discussed in Kedrina-Okutan et al., 2018). In non-*Vitis vinifera* genotypes, (-)-epigallocatechin was
539 additionally detected with increasing concentration during the vegetative season (Figure 6).
540 (Epi)gallocatechin-(epi)catechin was tentatively identified in *V. amurensis*, consistently with previously
541 published dimeric PA identification (Flamini, 2013). (Vagiri et al., 2017) reported a negative correlation
542 between (-)-epigallocatechin and the presence of septoria leaf spot caused by *Mycosphaerella ribis* in
543 black currant leaves. The high concentration of constitutive polyphenolic compounds throughout the
544 season in *V. amurensis* leaves contributes to explain this biotype excellent ability to protect against
545 adverse environmental condition (frost and diseases). Furthermore, the abundance of polyphenols in *V.*
546 *amurensis* leaves emphasizes that this species has the potential for further use as a source of natural
547 bioactive compounds for nutraceutical and pharmacological uses.

548 The polyphenolic composition of *V. candicans* leaves was studied here for the first time, to the
549 best of our knowledge. This American species has been described as highly vigorous, highly resistant
550 to downy mildew and tolerant to drought and salt (Ollat et al., 2016). *V. candicans* leaves displayed a
551 low concentration of total polyphenols, total flavonols, but a relatively high concentration of
552 proanthocyanidins and this genotype was among the few able to accumulate procyanidin B₁ in veins
553 (Figure 7). Procyanidin B₁, the dimeric proanthocyanidin consisting in units of (-)-epicatechin and (+)-
554 catechin, has an ecological significance in protecting plants against pathogens, insect pests and larger
555 herbivores (Dixon et al., 2005). For instance, previous studies demonstrated that *B. cinerea* remained
556 quiescent in immature strawberry until B proanthocyanidins are present at an essential high
557 concentration: this effect is related to the proanthocyanidin inhibitory potential of the fungal
558 polygalacturonase (Jersch et al., 1989).

559 The Eurasian wild species *V. v. sylvestris*, the ancestor of the worldwide cultivated *V. vinifera*
560 *sativa*, accumulated relatively high concentration of total polyphenols and total phenolic acids (Figure
561 2; Figure 5). Among phenolic acids, the predominant non-flavonoid polyphenols, caftaric acid was the
562 prevalent form, representing up to 90% of total phenolic acids (Figure 5C), similarly to what was
563 previously assessed in *V. vinifera sativa* leaves (Kedrina-Okutan et al., 2018). Caftaric acid is an

564 important bioactive component found in chicory, in the medical plant *Echinacea purpurea* (Bel-Rhliid
565 et al., 2012) and, in *V. coignetiae* juice, where it exhibited anti-mutagenic and anti-inflammatory
566 properties (Kamiya et al., 2018). In *V. v. sylvestris* leaves, we found astilbin (Figure 8B), in
567 concentrations that were in line with those found in Cabernet Sauvignon, Grenache, Shiraz and Barbera,
568 that were much lower respect to those found in Nebbiolo and Pinot noir (Kedrina-Okutan et al., 2018).
569 Additionally, *V. v. sylvestris* displayed the most complex flavonol profile (Figure 4). In *V. v. sylvestris*
570 veins we identified and quantified seven flavonols out of the eight flavonol compounds
571 chromatographically separated; *V. v. sylvestris* was the exclusive genotype that accumulated kaempferol
572 3-*O*-rhamnoside (Figure 4, Table 3) and, compared to *V. vinifera sativa*, it accumulated additional
573 flavonol rhamnosides (kaempferol- and quercetin-3-*O*-rhamnoside), showing a wider flavonol-profile
574 diversification.

575 The peculiar traits of individual *Vitis* species leaf polyphenolic compositions were highlighted
576 by the proposed PCA models. In particular, *V. riparia* and *V. rupestris* displayed similar eigenvectors
577 on PRIN1 and PRIN3, highlighting their similar concentrations of dimeric proanthocyanidins, similar
578 percentage incidence of caftaric acid over total phenolic acids (on PRIN1) and similar concentration of
579 flavan-3-ols on PRIN3 (Table 4; Figure 10). *V. riparia* sharply distinguished on PRIN2 due its high
580 flavonol content (Figure 3A) and the highest percentage incidence of myricetin derivatives respect to
581 the other analyzed genotypes (Figure 4). Flavonols allowed the species separation, also when
582 considering their concentration: in fact, flavonol glucosides, together with flavanonols, allowed a sharp
583 separation of *V. amurensis* samples and were negatively associated to the second PRIN of the vein PCA
584 model (Figure 11; Table 5). Oppositely, *V. berlandieri* and Börner associated to positive values of
585 PRIN2, with reduced amounts of flavonol-glucosides comparing to the other species. **Quercetin 3-*O*-**
586 **glucuronide was the prevalent flavonol in the analyzed genotypes, consistently with other studies on *V.***
587 ***vinifera* and *V. labrusca* leaves (Dresch et al., 2014; Kedrina-Okutan et al., 2018).** The differences in
588 the flavonol profile might be related to the genus *Vitis* evolution: in fact, evidences about the reduction
589 in structural complexity and in molecule diversification were reported (Moore and Giannasi, 1994), and

590 they were also ascribed to the flavonoid biosynthetic pathway, that was shown to have undergone a
591 simplification during evolution, particularly in domesticated genotypes. Interestingly, *V. v. sylvestris*,
592 which, being the wild form of *V. v. sativa*, underwent no or low selection pressure, showed the widest
593 flavonol profile diversification and comprised high concentration of quercetin 3-*O*-rhamnoside (22% of
594 total flavonols); besides, *V. v. sylvestris* also displayed a higher flavonol profile diversity respect to
595 cultivated varieties of *V. vinifera*. Quercetin 3-*O*-rhamnoside was also found in *V. berlandieri* and in
596 Börner. Hilbert et al. (2015) identified flavonol rhamnosides in berry skin extracts of *V. cinerea*, which
597 is native to the United States and belongs to the subgeneric group of *Cinereae* (Galet, 1988), the same
598 of *V. berlandieri* and Börner. Thus, flavonol rhamnosides seem to be genotype-specific compounds,
599 therefore they have the potential to be exploited as specific markers of some species and/or subspecies.

600 Vein flavanone concentration, associated to PRIN2 (Figure 11; Table 5) allowed the sharp
601 separation of *V. amurensis*, and, secondarily the separation of *V. coignetiae* and *V. v. sylvestris*. Since
602 flavanones accumulate prevalently in leaf veins, their concentration on a per whole leaf basis is
603 supposed to be low, considering the limited percentage incidence of veins on the total leaf area, but the
604 role of these tissue-specific compounds arouses the interest about their interaction with vein-located
605 pathogens. In some *V. vinifera sativa* varieties, we previously identified these molecules, particularly in
606 veins (Kedrina-Okutan et al., 2018) where astilbin was the predominant compound. We also associated
607 astilbin accumulation to a possible defense-mechanism in cv Nebbiolo (Ferrandino et al., 2019). In fact,
608 in Nebbiolo leaves of plants affected by *Flavescence dorée* (FD), we found a significantly higher
609 concentration of flavanones compared to healthy vines, hypothesizing a possible capability to limit the
610 pathogen development. Interestingly, together with *V. v. sylvestris*, also *V. amurensis*, whose capability
611 to defend against pathogens is detailed and *V. coignetiae*, known for its strong antiradical role,
612 accumulate flavanones.

613 As to the ecological role of leaf polyphenols, further studies would be of extreme interest to
614 understand: i) if genotype-related molecules such as flavanones and flavones have specific capacities

615 to limit diseases, starting from *in vitro* trials, ii) if grapevine plants can be induced to produce specific
616 molecules, once demonstrated that they can efficaciously serve to limit pathogen development.

617 As to the nutraceutical, pharmaceutical and feed/food ingredients, grapevine leaf are a source of
618 bioactive compounds that in future could be economically exploited, also exploiting their wide and still
619 little investigated, biodiversity.

620
621 ACKNOWLEDGMENTS. Authors gratefully acknowledge Tiziano Strano, DISAFA, University of
622 Turin, for managing the collection vineyard. Financial support from ‘Ricerca locale 2016
623 Caratterizzazione polifenolica di foglie di vite’, University of Turin.

624 REFERENCES

625 Abascal, K., Ganora, L., Yarnell, E., 2005. The effect of freeze-drying and its implications for botanical
626 medicine: a review. *Phyther. Res.* 19, 655–660. <https://doi.org/10.1002/ptr.1651>

627 Ahn, J.Y., Choi, S.E., Jeong, M.S., Park, K.H., Moon, N.J., Joo, S.S., Lee, C.S., Choi, Y.W., Kapsok,
628 L., Lee, M.-K., Lee, M.W., Seo, J.S., 2010. Effect of taxifolin glycoside on atopic dermatitis-like
629 skin lesions in NC/Nga mice. *Phyther. Res.* 24, 1071–1077. <https://doi.org/10.1002/ptr>

630 Ali, K., Maltese, F., Figueiredo, A., Rex, M., Margarida, A., Zyprian, E., Salomé, M., Verpoorte, R.,
631 Choi, Y.H., 2012. Alterations in grapevine leaf metabolism upon inoculation with *Plasmopara*
632 *viticola* in different time-points. *Plant Sci.* 191–192, 100–107.
633 <https://doi.org/10.1016/j.plantsci.2012.04.014>

634 Bak, M., Jun, M., Jeong, W., 2012. Procyanidins from wild grape (*Vitis amurensis*) seeds regulate ARE-
635 mediated enzyme expression via Nrf2 coupled with p38 and PI3K/Akt pathway in HepG2 cells
636 801–818. <https://doi.org/10.3390/ijms13010801>

637 Bak, M.J., Truong, V.-L., Ko, S.-Y., Nguyen, X.N.G., Ingkasupart, P., Jun, M., Shin, J.Y., Jeong, W.-
638 S., 2016. Antioxidant and hepatoprotective effects of procyanidins from wild grape (*Vitis*

639 *amurensis*) seeds in ethanol-induced cells and rats. Int. J. Mol. 17, 758.
640 <https://doi.org/10.3390/ijms17050758>

641 Bel-Rhlid, R., Page-Zoerkler, N., Fumeaux, R., Ho-Dac, T., Chuat, J.-Y., Sauvageat, J.L., Raab, T.,
642 2012. Hydrolysis of chicoric and caftaric acids with esterases and *Lactobacillus johnsonii* in vitro
643 and in a gastrointestinal model. J. Agric. Food Chem. 60, 9236–9241.
644 <https://doi.org/10.1021/jf301317h>

645 Chen, Q., Diao, L., Song, H., Zhu, X., 2018. *Vitis amurensis* Rupr: A review of chemistry and
646 pharmacology. Phytomedicine 49, 111–122. <https://doi.org/10.1016/j.phymed.2017.08.013>

647 Dani, C., Oliboni, L.S., Agostini, F., Funchal, C., Serafini, L., Henriques, J.A., Salvador, M., 2010.
648 Phenolic content of grapevine leaves (*Vitis labrusca* var. Bordo) and its neuroprotective effect
649 against peroxide damage. Toxicol. Vitro. 24, 148–153. <https://doi.org/10.1016/j.tiv.2009.08.006>

650 Di Stefano, R., Cravero, M.C., 1992. The separation of hydroxycinnamates in wine. Sci. des Aliment.
651 139–144.

652 Dixon, R., Xie, D.-Y., Sharma, S.B., 2005. Proanthocyanidins – a final frontier in flavonoid research?
653 New Phytol. 165, 9–28.

654 Dixon, R.A., 2001. Natural products and plant disease resistance. Nature 411, 843–847.
655 <https://doi.org/10.1038/35081178>

656 Dresch, R.R., Dresch, M.K., Guerreiro, A.F., Biegelmeyer, R., Holzschuh, M.H., Rambo, D.F.,
657 Henriques, A.T., 2014. Phenolic compounds from the leaves of *Vitis labrusca* and *Vitis vinifera* L.
658 as a source of waste byproducts: development and validation of LC method and antichemotactic
659 activity. Food Anal. Methods 7, 527–539. <https://doi.org/10.1007/s12161-013-9650-4>

660 Ferrandino, A., Guidoni, S., 2010. Anthocyanins, flavonols and hydroxycinnamates: An attempt to use
661 them to discriminate *Vitis vinifera* L. cv “Barbera” clones. Eur. Food Res. Technol. 230, 417–427.

662 <https://doi.org/10.1007/s00217-009-1180-3>

663 Ferrandino, A., Pagliarani, C., Kedrina-Okutan, O., Icardi, S., Bove, M., Lovisolo, C., Novello, V.,
664 Schubert, A., 2019. Non-anthocyanin polyphenols in healthy and Flavescence dorée infected
665 Barbera and Nebbiolo leaves. BIO Web Conf. 03003, 1–5.
666 <https://doi.org/https://doi.org/10.1051/bioconf/20191303003>

667 Flamini, R., 2013. Recent applications of mass spectrometry in the study of grape and wine polyphenols.
668 ISRN Spectrosc. 2013, 45. <https://doi.org/10.1155/2013/813563>

669 Galet, P., 1988. In *Cepages Et Vignobles De France* [Grapes and vineyards of France], 2nd ed.
670 Imprimerie Charles Dehan, Montpellier.

671 Gómez-Zeledón, J., Zipper, R., Spring, O., 2013. Assessment of phenotypic diversity of *Plasmopara*
672 *viticola* on *Vitis* genotypes with different resistance. Crop Prot. 54, 221–228.
673 <https://doi.org/10.1016/j.cropro.2013.08.015>

674 Hilbert, G., Tamsamani, H., Bordenave, L., Pedrot, E., Chaher, N., Cluzet, S., Delaunay, J.C., Ollat, N.,
675 Delrot, S., Mérillon, J.M., Gomès, E., Richard, T., 2015. Flavonol profiles in berries of wild *Vitis*
676 accessions using liquid chromatography coupled to mass spectrometry and nuclear magnetic
677 resonance spectrometry. Food Chem. 169, 49–58. <https://doi.org/10.1016/j.foodchem.2014.07.079>

678 Hmamouchi, M., Es-Safi, N., Lahrichi, M., Fruchier, A., Essassi, E.M., 1996. Flavones and flavonols in
679 leaves of some Moroccan *Vitis vinifera* cultivars. Am. J. Enol. Vitic. 47, 186–192.
680 <https://doi.org/10.1071/PP9960115>

681 Jeong, H.Y., Kim, J.Y., Lee, H.K., Ha, D.T., Song, K.S., Bae, K., Seong, Y.H., 2010. Leaf and stem of
682 *Vitis amurensis* and its active components protect against amyloid β protein (25-35)-induced
683 neurotoxicity. Arch. Pharm. Res. 33, 1655–1664. <https://doi.org/10.1007/s12272-010-1015-6>

684 Jersch, S., Scherer, C., Huth, G., Schlosser, E., 1989. Proanthocyanidins as basis for quiescence of

685 *Botrytis cinerea* in immature strawberry fruits. J. Plant Dis. Prot. 96, 365–378.

686 Jiang, N., Doseff, A., Grotewold, E., 2016. Flavones: from biosynthesis to health benefits. Plants 5, 27-
687 52. <https://doi.org/10.3390/plants5020027>

688 Jürges, G., Kassemeyer, H.H., Dürrenberger, M., Düggelin, M., Nick, P., 2009. The mode of interaction
689 between *Vitis* and *Plasmopara viticola* Berk. & Curt. Ex de Bary depends on the host species. Plant
690 Biol. 11, 886–898. <https://doi.org/10.1111/j.1438-8677.2008.00182.x>

691 Kamiya, T., Tanimoto, Y., Fujii, N., Negishi, T., Suzuki, T., Hatano, T., Arimoto-Kobayashi, S., 2018.
692 2,6-Dimethoxy-1,4-benzoquinone, isolation and identification of anti-carcinogenic, anti-mutagenic
693 and anti-inflammatory component from the juice of *Vitis coignetiae*. Food Chem. Toxicol. 122,
694 172–180. <https://doi.org/10.1016/j.fct.2018.10.028>

695 Kedrina-Okutan, O., Novello, V., Hoffmann, T., Hadersdorfer, J., Occhipinti, A., Schwab, W.G.,
696 Ferrandino, A., 2018. Constitutive polyphenols in blades and veins of grapevine (*Vitis vinifera* L.)
697 healthy leaves. J. Agric. Food Chem. 66, 10977–10990. <https://doi.org/10.1021/acs.jafc.8b03418>

698 Kim, B., Woo, S., Kim, M.J., Kwon, S.W., Lee, J., Sung, S.H., Koh, H.J., 2018. Identification and
699 quantification of flavonoids in yellow grain mutant of rice (*Oryza sativa* L.). Food Chem. 241,
700 154–162. <https://doi.org/10.1016/j.foodchem.2017.08.089>

701 Kim, Y.J., Choi, S.E., Lee, M.W., Lee, C.S., 2008. Taxifolin glycoside inhibits dendritic cell responses
702 stimulated by lipopolysaccharide and lipoteichoic acid. J. Pharm. Pharmacol. 60, 1465–1472.
703 <https://doi.org/10.1211/jpp/60.11.0007>

704 Li, D., Wan, Y., Wang, Y., He, P., 2008. Relatedness of resistance to anthracnose and to white rot in
705 Chinese wild grapes. Vitis 47, 213–215. <https://doi.org/10.1073/pnas.96.3.1146>

706 Li, Z.H., Guo, H., Xu, W. Bin, Ge, J., Li, X., Alimu, M., He, D.J., 2016. Rapid identification of flavonoid
707 constituents directly from PTP1B inhibitive extract of raspberry (*Rubus idaeus* L.) leaves by

708 HPLC-ESI-QTOF-MS-MS. J. Chromatogr. Sci. 54, 805–810.
709 <https://doi.org/10.1093/chromsci/bmw016>

710 Llorens, E., García-Agustín, P., Lapeña, L., 2017. Advances in induced resistance by natural
711 compounds: towards new options for woody crop protection. Sci. Agric. 74, 90–100.
712 <https://doi.org/10.1590/1678-992x-2016-0012>

713 McNally, D.J., Wurms, K. V., Labbé, C., Bélanger, R.R., 2003. Synthesis of C-glycosyl flavonoid
714 phytoalexins as a site-specific response to fungal penetration in cucumber. Physiol. Mol. Plant
715 Pathol. 63, 293–303. <https://doi.org/10.1016/j.pmpp.2004.03.005>

716 Moore, M.O., Giannasi, D.E., 1994. Foliar flavonoids of eastern North American *Vitis* (*Vitaceae*) north
717 of Mexico. Plant Syst. Evol. 193, 21–36.

718 Moore, M.O., Giannasi, D.E., 1987. Foliar flavonoids of selected *Vitis* taxa in the Southeastern United
719 States. Biochem Syst Ecol 15, 79–83.

720 Ollat, N., Bordenave, L., Tandonnet, J.P., Boursiquot, J.M., Marguerit, E., 2016. Grapevine rootstocks:
721 origins and perspectives. Acta Hortic. 1136, 11–22.
722 <https://doi.org/10.17660/ActaHortic.2016.1136.2>

723 Pastrana-Bonilla, E., Akoh, C.C., Sellappan, S., Krewer, G., 2003. Phenolic content and antioxidant
724 capacity of muscadine grapes. J. Agric. Food Chem. 51, 5497–5503.
725 <https://doi.org/10.1021/jf030113c>

726 Ruel, J.R., Walker, M.A., 2006. Resistance to Pierce’s disease in *Muscadinia rotundifolia* and other
727 native grape species. Am. J. Enol. Vitic. 32, 155–158.

728 Singleton, V.L., Orthofer, R., Lamuela-Raventós, R.M., 1999. Analysis of total phenols and other
729 oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods Enzymol. 299,
730 152–178. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1)

731 Staudt, G., 1997. Evaluation of resistance to grapevine powdery mildew (*Uncinula necator* [Scxw.]
732 Burr., anamorph *Oidium tuckeri* BERK.) in accessions of *Vitis* species. *Vitis* 36, 151–154.

733 Takayama, F., Nakamoto, K., Kawasaki, H., Mankura, M., Egashira, T., Ueki, K., Hasegawa, A., Okada,
734 S., Mori, A., 2009. Beneficial effects of *Vitis coignetiae* Pulliat leaves on nonalcoholic
735 steatohepatitis in a rat model. *Acta Med. Okayama* 63, 105–111.
736 <https://doi.org/10.18926/AMO/31835>

737 Vagiri, M., Johansson, E., Rumpunen, K., 2017. Phenolic compounds in black currant leaves – an
738 interaction between the plant and foliar diseases? *J. Plant Interact.* 12, 193–199.
739 <https://doi.org/10.1080/17429145.2017.1316524>

740 Wan, Y., Schwaninger, H., He, P., Wang, Y., 2007. Comparison of resistance to powdery mildew and
741 downy mildew in Chinese wild grapes. *Vitis* 46, 132–136.

742 Wan, Y., Schwaninger, H.R., Baldo, A.M., Labate, J.A., Zhong, G.Y., Simon, C.J., 2013. A
743 phylogenetic analysis of the grape genus (*Vitis* L.) reveals broad reticulation and concurrent
744 diversification during neogene and quaternary climate change. *BMC Evol. Biol.* 13, 141–161.
745 <https://doi.org/10.1186/1471-2148-13-141>

746 Xiao, J., Capanoglu, E., Jassbi, A.R., Miron, A., 2016. Advance on the flavonoid C-glycosides and
747 health benefits. *Crit. Rev. Food Sci. Nutr.* 56, 29–45.
748 <https://doi.org/10.1080/10408398.2015.1067595>

749 Yang, L., Wen, K.S., Ruan, X., Zhao, Y.X., Wei, F., Wang, Q., 2018. Response of plant secondary
750 metabolites to environmental factors. *Molecules* 23, 762–788.
751 <https://doi.org/10.3390/molecules23040762>

752 Yim, N., Thi, D., Nam, T., Pyo, J., Lee, S., Na, M., Jung, H., Su, H., Ho, Y., Bae, K., 2010. Bioorganic
753 & Medicinal Chemistry Letters The antimicrobial activity of compounds from the leaf and stem of
754 *Vitis amurensis* against two oral pathogens. *Bioorg. Med. Chem. Lett.* 20, 1165–1168.

755 <https://doi.org/10.1016/j.bmcl.2009.12.020>

756 Zhang, J., Wu, X., Niu, R., Liu, Y., Liu, N., Xu, W., Wang, Y., 2012. Cold-resistance evaluation in 25

757 wild grape species. *Vitis* - J. Grapevine Res. 4, 153–160.

758 <https://doi.org/10.1017/S0020743810001558>

759