AperTO - Archivio Istituzionale Open Access dell'Università di Torino

An assessment of the modulation of the population dynamics of pathogenic *Fusarium oxysporum* f. sp. *lycopersici* in the tomato rhizosphere by means of the application of *Bacillus subtilis* QST 713, *Trichoderma* sp. TW2 and two composts

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1721175> since 2020-04-03T19:17:48Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Research paper**

2 **An assessment of the modulation of the population dynamics of pathogenic *Fusarium***
3 ***oxysporum* f. sp. *lycopersici* in the tomato rhizosphere by means of the application of *Bacillus***
4 ***subtilis* QST713, *Trichoderma* sp. TW2 and two other composts**

5
6 **Maria Alexandra Cucu^{a*}, Giovanna Gilardi^a, Massimo Pugliese^{a,b,c}, Maria Lodovica Gullino^{a,b}**
7 **and Angelo Garibaldi^a**

8
9
10 *^aAGROINNOVA - Centre of Competence for the Innovation in the Agro-Environmental Sector,*
11 *Turin University, Largo P. Braccini 2, 10095, Grugliasco, Turin, Italy*

12 *^bDepartment of Agricultural, Forest and Food Sciences (DISAFA), Turin University, Largo P.*
13 *Braccini 2, 10095 Grugliasco, Turin, Italy*

14 *^cAgriNewTech srl, Via Livorno 60, 10144 Turin, Italy*

15
16
17
18 Running head: Biological Control of Tomato Fusarium Wilt with BCAs and composts

19
20 *Corresponding author:

21 Dr. Maria Alexandra Cucu

22 Phone: +39 011 670 8544

23 Email: mariaalexandra.cucu@unito.it

24

25

26

27 **Abstract**

28 Fusarium wilt, caused by *Fusarium oxysporum* f. sp. *lycopersici* (Fol), results in considerable yield
29 losses for tomato crops throughout the world. In this study, experiments were carried out “*in situ*”
30 over two consecutive years to evaluate the efficiency of two biological control agents (BCAs)
31 (*Bacillus subtilis* QST713 and *Trichoderma* sp. TW2) and two composts in controlling the disease.
32 In this context, the quantitative polymerase chain reaction was used along with soil chemical
33 parameters to study the general effects of treatments on the severity of the disease and on the non-
34 target microbial populations residing in the studied rhizosphere and bulk soil. The ecological fitness
35 of the BCAs was also evaluated. Furthermore, as BCAs produce elicitors which may activate plant
36 defense reactions, particular attention was paid to the induction of pathogenesis-related genes (PR)
37 in the roots of tomato plants.

38 The preventative nursery application of all three types of biocontrol agents, that is, *Bacillus subtilis*
39 QST713, *Trichoderma* sp. TW2 and compost, as separate treatments, induced a significant
40 reduction in the disease, compared to the untreated control, and reduced tomato Fusarium wilt by
41 70%. This result was confirmed by the significant negative correlations between the abundance of
42 biological control agents and the severity of the disease. In general, the BCA and compost
43 treatments did not induce a negative effect on the non-target microbial communities. The transcript
44 levels coding for the studied pathogenesis-related (PR) genes were always higher in the presence of
45 Fol on its own (untreated control) for all genes considered. However, the accumulation of
46 transcripts in the tomato roots was different, depending on the treatment. An important level of
47 disease reduction was shown by a decrease in Fol abundance, together with a greater abundance of
48 the inoculated BCA populations and an accumulation of transcripts encoding PR genes.

49 In short, the results of this study reinforce the concept of the sustainability of treatments based on
50 biological control agents and composts for the management of tomato Fusarium wilt.

51

52 **Keywords:** *Fusarium oxysporum* f.sp. *lycopersici* (Fol); BCAs; disease suppression; pathogenesis
53 related genes (PR)

54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77

1. Introduction

Tomato (*Lycopersicon esculentum* Mill.) is among the world's most important vegetables, with a global annual yield of approximately 160 million tons for all types of tomatoes (FAO, see the statistical database [FAOSTAT], 2014; available on: <http://fatstat.fao.org>). According to Lopes and Ávila (2005), about 200 biotic or abiotic diseases have been reported on tomato throughout the world. *Fusarium oxysporum* f. sp. *lycopersici* (Fol) is a soil-borne pathogen that causes the Fusarium wilt of tomato, which results in severe production losses (Gale et al., 2003; Huang et al., 2012; McGovern, 2015).

The eradication of soil-borne pathogens (including Fol) is generally based on an integrated management programme, consisting of chemical soil fumigation and the use of resistant cultivars when available (McGovern, 2015), but also on the use of organic amendments (Dukare et al., 2011; Pugliese et al., 2008; 2010) and biocontrol agents (BCAs) (Muslim et al., 2003; Gowtham et al., 2016; Huang et al., 2012; Jangir et al., 2018). *Bacillus* spp. and *Trichoderma* spp., which are available as commercial and experimental products, are among the most frequently used BCAs today (Jacobsen et al., 2004; Lorito and Woo, 2015). They are abundant in soil and have been investigated for their potential application as BCAs against several plant pathogens, including Fol (Cotxarrera et al., 2002; de Medeiros et al., 2017; Ghazalibiglar et al., 2016; Lorito et al., 2010; Zhao et al., 2014). The combination of BCAs with organic amendments (e.g., compost, manure, plant waste) has also been studied to control soil-borne plant pathogens (Borrero et al., 2004, 2012; Cotxarrera et al., 2002; Pugliese et al., 2011).

BCA activities include the secretion of secondary metabolites (such as cyclic lipopeptides and volatile organic compounds), and competition for nutrients and parasitism (Kim et al., 2015; Vitullo et al., 2012). In addition to these antagonistic effects on plant pathogens, BCAs produce elicitors that activate plant defence reactions (Walters and Daniell, 2007), such as the induction of pathogenesis-related (PR) genes (Edreva, 2005) encoding chitinase and β -1,3-glucanases. These two groups of PR proteins have attracted considerable interest because of their “*in vitro*” and “*in*

104 *vivo*” inhibitory activity against fungi (Van den Elzen et al., 1993; Bargabus et al., 2002). In
 105 addition, PR proteins (glucanases and chitinase) have been seen to increase concomitantly with the
 106 development of the disease (Rep et al., 2002). For example, it has been acknowledged that the
 107 treatment of sugar beet with *Bacillus mycoides* induces the expression of PR genes such as β -1,3-
 108 glucanase (Bargabus et al., 2002). As far as the inoculation of tomato plants with *F. oxysporum* f.
 109 sp. *Lycopersici* is concerned, increases in chitinase, β -1,3-glucanase, and β -1,4-glucosidase
 110 activities have been observed to be lower in resistant cultivars than in susceptible ones (Ferraris et
 111 al., 1987). However, relatively little is known about the effect of BCA and compost as pre-planting
 112 treatments on the accumulation status of PR genes in plants at the end of the cropping cycle in the
 113 correspondences with the peak of the disease (Borrero et al., 2004).
 114 BCAs are known to colonise the rhizosphere, thereby indirectly affecting the microbial processes
 115 that are essential for the general functioning of the soil ecosystem. This rhizosphere colonization
 116 includes effects on the resident microorganisms and in consequence on carbon, nitrogen and
 117 phosphorus cycling (Bonanomi et al., 2010; Gupta et al., 2012). This may result in adverse effects
 118 on crop yield and health. Hence, there is a need to study the potential side - effects of the applied
 119 treatments on the indigenous rhizosphere and soil microbial populations (Bonanomi et al., 2018).
 120 The main objectives of this study were: i) to evaluate the effect of two BCAs and two composts on
 121 the severity of the fusarium wilt of tomato and on the non-target rhizosphere and bulk soil microbial
 122 communities in the tomato - *Fusarium oxysporum* f.sp. *lycopersici* (Fol) pathosystem, in a naturally
 123 infested soil; ii) to obtain new insights into the induction of pathogenesis-related (PR) genes in pre-
 124 planting treated tomato roots at the end of the tomato crop cycle, which usually coincides with the
 125 highest disease incidence. Therefore, the differential expression of genes encoding *GluA* (acidic
 126 extracellular b-1,3-glucanase), *GluB* (basic intracellular b-1,3-glucanase), *Chi3* (acidic extracellular
 127 chitinase), *Chi9* (basic intracellular chitinase) and *PR-1a* was studied.

128
 129

130 2. Materials and Methods

131 2.1. Layout of the experiments

132 Two experiments (2016 and 2017) were carried out in plastic tunnels on a commercial farm in Asti
133 (Northern Italy), which had a history of several tomato cycles prior to the beginning of this study, in
134 a sandy loam soil (sand : silt : clay 50 : 40 : 10 %, with neutral pH and 1.5 % organic matter),
135 naturally infested with Fol.

136 The experimental trials were carried out over two consecutive years, 2016 and 2017, in order to test
137 the efficacy of pre-planting soil treatments with a commercially available formulation of *Bacillus*
138 *subtilis* QST713 - (SM, Serenade Max, Bayer Crop Science, Italy) and an experimental BCA
139 (*Trichoderma* sp., TW2, AgriNewTech, Italy) (Table 1) against Fol. In addition, two commercial
140 composts that is Ant's Compost V and Ant's Compost M; AgriNewTech, Italy, produced from
141 green wastes in a dynamic industrial treatment system, were used (Table 1). Ant's Compost M is in
142 fact Ant's Compost V inoculated with *Trichoderma* sp. TW2. An untreated control was used to
143 monitor Fusarium wilt development. Summarised details of these treatments can be found in Table
144 1.

145 Tomato seeds (cv. Ingrid, Seminis) were sown in 40-plug trays (53 x 42 cm; plug 10x10 cm, 4l peat
146 capacity) filled with a peat mixture substrate.

147 Four soil treatments with BCAs were carry out on the plug tray between sowing and transplanting
148 in a greenhouse in a commercial nursery (Table 1). The BCAs were applied by spraying them onto
149 the soil surface in a high volume of water (400 ml/tray) using a 1 l capacity hand sprayer. The
150 application times were: 37 days after sowing, which was defined as T1, for the first treatment, 44
151 days after sowing, defined as T2, for the second treatment, 51 days after sowing, defined as T3, for
152 the third treatment and 58 days after sowing, defined as T4, for the fourth treatment. The compost
153 products were applied twice: they were first mixed with the substrate at sowing, and this
154 combination was defined as T0, and they were then mixed with the soil one week before

transplanting at T3, that is, 51 days after sowing, according to dosages and application times reported in Table 1.

The tomato seedlings at the fourth true leaf stage were transplanted (T58) into naturally Fol infested soil at a density of 2.5 plants/m², drip irrigated and then grown according to the cultural practices adopted by commercial growers in the region. The plots were arranged in a completely randomised block design, with four replicates per treatment. The plants were monitored every 10-15 days to determine the development of symptoms. Disease severity (DS) data were recorded, starting from the appearance of the first symptoms (yellow leaves and reduced growth). The final disease rating was carried out 128 days after transplanting, on 16-20 plants per treatment by dissecting each plant. For this purpose, a DS scale was created as follows: 0 = healthy plant, 25% = initial leaf chlorosis, 50% = severe leaf chlorosis and initial symptoms of wilting during the hottest hours of the day, 75% = severe wilting and severe symptoms of leaf chlorosis; 100% = plant totally wilted, leaves completely necrotic (Borrero et al., 2004; Srinivasan et al., 2009).

2.2. Rhizosphere and bulk soil sampling

The effect of the two BCAs and composts on indigenous microbial communities was studied at the rhizosphere and bulk soil levels by sampling at the end of the trials (2016 and 2017), as described by Cucu et al., (2018). In brief, rhizosphere and bulk soil were collected, with three biological replicates per treatment. Each replicate consisted of five sub-replicates, which were pooled together. The replicate samples were collected in a W - shaped sampling pattern from each plot throughout the field, in order to ensure a good bulk soil homogeneity. The dry weight of the roots was also determined.

The fresh rhizosphere and bulk soil samples were accurately homogenized separately, sieved through a 2 mm sieve and stored at 4°C. All the samples were split for further geochemical analysis: humidity, pH, total N (TN), inorganic N as nitrate (NO₃⁻) and ammonium (NH₄⁺),

180 dissolved organic carbon (DOC), total phosphorus (Pt) and available phosphorus (Pav), as well as
181 for molecular investigations.

182

183 2.3. Molecular analyses

184 2.3.1. Rhizosphere and bulk soil DNA extraction

185 A NucleoSpin® Soil kit (Macherey-Nagel GmbH & Co. KG, Düren, Germany) was used,
186 according to the manufacturer's instructions, for the rhizosphere and bulk soil genomic DNA
187 extraction. The extraction was conducted on fresh samples (250-500 mg of sample material). The
188 quantity and purity of the DNA were measured spectrophotometrically using a NanoDrop ND-1000
189 device (NanoDrop Technologies, Wilmington, DE, USA).

190

191 2.3.2 Microbial abundance (quantitative PCR (qPCR) assays)

192 The abundance of the bacterial and archaeal 16S rRNA genes, fungal 18S rRNA genes and of three
193 functional genes, (i.e., bacterial ammonia monooxygenase *amoA* - AOB, archaeal *amoA* - AOA and
194 fungal chitinase - *chiA* was determined by means of qPCR, using a StepOnePlus™ Real-Time PCR
195 System (Applied Biosystems, Foster City, CA, USA) for the rhizosphere and bulk soil DNA
196 samples. In addition, the abundance of *Fol*, *Bacillus* spp. and *Trichoderma* spp. was also
197 investigated. A description of the primer sets and amplification details is given in Table 2.

198 Amplicons were generated from each target gene for the standard preparation, purified (Invisorb
199 Fragment CleanUp, Stratec Molecular GmbH, Berlin, Germany) and ligated in a Strata-Clone PCR
200 cloning vector pSC-A (Strataclone PCR Cloning Kit, Agilent Technologies Inc.); the ligation
201 products were then transformed using StrataClone SoloPack competent cells (Agilent Technologies
202 Inc.). The specificity of the clones used as qPCR standards was checked, via sequencing, at LGC
203 Genomics GmbH (Berlin, Germany), and through BLAST analysis. Plasmid DNA was isolated
204 (GenElute™ Plasmid Miniprep Kit, Sigma-Aldrich, St. Louis, MO, USA) from standard clones and
205 quantified as described above.

206 After the qPCR assay optimisation, the final volume of the qPCR reactions was established as 25 μ l
207 for the 16S and 18S rRNA genes, for the bacterial and archaeal *amoA*, and for the Fol and BCA
208 quantification, while it was 20 μ l for *chiaA*. All the qPCR assays were performed using 10 ng DNA
209 as a template, except for the bacterial 16S rRNA gene, for which 5 ng DNA was used (Cucu et al.,
210 2017; 2018). The reaction mixtures contained a 1x Power SYBR green master mix (Applied
211 Biosystems), with 0.12 μ M of each oligonucleotide (Table 2) for the total bacteria, archaea, fungi
212 and Fol, 0.32 μ M for the AOB, AOA and BCAs and 0.4 μ M for *chiaA*.
213 All the considered genes were quantified in triplicate across the plates, while standards were run in
214 duplicate in 10-fold serial dilutions. The amplification efficiency ranged from 96%, (archaeal 16S
215 rRNA, fungal 18S rRNA, BCAs and their functional genes) to 103% (bacterial 16S rRNA). The
216 amplification efficiency of the nitrifiers was 95.3% and 99.1% for AOB and AOA, respectively. R^2
217 was always ≥ 0.98 . Melting curves of the amplicons were generated to ensure that the fluorescence
218 signals originated from specific amplicons and not from primer dimers or other artifacts. This was
219 confirmed by checking the amplification products on 1% agarose gel. Gene copy numbers were
220 calculated with StepOne™ software, version 2.2 (Applied Biosystems). The data were normalised
221 and presented in figures as copies g^{-1} dry soil.

222

223 2.3.3 Race identification of *Fusarium oxysporum* f. sp. *lycopersici* from the experimental field

224 Real time PCR was used to identify Fol from the experimental field. Four primer/probe sets were
225 used to identify the tomato wilt pathogen race, on the basis of rDNA -intergenic spacer and
226 avirulence genes. Real - time PCR trials were conducted using genomic DNA from mycelia isolated
227 from the experimental site and soil DNA. Reaction conditions were established according to the
228 protocol described by Inami et al., (2010).

229

230

231

232 2.3.4 RNA extraction and real-time RT-PCR gene expression analysis

233 In order to study the differential expression of the selected genes in plants pre-planting treated with
234 biocontrol agents and compost, the real time RT-PCR assay was used on RNA samples from roots
235 collected at the end of the tomato crop cycle, which had been conserved at -80°C. The total RNA
236 was extracted by means of an RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) using 0.1 g of
237 tomato roots. The RNA was DNase treated using TURBO DNase (Ambion™, Thermo Fisher
238 Scientific, Waltham, USA), according to the manufacturer's protocol, and then reverse transcribed
239 (500 ng total RNA) using a high capacity cDNA reverse transcription kit (Applied Biosystems).
240 The actin gene was used as a reference gene. The primers for the *LeChi3*, *LeChi9*, *TomB13GluA*,
241 *TomB13GluB*, and *LePr1a* target genes were used according to Aimé et al., (2008). Real-time PCR
242 reactions were carried out with 25 ng of cDNA, 500 nM of each primer, 10 µl of the 1x Power
243 SYBR green master mix (Applied Biosystems) and RNase-free water in a final volume of 20 µl.
244 cDNA was replaced by RNase free water in the negative control. A programme consisting of 15
245 min at 95°C, followed by 40 denaturation cycles for 15 s at 95°C, annealing for 30 s at 58°C and
246 extension for 30 s at 72°C was used for the real-time PCR, at the end of which fluorescence was
247 measured. Real-time PCR reactions were carried out in duplicate for each sample. Primer titration
248 and dissociation experiments were performed to confirm there was no formation of primer dimers
249 or false amplicons that could interfere with the results. After the real-time PCR experiment, the Ct
250 number was extracted for both the reference gene and the target gene considering an auto baseline
251 and a manual threshold. Gene expression levels (relative to the actin gene) were calculated for each
252 cDNA sample using the following equation: $\text{relative gene / actin} = (E_{\text{gene}} - Ct_{\text{gene}}) / (E_{\text{actin}} - Ct_{\text{actin}})$ ratio.
253

254

255 2.4. Chemical properties of the rhizosphere and bulk soil samples

256 The pH values were measured in water suspensions at a solid:liquid ratio of 1:2.5. The total
257 nitrogen (TN) was quantified using a Leco Tru Spect CN automatic analyser. The dissolved organic

258 carbon (DOC) was determined after sample acidification in a TOC/TN analyser (Multi NC 2100S,
259 Analytic Jena GmbH, Jena, Germany). Ammonium (NH_4^+) and nitrate (NO_3^-) were measured
260 colorimetrically, by means of a continuous flow auto-analyser (Alliance Evolution II), using
261 standard colorimetric techniques. The total phosphorus (Pt) was determined by means of 'ICP
262 Varian mod. Liberty LR', after microwave digestion with hydrogen peroxide, hydrochloric acid and
263 nitric acid, filtration and dilution. The available P (P_{av}) was extracted using sodium bicarbonate and
264 determined by means of the molybdenum blue method (Olsen et al., 1954), modified for continuous
265 flow colorimetric analysis (Alliance Evolution II).

266

267 2.5. Statistical analyses

268 The data on gene abundance, on the chemical properties of the soil and on the disease index were
269 subjected to a linear mixed model, with time considered as the random variable, using R software
270 (Software R 3.0.1, R foundation for Statistical Computing, Vienna, Austria, [http://www.R-](http://www.R-project.org)
271 [project.org](http://www.R-project.org)). The effects of two different BCA and compost applications on the abundance of the
272 studied genes, on the chemical properties of the soil and on disease severity were evaluated. All the
273 data were subjected to a Levene test to check for the homogeneity of variance, and normality was
274 tested on the residuals using the Shapiro - Wilk test; when not normally distributed, data from
275 disease severity (DS) were arcsin transformed, while the data from microorganism abundance were
276 log-transformed and normality was checked again. The means were separated by means of the
277 Bonferroni test. The statistical analysis included treatment, year, treatment $\times$ year and rhizosphere $\times$
278 bulk soil.

279 Pearson's linear correlation coefficients were calculated to assess the relationships between
280 microbial abundance and disease severity as well as between BCA and F_{ol} abundance and the soil
281 chemical properties.

282

283

284 3. Results

285 3.1. Disease severity

286 In 2016, in the presence of a low level of disease severity (DS 29.7%) in the untreated control - C
287 tomato plots, the two tested BCAs and composts provided a disease reduction of 89.5 to 93%. The
288 green compost - CV provided statistically similar results to those of the compost - CM, which was
289 enriched with *Trichoderma* sp. TW2 (a disease reduction of 93% and 64%, respectively). However,
290 no significant differences were observed between treatments. In 2017, a disease severity of 39.9%
291 was observed in the untreated tomato control plants, but all the treatments significantly reduced
292 Fusarium wilt symptoms as much as 50%, and statistically similar results between treatments were
293 obtained (Table 3).

295 3.2. Microbial abundance

296 3.2a Race identification and the abundance of soil-borne *F. oxysporum* f.sp. *lycopersici* (Fol)

297 The amplification reaction for the Fol race identification was positive for both the P1 and R1 Race -
298 1 specific sets (Inami et al., 2010), on the basis of the rDNA-IGS and the avirulence genes,
299 respectively (data not shown).

300 In general, Fol abundance was significantly higher in the bulk soil than in the rhizosphere soil, $P(F)$
301 $Rhizo \times Bulk\ soil = 0.003$. All the treatments resulted in a significant pathogen reduction, compared
302 to the untreated control - C, in both the rhizosphere and bulk soil at the end of both trials (2016 and
303 2017) (Figure 1). The treatment $\times$ year interaction was not significant.

305 3.2b Total microbial community abundance

306 At the end of both trials (2016 and 2017), the bacterial, archaeal and fungal communities showed a
307 notable homogeneity, with no significant treatment $\times$ year interaction. Therefore, the data were
308 presented as the average of all the values of both trials.

Overall, at the end of both trials, the bacterial 16S rRNA abundance was significantly higher at the bulk soil level than at the rhizosphere level (Table 4). A higher bacterial abundance was observed in rhizosphere soil samples after the *Trichoderma* sp. TW2 - TW2 treatment, than in the untreated control -C, which was significantly similar to the green compost - CV treatment. On the other hand, the highest bacterial abundance at the soil level was observed after the inoculated compost - CM treatment, while the lowest was observed after the *B. subtilis* treatment. The archaeal 16S rRNA gene abundance in the bulk soil samples was more stable and was not influenced by the treatments. However, the green compost - CV, green compost - CM+TW2, and *B. subtilis* treatments resulted in an elevated archaeal presence, compared to the untreated control - C in the rhizosphere soil samples (Table 4). The fungal 18S rRNA abundance was significantly higher in the rhizosphere samples than in the bulk soil ones (Table 4). A significantly higher fungal abundance was observed in the rhizosphere after the inoculated compost - CM, *Trichoderma* sp. TW2 - TW2 treatments as well as in the untreated control - C. *B. subtilis* and green compost CV significantly reduced the total fungal population, while the compost - CM, *Trichoderma* sp. TW2 - TW2 treatments did not affect it. At the bulk soil level, the highest fungal abundance was observed after the green compost - CV treatment, while the untreated control - C was statistically similar to the *Trichoderma* sp. TW2 - TW2 treatment. *B. subtilis* and green compost CM+TW2 significantly reduced the total fungal population.

327

3.2c Functional gene abundance

In general, the ammonia-oxidising bacterial (AOB) gene abundance was not significantly influenced by the treatments, compared to the untreated control, and was significantly more abundant in the rhizosphere soil than in the bulk soil samples. The highest abundance of AOB was observed in the untreated control - C of the bulk soil and the lowest value was observed for the untreated control - C of the rhizosphere samples (Table 5). In general, the ammonia-oxidising archaeal (AOA) gene abundance was significantly more abundant in the bulk soil samples than in

the rhizosphere soil ones (Table 5). The highest abundance in the rhizosphere samples was observed after the compost - CM treatment, while the highest abundance in the bulk soil samples was observed after the compost - CV treatment (Table 5).

3.2d BCA abundance

Overall, at the end of both trials, an increase in the indigenous populations of *Bacillus* spp. and *Trichoderma* spp. was observed for all the treatments, compared to the untreated control - C. No significant treatment $\times$ year interaction was observed. In general, the abundance of *Bacillus* spp. was significantly higher for the *B. subtilis* - SM treatment than for the untreated control - C and for all the other treatments in both the rhizosphere and bulk soil. However, both compost treatments resulted in an increased resident *Bacillus* abundance in the rhizosphere and bulk soils after both trials. Nevertheless, the *Trichoderma* sp. TW2 - TW2 - treatment showed similar abundance to those of the untreated control - C (Table 6).

The inoculation with *Trichoderma* sp. significantly increased the overall *Trichoderma* spp. community in both the rhizosphere and in the bulk soils, compared to the untreated control - C. The highest abundance of *Trichoderma* spp. was observed in the rhizosphere after the compost - CM treatment and in the bulk soil after the *Trichoderma* sp. TW2 - TW2 treatment (Table 6). *B. subtilis* - SM resulted in a somewhat lower abundance than the *Trichoderma* sp. - TW2 treatment.

In general, the *chiA* gene (*Trichoderma* based chitinase) abundance was significantly influenced by the treatments, compared to the untreated control - C (Table 7). All the treatments at the rhizosphere soil level resulted in significant differences between the *chiA* gene abundances. The abundance of the *chiA* gene was higher in the rhizosphere soil samples for the *Trichoderma* sp. TW2 and the compost - CM treatments and lower and significantly similar to the untreated control - C after the *B. subtilis* - SM treatment. All the treatments at the bulk soil level also resulted in a higher *chiA* abundance than the untreated control - C. The highest *chiA* gene abundance at the bulk soil level was observed after the green compost - CV treatment.

361 3.2e Gene expression - analysis with RT-PCR

362 The levels of the transcripts encoding the studied pathogenesis-related (PR) genes were always
363 higher for the presence of Fol on its own (untreated control - C samples) for all the considered
364 genes. The transcript accumulation in the tomato roots differed according to the treatments (Figure
365 2); the transcript level profiles encoding *Chi3* and *Chi9* were rather similar, and were higher in the
366 *Trichoderma* sp. based treatments than the treatments with *B. subtilis* - SM and green compost -
367 CV. Overall, the expression level of *GluA* was lower than the expression level of *GluB*. The *B.*
368 *subtilis* - SM and both compost treatments showed a slightly higher expression of *GluA* than the
369 *Trichoderma* sp. TW2 - TW2 inoculation on its own. The *GluB* gene was expressed more in the *B.*
370 *subtilis* - SM inoculated compost and *Trichoderma* sp. TW2 - TW2 treatments. The expression
371 level of the PR-1 gene was very low for all the treatments.

372

373 3.3. Chemical properties

374 The pH was generally significantly lower ($p < 0.05$) at the rhizosphere level than at the bulk soil
375 level for both trials, with values ranging from 7.43 to 7.60 for the rhizosphere and between 7.95 and
376 8.18 for the bulk soil samples. The concentrations of TN, NH_4^+ , NO_3^- and DOC were generally
377 significantly higher in the rhizosphere samples than in the bulk soil ones. The untreated control - C
378 was characterised by higher NH_4^+ , NO_3^- and DOC concentrations than all the treatments at a bulk
379 level (Table S1). No significant treatment $\times$ year interaction was observed.

380 The dry weight of the roots was not significantly different between the SM, CM and CV treatments.
381 The highest dry weight was observed after the TW2 treatment, while the lowest was observed in the
382 untreated control - C (Figure S1).

383

384 3.5. Correlations between the total microbial communities (16S bacteria, 16S archaea and 18S
385 fungi), the functional genes (fungal *chiA* gene, bacterial and archaeal *amoA* genes,), the BCAs
386 (*Bacillus*, *Trichoderma*) and Fol abundances, and DS.

387 In general, the correlations were negative, with r coefficients ranging from -0.3 to -0.7 ($p < 0.001$)
388 at the rhizosphere level and from -0.3 to -0.8 ($p < 0.001$) at the bulk soil level (Table 8 and Table 9,
389 respectively).

390

391

392 **4. Discussion**

393 In the present study, pre-planting treatments with BCAs (*Bacillus subtilis* and *Trichoderma* sp. -
394 TW2) and two composts were used to control tomato Fusarium wilt caused by *Fusarium oxysporum*
395 f.sp. *lycopersici* (Race 1) in naturally infested soil. The evaluation was based on the two-year effect
396 of these treatments on the disease incidence, PR gene expression and on the rhizosphere/soil
397 microflora and chemical properties. The risk assessment of the BCAs and compost utilised within
398 the frame of this study pointed out the impact these treatments had on the pathogen population and
399 disease control corroborated with the PR gene expression status, as well as on the rhizosphere and
400 bulk soil microbial community dynamics. This impact is important as the resident microbes may
401 colonise similar niches as the newly introduced BCA microorganisms and thus compete for similar
402 resources in the rhizosphere and soil environment (Winding et al., 2004).

403

404 *4.1. Disease severity, and F. oxysporum f. sp. lycopersici (Fol) abundance*

405 The preventative application of two BCAs and composts, starting from the nursery, significantly
406 reduced Fol Race 1 abundance, compared to the untreated control. These findings were in accord
407 with the results of previous studies that showed that *Fusarium* abundance at the rhizosphere level of
408 different hosts, such as cucumber, banana and lettuce, was reduced after BCA treatments (Qiu et al.,
409 2012; Shen et al., 2015; Fu et al., 2017). In the present study, Race 1 of Fol abundance was low
410 after the treatment with the enriched compost (CM), thus indicating that an effective Fusarium wilt
411 control may be achieved on tomato with organic amendments, as previously pointed out (Borrero et
412 al., 2004; Cotxarerra et al., 2002). The compost utilisation probably resulted in changes in the

413 activity, density and structure of the non - target microbial community in the soil, thus enhancing
414 competition with the pathogen for space and nutrients, such as carbon (C) and nitrogen (N). This
415 may have resulted in a reduction in the Fol abundance and its potential activity, as already reported
416 by Bonilla et al., (2012), Cucu et al., (2018), Larkin et al., (2015).

417

418 4.2. Plant response to *Fol* infection

419 Plants usually have a complex defence system against pathogens that are well known as stress
420 factors, including different types of stress proteins with putative protective functions (Brown et al.,
421 2017). A high accumulation of transcripts encoding the studied PR genes in the plant roots from the
422 untreated control was observed in this study. This may be correlated to the colonisation of the
423 tomato vessels by Fol, as previously reported by Olivain and Alabouvette, (1999). The here
424 presented results are in line with those of Aime' et al., (2008), who found a higher PR gene
425 expression in tomato plants inoculated with a pathogenic strain, than in plants inoculated with a
426 non-pathogenic strain of *F. oxysporum*. Moreover, the pre-inoculation of plants with *Trichoderma*
427 sp. resulted in an even higher expression of PR genes encoding chitinase, thus indicating that
428 *Trichoderma* spp. induced a potentiated status in the plant that enabled it to be more resistant to
429 subsequent pathogen infection, as was also observed by Shores et al., (2005). This hypothesis is in
430 line with that of Shores et al., 2005 and Yedidia et al., 2003, who found that, in a *T. asperellum* -
431 cucumber system, PR protein chitinase, β -1,3-glucanase and peroxidase were induced by
432 *Trichoderma*. Jangir et al., (2018) showed that *Bacillus* sp. produced chitinase and β -1, 3-glucanase
433 that might play a role in the digestion of the pathogen's hyphae as carbon used than as energy
434 sources. Bargabus et al., (2002) showed that the treatment of sugar beet with *Bacillus mycoides*
435 induced the expression of PR genes that is new isoforms of β -1,3-glucanase. The potential
436 production of PR-proteins with chitinase and β -1,3- glucanase activity is presumably part of the
437 hypersensitive defence mechanism of tomato plants and may be responsible for the induction of the
438 resistance developed by the plant after treatment with the antagonists and infection with Fol.

439 4.3. Ecological feedback of the applied treatments and their effect on the total indigenous
440 prokaryotes

441 Rhizosphere microbial communities play important roles in plant health and disease prevention
442 (Bonanomi et al., 2018; Dudenhöffer et al., 2016). Analysing the abundance of the BCA - like
443 microorganisms, introduced into the soil as strains or through the compost treatments, showed a
444 very good feedback of the targeted populations (e.g., *Bacillus* spp. and *Trichoderma* spp.) as well as
445 excellent efficiency, as highlighted by the strong negative correlations with disease severity.
446 The abundance of *Trichoderma* spp. showed different trends in all the treatments. A higher
447 abundance of *Trichoderma* was detected in the rhizosphere of the inoculated compost treatment, but
448 also after the inoculation of the single strain. It is well known that composts represent an optimal
449 substrate for *Trichoderma* spp., as the organic matter composition and the associated biotic and
450 abiotic conditions can in general influence *Trichoderma* activity (Vinale et al., 2008; Lorito and
451 Woo, 2015). *Trichoderma* agents have been reported to be antagonistic to the *Fusarium* pathogens:
452 *F. oxysporum* f. sp. *ciceris*, *F. solani*, *F. oxysporum* f. sp. *lycopersici* (Dubey et al., 2007; Rojo et
453 al., 2007; Christopher et al., 2010). Gachomo and Kotchoni (2008) revealed that *Trichoderma*
454 isolates displayed various extracellular enzyme activities, for example amylase, chitinase, pectinase,
455 protease, lipase and cellulase activities, in order to compete with other microbes, and that the
456 production of volatile compounds by *Trichoderma* species could inhibit the growth of pathogenic
457 microorganisms. One of the main mechanisms of *Trichoderma* in the control of the soil-borne
458 pathogen is due to the presence of genes encoding cell wall degrading enzymes (Lorito et al., 1998,
459 Lorito and Woo, 2015). In this regard, the abundance of the *chiA* gene and its negative correlation
460 with disease severity has confirmed the potential antagonistic activity of the inoculated
461 *Trichoderma* strain TW2, but also of the inoculated compost - CM. This result has pointed out an
462 enhanced nitrogen efficiency as a result of the improvement of the nitrogen reduction and
463 assimilation mechanism, as also evidenced by Lorito et al., (2010) and Borrero et al., (2012). In
464 addition, the competition for carbon and other growing factors at the rhizosphere level helped to

sustain the control of Fol by *Trichoderma*, as highlighted by the very good correlation with dissolved organic carbon at the rhizosphere and bulk soil levels as well as with the ammonium content at the rhizosphere level. These results support the results of Sivan and Chet (1989), who demonstrated that the competition for nutrients is the major mechanism used by *T. harzianum* to control *F. oxysporum* f. sp. *melonis*.

The *Bacillus* sp. treatment also resulted to be effective in reducing the disease. The abundance results highlighted the excellent feedback of *Bacillus* spp. from the rhizosphere and soil levels. The significant negative relationships ($p < 0.001$) between *Bacillus* spp. and disease severity pointed out the antagonistic function of *Bacillus* against Fol, which may be due to volatile compounds and siderophore production, as reported by Jangir et al., (2018) after having characterised antagonistic *Bacillus* sp., isolated from a tomato rhizosphere, and to its control mechanisms against Fol.

Although a significant suppression of Fusarium wilt was achieved when the BCAs were employed individually (i.e., SM, TW2), alone or together with compost (i.e., CM), it is possible to assume, in agreement with Baker (1990), that the effects that were observed on antagonistic microbial communities after BCA applications may be the key to a successful and integrated approach for the control of tomato Fusarium wilt. In addition, as reported by Vinale et al., (2008), the biotic components of the soil environment also have relevant effects on the activity of the used biocontrol agents.

Overall, the release and successful proliferation of BCAs in the studied soil did not negatively alter the abundance of the functionally relevant indigenous soil microorganisms. The here presented results support previous reports which were based on the analysis of microbial densities (Ghini et al., 2000; Gullino et al., 1995; Mezzalana et al., 1998). A higher total fungal abundance was observed in the rhizosphere than in the bulk soil. The increased fungal community that was observed for all the treatments and for the untreated control was probably caused by the elevated level of organic carbon (Table S1). This may be linked to a competitive potential, based on an increased root exudation, which favours rhizosphere colonisation by fungi, including Fol and

491 *Trichoderma* spp. This result has been corroborated by considering the root dry weight (Figure S1).
492 According to Griffiths and Philippot (2013), the results suggested a great tolerance and resilience
493 potential of the native fungal community towards invading microorganisms (BCAs). Similarly,
494 Edel-Hermann et al., (2009) and Savazzini et al., (2009) only observed transient community shifts
495 in indigenous microbial populations in response to inoculation with antagonistic *Fusarium*
496 *oxysporum* and *Trichoderma atroviride*. However, a difference among treatments was observed at
497 both rhizosphere and bulk soil level, as *B. subtilis* and compost resulted in a decrease of total fungal
498 abundance. This result highlighted on one hand the rhizosphere competence of *B. subtilis* which
499 probably had a quick take over of space and nutrients. On the other hand the capacity of plant to
500 select the beneficial microorganisms as well as the influence of soil characteristics are well known
501 to be keys of soil biomass density and composition (Chapparo et al., 2012) Compost and
502 formulations of compost with bacteria or fungi (e.g., *Trichoderma* W2) may also dictate
503 competition for nutrients and the predominance of some microbial groups to another (Pugliese et al.
504 2011).

505 A high resilience of the total archaeal community was observed at both the rhizosphere and the soil
506 level, compared to the total bacterial community. Bacterial abundance was lower at the rhizosphere
507 level than at the bulk soil level, probably as a result of increased competition with the fungal
508 counterparts, including Fol. Archaeal nitrifiers (AOA) were more abundant in the bulk soil samples
509 than in the rhizosphere samples, which may be a result of the lower organic carbon content of the
510 soil and the higher pH (Wessén et al., 2010; Bates et al., 2011). A negative correlation was found
511 between the abundance of AOA and Race 1 of Fol at a soil level, thus suggesting a direct effect of
512 archaeal nitrifiers on the pathogen. However, these speculative interpretations require further
513 research to clarify the mode of actions of AOA responses to the development of Fol in the soil of
514 tomatoes. Nevertheless, the higher AOA abundance in the soil samples indicated that the AOA had
515 probably adapted to the decomposition of recalcitrant organic matter, as already described by Cucu
516 et al. (2017). The same pattern was observed by Cucu et al., (2018) for a lettuce - *Fusarium* wilt

517 pathosystem. This highlights the importance of abiotic factors, such as pH and soil type, in driving
518 not only the dynamics of prokaryotes, but also a higher level of suppressiveness towards *Fusarium*
519 spp. in the soil environment with pH ~8. This finding supports the findings of earlier studies which
520 correlated soil suppressiveness to *Fusarium* spp. with such abiotic soil characteristics as the clay
521 content and pH (Höper et al., 1995; Yergeau et al., 2010). The bacterial nitrifiers showed a higher
522 abundance and, as a consequence, a higher resilience at the rhizosphere level than the archaeal
523 counterparts. This result, as already pointed out by Wessén et al., (2010), points out that a low
524 ammonia environment is the key factor that determines a niche separation of AOA and AOB in
525 neutral soils.

526 The effect of *Bacillus* sp. and *Trichoderma* sp. on the nitrifiers was similar between treatments, thus
527 suggesting that *Bacillus subtilis* and *Trichoderma* sp. TW2 may not have any negative ecological
528 impact on other groups of microorganisms

529

530 **Conclusions and outlook**

531 The results have pointed out that the application of *Bacillus subtilis*, *Trichoderma* sp. TW2 and
532 compost treatments is safe for the rhizosphere and soil resident non-target microbial communities as
533 well as for the effective control of Fol on tomato. Moreover, the different response of the AOA
534 population to various treatments at a rhizosphere and a bulk level could indicate their suitability as
535 indicators to assess perturbations in soils, for example as the result of the introduction of new
536 microorganisms which act as biocontrol agents in the soil environment. Nevertheless, for a better
537 understanding of the impact that different treatments (e.g., biological or chemical) may have on
538 *Fusarium oxysporum* f.sp. *lycopersici*, it is necessary to consider not only non-target soil microbial
539 communities, but also different soil types characterised by different physical - chemical properties,
540 as abiotic factors.

541

542

543 **Acknowledgements**

544 This project has received funding from the European Union’s Horizon 2020 research and
545 innovation programme under grant agreement No. 633999 (EUCLID). The authors thank Professor
546 Marguerite Jones for the language revision.

549 **Conflict of Interest**

550 The authors declare that they have no conflict of interest. Massimo Pugliese declares he has a
551 financial interest (shareholder) in the AgriNewTech company that provided some of the products
552 tested in this study.

569 **References**

- 570 Aimé, S., Cordier, C., Alabouvette, C., Olivain, C., 2008. Comparative analysis of PR gene
571 expression in tomato inoculated with virulent *Fusarium oxysporum* f. sp. *lycopersici* and the
572 biocontrol strain *F. oxysporum* Fo47. *Physiol. Mol. Plant Pathol.*, 73, 9-15.
- 573 Baker, R., 1990. In *Biological control of soil-borne pathogens*. (Ed. Hormby, D.) CAB Inte.
574 Wellinford, UK., pp. 375-388.
- 575 Bargabus, R.L., Zidack, N.K., Sherwood, J.W., and Jacobsen., B.J., 2002. Characterization of
576 systemic resistance in sugar beet elicited by a non-pathogenic, phyllosphere-colonizing *Bacillus*
577 *mycooides*, biological control agent. *Physiol. Mol. Plant Pathol.*, 61, 289-298.
- 578 Bates, S.T., Berg-Lyons, D., Caporaso, J.G., Walters, W.A., Knight, R. and Fierer, N., 2011.
579 Examining the global distribution of dominant archaeal populations in soil. *ISME J.*, 5, 908-917.
- 580 Baysal, Ö., Çalışkan, M., Yeşilova, Ö., 2008. An inhibitory effect of a new *Bacillus subtilis* strain
581 (EU07) against *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Physiol. Mol. Plant. Pathol.*, 73, 25-
582 32.
- 583 Borrero C., Trillas M.I., Ordovas J., Tello J.C., Aviles M., 2004. Predictive factors for the
584 suppression of Fusarium wilt of tomato in plant growth media. *Phytopatol.*, 94, 1094-1101.
- 585 Borrero, C. , Trillas, M. , Delgado, A. and Avilés, M., 2012. Effect of ammonium/nitrate ratio in
586 nutrient solution on control of Fusarium wilt of tomato by *Trichoderma asperellum* T34. *Plant*
587 *Pathol.*, 61, 132-139.
- 588 Bonanomi, G., Antignani, V., Capodilupo, M. and Scala, F., 2010. Identifying the characteristics of
589 organic soil amendments that suppress soilborne plant diseases. *Soil. Biol. Biochem.*, 42, 136-144.
- 590 Bonanomi G., Lorito, M., Vinale F., Woo, S.L., 2018. Organic amendments, beneficial microbes,
591 and soil microbiota: toward a unified framework for disease suppression. *Annu. Rev. Phytopathol.*,
592 56, 1-20.
- 593 Bonilla, N., Gutiérrez-Barranquero, J.A., de Vicente, A. and Cazorla, F.M., 2012. Enhancing soil
594 quality and plant health through suppressive organic amendments. *Div.*, 4, 475-491.

595 Brown, A., Cowen, L.E., di Pietro, A., and Quinn, J., 2017. Stress adaptation. Microbiol. spectrum,
596 5(4), 1-38.

597 Chowdhury, S.P., Dietel, K., Randler, M., Schmid, M., Junge, H., Borriss, R., Hartmann, A.,
598 Grosch, R., 2013. Effects of *Bacillus amyloliquefaciens* FZB42 on lettuce growth and health under
599 pathogen pressure and its impact on the rhizosphere bacterial community. PLoS One 8,
600 doi.org/10.1371/journal.pone.0068818.

601 Chaparro, J.M., Sheflin, A.M., Manter, D.K. et al., (2012). Manipulating the soil microbiome to
602 increase soil health and plant fertility. Biol. Fertil. Soils, 48, 489. [https://doi.org/10.1007/s00374-](https://doi.org/10.1007/s00374-012-0691-4)
603 012-0691-4.

604 Christopher, D.J., Suthin, R.T., Rani, S.U., Udhayakumar, R., 2010. Role of defence enzymes
605 activity in tomato as induced by *Trichoderma virens* against Fusarium wilt caused by *Fusarium*
606 *oxysporum* f sp. *lycopersici*. J.biopesticides, 3, 158 - 162.

607 Cotxarrera, L., Trillas-Gay, M.I., Steinberg, C., Alabouvette, C., 2002. Use of sewage sludge
608 compost and *Trichoderma asperellum* isolates to suppress Fusarium wilt of tomato. Soil Biol.
609 Biochem., 34, 467-476.

610 Cucu, M.A., Marchan, S., Said-Pullicino, D., Celi, L., Kandeler, E., Rasche, F., 2017. Resource
611 driven community dynamics of NH_4^+ assimilating and N_2O reducing archaea in a temperate paddy
612 soil. Pedobiologia, 62, 16 -27.

613 Cucu, M.A., Gilardi, G., Pugliese, M., Matic, S., Ulrich, G., Gullino, M.L., Garibaldi, A. 2018.
614 Influence of different biological control agents and compost on total and nitrification driving
615 microbial communities at rhizosphere and soil level in a lettuce - *Fusarium oxysporum* f. sp.
616 *lactucae* pathosystem. J. Appl. Microbiol., doi: 10.1111/jam.14153.

617 de Medeiros, H.A., de Araújo Filho, J.V., de Freitas, L.G., Castillo, P., Rubio, M.B., Hermosa, R.
618 and Monte, E., 2017. Tomato progeny inherit resistance to the nematode *Meloidogyne javanica*
619 linked to plant growth induced by the biocontrol fungus *Trichoderma atroviride*. Sci. Rep., 7,
620 40216, 1-13.

621 Dubey, S.C., Suresh, M., Singh, B., 2007. Evaluation of *Trichoderma* species against *Fusarium*
 622 *oxysporum* f. sp. *ciceris* for integrated management of chickpea wilt. Biol. Control, 40, 118-127.
 623 Dudenhoffer, J.H., Scheu, S. and Jousse, A., 2016. Systemic enrichment of antifungal traits in the
 624 rhizosphere microbiome after pathogen attack. J. Ecol., 104, 1566-1575.
 625 Dukare, A.S., Prasanna, R. Chandra, Dubey, S., Nain, L., Chaudhary, V., Singh, R., Saxena, A.K.,
 626 2011. Evaluating novel microbe amended composts as biocontrol agents in tomato. Crop Prot., 30,
 627 436-442.
 628 Edel-Hermann, V., Brenot, S., Gautheron, N., Aimé, S., Alabouvette, C., Steinberg, C., 2009.
 629 Ecological fitness of the biocontrol agent *Fusarium oxysporum* Fo47 in soil and its impact on the
 630 soil microbial communities, FEMS Microbiol. Ecol., 68, 37-45.
 631 Edreva, A, 2005. Pathogenesis-related proteins: research progress in the last 15 years. Gen. Appl.
 632 Plant Physiol., 31, 105-124.
 633 Ferraris, L., Abbastista Gentile, I., Matta, A., 1987. Activation of glycosidases as a consequence of
 634 infection stress in *Fusarium* wilt of tomato. J Phytopathol, 118, 317-325.
 635 Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E. and Oakley, B.B., 2005. Ubiquity and
 636 diversity of ammonia oxidizing archaea in water columns and sediments of the ocean. Proc. Natl.
 637 Acad. Sci. USA, 102, 14683-14688.
 638 Fu, L., Penton, R., Ruan, Y., Shen, Z., Xue, C., Li, R. and Shen, Q., 2017. Inducing the rhizosphere
 639 microbiome by biofertilizer application to suppress banana *Fusarium* wilt disease. Soil Biol.
 640 Biochem., 104, 39-48.
 641 Gachomo, E., Kotchoni, S., 2008. The use of *Trichoderma harzianum* and *T. viride* as potential
 642 biocontrol agents against peanut microflora and their effectiveness in reducing aflatoxin
 643 contamination of infected kernels. Biotechnol., 7, 439-447.
 644 Gale, L.R., Katan, T., Kistler, H.C., 2003. The probable center of origin of *Fusarium oxysporum* f.
 645 sp. *lycopersici* VCG 0033. Plant Dis., 87, 1433-1438.

646 Gamliel, A., Austerweil, M., Kritzman, G., 2000. Non-chemical approach to soilborne pest
647 management - organic amendments. Crop Prot., 19, 847-853.

648 Gantner, S., Andersson, A.F., Alonso- Sáez, L. and Bertilsson, S., 2011. Novel primers for 16S
649 rRNA-based archaeal community analyses in environmental samples. J. Microbiol. Meth., 84, 12-
650 18.

651 Gao, W., Zhang, W. and Meldrum, D.R., 2011. RT-qPCR based quantitative analysis of gene
652 expression in single bacterial cells. J. Microbiol. Meth., 85, 221-227.

653 Garbeva, P., van Veen, J.A. and van Elsas, J.D., 2004. Microbial diversity in soil: selection of
654 microbial populations by plant and soil type and implications for disease suppressiveness. Ann.
655 Rev. Phytopathol., 42, 243-270.

656 Ghazalibiglar, H., Kandula, D.R.W. and Hampton, J.G., 2016. Biological control of Fusarium wilt
657 of tomato by *Trichoderma* isolates. New Zealand Plant Prot., 69, 57-63.

658 Ghini, R., Mezzalama, M., Ambrosoli, R., Barberis, E., Garibaldi, A. and De Stefano Piedade,
659 S.M., 2000. *Fusarium oxysporum* strains as biocontrol agents against Fusarium wilt: effects on soil
660 microbial biomass and activity. Pesqui. Agropecu. Bras., 35, 93-101.

661 Griffiths, B.S., Philippot, L., 2013. Insights into the resistance and resilience of the soil microbial
662 community. FEMS Microbiol. Rev., 37, 112-129.

663 Gowtham, H.G., Hariprasad, P., Nayak, S.C. and Niranjana, S.R., 2016. Application of
664 rhizobacteria antagonistic to *Fusarium oxysporum* f. sp. *lycopersici* for the management of
665 Fusarium wilt in tomato. Rhizosphere, 2, 72-74.

666 Gullino, M.L., Migheli, Q., Mezzalmaa, M., 1995. Risk analysis in the release of biological control
667 agents. Antagonistic *Fusarium oxysporum* as a case study. Plant Dis., Saint Paul, 79, 1193-1201.

668 Gupta, R., Bru, D., Bisaria, V.S., Philippot, L. and Sharma, S., 2012. Responses of *Cajanus cajan*
669 and rhizospheric N cycling communities to bioinoculants. Plant Soil, 358, 143-154.

670 Hagn, A., Wallisch, S., Radl, V., Munch, J.C. and Schloter, M., 2007. A new cultivation
 671 independent approach to detect and monitor common *Trichoderma* species in soils. J. Microbiol.
 672 Meth., 69, 86-92.

673 Höper, H., Steinberg, C., Alabouvette C., 1995. Involvement of clay type and pH in the mechanisms
 674 of soil suppressiveness to Fusarium wilt of flax. Soil Bio. Biochem., 27, 955-967.

675 Huang, C.-H., Roberts, P.D., Datnoff, L.E., 2012. Fusarium diseases of tomato. In: Gullino, M.L.,
 676 Katan, J., Garibaldi, A. (Eds.), Fusarium wilts of greenhouse vegetable and ornamental crops. APS
 677 Press, St. Paul, MN, USA, pp. 145-158.

678 Jangir, M., Pathak, R., Shar, S., Sharma, S., 2018. Biocontrol mechanisms of *Bacillus* sp. isolated
 679 from tomato rhizosphere against *Fusarium oxysporum* f. sp. *lycopersici*. Biol. Control., 123, 60-70.

680 Kim, Y.G., Kang, H.K., Kwon, K.D., Seo, C.H., Lee, H.B., Park, Y., 2015. Antagonistic activities
 681 of novel peptides from *Bacillus amyloliquefaciens* PT14 against *Fusarium solani* and *Fusarium*
 682 *oxysporum*. J. Agric. Food Chem., 63, 10380-10387.

683 Krober, M., Wibberg, D., Grosch, R., Eikmeyer, F., Verwaaijen, B., Chowdhury, S.P., Hartmann,
 684 A., Puhler, A., Schluter, A., 2014. Effect of the strain *Bacillus amyloliquefaciens* FZB42 on the
 685 microbial community in the rhizosphere of lettuce under field conditions analyzed by whole
 686 metagenome sequencing. Front. Microbiol., 5, 252.

687 Inami, K., Yoshioka, C., Hirano, Y., Kawabe, M., Tsushima, S., Teraoka, T., Arie., T., 2010. Real-
 688 time PCR for differential determination of the tomato wilt fungus, *Fusarium oxysporum* f. sp.
 689 *lycopersici*, and its races. J. Gen. Plant Pathol., 76, 116 -121.

690 Jones, J.B., Jones, J.P., Stall, J.P., Zitter, T.A., 1991. Compendium of tomato diseases. APS Press,
 691 St Paul, MN, USA.

692 Jacobsen, B.J., Zidack, N.K., Larson B.J., 2004. The role of *Bacillus*-based biological control
 693 agents in integrated pest management systems: Plant Diseases. Phytopathol., 94, 1272-1275.

694 Lane, D. 1991. 16S/23S rRNA sequencing. In Nucleic acid techniques systematics (1st edn) ed.
 695 Stackebrandt, A. and Goodfello, M. pp. 115-257. West Sussex: Wiley.

696 Larkin, R.P., 2015. Soil health paradigms and implications for disease management. *Annu. Rev.*
697 *Phytopathol.*, 53, 199-221.

698 Lopes, C.A., Ávila, A.C., 2005. Doenças do tomateiro. Embrapa Hortaliças, Brasília, p 151.

699 McGovern, R.J., 2015. Management of tomato diseases caused by *Fusarium oxysporum*. *Crop*
700 *Prot.*, 73, 78-92.

701 Lorito, M., Woo, S.L., Garcia, Fernandez I., Colucci, G., Harman, G.E., Pintor-Toro, J.A.,
702 Filippone, E., Muccifora, S., Lawrence, C.B., Zoina, A., Tuzun, S., Scala ,F., 1998. Genes from
703 mycoparasitic fungi as a novel source for improving plant resistance to fungal pathogens. *Proc.*
704 *Natl. Acad. Sci. USA*, 95, 7860-786.

705 Lorito, M., Woo, S. L., Harman, G. E., Monte, E., 2010. Translational research on *Trichoderma*:
706 from 'omics to the field. *Annu. Rev. Phytopathol.*, 48, 395-417.

707 Lorito. M., Woo, S.L., 2015. “*Trichoderma*: a multi - purpose toll for integrated pest management”,
708 in *Principles of plant-microbe interactions*, (Lugtenberg, B., ed.) Springer International Publishing,
709 pp., 345-353.

710 McSpadden Gardener, B.B., Fravel, D.R., 2002. Biological control of plant pathogens: research,
711 commercialization, and application in the USA. *Plant Health Prog.*, 3, 1-18.

712 Mezzalama, M., Ghini, R., Ruffa, P., Ambrosoli, R., Garibaldi, A., 1998. Effect of antagonistic
713 *Fusarium oxysporum* on functional groups of indigenous bacteria in soil. *Appl. Soil Ecol.*, 7, 31-40.

714 Muslim, A., Horinouchi, H., Hyakumaki, M., 2003. Control of *Fusarium* crown and root rot of
715 tomato with hypovirulent binucleate *Rhizoctonia* in soil and rock wool systems. *Plant Dis.*, 87, 739-
716 747.

717 Olivain, C, Alabouvette, C. 1999. Process of tomato root colonization by a pathogenic strain of
718 *Fusarium oxysporum* f. sp. *lycopersici* in comparison with a non-pathogenic strain. *New Phytol.*,
719 141, 497-510.

720 Muyzer, G., de Waal, E.C. and Uitterlinden, A.G., 1993. Profiling of complex microbial
 721 populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction
 722 amplified genes coding for 16S rRNA. Appl. Environ. Microbiol., 59, 695-700.

723 Olsen, S., Cole, C., Watanabe, F. and Dean, L., 1954. Estimation of available phosphorus in soils
 724 by extraction with sodium bicarbonate. USDA Circular Nr 939. Washington, DC: US Gov. Print.
 725 Office.

726 Pal, K.K., McSpadden Gardener, B., 2006. Biological control of plant pathogens. Plant Health Instr.
 727 <http://dx.doi.org/10.1094/PHI-A-2006-1117-02>.

728 Mishra, P.K., Fox, R.T.V., Culham, A., 2003. Development of a PCR-based assay for rapid and
 729 reliable identification of pathogenic *Fusaria*. FEMS Microbiol. Lett., 218, 329-332.

730 Pugliese, M., Liu, B.P., Gullino, M.L., Garibaldi A., 2008. Selection of antagonists from compost to
 731 control soil-borne pathogens. J. Plant Dis. Protect., 115, 220-228.

732 Pugliese, M., Gullino, M.L., Garibaldi, A., 2010. Efficacy of microorganisms selected from
 733 compost to control soil-borne pathogens. Commun. Agric. Appl. Biol. Sci., 75, 665-9.

734 Pugliese, M., Liu, B., Gullino, M.L., Garibaldi, A., 2011. Microbial enrichment of compost with
 735 biological control agents to enhance suppressiveness to four soil-borne diseases in greenhouse. J
 736 Plant Dis. Protect., 8, 45-50.

737 Qiu, M., Zhang, R., Xue, C., Zhang, S., Zhang, S.L. and Shen, Q., 2012. Application of bio-organic
 738 fertilizer can control Fusarium wilt of cucumber plants by regulating microbial community of
 739 rhizosphere soil. Biol. Fertil. Soils 48, 807-816.

740 Rep, M., Dekker, H.L., Vossen, J.H., de Boer, A.D., Houterman, P.M., Speijer, D., et al. 2002.
 741 Mass spectrometric identification of isoforms of PR proteins in xylem sap of fungus-infected
 742 tomato. Plant Physiol., 130, 904-917.

743 Rojo, F.G., Reynoso, M.M., Ferez, M., Chulze, S.N., Torres, A.M., 2007. Biological control by
 744 *Trichoderma* species of *Fusarium solani* causing peanut brown root rot under field conditions. Crop
 745 Prot., 26, 549-555.

746 Rotthauwe, J.H., Witzel, K.P. and Liesack, W., 1997. The ammonia monooxygenase structural gene
747 amoA as a functional marker: molecular fine-scale analysis of natural ammonia oxidizing
748 populations. Appl. Environ. Microbiol., 63, 4704-4712.

749 Savazzini, F., Oliveira Longa C.M., Pertot, I., 2009. Impact of the biocontrol agent *Trichoderma*
750 *atroviride* SC1 on soil microbial communities of a vineyard in northern Italy. Soil Biol. Biochem.,
751 41, 1457-1465.

752 Shen, Z., Ruan, Y., Xue, C., Zhong, S., Li, R. and Shen, Q., 2015. Soils naturally suppressive to
753 banana Fusarium wilt disease harbor unique bacterial communities. Plant Soil 1-2, 21-33.

754 Shores, M., Yedidia, I., Chet, I., 2005. Involvement of jasmonic acid/ethylene signalling pathway
755 in the systemic resistance induced in cucumber by *Trichoderma asperellum* T203. Phytopathol., 95,
756 76-84.

757 Sivan, A., Chet, I., 1989. The possible role of competition between *Trichoderma harzianum* and
758 *Fusarium oxysporum* on rhizosphere colonization. Phytopathol., 79, 198-203.

759 Srinivasan, K., Gilardi, G., Garibaldi, A. and Gullino, M.L., 2009. Bacterial antagonists from used
760 rockwool soilless substrates suppress Fusarium wilt on tomato. J Plant Pathol., 91, 147-154.

761 Suárez-Estrella, F., Vargas-García, C., López, M.J., Capel, C., and Moreno, J., 2007. Antagonistic
762 activity of bacteria and fungi from horticultural compost against *Fusarium oxysporum* f. sp.
763 *melonis*. Crop Prot., 26, 46-53.

764 Stein, T., 2005. *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. Mol.
765 Microbiol., 56, 845-857.

766 Vainio, E.J. and Hantula, J., 2000. Direct analysis of wood inhabiting fungi using denaturing
767 gradient gel electrophoresis of amplified ribosomal DNA. Mycol. Res., 104, 927-936.

768 Van den Elzen, PMJ., Jongendijk, E., Melchers, LS., and Cornelissen, BJC., 1993. Virus and fungal
769 resistance from laboratory to field. Phil. Trans. R. Soc. Lond. B., 342, 271-278.

770 Vinale, F., Sivasithamparan, K., Ghisalberti, E.L., Marra, R., Woo S.L., Lorito, M., 2008.
771 *Trichoderma* - plant - pathogen interactions. Soil Biol. Biochem., 40, 1-10.

772 Vitullo, D., Di Pietro, A., Romano, A., Lanzotti, V., Lima, G., 2012. Role of new bacterial
 773 surfactins in the antifungal interaction between *Bacillus amyloliquefaciens* and *Fusarium*
 774 *oxysporum*. Plant. Pathol., 61, 689-699.

775 Walters, D., and Daniell, T., 2007. Microbial induction of resistance to pathogens, p. 143-156. In
 776 Walters, D., Newton, A., and Lyon, G. (eds.), Induced resistance for plant defence. Blackwell
 777 Publishing, Oxford, UK.

778 Whipps, J.M., 2001. Microbial interactions and biocontrol in the rhizosphere. J. Exp. Bot., 52, 487-
 779 511.

780 Winding, A., Binnerup, S.J., Pritchard, H., 2004. Non-target effects of bacterial biological control
 781 agents suppressing root pathogenic fungi. FEMS Microbiol. Ecol., 47, 129-141.

782 Wessen, E., Nyberg, K., Jansson, J.K. and Hallin, S., 2010. Responses of bacterial and archaeal
 783 ammonia oxidizers to soil organic and fertilizer amendments under long-term management. Appl.
 784 Soil Ecol., 45, 193-200.

785 Wu, B., Wang, X., Yang, L., Yang, H., Zeng, H., Qiu, Y., Wang, C., Yu, J., Li, J., Xu, D., He, Z.,
 786 Chen, S., 2016. Effects of *Bacillus amyloliquefaciens* ZM9 on bacterial wilt and rhizosphere
 787 microbial communities of tobacco. Appl. Soil Ecol., 103, 1-12.

788 Yedidia, I., Shores, M., Kerem, K., Benhamou, N., Kapulnik, Y., Chet, I. 2003. Concomitant
 789 induction of systemic resistance to *Pseudomonas syringae* pv. lachrymans in cucumber by
 790 *Trichoderma asperellum* (T-203) and the accumulation of phytoalexins. Appl. Environ. Microbiol.,
 791 69, 7343-53.

792 Yergeau, E., Labour, K., Hamel, C., Vujanovic, V., Nakano-Hylander, A., Jeannotte, R., St-Arnaud,
 793 M., 2010. Patterns of *Fusarium* community structure and abundance in relation to spatial, abiotic
 794 and biotic factors in soil. FEMS Microbiol. Ecol., 71, 34-42.

795 Zhao, P., Quan, C., Wang, Y., Wang, J., Fan, S., 2014. *Bacillus amyloliquefaciens* Q-426 as a
 796 potential biocontrol agent against *Fusarium oxysporum* f. sp. *spinaciae*. J. Basic Microbiol., 54,
 797 448-456.

798 **Figure Captions**

799

800 **Figure 1** Abundance of *Fusarium oxysporum* f.sp. *lycopersici* (Fol) in the rhizosphere and bulk soil
801 for different treatments and the untreated control (SM - *Bacillus subtilis* - Serenade Max; CM -
802 Green compost plus *Trichoderma* sp. TW2; CV - green compost; TW2 - *Trichoderma* sp. TW2; C -
803 untreated control) (n = 6, means±standard errors). Different letters above the bars indicate
804 significant differences between treatments in the rhizosphere (uppercase letters) and bulk soil
805 (lowercase letters); *indicate significant differences between rhizosphere and bulk soil

806 **Figure 2** Real-time RT-PCR analysis of PR gene expression in the tomato roots for different
807 treatments and for the untreated control. Actin was used as the reference gene. The error bars show
808 standard deviations for triplicate assays.

809

Fusarium oxysporum f.sp. *lycopersici*

810

811
812 Figure 1
813
814
815
816
817
818
819
820
821

823 Figure 2

824 Tables

825 Table 1 General information on the trials and timing of the operations carried out in 2016 and 2017

Treatment	Microorganism/a.i.	Dosage	Tray treatment	Plot Treatment
Serenade max - SM	<i>Bacillus subtilis</i> QST 713	2.9X10 ¹⁰ cells/L water	T37; T44; T51;T58*	-
ANT'S COMPOST M - CM	<i>Green compost + Trichoderma</i> sp. TW2	8 g/seedling;1kg/0.1m ³ of soil	T0	- T51
ANT'S COMPOST V - CV	<i>Green compost</i>	8 g/seedling; 1kg/0.1m ³ of soil	T0	- T51
<i>Trichoderma</i> sp. TW2 -TW2	<i>Trichoderma</i> sp. TW2	1x10 ⁷ (conidia /ml)	T37; T44; T51;T58	
Untreated control - C	-	-		

826

827 *Treatments: T0 at sowing; T1, 37 days after sowing; T2, 44 days after sowing; T3: 51 days after sowing; T 4: 58 days after sowing and immediately
828 before transplanting.

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845 Table 2 Description of the primer sets and amplification details used for the quantitative PCR.

Target group	Primer (reference)	Amplification details
<i>Fusarium oxysporum</i> f.sp. <i>lycopersici</i> (Fol)	FO1(Prashant et al., 2003) 5'-ACATACCACTTGTTGCCTCG-3' FO2 (Prashant et al., 2003) 5'-CGCCAATCAATTTGAGGAACG-3'	40 cycles 95°C 15s, 60°C 60s, 72°C 45s
All bacteria (16S rRNA gene)	Eub338 (Lane 1991) 5'-ACTCCTACGGGAGGCAGCAGCAG-3' Eub518 (Muyzer et al. 1993) 5'-ATTACCGCGGCTGCTGG-3'	40 cycles 95°C 30s, 55°C 35s, 72°C 45s
All archaea (16S rRNA gene)	340F (Gantner et al., 2011) 5'-CCCTAYGGGGYGCASCAG-3' 1000R (Gantner et al., 2011) 5'-GGCCATGCACYWCYTCTC-3'	40 cycles 95°C 30s, 57°C 30s, 72°C 30s
All fungi (18S rRNA gene)	FR1 (Vainio and Hantula, 2000) 5'-AIC CAT TCA ATC GGT AIT-3' 390FF (Vainio and Hantula, 2000) 5'-CGA TAA CGA ACG AGA CCT-3'	45 cycles 95°C 30s, 50°C 30s, 70°C 60s
Ammonia oxidizing bacteria (AOB)	AmoA-1f (Rotthauwe et al. 1997) 5'-GGGGTTTCTACTGGTGGT-3' AmoA-2r (Rotthauwe et al. 1997) 5'-CCCCTCKGSAAAGCCTTCTTC-3'	45 cycles 95°C 30s, 57°C 45s, 72°C 45s, 78°C 20s
Ammonia oxidizing archaea (AOA)	Arch-amoAf (Francis et al., 2005) 5'-STAATGGTCTGGCTTAGACG-3' Arch-amoAr (Francis et al.,2005) 5'-GCGGCCATCCATCTGTATGT-3'	45cycles 95°C 30s, 53°C 45s, 72°C 45s, 78°C 20s
<i>chiA</i> gene	chiaxf (Cucu et al., 2018) 5'-ACCCTGCCGATGACACTCAG-3' chiaxr (Cucu et al., 2018) 5'-GGCAGCGATGGAGAGAAGGA-3'	35cycles 95°C 15s, 59.9°C 30s, 70°C 30s
<i>Bacillus subtilis</i>	Forward <i>B. subtilis</i> (Gao et al., 2011) 5'-TCGCGGTTTCGCTGCCCTTT-3' Reverse <i>B. subtilis</i> (Gao et al., 2011) 5'-AAGTCCCGCAACGAGCGCAA-3'	40 cycles 95°C 30s, 60°C 60s, 80°C 10s
<i>Trichoderma spp.</i>	uTf (Hagn et al., 2007) 5'-AACGTTACCAAACGTGTTG-3' uTr (Hagn et al., 2007) 5'-AAGTTCAGCGGTATTCCT-3'	35 cycles 95 °C 30 s, 55.5 °C 30 s, 72 °C 30 s

846

847

848

849

850

851

Table 3 Effect of the preventative soil treatments with BCAs and compost on disease severity at the end of trials 1 and 2, in 2016 and 2017, respectively

Treatment	Disease severity**			
	%			
	2016		2017	
<i>Bacillus subtilis</i> QST 713 - SM	6.3	±3.8 ^a	19.5	±3.0 ^a
<i>Green compost + Trichoderma</i> sp. TW2 - CM	10.4	±5.2 ^a	21.9	±2.8 ^a
<i>Green compost</i> - CV	1.9	±1.9 ^a	14.5	±2.6 ^a
<i>Trichoderma</i> sp. TW2 - TW2	3.1	±3.1 ^a	16.0	±1.3 ^a
Untreated control - C	29.7	±2.3 ^b	39.9	±2.4 ^b

*Treatments: T0 at sowing; T1, 37 days after sowing; T2, 44 days after sowing; T3: 51 days after sowing; T 4: 58 days after sowing and immediately before transplanting.

**Sixteen plants/treatment were evaluated for disease severity using the following rating scale: 0 = healthy plant, 25 = initial leaf chlorosis, 50 = severe leaf chlorosis and initial symptoms of wilting during the hottest hours of the day, 75 = severe wilting and severe symptoms of leaf chlorosis; 100 = plant totally wilted, leaves completely necrotic.

883 Table 4 Abundance of 16SrRNA (all bacteria and archaea) and 18S rRNA (fungi) in the rhizosphere
884 and bulk soil after different treatments and in the untreated control

885

Treatment	Rhizosphere soil	Bulk soil
<i>16S Bacteria (logcopy DNA⁻¹)</i>		
<i>Bacillus subtilis</i> (SM)	10.686 c	10.963 d
Green compost+TW2 (CM)	10.755 b	11.287 a
Green compost (CV)	10.454 d	10.904 d
<i>Trichoderma</i> sp. TW2 (TW2)	10.903 a	11.191 b
Untreated control - C	10.477 d	11.077 c
<i>P(F) Treat</i>	0.000	
<i>P (F) Year</i>	0.000	
<i>P(F) Treat*Year</i>	ns	
<i>P(F) Rhizo*Bulk soil</i>	0.000	
<i>16S archaea (logcopy DNA⁻¹)</i>		
<i>Bacillus subtilis</i> (SM)	7.728 a	7.995 a
Green compost+TW2 (CM)	7.748 a	7.982 a
Green compost (CV)	7.751 a	8.023 a
<i>Trichoderma</i> sp. TW2 (TW2)	7.592 b	7.974 a
Untreated control - C	7.549 b	7.988 a
<i>P(F) Treat</i>	0.000	
<i>P (F) Year</i>	0.000	
<i>P(F) Treat*Year</i>	ns	
<i>P(F) Rhizo*Bulk soil</i>	0.002	
<i>18S fungi (logcopy DNA⁻¹)</i>		
<i>Bacillus subtilis</i> (SM)	9.669 c	8.955 b
Green compost+TW2 (CM)	10.111 a	8.927 b
Green compost (CV)	9.813 b	9.226 a
<i>Trichoderma</i> sp. TW2 (TW2)	10.000 a	8.631 c
Untreated control - C	9.979 a	8.542 c
<i>P(F) Treat</i>	0.000	
<i>P (F) Year</i>	0.000	
<i>P(F) Treat*Year</i>	ns	
<i>P(F) Rhizo*Bulk soil</i>	0.000	

886

887

888 Table 5 Abundance of bacterial (AOB) and archaeal (AOA) *amoA* genes in the rhizosphere and
889 bulk soil after different treatments and in the untreated control

Treatment	Rhizosphere soil		Bulk soil	
<i>AOB (logcopy DNA⁻¹)</i>				
<i>Bacillus subtilis</i> (SM)	7.316	a	6.233	c
Green compost+TW2 (CM)	7.280	a	6.665	b
Green compost (CV)	7.321	a	6.686	b
<i>Trichoderma</i> sp. TW2 (TW2)	7.387	a	6.603	b
Untreated control - C	7.125	b	6.877	a
<i>P(F) Treat</i>	<i>0.000</i>			
<i>P (F) Year</i>	<i>0.000</i>			
<i>P(F) Treat*Year</i>	<i>ns</i>			
<i>P(F) Rhizo*Bulk soil</i>	<i>0.007</i>			
<i>AOA (logcopy DNA⁻¹)</i>				
<i>Bacillus subtilis</i> (SM)	6.657	b	7.915	b
Green compost+TW2 (CM)	6.974	a	7.884	b
Green compost (CV)	6.721	b	8.149	a
<i>Trichoderma</i> sp. TW2 (TW2)	6.674	b	7.935	b
Untreated control - C	6.681	b	7.926	b
<i>P(F) Treat</i>	<i>0.000</i>			
<i>P (F) Year</i>	<i>0.000</i>			
<i>P(F) Treat*Year</i>	<i>0.015</i>			
<i>P(F) Rhizo*Bulk soil</i>	<i>0.000</i>			

903 Table 6 Abundance of *Bacillus subtilis* and *Trichoderma* spp. genes in the rhizosphere and bulk soil
 904 after different treatments and the untreated control

905
 906

Treatment	Rhizosphere soil		Bulk soil	
	<i>Bacillus (logcopy DNA⁻¹)</i>			
<i>Bacillus subtilis</i> (SM)	6.024	a	6.458	a
Green compost +TW2 (CM)	4.702	c	5.798	b
Green compost (CV)	5.980	b	5.662	c
<i>Trichoderma</i> sp. TW2 (TW2)	4.673	d	4.865	d
Untreated control - C	4.685	d	4.931	d
<i>P(F) Treat</i>	0.000			
<i>P (F) Year</i>	0.000			
<i>P(F) Treat*Year</i>	ns			
<i>P(F) Rhizo*Bulk soil</i>	0.013			
	<i>Trichoderma (logcopy DNA⁻¹)</i>			
<i>Bacillus subtilis</i> (SM)	4.762	d	3.731	c
Green compost+TW2 (CM)	6.922	a	4.483	a
Green compost (CV)	5.494	c	3.652	b
<i>Trichoderma</i> sp. TW2 (TW2)	6.832	b	4.632	a
Untreated control - C	4.791	d	3.977	c
<i>P(F) Treat</i>	0.000			
<i>P (F) Year</i>	0.000			
<i>P(F) Treat*Year</i>	ns			
<i>P(F) Rhizo*Bulk soil</i>	0.016			

907
 908
 909
 910
 911
 912
 913
 914
 915

916 Table 7 Abundance of the *chiaA* gene in the rhizosphere and bulk soil after different treatments and
 917 the untreated control

918
 919

Treatment	Rhizosphere soil		Bulk soil	
<i>chiA</i> gene (logcopy DNA ⁻¹)				
<i>Bacillus subtilis</i> (SM)	3.632	c	4.384	b
Green compost+TW2 (CM)	5.736	a	4.363	b
Green compost (CV)	4.338	b	4.403	a
<i>Trichoderma</i> sp. TW2 (TW2)	5.789	a	4.321	b
Untreated control - C	3.373	c	3.890	c
<i>P</i> (<i>F</i>) Treat	0.000			
<i>P</i> (<i>F</i>) Year	0.000			
<i>P</i> (<i>F</i>) Treat*Year	ns			
<i>P</i> (<i>F</i>) Rhizo*Bulk soil	0.015			

920
 921

Table 8 Pearson's correlation coefficient for the BCAs, Fol race 1 abundance and chemical properties in the rhizosphere (rhizo) and bulk soil samples

Property	<i>Bacillus subtilis</i>		<i>Trichoderma</i> spp.		Fol	
	rhizo	bulk soil	rhizo	bulk soil	rhizo	bulk soil
pH	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
TN	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
NH ₄ ⁺	<i>ns</i>	<i>ns</i>	0.78***	<i>ns</i>	<i>ns</i>	<i>ns</i>
NO ₃ ⁻	-0.5	<i>ns</i>	-0.75***	<i>ns</i>	<i>ns</i>	<i>ns</i>
TP	-0.60***	-0.40**	0.5**	<i>ns</i>	-0.4**	-0.5**
AP	0.40**	-0.50**	0.67**	0.50**	0.575**	0.468*
DOC	-0.7***	0.5 *	-0.8***	0.67***	0.676***	0.67**

Significance levels: not significant-ns: $p > 0.05$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

939 Table 9 Pearson’s correlation coefficients between microbial gene abundance and disease severity
940 (DS) in rhizosphere (rhizo) and bulk soil samples

941

942

Microbial abundance	Disease severity (DS)	
	rhizo	bulk soil
Fol	0.75***	0.63***
16S Bacteria	ns	ns
16S Archaea	ns	ns
18S Fungi	0.67	ns
AOB	-0.55**	0.65***
AOA	-0.57***	-0.76***
<i>Bacillus</i>	-0.45**	-0.47*
<i>Trichoderma</i>	-0.75***	-0.60***
<i>chiA</i> gene	-0.55***	-0.65***

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960 Significance levels: not significant-ns: p>0.05; *p<0.05; **p<0.01; ***p<0.001

961

962