

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Biodiversity of leaf litter fungi in streams along a latitudinal gradient

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1728420> since 2020-02-19T09:23:18Z

Published version:

DOI:10.1016/j.scitotenv.2019.01.122

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Biodiversity of leaf litter fungi in streams along a latitudinal gradient

Sahadevan Seenaa^{1,2*}, Felix Bärlocher³, Olímpia Sobral^{1,2}, Mark O. Gessner^{4,5,6} David Dudgeon⁷
Brendan G. McKie⁸, Eric Chauvet⁹, Luz Boyero^{10,11,12}, Verónica Ferreira^{1,13}, André
Frainer^{14,15}, Andreas Bruder¹⁶, Christoph D. Matthaei¹⁶, Stefano Fenoglio¹⁷, Kandikere R.
Sridhar¹⁸, Ricardo J. Albariño¹⁹, Michael M. Douglas^{20,21}, Andrea C. Encalada^{22,1}, Erica
Garcia²¹, Darren P. Giling⁵, Sudeep D. Ghate¹⁸, Vítor Gonçalves²³, Tomoya Iwata²⁴, Andrea
Landeira-Dabarca²², Damien McMaster²¹, Adriana O. Medeiros²⁵, Josheena Naggea²⁶, Jesús
Pozo¹⁰, Pedro M. Raposeiro²³, Christopher M. Swan²⁷, Nathalie S.D. Tenkiano²⁸, Catherine M.
Yule²⁶, Manuel A.S. Graça^{1,13}

*Correspondence should be addressed to S. Seenaa (seenaa.sahadevan@gmail.com)

¹MARE-Marine and Environmental Sciences Centre, Department of biology, University of Coimbra, PT-3004-517 Coimbra, Portugal.

²Centre of Molecular and Environmental Biology (CBMA), Department of Biology, University of Minho, Campus de Gualtar, 4710-057, Braga, Portugal.

³Department of Biology, Mt. Allison University, Sackville, NB, E4L 1G7, Canada.

⁴Department of Experimental Limnology, Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Alte Fischerhütte 2, 16775 Stechlin, Germany.

⁵Department of Ecology, Berlin Institute of Technology (TU Berlin), Ernst-Reuter-Platz 1, 10587 Berlin, Germany.

⁶Berlin Brandenburg Institute of Advanced Biodiversity Research (BBIB), Altensteinstraße 34, 14195 Berlin, Germany.

⁷School of Biological Sciences, The University of Hong Kong, Pokfulam, Hong Kong.

⁸Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, P.O. Box 7050, SE-75007, Uppsala, Sweden.

⁹EcoLab, Université de Toulouse, CNRS, INPT, UPS, Toulouse, France.

¹⁰Faculty of Science and Technology, University of the Basque Country (UPV/EHU), Leioa, Spain.

¹¹IKERBASQUE, Basque Foundation for Science, Bilbao, Spain.

¹²College of Science and Engineering, James Cook University, Townsville, Qld, Australia.

¹³Department of Life Sciences, University of Coimbra, PT-3004-517, Coimbra, Portugal.

¹⁴Norwegian College of Fishery Science, UiT The Arctic University of Norway, 9037, Tromsø, Norway.

¹⁵Department of Arctic and Marine Biology, UiT The Arctic University of Norway, 9037, Tromsø, Norway.

¹⁶Department of Zoology, University of Otago, PO Box 56, Dunedin 9054, New Zealand.

¹⁷Department of Science and Technological Innovation, University of Piemonte Orientale, Alessandria, I-15121, Italy.

¹⁸Department of Biosciences, Mangalore University, Mangalagangothri, Mangalore 574 199, Karnataka, India.

¹⁹Fotobiology Lab, INIBIOMA, UNComahue-CONICET, 8400 Bariloche, Argentina.

²⁰School of Earth and Environment, University of Western Australia, Perth, WA 6009, Australia. ²¹Research Institute for the Environment and Livelihoods, Charles Darwin University, Darwin, NT 0909, Australia.

²²Laboratorio de Ecología Acuática, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Diego de Robles y Vía Interoceánica, 17-1200-841 Quito, Ecuador.

²³Centro de Investigação em Biodiversidade e Recursos Genéticos, CIBIO, InBIO Laboratório Associado, Pólo dos Açores & Departamento de Biologia da Universidade dos Açores, Ponta Delgada, Açores, Portugal.

²⁴Faculty of Life and Environmental Sciences, University of Yamanashi, Takeda, Kofu 400-8510, Japan.

²⁵Laboratório de Microbiologia Ambiental Departamento de Botânica, Instituto de Biologia, Universidade Federal da Bahia, Campus Ondina, 40170-115 Salvador, BA, Brazil.

²⁶School of Science, Monash University Malaysia, Bandar Sunway, Selangor 47500, Malaysia.

²⁷Department of Geography & Environmental Systems, University of Maryland, Baltimore County, 211 Sondheim Hall, 1000 Hilltop Circle, Baltimore, MD 21250 USA.

²⁸Université Julius N'Yerere de Kankan, Kankan, Guinea.

Abstract

Global patterns of biodiversity have emerged for soil microorganisms and particularly for macroorganisms. The significance of microbial functions in ecosystems is also well established. Virtually unknown, however, are large-scale patterns of microbial diversity in freshwaters, despite the importance of these ecosystems as hotspots of biodiversity and biogeochemical processes. Here we report on the first large-scale study of biodiversity of leaf litter fungi in streams along a latitudinal gradient unravelled by Illumina sequencing. The study is based on fungal colonization experiments of standardised plant litter in 19 globally distributed stream locations between 69°N and 44°S. Fungal richness showed a hump-shaped distribution along the absolute latitude. Strikingly, community composition of fungi was more clearly grouped according to thermal preferences than to biogeography. Our results suggest that identifying differences in key environmental drivers, such as temperature, among taxa and ecosystem types is critical to developing an overarching theory of global diversity distribution of aquatic fungi.

Introduction

The role of freshwater ecosystems as components of the global carbon cycle is being increasingly acknowledged (Perkins et al. 2012; Martinez et al. 2014). Streams are hotspots of CO₂ emission (Perkins et al. 2012; Battin et al. 2009) as they receive organic carbon of terrestrial origin and its in-stream decomposition releases CO₂ to the atmosphere. The underlying processes, including the decomposition of terrestrial plant litter, are mainly driven by the activities of fungi (Boyero et al. 2011; Duarte et al. 2016b; Duarte et al. 2015; Seena et al. 2017). Importantly, decomposition rates of litter in streams can be affected by fungal diversity, either directly or through trophic effects (Jabiol et al. 2013b). Therefore, it is essential

to understand the global distribution and richness of aquatic fungi, to allow predictions of ecosystem responses to global changes (Violle et al. 2014).

There has been considerable debate as to whether microbes follow similar global distribution patterns as plants and animals (Fuhrman et al. 2008; Fierer et al. 2011; Azovsky and Mazei 2013; Tedersoo et al. 2014). Currently discussed topics include whether microbes follow a latitudinal diversity gradient characterized by increasing richness from the poles to the tropics (Hillebrand 2004; Mittelbach et al. 2007; Fuhrman et al. 2008; Andam et al. 2016) and to what extent biogeographic history structures present-day species distributions. The importance of plate tectonics as a significant factor governing the global distribution of plants and animals is well established (Holden 2012; Cox et al. 2016a; Briggs 1995), with many taxa following either the Laurasian or Gondwanan distribution patterns (Holden 2012). In contrast, one of the most enduring tenets in microbial ecology is Baas-Becking's hypothesis that "*everything is everywhere, but the environment selects*" (Baas-Becking 1934; O'Malley 2007), implying that latitudinal and biogeographical patterning of microbial diversity is weak or non-existent. Nonetheless, Taylor et al. (2006) concluded that microfungi do have a biogeography and that fungal species do not necessarily occur everywhere.

The richness of soil fungi (with some exceptions) generally decreases towards the poles (Tedersoo et al. 2014; Sul et al. 2013), with soil fungi that show strong dispersal abilities dominating at high latitudes (Cox et al. 2016b). Large-scale studies have also demonstrated that soil fungal richness is influenced by climate as well as by chemical and physical soil properties (Tedersoo et al. 2014; Zhou et al. 2016). Lastly, evidence exists for strong geographic structure and dispersal limitation of soil fungi (Talbot et al. 2014; Tedersoo et al. 2014). In a previous study, species richness of aquatic hyphomycetes, a polyphyletic group of stream fungi that assume dominant roles in litter decomposition, peaked at mid-latitudes and was lowest towards

the extremes of the latitudinal gradient on the northern hemisphere, with the exact shape of the diversity pattern modified by the chemistry of the litter examined (Jabiol et al. 2013a).. However, the study was based on only five streams and therefore of limited statistical power. A globally congruent pattern of stream microbial diversity has yet to emerge.

The objective of this study was to determine the community composition and diversity of litter-associated fungi, assessed as richness of operational taxonomic units (OTUs), in a coordinated global-scale study stretching over a 113° latitudinal gradient of 19 stream locations on five continents. Given that latitude is widely accepted as an overarching climate surrogate (Parmesan and Yohe 2003; Jetz et al. 2008; Boyero et al. 2011; Jabiol et al. 2013a), we tested the hypothesis that fungal richness decreases with latitude, similar to the pattern described for macroorganisms. Furthermore, we predicted that the global distribution of microbial taxa would follow the terrestrial biogeographic realms, eight of which are generally recognized to be based on distributional patterns of terrestrial species that result from isolation of populations by continental drift (Olson et al. 2001).

Materials and methods

Study sites and field work

A total of 19 study streams were distributed across both hemispheres with locations extending from 69° N to 44° S (Table 1 and Fig. 1). Mean annual air temperature (°C) and rainfall (mm) data were obtained from climate-data.org (<http://en.climate-data.org/>; accessed February 2016) and AIC (Autoridad Interjurisdiccional de Cuencas de los Ríos Limay, Neuquén y Negro, Bureau of Water Resources Management, Argentina; <http://www.aic.gov.ar/aic/default.aspx#v>; accessed February 2016). Characteristic properties of the experimental streams were: (1) low order (1-3 according to Strahler (Strahler 1957)); (2) <50 cm depth and <5 m wide; (3) coarse substrates, generally dominated by cobbles; (4) no

major anthropogenic impacts; and (5) no invasive tree species. Stream physico-chemical characteristics including concentrations of nutrients (nitrogen [N] and phosphorus [P]) were determined (APHA 1995) when the litter was deployed and retrieved.

Alnus glutinosa (L.) Gaertn. (black alder: Betulaceae) leaves were collected at a single site on the banks of the Mondego River at Lages, Coimbra (40°11'21"N, 8°25'30"W"). Black alder was chosen because the genus is widespread in the Holarctic and also occurs in the Neotropics (Boyero et al. 2011) and has high-quality leaves (Royer and Minshall 2001; Hladysz et al. 2009; Fernandes et al. 2014). Although alder trees do not occur in some of the study regions, their high-quality leaves (i.e., soft, high nitrogen concentrations) do not impose any colonization challenge to microbial communities (Fernandes et al. 2014; Royer and Minshall 2001; Hladysz et al. 2009; Chauvet et al. 2016). Moreover, the species has been previously used as a standard litter in global decomposition studies (Boyero et al. 2011), and the leaves are readily consumed by tropical detritivores (Encalada et al. 2010).

Kits containing air-dried alder leaves (three replicates, each containing 2 g of leaves), fine-mesh (0.5 mm) nylon bags, DNAGard[®] (Biomatrica, San Diego, CA, USA) and protocols were shipped from Coimbra, Portugal, to the other 18 locations distributed on all continents. DNA was extracted and pooled from 10 leaf discs (12 mm diameter) cut from randomly selected alder leaves and composition of the initial microbial community was determined by Illumina MiSeq sequencing before shipping. The leaves were not sterilized before colonization to avoid changes in litter chemistry (Howard and Frankland 1974). Microbes initially present on the litter have little influence on microbial colonization dynamics (Bärlocher and Kendrick 1974), since stream microbes are rapid colonizers that quickly outcompete terrestrial taxa (Nikolcheva et al. 2005; Frossard et al. 2013).

Strictly standardized litter colonization experiments were conducted in the 19 study streams during the dry season in tropical and during autumn in temperate and arctic streams. At each site, litter bags were deployed in riffle areas rich in oxygen (see Table S2) at water depths of less than 30 cm. Experiments were terminated when an estimated 40-50 % of the initial litter mass was lost, as inferred from previous decomposition studies (Boyero et al. 2011; Boyero et al. 2015). The exact colonization period at each site is given in Table S1. Three litter bags were retrieved and 10 leaf discs (12 mm diameter) were cut per bag with a sterile cork borer, immediately placed in 3 sterile screw-cap tubes containing 1 ml of DNAgard[®] solution and sent to the laboratory at the University of Coimbra, Portugal, for DNA extraction.

DNA extraction, Miseq sequencing and diversity analysis

From each replicate set of 10 leaf discs, microbial DNA was extracted with the PowerSoil[®] DNA isolation kit (MoBio laboratories, Carlsbad, CA, USA) according to the manufacturer's instructions. The concentration of extracted DNA (>20 ng/ μ l) was confirmed with a NanoDrop 2000c spectrophotometer (Wilmington, DE, USA) before storing the DNA at -20 °C. The DNA of all replicate samples from each country were pooled and sent to Research and Testing Laboratory (Lubbock, TX, USA; www.researchandtesting.com) for Illumina MiSeq sequencing using ITS primer pairs for fungi (ITS3 F: 5'GCATCGATGAAGAACGCAGC 3' and ITS4R: 5'TCCTCCGCTTATTGATATGC3') (White et al. 1990).

Quality trimming was performed on the fastq using the SolexaQA++ dynamictrim utility (Cox et al. 2010) with a minimum threshold of 25 and if after quality trimming the length of the reads became less than 50 bp it was removed. The forward and reverse reads of our dataset was merged by using FLASH (Magoč and Salzberg 2011). Dereplication step was used to remove duplicated reads by using VSEARCH software (Rognes et al. 2016). Chimera was removed using Mothur (version 1.39.5; Schloss et al. 2009) using the unified system for DNA based

fungus species (UNITE) and international nucleotide sequence databases (INSDC) fungus ITS databases (version released on 28.06.2017) as reference (Koljalg et al. 2013). The clustering of the OTUs was achieved by using Swarm (version 2.2.2; Mahé et al. 2014) and the longest sequence from each OTU was selected as representatives of each OTU. The singletons were removed from the analyses and assignment of taxonomy was performed with a BLAST of the OTU representatives against UNITE+INSDC fungus ITS databases (version released on 28.06.2017, <https://unite.ut.ee>). The robust assignment to the fungus kingdom was made for the sequences with E-values $< e^{-50}$ and sequence similarity $> 75\%$. Moreover, query sequences with an E-value between e^{-08} and e^{-50} with sequence similarity $> 75\%$ were manually checked against the 100 best-matching sequences for accurate assignment (Li et al. 2016). Raw sequences from the alder leaves before exposure in the stream and after retrieval of the leaves were deposited in the National Center for Biotechnology Information (NCBI) database (Sequence Read Archives; SRA) under accession numbers SRP072752 and SRP100503, respectively.

Data analysis

Rarefaction curves showing the number of sequences versus the number of fungus OTUs in all locations were computed to check whether OTU richness was close to saturation. The control OTUs were removed from the OTUs of the samples and the rarefied fungus richness and relative abundances were calculated. All analyses were conducted in R (R Development Core Team, 2011), using vegan package (Oksanen et al. 2011). Barplots of the fungus phylum were generated using average OTU relative abundance.

We conducted linear regressions (Zuur et al. 2007) to assess the relationship between the physico-chemical characteristics of the stream and the locations with latitude. A quadratic term was added to the linear regression model to assess the relationship between OTU richness and

Latitude. All regressions were performed using (R Development Core Team 2011). All data were checked for normality and homoscedasticity; where necessary, a natural log transformation was applied to meet these assumptions. A Bray–Curtis similarity matrix (Bray and Curtis 1957) was calculated based on the log transformed relative abundance data. Influence Differences in fungal community structure were displayed using Canonical Analysis of Principal Coordinates (CAP; Anderson and Willis 2003) based on relative abundances of OTUs. Permutational multivariate analysis of variance (PERMANOVA; Anderson et al. 2008) tested for differences between sample clouds which were separated by two CAP axes. Unrestricted permutation of the raw data at 9999 permutations was used for PERMANOVA. PRIMER 6 was used to perform PERMANOVA and CAP analyses (Version 6, Primer-E Ltd, Plymouth, UK; Clarke and Gorley 2006) A Venn diagram was generated with Venny 2.1 (Oliveros 2015) to determine the shared OTUs between the sample clouds which were separated by two CAP axes.

Results

Physico-chemical characteristics of streams

Mean stream water temperature during the study (3.1–26.2 °C), mean annual air temperature (0.9–27.4 °C), and mean annual rainfall (527–4273 mm) at the 19 locations of our study (Table 2) were all significantly related to absolute latitude (Fig. 2). The site in Ecuador appears to contradict this trend, however, it was located at an unusually high altitude (3061 m asl). There were no latitudinal trends in mean pH, conductivity or concentrations of dissolved oxygen, phosphate or nitrate.

Diversity of litter-associated fungi

Illumina MiSeq sequencing of colonized alder leaves revealed 1,093,416 reads. These were reduced to 859,892 after filtering for quality and length after joining the forward and reverse

reads it generated in total 229,194 reads. After dereplication this was further reduced to 44138 reads; after performing the chimera 42,618 reads remained (Table S2). The no hits and OTUs belonging to the kingdom Plantae (14.6%) were removed. The removal of the control OTUs from the samples yielded 1311 OTUs (Table S3). Collectively, fungal OTUs were assigned primarily to the phylum Ascomycota (79.71 %), followed by Mucoromycota (16.8 %), Basidiomycota (0.43%) and Chytridiomycota (0.2%). The remaining 10.62 % of the fungal OTUs were unknown. The relative abundances of fungal classes varied among locations: streams with water temperature (Fig. 3) above 8.6°C (except in Japan) consisted mainly filamentous fungi of moulds (Azores, Brazil, Canada, Ecuador, Hong Kong, Malaysia, Portugal, Spain and USA), Yeasts (India and Guinea) and unknown fungi and chytrids (Australia).

Fungal richness

Rarefaction curves of the streams had clearly distinct and separate trajectories, suggesting that they realistically represented inherent diversity patterns (Fig. S1). OTU richness was lowest at latitudinal extremes, peaked at mid-latitude regions (Fig. 4), and was insensitive to major climatic factors like temperature or rainfall.

Endemism and ubiquity

A total of 950 OTUs (out of 1311 OTUs) were unique to single locations and classified as exclusive endemics. The richness of endemic fungal species did not follow any specific geographical trend (Fig. S2). India and Hong Kong (97 %) had the highest fungal endemism, closely followed by Malaysia (72%), Portugal (55%) and Japan (54%). Australia and Guinea did not have any exclusive endemics. 93% of the total OTUs were restricted to two localities, showing moderate endemism. The majority of endemic fungi were in the Neotropic realm (55%) followed by Indo-Malay realm (40 %), the Palearctic (35 %), Australasia (21%) and

Nearctic realm (19 %). In our study, the Afrotropic realm did not have endemic fungi. Out of all fungi, 14.5% were aquatic hyphomycetes (litter decomposers), however, in France (48%), Germany (47.5) and Italy (46.8%) almost half of the fungi were aquatic hyphomycetes. The aquatic hyphomycete, *Lemonniera aquatica* was the most widespread species and occurred in 8 locations.

Influence of water temperature

To discriminate among drivers of community structure, canonical analysis of principal coordinates (CAP) using Bray–Curtis similarities from log-transformed relative species abundance data of OTUs was used. The resulting pattern was clear and the communities were separated into three groups ($\leq 8.1^\circ\text{C}$, $8.6\text{-}19.8^\circ\text{C}$ and $\geq 21.3^\circ\text{C}$) based on stream temperature, except for Japan (Fig. 5). The overall classification accuracy rate based on $\leq 8.1^\circ\text{C}$, $8.6\text{-}19.8^\circ\text{C}$ and $\geq 21.3^\circ\text{C}$ is 83.3 %, 87.5% and 80% respectively. The results demonstrated that the first ($\delta^2=0.9371$) and the second ($\delta^2=0.7978$) squared canonical correlation was large. The first canonical axis separated the communities of the tropical streams from others and the second axis separated the communities from the stream with the $8.6\text{-}19.8^\circ\text{C}$ temperature range to the others. A permutational multivariate analysis of variance (PERMANOVA) test demonstrated a significant effect of the stream temperature ($F_{2,18} = 2.19$, $P = 0.0001$). 773, 318 and 350 OTUs exclusively belonged to $\leq 8.1^\circ\text{C}$, $8.6\text{-}19.8^\circ\text{C}$ and $\geq 21.3^\circ\text{C}$, respectively. Streams with water temperature $\leq 8.1^\circ\text{C}$ and $8.6\text{-}19.8^\circ\text{C}$ shared 63 OTUs while streams with $\leq 8.1^\circ\text{C}$ and $\geq 21.3^\circ\text{C}$, and $8.6\text{-}19.8^\circ\text{C}$ and $\geq 21.3^\circ\text{C}$ water temperature shared 57 and 21 OTUs respectively (Fig. 6).

Discussion

One of the most studied and widely documented macroecological pattern is the latitudinal gradient in species richness. A decrease in species richness with latitude is the oldest and most

fundamental pattern of the distribution of life on earth, particularly in plants and animals (Peay et al 2016). As many factors (such as temperature, rainfall) change with latitude it is important to determine if litter associated fungi correspond to macroecological patterns, in order to recognize the general drivers of fungal biodiversity. A key discovery of our large-scale coordinated study is that the fungal richness diverges from the conventional macroecological pattern. In our study, it follows a hump-shaped relationship with absolute latitude with a peak at mid-latitude regions (temperate streams) (Fig. 4). This could be possibly due to more varied niches owing to the strong seasonality in the temperate streams (Shearer et al., 2007; Jabiol et al). In our study, stream temperature, air temperature and rainfall decreased with latitude, implying that extreme conditions were not favorable for litter associated fungi. A hump-shaped curve along the latitudinal gradient was also described in previous studies pertaining to aquatic hyphomycetes, the major litter decomposers (Jabiol et al. 2013a; Wood-Eggenschwiler and Bärlocher 1983; Shearer et al. 2007). This discrepancy differs strikingly from the global soil fungal study which demonstrated macroecological patterns similar to those of other organisms, albeit with several notable exceptions.

Stream and soil environments and their microbial communities differ in many fundamental ways (Vinson and Hawkins 2003, Bärlocher and Boddy 2016), implying that global microbial diversity patterns need not necessarily match. For instance, precipitation events affect fungal species richness in streams through hydrological disturbances such as droughts and floods, whereas the main control in terrestrial ecosystems is primary productivity (Hawkins and Diniz-Filho 2004; Hawkins et al. 2003). Similarly, moisture is a limiting factor in soils but not in (permanent) streams.

Our study demonstrates the tendency of litter associated fungi to occur in thermal bands. We identified three fungal clusters encompassing locations from different biogeographic realms.

These clusters corresponded to different stream water temperature bands, indicating that water temperature plays an essential role in determining the occurrence and composition of litter-associated microbes in streams across the globe. Given that aquatic fungi play a key role in litter decomposition in streams (Baldy et al. 1995; Weyers and Suberkropp 1996), and species differ in their preferred thermal ranges for growth (Krauss et al. 2011), reproduction (Chauvet and Suberkropp 1998) and activity (Ferreira and Chauvet 2011), large-scale variation and changes in temperature are likely to influence not only the composition of fungal communities (i.e. biodiversity patterns) but also rates of carbon and nutrient cycling (Martinez et al. 2014; Dang et al. 2009).

Our results show that even the most common fungi, *Lemonniera aquatica* was limited to 8 locations. Thus, freshwater fungi associated with decomposing litter appear to follow the 'moderate endemicity model' of microbial biogeography initially proposed for small eukaryotes (Foissner 1999). However, this conclusion needs further backing by empirical data on a wide range of substrate and habitats before generalisations about global distribution patterns can be confidently made. This includes comparisons of different locations under similar climates and/or at similar latitudes to substantiate whether community similarity across regions is typically temperature-driven, as our results suggest.

Some published evidence indicates that the biogeography of freshwater fungi classified as aquatic hyphomycetes is species-specific (Duarte et al. 2012), and that, as confirmed here, community composition in geographically distant locations within comparable climatic zones may be similar (for review, see Duarte et al. 2016a). A caveat to generalisations about biogeographic patterns in microbial diversity is the fact that fungi undergo distinct successions during litter decomposition (Suberkropp 1984; Gessner et al. 1993).

The fungal communities in our globally distributed study streams were invariably dominated by Ascomycota as has been observed in soils (Schneider et al. 2012). In particular, fast-growing moulds like *Penicillium* and *Mucor*, yeasts, Chytrids were the most abundant fungi in streams with water temperature ≥ 8.6 °C (Fig. 3). The one exception is the stream in Japan, located at a higher altitude (1076 m) than the other locations (75 m - 300 m) within the same water temperature range (Table 1). Altitude is also known to play a significant role in structuring fungal communities in streams (Chauvet 1991), sediments (Wu et al. 2013) and soils (Siles et al. 2017).

Conclusions

Our results point to an overwhelming influence of temperature on the global distribution of litter-associated fungi in streams. This suggests that global warming might induce shifts in microbial communities colonizing decomposing litter in streams, particularly in communities with species adapted to cool environments (Christiansen et al. 2017) or those that currently experience minimal temperature fluctuations, such as in streams near the equator (Perez et al. 2016). A corollary of such community shifts that may involve the loss of key species is that expression patterns of enzymes essential in litter decomposition may lead to cascading adverse effects on food webs, alter biogeochemical cycles (Christiansen et al. 2017) and compromise ecosystem services and human well-being (Chapin et al. 2000; Sandifer et al. 2015; Wall et al. 2015).

Figure Legends

Figure 1: Study streams distribution across the globe. Norway (NOR), Germany (DEU), Canada (CAN), Italy (ITA), France (FRA), Spain (ESP), Portugal (PRT), United States of America (USA), Azores (AZO), Japan (JPN), Hong Kong (HKN), India (IND), Guinea (GIN)

Malaysia (MYS), Ecuador (ECU), Brazil (BRA), Australia (AUS), Argentina (ARG), New Zealand (NZL).

Figure 2: Linear regression model (n=19) of the relationships between absolute latitude and the natural logarithm of the mean annual rainfall (a), mean stream water temperature (b) and mean annual air temperature (c).

Figure 3: Relative abundance of fungal operational taxonomic units. For locations of stream sites see Fig. 1.

Figure 4: Polynomial regression model between absolute latitude and fungal OTU richness.

Figure 5: Canonical analysis of principal coordinates (CAP) ordinations of the relative abundances of fungal OTUs. Locations are colour coded according to mean water temperature. Mean water temperature bands were ≤ 8.1 , 8.6-19.8 and ≥ 21.3 °C. For locations of stream sites see Fig. 1.

Figure 6: Venn diagram showing the unique and shared OTUs between the stream water temperature groups. Mean water temperature bands were ≤ 8.1 , 8.6-19.8 and ≥ 21.3 °C.

Supplementary Figures

Figure S1: Rarefaction curves of fungal samples in all the streams across the globe.

Figure S2: Percentage of endemic species in the 19 streams across the globe. For locations of stream sites see Fig. 1

Acknowledgements

We are grateful to the Ecuadorian Ministry of Environment (Permits: #56-IC-FAU/FLO-DPN/MA, MAE-DNB-CM-2015-0017), and the village of Oyacachi for access to sites. Thank you also to A. Eguiguren and J. Martinez (Ecuador), S. Lamothe (France), M. Degebrodt (Germany), L.C.Y. Ng (Hong Kong) and C. Wahl (USA) for assistance in the field. Finally, we thank I. Abrantes for providing laboratory facilities.

Funding

This research was supported by the European Regional Development Fund-Operational Competitiveness Programme (FEDER-POFC- COMPETE; FCOMP-01-0124-FEDER-027793) and by national funds of the Portuguese Foundation for Science and Technology (FCT) through the project ‘DiverseAquaFun-Molecular profiling of taxonomic, functional and genetic diversity of aquatic fungi along a pollution gradient’ (PTDC/AAG-GLO/3896/2012) granted to S. Seena, PEst-OE/BIA/ UI4050/2014 granted to CBMA, and UID/MAR/04292/2013 granted to MARE. FCT also supported S. Seena and O. Sobral (PTDC/AAG-GLO/3896/2012), S. Seena (SFRH/BPD/103865/2014) and V. Ferreira (IF/00129/2014). The fieldwork performed in the Azores was funded by FCT (EXPL/AAG-GLO/0189/2013). The fieldwork conducted in Ecuador was supported by the projects DEB-1046408, DEB-1045960, and DEB-1045991. The funding bodies did not influence the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest, financial or others.

References

- Andam CP, Doroghazi JR, Campbell AN, Kelly PJ, Choudoir MJ, Buckley DH (2016) A Latitudinal Diversity Gradient in Terrestrial Bacteria of the Genus *Streptomyces*. *Mbio* 7:e02200-15(2). doi:ARTN e02200-1510.1128/mBio.02200-15
- Anderson M, Gorley RN, Clarke KR (2008) PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods. PRIMER-E. doi:citeulike-article-id:7955507
- Anderson MJ, Willis TJ (2003) Canonical analysis of principal coordinates: A useful method of constrained ordination for ecology. *Ecology* 84 (2):511-525. doi:Doi 10.1890/0012-9658(2003)084[0511:Caopca]2.0.Co;2
- APHA (1995) Standard Methods for the Examination of Water and Waste Water. Byrd Prepress Springfield, Washington, USA
- Azovsky A, Mazei Y (2013) Do microbes have macroecology? Large-scale patterns in the diversity and distribution of marine benthic ciliates. *Global Ecol Biogeogr* 22 (2):163-172. doi:10.1111/j.1466-8238.2012.00776.x
- Baldy V, Gessner MO, Chauvet E (1995) Bacteria, Fungi and the Breakdown of Leaf Litter in a Large River. *Oikos* 74 (1):93-102. doi:10.2307/3545678

- Bärlocher F, Boddy L (2016) Aquatic fungal ecology - how does it differ from terrestrial? *Fung Ecol* 19:5-13. doi.org/10.1016/j.funeco.2015.09.001
- Bärlocher F, Kendrick B (1974) Dynamics of the Fungal Population on Leaves in a Stream. *J Ecol* 62 (3):761-791. doi:10.2307/2258954
- Bass-Becking LGM (1934) *Giobiologie of Inleiding Tot de Milieukunde* W.P. Van Stockum & Zoon, The Hague, The Netherlands
- Battin TJ, Luysaert S, Kaplan LA, Aufdenkampe AK, Richter A, Tranvik LJ (2009) The boundless carbon cycle. *Nat Geosci* 2 (9):598-600. doi:10.1038/ngeo618
- Boyero L, Pearson RG, Gessner MO, Barmuta LA, Ferreira V, Graca MA, Dudgeon D, Boulton AJ, Callisto M, Chauvet E, Helson JE, Bruder A, Albarino RJ, Yule CM, Arunachalam M, Davies JN, Figueroa R, Flecker AS, Ramirez A, Death RG, Iwata T, Mathooko JM, Mathuriau C, Goncalves JF, Jr., Moretti MS, Jingtut T, Lamothe S, M'Erimba C, Ratnarajah L, Schindler MH, Castela J, Buria LM, Cornejo A, Villanueva VD, West DC (2011) A global experiment suggests climate warming will not accelerate litter decomposition in streams but might reduce carbon sequestration. *Ecol Lett* 14 (3):289-294. doi:10.1111/j.1461-0248.2010.01578.x
- Boyero L, Pearson RG, Gessner MO, Dudgeon D, Ramirez A, Yule CM, Callisto M, Pringle CM, Encalada AC, Arunachalam M, Mathooko J, Helson JE, Rincón J, Bruder A, Cornejo A, Flecker AS, Mathuriau C, M'Erimba C, Goncalves Jr JF, Moretti M, Jingtut T (2015) Leaf-litter breakdown in tropical streams: is variability the norm? *Freshw Sci* 34:759-769
- Bray JR, Curtis JT (1957) An ordination of the upland forest communities of southern Wisconsin. *Ecological monographs* 27 (4):325-349
- Briggs JC (1995) *Global Biogeography. Developments in Palaeontology and Stratigraphy*. Elsevier, Amsterdam, The Netherlands
- Bruder A, Schindler MH, Moretti MS, Gessner MO (2014) Litter decomposition in a temperate and a tropical stream: the effects of species mixing, litter quality and shredders. *Freshwater Biol* 59 (3):438-449. doi:10.1111/fwb.12276
- Chapin FS, 3rd, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, Hooper DU, Lavorel S, Sala OE, Hobbie SE, Mack MC, Diaz S (2000) Consequences of changing biodiversity. *Nature* 405 (6783):234-242. doi:10.1038/35012241
- Chauvet E (1991) Aquatic Hyphomycete Distribution in South-Western France. *J Biogeogr* 18 (6):699-706. doi:10.2307/2845551
- Chauvet E, Ferreira V, Giller PS, McKie BG, Tiegs SD, Woodward G, Elosgi A, Dobson M, Fleituch T, Graça MAS, Gulis V, Hladysz S, Lacoursière JO, Lecerf A, Pozo J, Preda E, Riipinen M, Rîşnoveanu G, Vadineanu A, Vought LBM, Gessner MO (2016) Chapter Three - Litter Decomposition as an Indicator of Stream Ecosystem Functioning at Local-to-Continental Scales: Insights from the European RivFunction Project. In: Alex J. Dumbrell RLK, Guy W (eds) *Advances in Ecological Research*, vol Volume 55. Academic Press,, Oxford, UK, pp 99-182. doi:http://dx.doi.org/10.1016/bs.aecr.2016.08.006
- Chauvet E, Suberkropp K (1998) Temperature and sporulation of aquatic hyphomycetes. *Appl Environ Microb* 64 (4):1522-1525
- Christiansen CT, Haugwitz MS, Priemé A, Nielsen CS, Elberling B, Michelsen A, Grogan P, Blok D (2017) Enhanced summer warming reduces fungal decomposer diversity and litter mass loss more strongly in dry than in wet tundra. *Glob Change Biol* 23 (1):406-420. doi:10.1111/gcb.13362
- Clarke KR, Gorley RN (2006) *Primer V6: User Manual - Tutorial*. Plymouth Marine Laboratory, Plymouth, UK

- Cox CB, Moore PD, Ladle RJ (2016a) *Biogeography: An ecological and evolutionary approach*. Wiley-Blackwell, Oxford, UK
- Cox F, Newsham KK, Bol R, Dungait JAJ, Robinson CH (2016b) Not poles apart: Antarctic soil fungal communities show similarities to those of the distant Arctic. *Ecol Lett* 19 (5):528-536. doi:10.1111/ele.12587
- Cox MP, Peterson DA, Biggs PJ (2010) SolexaQA: At-a-glance quality assessment of Illumina second-generation sequencing data. *BMC Bioinformatics* 11 (1):485. doi:10.1186/1471-2105-11-485
- Dang CK, Schindler M, Chauvet E, Gessner MO (2009) Temperature oscillation coupled with fungal community shifts can modulate warming effects on litter decomposition. *Ecology* 90 (1):122-131
- Duarte S, Bärlocher F, Pascoal C, Cassio F (2016a) Biogeography of aquatic hyphomycetes: Current knowledge and future perspectives. *Fungal Ecol* 19:169-181. doi:10.1016/j.funeco.2015.06.002
- Duarte S, Bärlocher F, Pascoal C, Cássio F (2016b) Biogeography of aquatic hyphomycetes: Current knowledge and future perspectives. *Fungal Ecol* 19 (Supplement C):169-181. doi:https://doi.org/10.1016/j.funeco.2015.06.002
- Duarte S, Bärlocher F, Trabulo J, Cássio F, Pascoal C (2015) Stream-dwelling fungal decomposer communities along a gradient of eutrophication unraveled by 454 pyrosequencing. *Fungal Divers* 70 (1):127-148. doi:10.1007/s13225-014-0300-y
- Duarte S, Seena S, Bärlocher F, Cassio F, Pascoal C (2012) Preliminary Insights into the Phylogeography of Six Aquatic Hyphomycete Species. *PLOS ONE* 7(9): e45289 (9). doi:ARTN e4528910.1371/journal.pone.0045289
- Encalada AC, Calles J, Ferreira V, Canhoto CM, GraÇA MAS (2010) Riparian land use and the relationship between the benthos and litter decomposition in tropical montane streams. *Freshwater Biol* 55 (8):1719-1733. doi:10.1111/j.1365-2427.2010.02406.x
- Fernandes I, Seena S, Pascoal C, Cássio F (2014) Elevated temperature may intensify the positive effects of nutrients on microbial decomposition in streams. *Freshwater Biol* 59 (11):2390-2399. doi:10.1111/fwb.12445
- Ferreira V, Chauvet E (2011) Synergistic effects of water temperature and dissolved nutrients on litter decomposition and associated fungi. *Glob Change Biol* 17 (1):551-564. doi:10.1111/j.1365-2486.2010.02185.x
- Fierer N, McCain CM, Meir P, Zimmermann M, Rapp JM, Silman MR, Knight R (2011) Microbes do not follow the elevational diversity patterns of plants and animals. *Ecology* 92 (4):797-804
- Foissner W (1999) Protist diversity: Estimates of the near-imponderable. *Protist* 150 (4):363-368
- Frossard A, Gerull L, Mutz M, Gessner MO (2013) Litter Supply as a Driver of Microbial Activity and Community Structure on Decomposing Leaves: a Test in Experimental Streams. *Appl Environ Microb* 79 (16):4965-4973. doi:10.1128/Aem.00747-13
- Fuhrman JA, Steele JA, Hewson I, Schwalbach MS, Brown MV, Green JL, Brown JH (2008) A latitudinal diversity gradient in planktonic marine bacteria. *P Natl Acad Sci USA* 105 (22):7774-7778. doi:10.1073/pnas.0803070105
- Gessner MO, Thomas M, Jean-Louis A-M, Chauvet E (1993) Stable successional patterns of aquatic hyphomycetes on leaves decaying in a summer cool stream. *Mycol Res* 97 (2):163-172. doi:http://dx.doi.org/10.1016/S0953-7562(09)80238-4
- Hawkins BA, Diniz-Filho JAF (2004) 'Latitude' and geographic patterns in species richness. *Ecography* 27 (2):268-272. doi:10.1111/j.0906-7590.2004.03883.x

- Hawkins BA, Porter EE, Diniz-Filho JAF (2003) Productivity and history as predictors of the latitudinal diversity gradient of terrestrial birds. *Ecology* 84 (6):1608-1623. doi:10.1890/0012-9658(2003)084[1608:PAHAPO]2.0.CO;2
- Hillebrand H (2004) On the generality of the latitudinal diversity gradient. *Am Nat* 163 (2):192-211. doi:10.1086/381004
- Hladyz S, Gessner MO, Giller PS, Pozo J, Woodward GUY (2009) Resource quality and stoichiometric constraints on stream ecosystem functioning. *Freshwater Biol* 54 (5):957-970. doi:10.1111/j.1365-2427.2008.02138.x
- Holden J (2012) *An Introduction to Physical Geography and the Environment*, vol 3rd Edn Pearson education limited, Essex, England
- Howard PJA, Frankland JC (1974) Effects of certain full and partial sterilization treatments on leaf litter. *Soil Biol Biochem* 6 (2):117-123. doi:http://dx.doi.org/10.1016/0038-0717(74)90070-4
- Jabiol J, Bruder A, Gessner MO, Makkonen M, McKie BG, Peeters ETHM, Vos VCA, Chauvet E (2013a) Diversity patterns of leaf-associated aquatic hyphomycetes along a broad latitudinal gradient. *Fungal Ecol* 6 (5):439-448. doi:https://doi.org/10.1016/j.funeco.2013.04.002
- Jabiol J, McKie BG, Bruder A, Bernadet C, Gessner MO, Chauvet E (2013b) Trophic complexity enhances ecosystem functioning in an aquatic detritus-based model system. *J Anim Ecol* 82 (5):1042-1051. doi:10.1111/1365-2656.12079
- Jetz W, Sekercioglu CH, Bohning-Gaese K (2008) The worldwide variation in avian clutch size across species and space. *PLOS Biol* 6 (12):2650-2657. doi:10.1371/journal.pbio.0060303
- Koljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Duenas M, Grebenc T, Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lucking R, Martin MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Poldmaa K, Saag L, Saar I, Schussler A, Scott JA, Senes C, Smith ME, Suija A, Taylor DL, Telleria MT, Weiss M, Larsson KH (2013) Towards a unified paradigm for sequence-based identification of fungi. *Mol Ecol* 22 (21):5271-5277. doi:10.1111/mec.12481
- Krauss GJ, Solé M, Krauss G, Schlosser D, Wesenberg D, Bärlocher F (2011) Fungi in freshwaters: ecology, physiology and biochemical potential. *FEMS Microbiol Rev* 35 (4):620-651
- Li W, Wang M, Bian X, Guo J, Cai L (2016) A High-Level Fungal Diversity in the Intertidal Sediment of Chinese Seas Presents the Spatial Variation of Community Composition. *Frontiers in Microbiology* 7:2098
- Magoč T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27 (21):2957-2963. doi:10.1093/bioinformatics/btr507
- Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M (2014) Swarm: robust and fast clustering method for amplicon-based studies. *PeerJ* 2:e593. doi:10.7717/peerj.593
- Martinez A, Larranaga A, Perez J, Descals E, Pozo J (2014) Temperature affects leaf litter decomposition in low-order forest streams: field and microcosm approaches. *FEMS Microbiol Ecol* 87 (1):257-267. doi:10.1111/1574-6941.12221
- Mittelbach GG, Schemske DW, Cornell HV, Allen AP, Brown JM, Bush MB, Harrison SP, Hurlbert AH, Knowlton N, Lessios HA, McCain CM, McCune AR, McDade LA, McPeck MA, Near TJ, Price TD, Ricklefs RE, Roy K, Sax DF, Schluter D, Sobel JM, Turelli M (2007) Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecol Lett* 10 (4):315-331. doi:10.1111/j.1461-0248.2007.01020.x

- Nikolcheva LG, Bourque T, Barlocher F (2005) Fungal diversity during initial stages of leaf decomposition in a stream. *Mycol Res* 109:246-253. doi:10.1017/S0953756204001698
- O'Malley MA (2007) The nineteenth century roots of 'everything is everywhere'. *Nat Rev Microbiol* 5 (8):647-651. doi:10.1038/nrmicro1711
- Oksanen J, Blanchet FG, Kindt R, Legendre P, O'hara RB, Simpson GL, Solymos P, Stevens H, Wagner H (2011) *Vegan: Community Ecology Package*. R package version 1.17-8.
- Oliveros J (2015) VENNY. An interactive tool for comparing lists with Venn diagrams. *BioinfoGP, CNB-CSIC*. doi:citeulike-article-id:6994833
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, D'Amico JA, Itoua I, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Ricketts TH, Kura Y, Lamoreux JF, Wettengel WW, Hedao P, Kassem KR (2001) Terrestrial Ecoregions of the World: A New Map of Life on Earth. *BioScience* 51 (11):933-938. doi:10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2
- Parmesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421 (6918):37-42. doi:10.1038/nature01286
- Peay KG, Kennedy PG, Tablot JM (2016) Dimensions of biodiversity in the Earth mycobiome. *Nature Reviews/Microbiology* 14:434-447. doi:10.1038/nrmicro.2016.59
- Perez TM, Stroud JT, Feeley KJ (2016) Thermal trouble in the tropics. *Science* 351 (6280):1392
- Perkins DM, Yvon-Durocher G, Demars BOL, Reiss J, Pichler DE, Friberg N, Trimmer M, Woodward G (2012) Consistent temperature dependence of respiration across ecosystems contrasting in thermal history. *Glob Change Biol* 18 (4): 1300-1311. doi:10.1111/j.1365-2486.2011.02597.x
- Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584. doi:10.7717/peerj.2584
- Royer TV, Minshall GW (2001) Effects of nutrient enrichment and leaf quality on the breakdown of leaves in a hardwater stream. *Freshwater Biol* 46 (5):603-610. doi:10.1046/j.1365-2427.2001.00694.x
- Sandifer PA, Sutton-Grier AE, Ward BP (2015) Exploring connections among nature, biodiversity, ecosystem services, and human health and well-being: Opportunities to enhance health and biodiversity conservation. *Ecosyst Serv* 12:1-15. doi:10.1016/j.ecoser.2014.12.007
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75 (23):7537-7541. doi:10.1128/AEM.01541-09
- Schneider T, Keiblinger KM, Schmid E, Sterflinger-Gleixner K, Ellersdorfer G, Roschitzki B, Richter A, Eberl L, Zechmeister-Boltenstern S, Riedel K (2012) Who is who in litter decomposition[quest] Metaproteomics reveals major microbial players and their biogeochemical functions. *ISME J* 6 (9):1749-1762. doi:http://www.nature.com/ismej/journal/v6/n9/supinfo/ismej201211s1.html
- Seena S, Carvalho F, Cássio F, Pascoal C (2017) Does the developmental stage and composition of riparian forest stand affect ecosystem functioning in streams? *Science of The Total Environment* 609 (Supplement C):1500-1511. doi:https://doi.org/10.1016/j.scitotenv.2017.07.252
- Shearer CA, Descals E, Kohlmeyer B, Kohlmeyer J, Marvanová L, Padgett D, Porter D, Raja HA, Schmit JP, Thorton HA, Voglymayr H (2007) Fungal biodiversity in aquatic habitats. *Biodiversity and Conservation* 16 (1):49-67. doi:10.1007/s10531-006-9120-z

- Siles JA, Cajthaml T, Filipová A, Minerbi S, Margesin R (2017) Altitudinal, seasonal and interannual shifts in microbial communities and chemical composition of soil organic matter in Alpine forest soils. *Soil Biol Biochem* 112:1-13. doi:<https://doi.org/10.1016/j.soilbio.2017.04.014>
- Strahler AN (1957) Quantitative analysis of watershed geomorphology. *Eos, Trans Amer Geophys Union* 38 (6):913-920. doi:10.1029/TR038i006p00913
- Suberkropp K (1984) Effect of temperature on seasonal occurrence of aquatic hyphomycetes. *T Brit Mycol Soc* 82 (1):53-62. doi:[http://dx.doi.org/10.1016/S0007-1536\(84\)80211-9](http://dx.doi.org/10.1016/S0007-1536(84)80211-9)
- Sul WJ, Oliver TA, Ducklow HW, Amaral-Zettler LA, Sogin ML (2013) Marine bacteria exhibit a bipolar distribution. *P Natl Acad Sci USA* 110 (6):2342-2347. doi:10.1073/pnas.1212424110
- Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI, Erlandson S, Vilgalys R, Liao HL, Smith ME, Peay KG (2014) Endemism and functional convergence across the North American soil mycobiome. *P Natl Acad Sci USA* 111 (17):6341-6346. doi:10.1073/pnas.1402584111
- Taylor JW, Turner E, Townsend JP, Dettman JR, Jacobson D (2006) Eukaryotic microbes, species recognition and the geographic limits of species: examples from the kingdom Fungi. *Philos T R Soc B* 361 (1475):1947-1963. doi:10.1098/rstb.2006.1923
- Team RDC (2011) A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <http://www.r-project.org/>
- Tedersoo L, Bahram M, Polme S, Koljalg U, Yorou NS, Wijesundera R, Villarreal Ruiz L, Vasco-Palacios AM, Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Poldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Partel K, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH, Morgado LN, Mayor J, May TW, Majuakim L, Lodge DJ, Lee SS, Larsson KH, Kohout P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo LD, Greslebin A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, De Kesel A, Dang T, Chen X, Buegger F, Brearley FQ, Bonito G, Anslan S, Abell S, Abarenkov K (2014) Fungal biogeography. Global diversity and geography of soil fungi. *Science* 346 (6213):1256688. doi:10.1126/science.1256688
- Vinson MR, Hawkins CP (2003) Broad-scale geographical patterns in local stream insect genera richness. *Ecography* 26 (6):751-767. doi:10.1111/j.0906-7590.2003.03397.x
- Violle C, Reich PB, Pacala SW, Enquist BJ, Kattge J (2014) The emergence and promise of functional biogeography. *P Natl Acad Sci USA* 111 (38):13690-13696. doi:10.1073/pnas.1415442111
- Wall DH, Nielsen UN, Six J (2015) Soil biodiversity and human health. *Nature* 528 (7580):69-76. doi:10.1038/nature15744
- Weyers HS, Suberkropp K (1996) Fungal and Bacterial Production during the Breakdown of Yellow Poplar Leaves in 2 Streams. *J N Am Benthol Soc* 15 (4):408-420. doi:10.2307/1467795
- White T, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Shinsky JJ, White TJ (eds) *PCR Protocols: A Guide to Methods and Applications*. Academic Press, San Diego, California, USA, pp 315-322. doi:citeulike-article-id:671166
- Wood-Eggenschwiler S, Bärlocher F (1983) Aquatic hyphomycetes in sixteen streams in France, Germany and Switzerland. *Transactions of the British Mycological Society* 81 (2):371-379. doi:[https://doi.org/10.1016/S0007-1536\(83\)80089-8](https://doi.org/10.1016/S0007-1536(83)80089-8)
- Wu B, Tian J, Bai C, Xiang M, Sun J, Liu X (2013) The biogeography of fungal communities in wetland sediments along the Changjiang River and other sites in China. *ISME J* 7 (7):1299-1309. doi:10.1038/ismej.2013.29

- Zhou J, Deng Y, Shen L, Wen C, Yan Q, Ning D, Qin Y, Xue K, Wu L, He Z, Voordeckers JW, Nostrand JDV, Buzzard V, Michaletz ST, Enquist BJ, Weiser MD, Kaspari M, Waide R, Yang Y, Brown JH (2016) Temperature mediates continental-scale diversity of microbes in forest soils. *Nat Commun* 7:12083. doi:10.1038/ncomms12083. <http://www.nature.com/articles/ncomms12083> - supplementary-information
- Zuur A, Ieno EN, Smith GM (2007) *Analyzing ecological data. Statistics for biology and health.* Springer Science & Business Media, New York, USA