

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Structural characterization of the third scavenger receptor cysteine-rich domain of murine neurotrypsin**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1742221> since 2020-06-24T10:37:03Z

*Published version:*

DOI:10.1002/pro.3587

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1                                   **Structural characterization of the**  
2                                   **third scavenger receptor cysteine-rich domain**  
3                                   **of murine Neurotrypsin**

4  
5                                   Anselmo Canciani<sup>1</sup>, Gianluca Catucci<sup>2</sup>, Federico Forneris<sup>1,\*</sup>

6  
7                                   <sup>1</sup>*The Armenise-Harvard Laboratory of Structural Biology, Department of Biology and*  
8                                   *Biotechnology, University of Pavia, Via Ferrata 9/A, 27100 Pavia (Italy).*

9                                   *<http://fornerislab.unipv.it>; <sup>2</sup>Department of Life Sciences and Systems Biology,*  
10                                   *University of Torino, 10123 Turin, Italy*

11  
12 \* Correspondence to: Federico Forneris, Tel. +39 0382 985228, Fax +39 0382 528496, e-  
13 mail: [federico.forneris@unipv.it](mailto:federico.forneris@unipv.it)

14  
15 ***Running title:*** *Crystal structure of Neurotrypsin SRCR3*

16  
17 **Total Number of Manuscript Pages:** 26

18 **Total Number of Supplementary Material Pages:** 5

19 **Total Number of Tables:** 3

20 **Total Number of Figures:** 4

21  
22 **Description of Supplementary Materials:**

23 Supplementary materials consist of a single PDF file including 4 supplementary figures. All  
24 supplementary materials are cited within the manuscript.

25 **Abstract**

26 Neurotrypsin (NT) is a multi-domain serine-protease of the nervous system with only one  
27 known substrate: the large proteoglycan Agrin. NT has been seen to be involved in the  
28 maintenance/turnover of neuromuscular junctions and in processes of synaptic plasticity in  
29 the central nervous system. Roles which have been tied to its enzymatic activity, localized in  
30 the C-terminal serine-protease (SP) domain. However the purpose of NT's remaining 3-4  
31 scavenger receptor cysteine-rich (SRCR) domains is still unclear. We have determined the  
32 crystal structure of the third SRCR domain of murine NT (mmNT-SRCR3), immediately  
33 preceding the SP domain and performed a comparative structural analysis using homologous  
34 SRCR structures. Our data and the elevated degree of structural conservation with  
35 homologous domains highlight possible functional roles for NT SRCRs. Computational and  
36 experimental analyses suggest the identification of a putative binding region for Ca<sup>2+</sup> ions,  
37 known to regulate NT enzymatic activity. Furthermore, sequence and structure comparisons  
38 allow to single out regions of interest that, in future studies, might be implicated in Agrin  
39 recognition/binding or in interactions with as of yet undiscovered NT partners.

40

41 **Keywords**

42 Scavenger receptor cysteine-rich domain; SRCR; Protease; Neurotrypsin; Neuromuscular  
43 Junctions.

44

45

46 **Introduction**

47 Neurotrypsin (NT), also known as PRSS12 and motopsin, is a multi-domain extracellular  
48 serine-protease of the nervous system first described in the late 90's (1; 2). Spanning 761  
49 amino acids, the murine variant encompasses an N-terminal proline rich segment, a kringle  
50 domain (Kr), three scavenger receptor cysteine rich (SRCR) domains and a C-terminal  
51 serine-protease domain (SP) (1; 2). In comparison to the human homolog (875 a.a.), sharing  
52 an overall sequence identity of 87%, the murine ortholog is shorter in virtue of an additional  
53 N-terminal SRCR domain (3) (Figure 1 A, Supplementary Fig. 1).

54 Produced predominantly by neurons of the central and peripheral nervous systems, NT is  
55 stocked in synaptic vesicles and released in an activity-dependant manner (4; 5). Upon  
56 secretion, NT cleaves the large proteoglycan Agrin, its only known substrate, generating two  
57 C-terminal fragments. This event is likely responsible for the roles played by NT in  
58 neuromuscular junction maintenance/turnover and in neuronal plasticity (4-7). Such is the  
59 importance of this cleavage event that de-regulation and/or inactivation result, respectively,  
60 in Sarcopenia (muscle wasting disease of the elderly) (7) and non-syndromic mental  
61 retardation (4).

62 While the significance of the catalytic domain has been touched upon, the role and relevance  
63 of the accessory domains remain unclear. Nonetheless it would be reasonable to attribute  
64 them with a role in protein-protein interaction and/or substrate recognition. Indeed, literature  
65 reports two potential NT interactors, in addition to the substrate Agrin, identified through  
66 yeast two-hybrid screening experiments: the product of seizure related gene 6 (Sez-6) (8) and  
67 the Integral Membrane Protein 2A (Itm2a) (8). These have been seen to influence,  
68 respectively, neuronal development and function (9; 10), and the differentiation of different  
69 tissue/cell types (11-13), among which skeletal muscle (14). NT's interaction with Sez-6 has

70 been tied to one of the accessory domains: the Kr domain (8). Conversely, the domain(s)  
71 responsible for interacting with Itm2a has yet to be identified.

72 The scavenger receptor cysteine-rich (SRCR) domain is a small (90-110 residues)  
73 distinguishing element of a broad superfamily of proteins whose members span different  
74 functional roles (15-17). It displays a highly conserved structure with five anti-parallel beta  
75 strands cradling an alpha helix and a distinct disulfide bridge architecture (18; 20-23). A  
76 feature which has been used to identify two types of SRCR domains, A and B, bearing  
77 respectively 6 and 8 cysteines. While the number of cysteines differs, the relative pairing is  
78 surprisingly consistent and tends to form as follows: C1-C4, C2-C7, C3-C8 and C5-C6. The  
79 first pair distinguishes types A and B as it is present exclusively in the latter (16). Members  
80 of the SRCR superfamily can share little functional similarity playing roles in immune  
81 response (18), cell differentiation (24), apoptosis (25) and tumor suppression alike. While it  
82 has been difficult to assign a univocal function to SRCR domains, a commonality can be  
83 drawn across several SRCR superfamily members, broadly attributing the SRCR domain with  
84 a role in protein-protein/protein-substrate interaction (26). It is therefore likely NT SRCR  
85 domains may mediate interactions with the surrounding environment. They could be  
86 responsible for the reported glycosaminoglycan and heparin binding capabilities (27),  
87 contribute to its specificity, mediate its localized lingering at the synapse (19), contribute to  
88 its specificity, mediate its localized lingering at the synapse (20) and/or be involved in  
89 binding to, as of yet, unidentified partners. Of NT's accessory domains only the structure of  
90 the Kr domain has been successfully determined and assigned a possible role (21; 8). The  
91 lack of structural information pertaining to the remaining domains limits our functional  
92 understanding of NT. As such new structural data on the SP or SRCR domains might provide  
93 key insights on the mechanisms underlying NT's biological role.

94 Here we report the recombinant production, purification, crystallization and crystal structure  
95 determination of the third SRCR domain of murine NT (mmNT-SRCR3), and analysis of its  
96 molecular architecture based on comparisons with homologous SRCR domains. Our  
97 structural data constitute a solid additional step towards the understanding of NT's molecular  
98 interactions and biological functions.

99

## 100 **Results**

### 101 *Purification of mmNT-SRCR3*

102 Owing to the high cysteine content of extracellular SRCR domains, we produced the mmNT-  
103 SRCR3 in soluble form using a *E. coli* strain particularly adapted to facilitate disulfide bond  
104 formation. Purification of mmNT-SRCR3 was performed using a two-step immobilized Ni-  
105 affinity chromatography (Ni-IMAC) followed by size-exclusion chromatography (SEC)  
106 (Supplementary Fig. 2). The purified material was used for crystallization experiments and  
107 protein characterization studies in solution.

108

### 109 *Crystal structure of mmNT-SRCR3*

110 A single mmNT-SRCR3 crystal allowed complete X-ray data collection and structure  
111 determination (Figure 1 B); data processing and structure refinement statistics are reported in  
112 table 1. This process highlighted the presence of two mmNT-SRCR3 monomers in the  
113 asymmetric unit assembling into what appeared to be a crystallographic dimer (Figure 1 C).  
114 Each monomer presents a very compact fold with a central alpha helix ( $\alpha 1$ ) nested in five  
115 beta strands ( $\beta 1$ - $\beta 5$ ) forming a twisted anti-parallel cradle (Figure 1 B). The B-factor  
116 distribution across the structure highlights the high stability of the central core and  
117 progressive flexibility of the peripheral loops interconnecting the secondary structure  
118 elements (Figure 1 C). Of particular interest is the Val466–Asn474 segment, whose flexibility

119 is such that the electron density map of that region is very poor despite the 1.7-Å resolution  
120 (Supplementary Fig. 3).

121 Each mmNT-SRRCR3 model spans 107 residues and lacks the final C-terminal “KKASS”  
122 sequence, owing to the high mobility of this five-residue tail. Superposition of the two  
123 monomers found in the asymmetric unit and analysis of the root mean square deviation  
124 (r.m.s.d) shows that the only visible significant difference is located in the aforementioned  
125 Val466–Asn474 loop, further supporting its flexible nature (Figure 1 D). Calculations of  
126 surface electrostatic potentials using CCP4mg (22) displayed a disc-like shape with mixed  
127 charge distribution along its sides and two opposite faces, one almost fully hydrophobic and  
128 the other displaying a patch of negative charges (Figure 1 E).

129

### 130 *SEC-SAXS analysis*

131 In order to assess the oligomeric state of the purified protein sample in solution, and thus  
132 ascertain the nature of the mmNT-SRRCR3 dimer observed in the crystal packing (Figure 1 C),  
133 we performed a SEC-small angle x-ray scattering (SEC-SAXS) experiment. The initial  
134 chromatogram displayed a well-defined peak with good scattering intensities and a very  
135 homogeneous mass distribution (Figure 2 A). mmNT-SRRCR3 presents as a monodisperse  
136 species with an MW of 12 kDa calculated from averaged peak intensities (Table 2). This is in  
137 line with the 12.5 kDa calculated for a monomer of this construct and incompatible with  
138 higher order oligomeric species. *CRY SOL* (23) was used to compare the averaged scattering  
139 curve of the SEC-SAXS peak to a theoretically calculated scattering curve for our  
140 crystallographic structure. The single mmNT-SRRCR3 monomer displayed a very good fit ( $\chi^2$   
141 = 1.4) with the in-solution data (Figure 2 B), ruling the dimer as induced by the  
142 crystallization process.

143

144 *Comparison with other SRCR domains*

145 To better understand the functional role of NT's SRCR domains, we used the *DALI* server  
146 (24) to compare this structure with other entries in the Protein Data Bank (PDB). This search  
147 returned several high scoring matches which were manually filtered to remove redundancies  
148 resulting in a list of only 8 hits (Table 3). While limited in sequence identity, structural  
149 superpositions with mmNT-SRCR3 (Figure 3 A) evidenced a high degree of structural  
150 conservation, strongest within the central core and with greater variability in the peripheral  
151 loops. Comparison with an expanded pool of homologs using *ConSurf* (25) highlighted a  
152 similar trend for sequence conservation. When mapped to the structure of mmNT-SRCR3, it  
153 was possible to see how the areas of greater sequence identity, for the most part,  
154 corresponded to highly conserved secondary structure elements forming the domain core  
155 (Figure 3 B). Conversely the more flexible external loops displayed a significantly greater  
156 compositional variability. Of particular note is the  $\beta$ 4 strand that, while being one of five core  
157  $\beta$  strands displays little to no amino acid conservation (Figure 3 C).

158 *DALI* matches with > 30% identity evidenced several perfectly conserved residues which  
159 were mirrored in ConSurf alignments. Among these is a six-cysteine network (residues 411,  
160 424, 455, 465, 475, and 485) responsible for the archetypical SRCR domain disulfide bridges  
161 (Figure 3 B). This canonical pairing is expressed relative to type B SRCR domains and while  
162 type A domains lack the C1-C4 pair, the relative numbering of the other pairs remains  
163 conventionally and structurally unaltered. Therefore, in mmNT-SRCR3 we found that  
164 Cys411-Cys475 corresponds to the C2-C7 pair, while Cys424-Cys485 and Cys455-Cys465  
165 correspond to the C3-C8 and C5-C6 pairs, respectively. Residues Gly391, Gly397, Glu400,  
166 Ala420, Val422, Leu427, Gly457, Glu459 and Val483 are also fully conserved (Figure 3 C).  
167 Of these, Ala420, Val422 and Leu427 can be mapped to the central helix while Gly397,  
168 Glu400 and Val483 can be found on strands  $\beta$ 2 and  $\beta$ 5, respectively (Figure 3 B).



169 Conversely, the remaining Gly391, Gly457 and Glu459 locate on the more flexible, and  
170 generally less conserved, external loops (Figure 3 B).

171 Comparison of mmNT-SRCR3 with other mmNT SRCR domains evidenced a certain extent  
172 of conservation (44-58% sequence identity) comparable to that of the highest *DALI* matches  
173 (Supplementary Fig. 1 B). Extending this analysis to include human NT evidenced how its  
174 SRCR domains 2-4 correspond, respectively, to mmNT SRCR domains 1-3, while the first  
175 human SRCR domain is not conserved across the two homologs (Supplementary Fig. 1B).

176

### 177 *Evidence for Ca<sup>2+</sup> binding*

178 Given the significant contributions of Ca<sup>2+</sup> to NT activity (5), and the reported Ca<sup>2+</sup>-based  
179 modulation of SRCR-mediated interactions (18; 26), we investigated whether mmNT-SRCR3  
180 might contribute in similar fashion. Therefore, an Isothermal Titration Calorimetry (ITC)  
181 experiment was conducted in presence of CaCl<sub>2</sub>. This evidenced a Ca<sup>2+</sup> binding to mmNT-  
182 SRCR3 (Figure 4 A), with  $K_d = 10.1$  mM,  $\Delta H = -7.7$  Kcal/mol,  $\Delta S = -16.8$  cal/mol/deg.

183 A comparison of mmNT-SRCR3 with the structure of the monomeric MARCO SRCR  
184 domain (18) highlighted a cluster of highly conserved residues coordinating a bivalent metal-  
185 ion in MARCO (Figure 4 B). These correspond to amino acids Asp412-Asp413, His473,  
186 Asn474 and Glu479, that significantly contribute mmNT-SRCR3's negatively charged patch.  
187 Interestingly the same residues were seen to be perfectly conserved in the DMBT1 SRCR-1  
188 domain, known as well to display Ca<sup>2+</sup> binding (27) (Figure 4 C).

189

## 190 **Discussion**

191 The scavenger receptor cysteine-rich domain groups numerous proteins with different  
192 functions in a large superfamily (15). However, the understanding of the specific function of  
193 SRCR domains is, to date, limited to a general consensus of protein-protein/protein-ligand

194 interaction. The solution of the structure of the third SRCR domain of murine Neurotrypsin  
195 (mmNT-SRCR3) and its analysis highlighted several interesting features that might provide a  
196 functional insight both in regards to NT and SRCR domains in general. Most striking of all  
197 was the degree of similarity with the available homologous structures despite a generally  
198 poor sequence conservation (Table 3).

199 At the core of mmNT-SRCR3 is an extremely stable secondary structure element  
200 organization which seems to be perfectly conserved across available structures. Such  
201 variability could be traced to the more flexible external loops, most notably the Val466-  
202 Asn474 loop. This region not only displayed the highest mobility across homologs but also  
203 within the mmNT-SRCR3 monomers observed in the crystallographic dimer. Mapping of  
204 sequence conservation to the structure of mmNT-SRCR3 showed that the least conserved  
205 residues were located on this and other peripheral flexible loops, while the core secondary  
206 structure elements bore the majority of highly conserved amino acids. Of particular note are:  
207 a six-cysteine network responsible for the typical disulfide bond architecture of SRCR  
208 domains, and several perfectly conserved residues (Gly397, Glu400, Ala420, Val422,  
209 Leu427, and Val483) mapped to the highly stable core. These, owing to their localization and  
210 conservation, are likely to provide essential contributions to the fold of SRCR domains.  
211 Conversely, three additional highly conserved residues (Gly391, Gly457 and Glu459),  
212 mapped to the more flexible loop regions, and the  $\beta$ 4 secondary structure element, which  
213 displays structural but not compositional conservation, might contribute to SRCR domain  
214 function.

215 Among the more conserved regions two stretches, spanning Gly397–Val401 and Trp407–  
216 Asp412 respectively, should be mentioned owing to their documented evolutionary  
217 conservation (28). The first likely contributes functionally, as similar consensus sequences  
218 were seen to mediate protein-target interaction in other SRCR-SF members including

219 MARCO (29), DMBT1 (35), CD163 (30), as well as structurally, given its mapping to the  
220 highly conserved  $\beta$ 1 secondary structure element. Conversely, the second has yet to be  
221 assigned with a functional role but it is thought that the conservation of this stretch is  
222 structurally related.

223 Owing to structural conservation it is possible to suppose a protein-target role for mmNT-  
224 SRCR3. Additional observations drawn from that structure and other SRCR-SF members  
225 might hint at its more specific function in relation to NT. Notably, analysis of surface charge  
226 distribution of mmNT-SRCR3 evidenced a negatively charged patch corresponding to a  
227 cluster of residues (Asp412, Asp413, His473, Asn474 and Glu479), conserved in the metal-  
228 ion binding MARCO SRCR domain (18), that might be involved in  $\text{Ca}^{2+}$  ion binding known  
229 to be important for NT enzymatic activity (31). A plausible hypothesis, given that ITC  
230 experiments showed  $\text{Ca}^{2+}$  binding for mmNT-SRCR3 and that other SRCR-SF members  
231 (MARCO, DMBT1 and CD163) also display a  $\text{Ca}^{2+}$  dependant modulation mapped to their  
232 SRCR domains (32; 30; 18). Further analogies can be speculated regarding the nature of  
233 possible binding partners. As several SRCR-SF members are known to bind to bacteria (32;  
234 27), lipopolysaccharides (LPS) directly (29; 33) or glycoproteins (34) via their SRCR domains.  
235 Thus it is possible that NT SRCR domains might mediate similar interactions between NT  
236 and its heavily glycosylated substrate Agrin (19).

237 Finally, cross-comparison of human and murine NT evidenced how the first SRCR domain in  
238 the human NT sequence represents the principal source of divergence between the two  
239 homologs. Surprisingly, also the residues putatively coordinating  $\text{Ca}^{2+}$  in mmNT-SRCR3 are  
240 well conserved across all NT SRCR domains, except for human NT-SRCR1 (Supplementary  
241 Fig. 4). An observation which hints at a possible lack of  $\text{Ca}^{2+}$  binding for that SRCR domain  
242 and that, in conjunction with its absence in murine NT, opens to speculation regarding the  
243 function and relevance of the human NT SRCR1.

244 In conclusion, it seems plausible that the broad protein-target interaction attributed to SRCR  
245 domains might be mediated by the highly conserved nature of their core structure. While the  
246 more structurally and compositionally variable regions could be responsible for more  
247 nuanced and specific functional aspects such as ligand recognition. Finally in regards to NT,  
248 our characterization of mmNT-SRCR3 allowed us to identify a  $\text{Ca}^{2+}$  binding site and infer, by  
249 comparison, a possible role for this domain in Agrin recognition.

250

251 **Materials and methods**

252 *Molecular cloning, recombinant expression and purification*

253 The cDNA encoding for the third SRCR domain of murine Neurotrypsin (UniProt id O08762,  
254 residues 383-494) was obtained from Source Bioscience (I.M.A.G.E. clone ID 3665834),  
255 amplified with a Phusion DNA polymerase (Thermo Fisher Scientific) using oligos  
256 AAAGATCTGGTTTTCCCATCAGACTAGTGGATG (forward) and  
257 AAGCGGCCGCACTTGATGCTTTCTTCTCTAAATAG (reverse), and inserted into the  
258 pCIOX recombinant expression vector (Addgene), yielding the final construct bearing an N-  
259 terminal 8-His-SUMO tag. This plasmid was transformed into chemically competent  
260 *Escherichia coli* SHuffle T7 cells (New England Biolabs), which were used for protein  
261 production.

262 A small culture of transformed cells was grown over night shaking at 30 °C in LB +  
263 kanamycin (50 µg/ml) and used to inoculate, in a 1:100 ratio, a larger volume (1 - 6 L) of  
264 autoinducing ZYP-5052 media (35) for large-scale production. Inoculated media was  
265 incubated shaking at 30 °C for 5 h, after which the temperature was lowered to 20 °C to  
266 induce recombinant protein expression, and further incubated for 20 h.

267 Cells were harvested by centrifugation (5000 x g, 15 min, 4 °C) with a swinging bucket  
268 centrifuge (Beckman Coulter). The supernatant was discarded and the cells were resuspended  
269 in buffer A (25 mM 2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid  
270 (HEPES)/NaOH, 0.5 M NaCl, pH 8) in a 1:5 (w/v) wet cell pellet-to-buffer ratio. This  
271 suspension was placed on ice for 30 minutes and then sonicated (80% amplitude, 8 cycles  
272 with 40 s on/20 s off pulses) to lyse the cells. Cell debris was removed by centrifugation  
273 (10'000 x g, 50 min, 4 °C) and the supernatant was filtered with a 0.45 µm syringe filter  
274 (Sartorius). The clarified lysate was loaded on a 5 mL HisTrap (GE Healthcare) column, pre-  
275 conditioned with buffer A, at a rate of 1 mL/min. Unbound material was washed from the

276 column with buffer A and non-specifically bound contaminants were removed with buffer A  
277 supplemented with 25 mM imidazole. mmNT-SRCR3 was eluted by further increasing the  
278 imidazole concentration to 250 mM. The eluted fractions were pooled, supplemented with  
279 SUMO protease (1.2 mg/mL stock, 1:300 v/v) and dialyzed overnight at 4 °C against buffer  
280 A. Following dialysis the sample was subject to centrifugation (5000 x g, 4 °C, 15 min) and  
281 the supernatant was loaded onto a 1 mL HisTrap column (GE Healthcare). Successful  
282 removal of the His-SUMO tag was evaluated through sample recovery in the flow through  
283 fractions, which were further purified by gel filtration on a Superdex 75 10/300 GL column  
284 (GE Healthcare) equilibrated with 25 mM HEPES/NaOH, 100 mM NaCl, pH 8. At each step  
285 of protein purification, samples were collected and analysed using SDS-PAGE. The final  
286 yield was  $\approx$  2 mg of pure protein per gram of bacterial cells.

287

#### 288 *Crystallization, X-ray Data collection and processing*

289 For crystallization screening the protein was concentrated to 20 mg/mL with a 5 kDa MWCO  
290 Vivaspin12 concentrator (Sartorius). Crystallization was performed using the sitting-drop  
291 vapor diffusion method at both 4 and 20 °C. Drops were set up in a volumetric 1:1 protein-to-  
292 precipitant solution ratio using an Oryx8 crystallization robot (Douglas Instruments) with  
293 MRC 96-well PS plates (SwissSci). Initial screening performed with commercial screens  
294 yielded thin microcrystals, not suitable for diffraction testing, in numerous conditions.  
295 Optimization of initial crystallisation hits was performed by hand using the sitting drop  
296 method. The best crystals were obtained at 4 °C, over the span of a month, by mixing  
297 mmNT-SRCR3 concentrated at 19 mg/mL with a solution composed of 16% PEG3350, 0.1  
298 M Tri-Sodium citrate, pH 7.5. Prior to flash-cooling in liquid nitrogen, crystals were  
299 harvested with nylon cryoloops (Hampton Research) and briefly soaked into a cryo-  
300 protectant solution (19% PEG3350, 0.1 M Tri-Sodium citrate, 20% Glycerol). X-ray

301 diffraction data were collected at 100 K at the ID30A-3 beamline of the ESRF synchrotron.  
302 Data were indexed and integrated using *XDS* (36) and scaled using *Aimless* (37). Data  
303 collection statistics are summarized in Table 1.

304

#### 305 *Structure determination and refinement*

306 The structure of mmNT-SRCR3 was solved at 1.7 Å by molecular replacement using the  
307 structure of the Mac-2 binding protein (M2BP) SRCR domain (pdb: 1BY2) (38) as search  
308 model. This model (50% sequence identity to mmNT-SRCR3) was selected based on  
309 conservation of sequence identity as evaluated using *NCBI BLAST* (39). Prior to molecular  
310 replacement, the sequences of mmNT-SRCR3 and M2BP SRCR were aligned using *EBI*  
311 *MUSCLE* (40) and non-conserved residues were adjusted using *CHAINSAW* (41). The  
312 resulting model was used in molecular replacement with *PHASER* (42). Two copies of the  
313 search model were found constituting the asymmetric unit, with a  $V_m$  of 3.11 Å<sup>3</sup> and 60%  
314 solvent content. The structure was refined with successive steps of manual building in *COOT*  
315 (43) and automated refinement with *REFMAC5* (44). Model validation was performed with  
316 *MolProbity* (45). Refinement statistics for the final model are reported in Table 1 as  
317 deposited to the PDB under accession code 6H8M. Electrostatic surface calculations and  
318 representations were generated with *CCP4mg* (22). Other structural images were generated  
319 with *UCSF Chimera* (46).

320

#### 321 *SEC-SAXS analysis*

322 Size exclusion chromatography coupled to small-angle X-ray scattering (SEC-SAXS)  
323 analysis was performed at the BM29 beamline of the ESRF synchrotron in Grenoble (France)  
324 using a protocol adapted from (47). The protein was concentrated to 15 mg/mL and a 15 µL  
325 sample was run on a Superdex 75 Increase 3.2/300 column (GE Healthcare) equilibrated in

326 25 mM HEPES/NaOH, 0.1 M NaCl, pH 8 and mounted on a Nexera High Pressure  
327 Liquid/Chromatography (HPLC; Shimadzu) system. SAXS data were collected from the  
328 sample capillary mounted on line with the HPLC system, using a Pilatus 1 M detector  
329 (Dectris) positioned at distance of 2.87 m allowing a global  $q$  range of 0.03–4.5 nm with 12.5  
330 keV energy. Analysis of scattering intensities was performed using *CHROMIXS* (48) and the  
331 *ATSAS* suite (49). Comparison of in-solution scattering to crystallographic data was carried  
332 out with CRY SOL (23). Details of SEC-SAXS data collection and analysis are summarized  
333 in Table 2 as deposited in SASBDB under accession code SASDES5.

334

### 335 *Isothermal Titration Calorimetry*

336 Titrations were carried out at 25°C in 100 mM NaCl 25 mM HEPES/NaOH pH8 using the  
337 high feedback mode of a MicroCal ITC200 instrument (Malvern Instruments, Worcestershire,  
338 UK). The concentration of mmNT-SRRCR3 in the cell was 1.2 mM, whereas the CaCl<sub>2</sub>  
339 solution in the syringe was at 120 mM. The first injection was kept to the minimum volume  
340 of 0.1 uL to allow complete equilibration of the cell. The following 19 titrations of CaCl<sub>2</sub>  
341 were performed using a 2 uL injection volume with a 120 sec. time interval while  
342 maintaining stirring at 750 rpm. The heat of dilution of 120 mM CaCl<sub>2</sub> was determined by  
343 performing a second titration in which the cell was filled only with buffer while all other  
344 experimental parameters remained unaltered. The net contribution of binding of CaCl<sub>2</sub> to  
345 mmNT-SRRCR3 was calculated by subtracting the heat of dilution of CaCl<sub>2</sub>. Data were  
346 analyzed using the “One Set of Sites” curve fitting model (MicroCal ITC200 Origin).

347

### 348 **Supplementary Material**

349 Supplementary figures and tables are available and linked through the text.

350



351 **Acknowledgements**

352 We thank the European Synchrotron Radiation Facility (ESRF) for the provision of synchrotron  
353 radiation facilities and beamline scientists of the ESRF and the European Molecular Biology  
354 Laboratory for assistance. This work was supported by a Cariplo Foundation Grant (id. 2014-  
355 0881), the Giovanni Armenise-Harvard Career Development Award (2013), the “Programma  
356 Rita Levi-Montalcini” from the Italian Ministry of University and Research (MIUR), and by  
357 the Italian Ministry of Education, University and Research (MIUR): Dipartimenti di  
358 Eccellenza Program (2018–2022) - Dept. of Biology and Biotechnology "L. Spallanzani",  
359 University of Pavia.

360

361 **Conflict of interest statement**

362 The authors declare no competing financial interests.

363

364 **Author Contributions**

365 FF conceived and supervised the project. AC cloned, expressed, purified, crystallized and  
366 solved the structure of mmNT-SRCR3. AC and FF analyzed crystallographic and SEC-SAXS  
367 data. GC performed ITC measurements and data processing. AC prepared the figures. AC  
368 and FF wrote the paper. All authors read and approved the final manuscript.

369

370 **Table 1. X-ray data collection and refinement statistics**

<b>Data Collection</b>	
Diffraction source	ESRF ID30A-3
Wavelength (Å)	0.983
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>
Cell parameters <i>a</i> , <i>b</i> , <i>c</i> (Å)	<i>a</i> = 57.8 Å, <i>b</i> = 62.9 Å, <i>c</i> = 84.5 Å, $\alpha = \beta = \gamma = 90^\circ$
Resolution range (Å) <sup>1,2</sup>	42.54 - 1.70 (1.73 - 1.70)
Unique reflections	34576 (1785)
Completeness (%)	100.0 (100.0)
Multiplicity	6.9 (6.9)
Mean ( <i>I</i> /σ( <i>I</i> ))	15.5 (0.5)
CC(1/2)	0.999 (0.434)
<i>R</i> <sub>sym</sub> <sup>3</sup>	0.049 (3.37)
<b>Refinement</b>	
<i>R</i> <sub>work</sub> / <i>R</i> <sub>free</sub>	0.19 / 0.21
No. of non-H atoms	1747
Protein	1662
Ligands	0
Waters	85
r.m.s. deviations	
Bonds (Å)	0.016
Angles (°)	1.699
Average <i>B</i> factors (Å <sup>2</sup> )	
Protein	65.3
Water	54.0
Ramachandran favoured (%)	97.6
Ramachandran allowed (%)	2.4

371

372 <sup>1</sup> Values for the outer shell are given in parentheses.373 <sup>2</sup> Resolution limits were determined by applying a cut-off based on the mean intensity correlation coefficient of  
374 half-datasets (CC1/2) approximately of 0.5 (50).375 <sup>3</sup>  $R_{sym} = \Sigma |I - \langle I \rangle| / \Sigma I$ , where *I* is the observed intensity for a reflection and  $\langle I \rangle$  is the average intensity  
376 obtained from multiple observations of symmetry-related reflections.

377

378 **Table 2. Summary of SEC-SAXS data collection and analysis**

---

<b><i>Data Collection</i></b>	
Light source and beamline	ESRF BM29
Beam energy (keV)	12.5
Sample-detector distance (m)	2.867
Exposure time (s)	1
Sample cell thickness (mm)	1
Temperature (°)	20
Final $q$ range ( $\text{nm}^{-1}$ )	0.01 - 4
<b><i>Data Analysis</i></b>	
Points used for Guinier analysis	27-173
Guinier $qR_g$ limits	0.24
Guinier $R_g$ (nm)	$1.54 \pm 0.03$
$I(0)$ ( $\text{nm}^{-1}$ )	$13.48 \pm 0.02$
Dmax (nm)	5.8
MW estimation ( $V_c$ based) (kDa)	12

---

379

380

381 **Table 3** List of homologous SRCR structures identified by *DALI*.

<i>Protein</i>	<i>PDB</i>	<i>Z-Score</i>	<i>r.m.s.d.</i>	<i>% id.</i>
Lysyl oxidase homolog 2	5ZE3	18.2	2.1	49
CD6	5A2E	18.0	1.5	49
Mac-2 binding protein (M2BP)	1BY2	18.0	1.8	50
Scavenger receptor cysteine-rich type 1 protein m	5JFB	16.6	1.8	38
Macrophage receptor (MARCO)	2OYA	13.7	1.7	52
Human complement factor I	2XRC	11.2	2.2	31
T-cell surface glycoprotein CD5	2JA4	7.5	2.4	27
Serine protease Hepsin	3T2N	7.2	3.0	16

382

383 **Figure Legends:**

384 *Figure 1* - Three-dimensional structure of the third SRCR domain of murine NT. (A) domain  
385 organization of mouse and human NT, highlighting the localization of mmNT-SRCR3 within  
386 the multidomain enzyme architecture. Colouring of the domains highlights sequence  
387 conservation between homologous enzymes, as described in Supplementary Fig. 1. (B)  
388 Cartoon representation of mmNT-SRCR3 displaying its main secondary structure elements,  
389  $\alpha$ -helix ( $\alpha 1$ ) in blue and  $\beta$ -strands ( $\beta 1 - \beta 5$ ) in cyan. (C) Cartoon “putty” representation of  
390 the two mmNT-SRCR3 monomers found in the crystallographic asymmetric unit, coloured  
391 blue-red by B-factors ( $34.2 - 162.5 \text{ \AA}^2$ ). (D) Superposition of the two mmNT-SRCR3  
392 molecules found in the asymmetric unit. chain A is shown in red, chain B is shown in cyan.  
393 (E) Surface representation with colouring by electrostatic potential ( $-2.0 - +2.0 \text{ V}$ ), negative  
394 charges in red, positive charges in blue and neutral areas in white. Monomer is presented  
395 orientations highlighting main features: uncharged bottom, mixed charges on sides and  
396 negatively charged patch on top.

397

398 *Figure 2* – mmNT-SRCR3 SEC-SAXS analysis. (A) SEC-SAXS chromatogram showing  
399 mmNT-SRCR3 scattering intensities and profile of mass distribution across the peak. (B)  
400 Averaged experimental peak scattering curve, after buffer subtraction, plotted superimposed  
401 to a theoretical, *CRY SOL*-derived, scattering curve for an mmNT-SRCR3 monomer ( $\chi^2 =$   
402 1.4).

403

404 *Figure 3* - Structural comparison of mmNT-SRCR3 with other SRCR domains. (A)  
405 Superposition with non-redundant *DALI* matches (Table 3) coloured red-yellow by r.m.s.d.  
406 ( $0.39 - 6.32 \text{ \AA}$ ). (B) Cartoon representation of mmNT-SRCR3 in two orientations, coloured  
407 cyan-brown ( $0 - 100\%$ ) by residue conservation after *ConSurf* analysis. The left panel maps

408 highly conserved residues from SRCR structure sequence alignments, while the right panel  
409 maps the principal secondary structure elements and the archetypical SRCR disulfide bridge  
410 organization. The C1-C4 pair is absent in SRCR type A domains, residues Cys411-Cys475  
411 correspond to the C2-C7 pair, Cys424-Cys485 to the C3-C8 pair and Cys455-Cys465 to the  
412 C5-C6 pair. (C) Sequence alignment of SRCR structures identified by *DALI* analysis of  
413 mmNT-SRCR3. Identical residues boxed in red, residues with 70% conservation boxed in  
414 yellow. mmNT-SRCR3 secondary structure elements displayed above alignment and  
415 consensus sequence with 70% conservation under alignment. Image created using *ESPRIT3*  
416 (51).

417

418 *Figure 4* – Evidence for mmNT-SRCR3 Ca<sup>2+</sup> binding. (A) Isothermal calorimetric titration of  
419 CaCl<sub>2</sub> (120 mM) with purified mmNT-SRCR3. Heat variation generated by each injection of  
420 titrant at each time interval (top panel) and the integration of each peak and the amount of  
421 heat produced (lower panel). (B) Structural inference of mmNT-SRCR3 binding of metal  
422 ions. Top panel, residues contributing to the formation of the negatively charged patch on  
423 mmNT-SRCR3 likely bind metal ions. Bottom panel, superposition of mmNT-SRCR3 (blue)  
424 with monomeric MARCO (cyan, PDB: 2OY3) coordinating Mg<sup>2+</sup> (green sphere) via those  
425 same amino acids. Side chains of residues Asp412, Asp413 and Glu479 display almost  
426 identical conformations, while Asn474 and His473 show higher flexibility. (C) Sequence  
427 alignment of mmNT-SRCR3 against MARCO and DMBT1 SRCR domains known to bind  
428 Ca<sup>2+</sup>. Blue arrows and boxes indicate residues evidenced in (B) which are perfectly conserved  
429 across all three proteins. Perfectly conserved residues boxed in red, residues with 70%  
430 conservation are boxed in yellow. Evaluation of sequence conservation is shown under  
431 alignment; symbols indicate: ! = Ile or Val, \$ = Leu or Met, % = Phe or Tyr, # = Asn, Asp,  
432 Glu or Gln. Residue numbering refers to mmNT-SRCR3.

433

434 **References**

- 435 1. Gschwend TP, Krueger SR, Kozlov SV, Wolfer DP, Sonderegger P (1997) Neurotrypsin, a  
436 novel multidomain serine protease expressed in the nervous system. *Mol Cell Neurosci*  
437 9:207-219. PMID: 9245503
- 438 2. Yamamura Y, Yamashiro K, Tsuruoka N, Nakazato H, Tsujimura A, Yamaguchi N (1997)  
439 Molecular cloning of a novel brain-specific serine protease with a kringle-like structure and  
440 three scavenger receptor cysteine-rich motifs. *Biochem Biophys Res Commun* 239:386-392.  
441 PMID: 9344839
- 442 3. Proba K, Gschwend TP, Sonderegger P (1998) Cloning and sequencing of the cDNA  
443 encoding human neurotrypsin. *Biochim Biophys Acta* 1396:143-147. PMID: 9540828
- 444 4. Molinari F, Rio M, Meskenaite V, Encha-Razavi F, Auge J, Bacq D, Briault S, Vekemans  
445 M, Munnich A, Attie-Bitach T, Sonderegger P, Colleaux L (2002) Truncating neurotrypsin  
446 mutation in autosomal recessive nonsyndromic mental retardation. *Science* 298:1779-1781.  
447 PMID: 12459588
- 448 5. Reif R, Sales S, Hettwer S, Dreier B, Gisler C, Wolfel J, Luscher D, Zurlinden A, Stephan  
449 A, Ahmed S, Baici A, Ledermann B, Kunz B, Sonderegger P (2007) Specific cleavage of  
450 agrin by neurotrypsin, a synaptic protease linked to mental retardation. *FASEB J* 21:3468-  
451 3478. PMID: 17586728
- 452 6. Stephan A, Mateos JM, Kozlov SV, Cinelli P, Kistler AD, Hettwer S, Rulicke T, Streit P,  
453 Kunz B, Sonderegger P (2008) Neurotrypsin cleaves agrin locally at the synapse. *FASEB J*  
454 22:1861-1873. PMID: 18230682
- 455 7. Butikofer L, Zurlinden A, Bolliger MF, Kunz B, Sonderegger P (2011) Destabilization of  
456 the neuromuscular junction by proteolytic cleavage of agrin results in precocious sarcopenia.  
457 *FASEB J* 25:4378-4393. PMID: 21885656

- 458 8. Mitsui S, Hidaka C, Furihata M, Osako Y, Yuri K (2013) A mental retardation gene,  
459 motopsin/prss12, modulates cell morphology by interaction with seizure-related gene 6.  
460 Biochem Biophys Res Commun 436:638-644. PMID: 23770366
- 461 9. Shimizu-Nishikawa K, Kajiwara K, Sugaya E (1995) Cloning and characterization of  
462 seizure-related gene, SEZ-6. Biochem Biophys Res Commun 216:382-389. PMID: 7488116
- 463 10. Gunnensen JM, Kim MH, Fuller SJ, De Silva M, Britto JM, Hammond VE, Davies PJ,  
464 Petrou S, Faber ES, Sah P, Tan SS (2007) Sez-6 proteins affect dendritic arborization patterns  
465 and excitability of cortical pyramidal neurons. Neuron 56:621-639. PMID: 18031681
- 466 11. Boeuf S, Borger M, Hennig T, Winter A, Kasten P, Richter W (2009) Enhanced ITM2A  
467 expression inhibits chondrogenic differentiation of mesenchymal stem cells. Differentiation  
468 78:108-115. PMID: 19541402
- 469 12. Lagha M, Mayeuf-Louchart A, Chang T, Montarras D, Rocancourt D, Zalc A, Kormish J,  
470 Zaret KS, Buckingham ME, Relaix F (2013) Itm2a is a Pax3 target gene, expressed at sites of  
471 skeletal muscle formation in vivo. PLoS One 8:e63143. PMID: 23650549
- 472 13. Tai TS, Pai SY, Ho IC (2014) Itm2a, a target gene of GATA-3, plays a minimal role in  
473 regulating the development and function of T cells. PLoS One 9:e96535. PMID: 24831988
- 474 14. Van den Plas D, Merregaert J (2004) Constitutive overexpression of the integral  
475 membrane protein Itm2A enhances myogenic differentiation of C2C12 cells. Cell Biol Int  
476 28:199-207. PMID: 14984746
- 477 15. Freeman M, Ashkenas J, Rees DJ, Kingsley DM, Copeland NG, Jenkins NA, Krieger M  
478 (1990) An ancient, highly conserved family of cysteine-rich protein domains revealed by  
479 cloning type I and type II murine macrophage scavenger receptors. Proc Natl Acad Sci U S A  
480 87:8810-8814. PMID: 1978939
- 481 16. Resnick D, Chatterton JE, Schwartz K, Slayter H, Krieger M (1996) Structures of class A  
482 macrophage scavenger receptors. Electron microscopic study of flexible, multidomain,



483 fibrous proteins and determination of the disulfide bond pattern of the scavenger receptor  
484 cysteine-rich domain. *J Biol Chem* 271:26924-26930. PMID: 8900177

485 17. Martinez VG, Moestrup SK, Holmskov U, Mollenhauer J, Lozano F (2011) The  
486 conserved scavenger receptor cysteine-rich superfamily in therapy and diagnosis. *Pharmacol*  
487 *Rev* 63:967-1000. PMID: 21880988

488 18. Ojala JR, Pikkarainen T, Tuuttila A, Sandalova T, Tryggvason K (2007) Crystal structure  
489 of the cysteine-rich domain of scavenger receptor MARCO reveals the presence of a basic  
490 and an acidic cluster that both contribute to ligand recognition. *J Biol Chem* 282:16654-  
491 16666. PMID: 17405873

492 19. Gisler C, Luscher D, Schatzle P, Durr S, Baici A, Galliciotti G, Reif R, Bolliger MF,  
493 Kunz B, Sonderegger P (2013) Zymogen activation of neurotrypsin and neurotrypsin-  
494 dependent agrin cleavage on the cell surface are enhanced by glycosaminoglycans. *Biochem*  
495 *J* 453:83-100. PMID: 23560819

496 20. Frischknecht R, Fejtova A, Viesti M, Stephan A, Sonderegger P (2008) Activity-induced  
497 synaptic capture and exocytosis of the neuronal serine protease neurotrypsin. *J Neurosci*  
498 28:1568-1579. PMID: 18272678

499 21. Ozhogina OA, Grishaev A, Bominaar EL, Patthy L, Trexler M, Llinas M (2008) NMR  
500 solution structure of the neurotrypsin Kringle domain. *Biochemistry* 47:12290-12298. PMID:  
501 18956887

502 22. McNicholas S, Potterton E, Wilson KS, Noble ME (2011) Presenting your structures: the  
503 CCP4mg molecular-graphics software. *Acta Crystallogr D Biol Crystallogr* 67:386-394.  
504 PMID: 21460457

505 23. Svergun D, Barberato C, Koch MHJ (1995) CRY SOL - a Program to Evaluate X-ray  
506 Solution Scattering of Biological Macromolecules from Atomic Coordinates. *Journal of*  
507 *Applied Crystallography* 28:768-773.

508 24. Holm L, Rosenstrom P (2010) Dali server: conservation mapping in 3D. *Nucleic Acids*  
509 *Res* 38:W545-549. PMID: 20457744

510 25. Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, Ben-Tal N (2016)  
511 ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation  
512 in macromolecules. *Nucleic Acids Res* 44:W344-350. PMID: 27166375

513 26. Purushotham S, Deivanayagam C (2014) The calcium-induced conformation and  
514 glycosylation of scavenger-rich cysteine repeat (SRCR) domains of glycoprotein 340  
515 influence the high affinity interaction with antigen I/II homologs. *J Biol Chem* 289:21877-  
516 21887. PMID: 24923446

517 27. Bikker FJ, Ligtenberg AJ, End C, Renner M, Blaich S, Lyer S, Wittig R, van't Hof W,  
518 Veerman EC, Nazmi K, de Blicq-Hogervorst JM, Kioschis P, Nieuw Amerongen AV,  
519 Poustka A, Mollenhauer J (2004) Bacteria binding by DMBT1/SAG/gp-340 is confined to  
520 the VEVLXXXXW motif in its scavenger receptor cysteine-rich domains. *J Biol Chem*  
521 279:47699-47703. PMID: 15355985

522 28. Yap NV, Whelan FJ, Bowdish DM, Golding GB (2015) The Evolution of the Scavenger  
523 Receptor Cysteine-Rich Domain of the Class A Scavenger Receptors. *Front Immunol* 6:342.  
524 PMID: 26217337

525 29. Brannstrom A, Sankala M, Tryggvason K, Pikkarainen T (2002) Arginine residues in  
526 domain V have a central role for bacteria-binding activity of macrophage scavenger receptor  
527 MARCO. *Biochem Biophys Res Commun* 290:1462-1469. PMID: 11820786

528 30. Madsen M, Moller HJ, Nielsen MJ, Jacobsen C, Graversen JH, van den Berg T, Moestrup  
529 SK (2004) Molecular characterization of the haptoglobin.hemoglobin receptor CD163.  
530 Ligand binding properties of the scavenger receptor cysteine-rich domain region. *J Biol*  
531 *Chem* 279:51561-51567. PMID: 15448162

532 31. Reif R, Sales S, Dreier B, Luscher D, Wolfel J, Gisler C, Baici A, Kunz B, Sonderegger P  
533 (2008) Purification and enzymological characterization of murine neurotrypsin. *Protein Expr*  
534 *Purif* 61:13-21. PMID: 18577456

535 32. Bikker FJ, Ligtenberg AJ, Nazmi K, Veerman EC, van't Hof W, Bolscher JG, Poustka A,  
536 Nieuw Amerongen AV, Mollenhauer J (2002) Identification of the bacteria-binding peptide  
537 domain on salivary agglutinin (gp-340/DMBT1), a member of the scavenger receptor  
538 cysteine-rich superfamily. *J Biol Chem* 277:32109-32115. PMID: 12050164

539 33. Sankala M, Brannstrom A, Schulthess T, Bergmann U, Morgunova E, Engel J,  
540 Tryggvason K, Pikkarainen T (2002) Characterization of recombinant soluble macrophage  
541 scavenger receptor MARCO. *J Biol Chem* 277:33378-33385. PMID: 12097327

542 34. Bowen MA, Bajorath J, Siadak AW, Modrell B, Malacko AR, Marquardt H, Nadler SG,  
543 Aruffo A (1996) The amino-terminal immunoglobulin-like domain of activated leukocyte cell  
544 adhesion molecule binds specifically to the membrane-proximal scavenger receptor cysteine-  
545 rich domain of CD6 with a 1:1 stoichiometry. *J Biol Chem* 271:17390-17396. PMID:  
546 8663238

547 35. Studier FW (2005) Protein production by auto-induction in high density shaking cultures.  
548 *Protein Expr Purif* 41:207-234. PMID: 15915565

549 36. Kabsch W (2010) Xds. *Acta Crystallogr D Biol Crystallogr* 66:125-132. PMID:  
550 20124692

551 37. Evans PR (2011) An introduction to data reduction: space-group determination, scaling  
552 and intensity statistics. *Acta Crystallogr D Biol Crystallogr* 67:282-292. PMID: 21460446

553 38. Hohenester E, Sasaki T, Timpl R (1999) Crystal structure of a scavenger receptor  
554 cysteine-rich domain sheds light on an ancient superfamily. *Nat Struct Biol* 6:228-232.  
555 PMID: 10074941

- 556 39. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment  
557 search tool. *J Mol Biol* 215:403-410. PMID: 2231712
- 558 40. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high  
559 throughput. *Nucleic Acids Res* 32:1792-1797. PMID: 15034147
- 560 41. Stein N (2008) CHAINSAW: a program for mutating pdb files used as templates in  
561 molecular replacement. *Journal of Applied Crystallography* 41:641-643.
- 562 42. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ (2007)  
563 Phaser crystallographic software. *J Appl Crystallogr* 40:658-674. PMID: 19461840
- 564 43. Emsley P, Lohkamp B, Scott WG, Cowtan K (2010) Features and development of Coot.  
565 *Acta Crystallographica Section D* 66:486-501.
- 566 44. Murshudov GN, Skubak P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, Winn  
567 MD, Long F, Vagin AA (2011) REFMAC5 for the refinement of macromolecular crystal  
568 structures. *Acta Crystallogr D Biol Crystallogr* 67:355-367. PMID: 21460454
- 569 45. Chen VB, Arendall WB, 3rd, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, Murray  
570 LW, Richardson JS, Richardson DC (2010) MolProbity: all-atom structure validation for  
571 macromolecular crystallography. *Acta Crystallogr D Biol Crystallogr* 66:12-21. PMID:  
572 20057044
- 573 46. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE  
574 (2004) UCSF Chimera--a visualization system for exploratory research and analysis. *J*  
575 *Comput Chem* 25:1605-1612. PMID: 15264254
- 576 47. Brennich ME, Round AR, Hutin S (2017) Online Size-exclusion and Ion-exchange  
577 Chromatography on a SAXS Beamline. *J Vis Exp*. PMID: 28117806
- 578 48. Panjkovich A, Svergun DI (2017) CHROMIXS: automatic and interactive analysis of  
579 chromatography-coupled small angle X-ray scattering data. *Bioinformatics*. PMID: 29300836

580 49. Franke D, Petoukhov MV, Konarev PV, Panjkovich A, Tuukkanen A, Mertens HDT,  
581 Kikhney AG, Hajizadeh NR, Franklin JM, Jeffries CM, Svergun DI (2017) ATSAS 2.8: a  
582 comprehensive data analysis suite for small-angle scattering from macromolecular solutions.  
583 J Appl Crystallogr 50:1212-1225. PMID: 28808438

584 50. Karplus PA, Diederichs K (2012) Linking crystallographic model and data quality.  
585 Science 336:1030-1033. PMID: 22628654

586 51. Robert X, Gouet P (2014) Deciphering key features in protein structures with the new  
587 ENDscript server. Nucleic Acids Res 42:W320-324. PMID: 24753421