

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The Functional Versatility of Transferrin Receptor 2 and Its Therapeutic Value

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1742689> since 2020-07-01T10:37:53Z

Published version:

DOI:10.3390/ph11040115

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

The functional versatility of Transferrin Receptor 2 and its therapeutic value

Antonella Roetto, PhD, Mariarosa Mezzanotte BsD, Rosa Maria Pellegrino, PhD

University of Torino, Department of Clinical and Biological Sciences, 10043 Orbassano, Torino, Italy

TFR2 GENE AND PROTEINS

Tfr2 is a type II transmembrane glycoprotein, member of the transferrin receptor family and homologous to Tfr1 [1].

It is **encoded** by *TFR2*, a 2471 bp long gene localized on the long arm of human chromosome 7 (7q22.1), that consists of 18 exons and gives origin to two main variants regulated by different specific promoters: Tfr2 α and Tfr2 β (**Figure 1**).

Tfr2 α results from the transcription of all exons and is prevalently and highly expressed in hepatocytes and erythroid cell lines. Tfr2 α cDNA is 2.3 kb long (AF067864) and the Tfr2 α is a protein of about 89 kDa encompassing 801 amino acids [2]. As Tfr1, Tfr2 α has a short cytoplasmic tail (aa 1-80) that contains a consensus sequence YQRV for endocytosis, a transmembrane domain (aa 81-104) with four cysteines (aa 89-98 and 108-111), involved in disulphide bonds, likely responsible for *TFR2* homodimerization, and a large extracellular domain (aa 105-801) comprising a protease-associated domain and two RGD motifs that bind **di-ferric Tf (Fe₂Tf)**. Furthermore, an N-terminal mitochondrial targeting sequence (MTS) has been found in Tfr2 intracellular domain [3]. *In vitro* analysis demonstrated that Tfr2 α on cell membrane can be shed and give origin to a soluble form and that this process is inhibited by **Fe₂Tf** [4]; however, this form could not be found in animal or human sera.

Tfr2 α transcription is up-regulated in **mouse embryonic fibroblast cells** (NIH3T3) by erythroid GATA1, EKLF and cEBP/ α transcriptional factors, while FOG1 seems to inhibit GATA1 enhancement [5]. Also, hepatic Hnf4 α stimulates Tfr2 α transcription since it is significantly decreased in liver-specific HNF4 α -null mice [6]. There is no Tfr2 α IRE-IRP dependent posttranscriptional regulation [7], while the hepatic tetraspanin CD81 is able to interact with Tfr2 α and induce its degradation [8].

Tfr2 β has an in-frame transcription start site in exon 4, so Tfr2 β cDNA (NM_001206855.1) lacks exons 1-3 transcript and presents 142 additional untranslated base pairs at its 5' end. Tfr2 β is ubiquitously expressed at low level, mostly expressed in spleen, heart and brain. The resulting protein lacks the cytoplasmic and the transmembrane domain [2]. Since no signal peptide involved in secretory pathway could be evidenced in Tfr2 β isoform, it is supposed to be a cytosolic 60 kDa protein identical to the Tfr2 α extracellular domain. At the moment, no transcriptional/translational regulatory pathway is known for Tfr2 β isoform (**Figure 1**).




Figure 1. Schematic representation of A) *TFR2* gene structure. *TFR2* 18 exons are in bright yellow *TFR2* α promoter is shown upstream the gene and transcriptional factors binding sites are highlighted by black arrows. 5' untranslated region (152 bp long) of *TFR2* β transcript is shown in orange; B) the two main transcripts and of Tfr2 α and Tfr2 β isoforms, that are identical in the common sequence and the two proteins localization, on the cell surface or in the cytosol respectively.

TFR2 and HFE3

Inactivating mutations of *TFR2* gene (OMIM: 604720) lead to type 3 **hereditary** hemochromatosis (*TFR2*-HHC or HFE3), a rare recessive disorder characterized by increased transferrin saturation and serum ferritin concentration and iron overload [9].

HFE3 is one of the 5 different forms of hereditary hemochromatosis, a genetically heterogeneous disorder due to the deregulation of iron protein Heparin (Hepc) [10] (**Table1**). *TFR2*-HHC presents an earlier age of onset than **type 1 hereditary hemochromatosis (HFE-HHC)** and some pediatric patients have been reported so far. However, the majority of the affected individuals are young adults with abnormal serum iron indices [11].

Most of the mutations involved in HFE3 pathogenesis cause an inactivation of both Tfr2 isoforms, but some of them, occurring in exons from 1 to 3, impair the production of only the Tfr2 α isoform [12-14]. Three patients with homozygote mutation M172K, that impairs Tfr2 β translation initiation codon were identified, all presenting typical hemochromatosis symptoms (cirrhosis, hypogonadism, **cardiomyopathy**, arthritis) **on average age of 38 \pm 5 years** [13,15].

Unfortunately, few clinical data are available on patients with these mutations to allow an exhaustive genotype/phenotype analysis.

Table 1. HHC types and their relationship with Hepcidin.

HH type	Acronym	Inheritance	Gene	Protein	Function
HFE1	HFE-HHC	AR	HFE	Hfe	Hepc regulator
HFE2a	HJV-HHC	AR	HJV	Hemojuvelin	Hepc regulator
HFE2b	HEPC-HHC	AR	HAMP	Hepc	Fe absorption inhibitor
HFE3	TFR2-HHC	AR	TFR2	Tfr2	Hepc regulator
HFE4	FPN1-HHC	AD	SLC40A1	Fpn1	Hepc receptor

Systemic Iron Metabolism: the Hepc-Fpn1 axis and the proteins involved in Hepc regulation

In mammals, the hepatocyte-secreted hormone Hepc regulates systemic iron homeostasis [16]. Hepc is codified by *HAMP* gene, which encodes for an 84 amino acids precursor protein from which active 20–25 amino acids peptides are generated [17]. It is expressed primarily in the liver although low levels have been also reported in other organs [18].

How *HAMP* gene expression is regulated is mostly unknown. There are no IRE elements in its transcript and the transcriptional factor CCAAT/enhancer binding protein- α is highly expressed in the liver and seems to stimulate *HAMP* expression, while the hepatocyte nuclear factor 4- α (HNF-4) represses Hepc expression [19].

The molecular processes involved in hepatic Hepc regulation are quite complex. Basal Hepc expression is regulated through the bone morphogenetic protein 6 (Bmp6) and **Smad** protein signalling pathway. In iron excess condition, Bmp6, produced and secreted by liver sinusoidal endothelial cells (LSECs) [20], binds to bone morphogenetic protein receptors, ALK2 and/or ALK3 [21], activin receptor type 2A (Actr2a) [22], Hemojuvelin (Hjv) and Neogenin [23]. The protein complex activates signals transducers Smad1/5/8 leading to their interaction with the common mediator **Smad4**. As a consequence of this interplay, Smad4 translocates into the nucleus and promotes Hepc transcription [16].

More recently, it has been demonstrated that bone morphogenetic protein 2 (Bmp2), expressed in LSECs, can **also** trigger Hepc transcription increase [24].

A second Hepc regulatory pathway involves **di-ferric Tf (Fe2Tf)** as the **signalling** of increased iron availability, transferrin receptor 1 (Tfr1), hemochromatosis type 1 protein (Hfe) and transferrin receptor 2 (Tfr2). It has been demonstrated that **Fe2Tf** competes with Hfe for binding Tfr1 then, when circulating **Fe2Tf** increases as a consequence of iron **raising**, Hfe dissociates from Tfr1 and binds Tfr2 [25]. Hfe/Tfr2 complex **is then** responsible for Hepc response to iron increase, through the activation of **Erk1/2** and MAPK cascade that has been proposed to potentially converge on the Bmps/Smad1/5/8 mediated pathway [26].

The hierarchy of the two pathways activation and their relationship are still not completely defined. *In vitro* data support the hypothesis that the complex Hfe/Tfr2 interacts with membrane Hvj (mHjv) on cell surface, **thereby** the link between the two signalling pathways occurs [27]. **It has been found** found *in vivo* that both Hfe and Tfr2 KO mice present lower pErk1/2 [28] and pSmad1/5/8 proteins [29,30], meaning that these two proteins **regulate** both signal translation pathways. **Hepatic Hcpc up-regulation** is inhibited by matriptase 2 (MT-2 or Tmprss6), that acts as Hcpc inhibitor cleaving mHjv expressed on the plasma membrane [31]. *TMPRSS6* gene expression has been found to be induced by chronic dietary iron loading and Bmp6 injection [32] and its interaction with Neogenin facilitates mHjv cleavage and inactivation in transfected cells [23] (**Figure 2**).




Figure 2. Graphic representation of the hepatic Hcpc pathway: In condition of systemic iron increase. The iron signalling proteins (Fe2Tf and Bmps) interact with iron sensors (Tfr1, Hfe, Tfr2, BmpRs) and their co-activator (mHjv) to promote Hcpc production. Hcpc secreted by hepatocytes is transported in plasma and binds iron exporter Fpn1 on duodenal and reticuloendothelial cells surface causing its internalization and intracellular degradation. As a consequence, iron remains entrapped in these cells, systemically reducing the metal availability.

On the contrary, in iron deficiency condition this signalling pathway is inhibited by soluble Hjv (sHjv) and Tmprss6, which physically interacts with mHjv causing its fragmentation.

Hcpc expression in hepatocytes is systemically regulated by multiple signals: systemic iron availability, such as iron loaded transferrin and hepatic iron stores, erythropoietic activity, hypoxia

and inflammation [33]. Hepc secreted by hepatocytes regulates iron release from duodenal enterocytes, splenic macrophages and hepatocytes, which are responsible for dietary iron absorption, contain large amounts of iron from erythrocyte recycling and act as an iron reservoir and export iron when needed, respectively. Hepc exerts its function binding the iron exporter Ferroportin 1 (Fpn1) [34] and stimulating the complex internalization and degradation, leading *de facto* to cellular retention of iron [35]. Elevated plasma Hepc, as in inflammatory state, downregulates iron efflux from several cell types and this leads to an overall reduction in plasma iron. On the contrary, low Hepc, as seen in iron-depleted or erythropoietic expansion conditions, causes an increased iron release by macrophages and by the basolateral site of villi duodenal cells.

A potent Hepc inhibitor signal is iron demand for erythropoiesis, mediated by three Hepc modulators (Gdf15, Twgs1, Erfe). Their roles and precise mechanisms in Hepc regulation are still not completely clear, but Erfe in particular has emerged as a potent Hepc negative regulator in condition of acute erythropoietic demand acting in conjunction with erythropoietin (Epo) signalling as well as in anemia of inflammation condition [36].

HAMP expression is induced by inflammation and infection. This acute phase response involves a different pathway from the above described one and is mainly mediated by interleukin 6 (IL-6) inflammatory cytokine and requires the signal transducer and activator of transcription 3 (STAT3) activation and the binding of STAT3 to a STAT3-binding motif in the Hepc promoter [37]. In addition, cytokine IL-22, involved in immunological response to extracellular infections, as well as Toll-like receptor 5 (TLR5) agonist flagellin, seem to upregulate Hepc, strengthening the hypothesis of a possible Hepc role in innate immunity [38].

Conversely, *HAMP* expression is repressed by hypoxia both *in vitro* and *in vivo* in animal models [33] and humans [39,40]. The mediator of Hepc response to hypoxia seems to be the hypoxia inducible factor (HIF), even though it is not clear if it acts directly or indirectly on Hepc regulation [16]. The fact that Hepc inhibitor *Tmprss6* presents a hypoxia responsive element (HRE) in its promoter [41] might make *Tmprss6* the linking protein between hypoxia and iron homeostasis.

Mutations in *HAMP* gene and in genes codifying for Hepc regulatory proteins (*HFE*, *TFR2* and *HJV*) cause the lack of up-regulation of Hepc as a response to increased liver iron stores. In fact, inappropriately low levels of liver Hepc are observed in patients and mouse models of hereditary hemochromatosis [42,43]. These conditions promote a continuous dietary iron absorption that leads to iron overload. On the contrary, inappropriately high Hepc has been found in animal models and patients with iron resistant iron deficient anemia (IRIDA), a genetic disorder due to mutations in *TMPRSS6* gene [44-46].

TFR2 IN LIVER

In the liver, *Tfr2α* is a sensor of circulating iron, but the knowledge about the *Tfr2α* hepatic function is still incomplete. It is known that *Tfr2α* localizes in caveolar microdomains [47], membrane structures involved in the recruitment of receptors that can be activated by ligand binding [48]. Also, *Tfr2α* localizes in lipid raft domains on the exosomal cell membrane where it is internalized by clathrin-mediated endocytosis, if transferrin saturation (TS) is low [49].

Tfr2α protein regulation occurs mainly through its stabilization on the cell membrane as a consequence of the binding to **Fe2Tf** [50,51]. An *in vitro* study showed that in presence of **Fe2Tf**, *Tfr2α* has an increased half-life and is recycled, while in presence of apo-Tf membrane *Tfr2α* is

mainly subjected to lysosomal degradation [52]. It has been recently demonstrated that CD81 is also able to induce Tfr2 α degradation, but the correlation between this Tfr2 α regulatory route and Hpc pathway is still obscure [8].

Therefore, increased TS has an opposite effect on the two Tfrs via two different mechanisms: it causes a decrease of Tfr1, regulated by the IRE-IRP system, but a stabilization of Tfr2 α on the cell surface [1].

This supports the hypothesis that Tfr2 α exerts its function(s) as a signalling receptor more than as an iron importer.

According to the available *in vitro* data, hepatic Tfr2 α interacts on the cell membrane with two main iron proteins, Tfr1 and Hfe.

The current model assumes that Tfr2 α , in conjunction with HFE and Tfr1, is partner of a sensor complex of circulating iron, that activates Hpc in response to elevated TS [53]. In physiological conditions (TS 30-35%) Hfe and the complex Tf/Tfr1 are bound on the plasma cells; when TS increases in response to increased iron availability, loaded Tf impairs Hfe binding to Tfr1/Tf complex, leading it to bind Tfr2 α that is stabilized on the membrane by the same Fe²Tf. The resulting complex Tfr2 α /Fe²Tf/Hfe causes the activation of Hpc transcription [25]. In the HuH7 hepatoma cells surface, this Tfr2 α /HFE interaction occurs within a multiprotein complex, that includes also mHjv, [27]. It remains to be demonstrated if this complex activates the intracellular signalling to up-regulate Hpc expression also *in vivo*.

In presence of Fe²Tf, Tfr2 α is able to activate Erk1/2 and p38 MAPK kinase signalling transduction pathway [47], since Tfr2 KO mice present a decrease of pErk1/2 [28]. Furthermore, also Smad1/5/8 pathway seems to be involved in Tfr2 α mediated signal transduction, since pSmad1/5/8 is decreased in Tfr2 KO animals as well [29].

Erk 1/2 phosphorylation could be increased also by Hfe overexpression and both Tfr2 and Hfe cause an increase of the pro-hormone convertase Furin [29], previously demonstrated to be involved in Hpc regulation [54]. Whether it is the sole and/or the main Tfr2 dependent Hpc regulatory pathway is still not clear.

TFR2 MOUSE MODELS

The first Tfr2 KO animal model was generated by targeted mutagenesis, introducing a premature stop codon (Y245X) in the murine Tfr2 coding sequence [55]. This mutation is homologous to the Y250X variant originally detected in humans and responsible for HFE3 [9]. Even young homozygous Y245X mice maintained on a standard diet had high liver iron concentration, in agreement with the observation of early iron overload in HFE3 patients [11]. As in humans, heterozygous animals were normal. The histological distribution of iron resembles the features of HFE3, with the typical liver periportal accumulation.

Subsequently, different murine models of Tfr2 inactivation were developed, including Tfr2 total (Tfr2 KO) and liver-specific (Tfr2 LCKO) knock-out [56,57] as well as Tfr2/Hfe double KO [28]. All these models are characterized by an inadequate hepatic Hpc expression and liver iron overload with variable severity. However, when generated in the same genetic background, Tfr2 KO mice were shown to have a more severe iron overload than Hfe KO although less severe than Tfr2/Hfe double KO [58]. These observations are in agreement with the model of Tfr2/Hfe proteins cooperation in the liver.

In a double Tfr2/Hjv KO mouse model, **plasma** Hepc and Hepc transcription was lower than in Tfr2 and similar to Hjv single KOs respectively. The same was true for the **Tfr2/Hfe** double KO, [59]. Also, a recent study on a mouse model with inactivation of both Bmp6 and Tfr2 (Tfr2/Bmp6 double KO) demonstrated that loss of functional Tfr2 further represses Hamp expression, **Smad5** phosphorylation and **plasma** Hepc amount in Bmp6 KO mice. The same results were obtained in the Hfe/Bmp6 double KO and the Hfe/Bmp6/Tfr2 triple KO [60]. All these data support the hypothesis that Tfr2 and Hfe act downstream Bmp6 and upstream Hjv in Hepc regulatory pathway.

Last, Tfr2 germinal vs liver specific KO animals highlighted a distinct function of Tfr2 outside the liver in maintaining iron balance. In fact, Tfr2 KO mice have less severe iron overload, slightly higher hemoglobin (Hb) levels [57,61] and moderate macrocytosis than Tfr2 LCKO [56,57].

To study the specific function of Tfr2 β isoform in iron metabolism regulation, a specific mouse model was generated introducing the M167K substitution in the Tfr2 protein [57]. This mutation, homologous to the one found in naturally mutant individuals with HFE3, substitutes the start codon methionine of the Tfr2 β isoform with a lysine. Interestingly, this knock-in mouse model (Tfr2 KI), specifically lacking the Tfr2 β -isoform ($\alpha^+\beta^0$) is characterized by normal transferrin saturation, liver iron concentration, Hepc and Bmp6 levels, but shows a transient anemia at young age. In additions, adult Tfr2 KI animals accumulate iron in the spleen, due to a significant reduction of iron exporter Fpn1 mRNA, thus suggesting a possible regulatory effect of Tfr2 β isoform on splenic Fpn1 expression. These data are further supported by the results obtained in Tfr2 macrophage-specific KO mouse model. These animals present normal systemic iron parameters but lower Fpn1 transcript and protein in peritoneal macrophages [62]. Recent studies demonstrated that Tfr2 β is well expressed in reticuloendothelial cells of different tissues, where it exerts its role in modulating iron availability in these tissues acting on Fpn1 transcription (see below). Since Fpn1 protein has several regulatory systems both at transcriptional [63,64] and post-transcriptional level through IRE/IRP system [7] and origins from different Fpn1 transcripts with or without IREs [65], it remains to be clarified how and when Tfr2 β acts on Fpn1 regulation.

TFR2 IN EXTRA-HEPATIC TISSUES

Tfr2 in the erythropoietic compartment

A Tfr2 α erythropoietic role was firstly hypothesized in genome-wide association studies that identified Tfr2 α polymorphisms affecting hematologic parameters [66,67]. These data were further strengthened by the identification of Tfr2 α as a component of the Erythropoietin receptor (EpoR) complex in erythroid progenitor cells. Tfr2 α results to be crucial for efficient transport of EpoR to the cell surface and for its terminal differentiation since human erythroid progenitors with silenced Tfr2 α show a delayed differentiation [68]. Another hint was provided by the increased Hb content present only in Tfr2 germinal KO but not in liver-specific KO mice. Since both mouse models manifest comparable iron overload, the lack of enhanced hemoglobinization in Tfr2 LCKO mice suggests that the erythroid function of Tfr2 α is preserved [68]. Also, double Tmprss6/Tfr2 KO mice

develop erythrocytosis, while in double *Tmprss6/Tfr2* LCKO mice, where *Tfr2 α* is functional in erythroid cells, RBC number is normal [61].

Recently, a mouse model lacking *Tfr2* in bone marrow cells (*Tfr2^{BMKO}*) was developed injecting BM cells from *Tfr2* KO mice in lethally irradiated C57/BL6 mice. *Tfr2^{BMKO}* mice manifest reduced MCV and low Hepc levels as typical response to iron deficiency, but an enhanced terminal erythropoiesis showed by increased RBC and Hb content [69]. Interestingly, erythropoiesis and Epo level in these mice do not change in a mild dietary restriction setting, as it happens for WT animals, where Epo level is drastically increased.

As a whole, these data suggest that the lack of *Tfr2* confers increased Epo sensitivity to erythroid progenitor cells, a hypothesis that is further supported by the induction of Epo target genes like Hamp regulator *Erfe* [70] in these animals.

A similar animal model was recently developed crossing *Tfr2* floxed mice with Vav-Cre expressing mice to obtain *Tfr2* silencing in erythroid compartment [62]. Results differ from the previous work since decreased RBC and splenomegaly were observed, but these discrepancies might be explained by the different procedures used to create the two mouse models, since in the first case *Tfr2* is silenced in all bone marrow (BM) cell lines after a BM transplant procedure, while in the latter only the erythroid cell line are *Tfr2* null.

In another study, *Tfr2* erythropoietic role was further investigated studying the erythropoiesis of two *Tfr2* mice with one or both *Tfr2* isoforms silenced (*Tfr2* KI and *Tfr2* KO) and with normal or increased iron availability [57]. The evaluations were performed in bone marrow and spleen, in young and adult animals to unravel the erythropoietic role of *Tfr2* isoforms at different ages and in the two main erythropoietic organs. It resulted that the lack of *Tfr2* in *Tfr2* KO mice leads to macrocytosis with low reticulocyte number and increased Hb value, together with an anticipation of erythropoiesis in young mice both in BM and in the spleen [71], probably because the increased systemic iron amount present in these animals allows them to reach mature erythropoiesis even at young age.

Although different animals and approaches were used in these studies and the partially contradictory results obtained, they all demonstrate that erythropoiesis is impaired by lack of *Tfr2* in BM independently from its activity in hepatic tissues.

Moreover, results obtained studying $\alpha^+\beta^0$ *Tfr2* KI mice [57] demonstrated for the first time the involvement of *Tfr2 β* in favouring iron availability for erythropoiesis. In fact, the sole lack of *Tfr2 β* , in normal systemic iron condition, causes an increased but immature splenic erythropoiesis seen only in young mice, as if they had insufficient iron availability during animal massive growth, that is normalized in animal adult age. Decreased iron availability for erythropoiesis in *Tfr2* KI young mice is demonstrated by increased ferritin (Ft) and decreased divalent metals transporter 1 (DMT1) in their splenic monocyte, the increase of *Erfe* transcription in BM and spleen and the low hepatic Hepc transcription that could in turn be responsible of increased splenic *Fpn1* amount in these animals [71].

This effect, due to *Tfr2 β* absence, in aged matched *Tfr2* KO ($\alpha^0\beta^0$) mice was compensated by the increased amount of circulating iron available that may be used for erythrocyte production. (Figure 3A).




Figure 3. Schematic model illustrating Tfr2 β function in regulation iron export from reticulo-endothelial (REL) cells. Lack of Tfr2 β causes an increased iron retention in REL cells that causes (A) the onset of an immature erythropoiesis in the spleen and (B) induces a cardio-protection against the effect of the reperfusion of oxygenated solutions after an ischemic event in heart (see text).

Tfr2 β in the Heart

The cardiac muscle is a major site of oxygen consumption, so an adequate intracellular iron pool is essential to its aerobic activity. This is demonstrated by the finding that deletion of cardiac Tfr1 in mice causes fatal energetic failure in cardiomyocytes [72]. Cardiomyocytes express relatively high levels of Hpc and Fpn1, despite the fact that these cells have no role in systemic iron control [73]. Studies on mouse models **demonstrated** that the cardiac Hpc/Fpn1 axis is essential for heart cells autonomous control of the intracellular iron pool that guaranties a normal cardiac functionality [73] and that Hpc/Fpn1 appears to protect the heart from the effects of systemic iron deficiency [74]. On the other side, cardiomyocytes are particularly susceptible to ROS-mediated damage because they are rich in mitochondria and consume large amounts of oxygen [75]. Therefore, when labile iron pool (LIP) expansion occurs, oxidative stress can affect cardiac functions, as it happens in severe juvenile HH forms [10].

Although ubiquitously expressed, Tfr2 β resulted to be highly transcribed in heart [2], so that a role for Tfr2 β isoform in cardiac iron management **has been** postulated.

Indeed, in the hearts of two Tfr2 $\alpha^+\beta^0$ mice with normal or increased systemic iron amount, Tfr2 KI and Tfr2 LCKO[57], the silencing of Tfr2 β induces a selective activation of different proteins involved in cell survival, antioxidant enzymes, and kinases involved in cardio-protective pathways that are usually activated **by** stressful stimuli.

In particular, Tfr2 KI and Tfr2 LCKO mice develop a greater resistance against acute Ischemia/Reperfusion (I/R) challenge, irrespective of animals' systemic iron content, via the activation of the RISK or SAFE/GSK3 β cardio-protective pathway respectively. The iron imbalance present in these mice hearts was demonstrated by the finding that both the models present the activation of antioxidant proteins, pro-apoptotic markers and catalase even before I/R [76]. They also have a slightly increased synthesis of cardiac ferritins, similarly to what happens in ischemic preconditioning, in which a small increase of ferritin protects cardiac cells from iron-mediated oxidative damage associated with ischemia-reperfusion injury [77].

Since previous data demonstrated a significant decrease of Fpn1 and an increased iron deposit in splenic macrophages in Tfr2 β null mice [57,71], one might hypothesize that Tfr2 β isoform inactivation in the heart causes an iron retention in cardiac reticuloendothelial cells, that is able to induce cardio-protective pathways activation and to reduce iron availability to form free oxygen radicals during the reperfusion phase (**Figure 3B**).

Tfr2 in the Central Nervous System (CNS)

Iron levels in the brain varies during life: iron amount increase with aging in the striatum and the brain stem [78]; it is present in most CNS cell types, neurons, oligodendrocytes, microglia and astrocytes [79]. A well-regulated iron homeostasis is important for brain development and function. Iron deficiency negatively impacts neurodevelopmental processes [80] and is also implicated in a number of psychiatric and neurological conditions, learning disabilities, attention deficit hyperactivity disorder (ADHD) and pediatric restlessleg–syndrome (RLS) [81,82]. On the contrary, brain iron overload is present in Alzheimer's and Huntington's neurodegenerative disorders as well as in Parkinson disease (PD). Nevertheless, the exact role of iron in these diseases onset/worsening is still debated and it remains to be clarified whether brain iron overload is directly involved in their pathogenesis or it is a secondary effect that contributes to their clinical symptoms progression.

The main sites of brain iron uptake are the brain vascular endothelial cells (BVECs) present in the blood-brain barrier (BBB) [84]. As in other organs, there are two main pathways responsible for CNS cells iron uptake, the Tf-Tfr1 pathway, and the NTBI transport pathway. Traditionally, Tf-Tfr1 is considered a major pathway and works as in all the other cell types of the organism, through a receptor mediated endocytosis of **plasma** Tf circulating in the ventricles [85].

The NTBI transport pathway has been recently reevaluated as a significant way to introduce iron in CNS and it could be done through vesicular or non-vesicular mechanism. In the first case Tf homologues, as lactoferrin and melanotransferrin, might be involved in Tfr mediated iron transport; moreover, the newly characterized Ft receptors Tim 2 and Scara5 can introduce iron inside the cells through a Ft-FtR pathway. Non vesicular iron uptake can be exerted by iron importer DMT1, that is present in endothelial cells of the brain microvasculature, as well as other importer like IN4/5/6 [86].

In CNS, iron exporter Fpn1 is found in BVECs, neurons, oligodendrocytes, astrocytes, the choroid plexus, and ependymal cells and microglia together with ceruloplasmin (CP) or hephaestin (Hp), the two ferroxidases that cooperate with Fpn1 to facilitate iron export [87]. Fpn1 could be the main responsible for iron release from CNS cell, even if other proteins and mechanism have been brought into play for these processes [86].

Inside brain cells, the majority of iron is bound to ferritin heteropolymers (Ft H/L) [88]. Their cellular distribution and ratio varies with iron status, age, and disease conditions [89].

CNS iron homeostasis is intracellularly modulated by (IRE/IRPs) system [90] and by local and systemic Hephc. Injection of Hephc into the mouse lateral cerebral ventricle **decreases** Fpn1 protein levels and treatment of primary cultured rat neurons with Hephc **decreases** Fpn1 expression and results in the subsequent release of iron [91]. More recently, it was demonstrated that injection of adenovirus expressing Hephc (ad-hepcidin) in brain ventricles reduces brain iron in iron-overloaded rats through the downregulation of iron transporter [92]. This data indicates that Hephc/Fpn1 axis is present and acts in CNS as in the other districts of the organism.

It remains a matter of debate whether Hephc acting on brain is locally produced or comes from the systemic circulation crossing the BBB or both [93].

Similar to other Hephc regulatory proteins, Tfr2 gene expression has been shown in total brain extracts [2,94], in brain tumor cell lines [95] or in specific neuronal subtypes as dopaminergic neurons [3]. Furthermore, a transcriptome study on Tfr2 null mice revealed that several genes involved in the control of neuronal functions are abnormally transcribed [96]. **Noteworthy**, the same experimental approaches applied to Hfe null mice, revealed that a consistent percentage of transcripts are modified in the same way in the two models [96]. This highlights the possibility of a cooperation between Tfr2 α and Hfe protein in CNS iron regulation as in the rest of the organism.

Immunofluorescence **studies** using a Tfr2 α specific antibody demonstrate that the protein is significantly produced in mouse hippocampus, amygdala, central nucleus, and in the hypothalamic paraventricular nucleus [97].

A recent study assessed the situation of iron in the brain **of** Tfr2 KO mouse model vs WT sib pairs submitted to an iron enriched diet (IED). They both are iron overloaded animals so one could distinguish the effects of Tfr2 silencing from those due to Tfr2-independent iron load modifications. It has been demonstrated that Tfr2 causes a lack of brain Hephc response to the systemic raise of iron levels, with altered iron mobilization and/or cellular distribution in the nervous tissue [98].

Moreover, Tfr2 KO mice present a selective over activation of neurons in the limbic circuit and the emergence of an anxious-like behaviour.

Also, microglial cells showed sensitivity to iron perturbations of Tfr2 KO mice, being more reactive, dystrophic and with a high level of apoptosis [97]. In light of these data, Tfr2 appears to be a key regulator of brain iron homeostasis and could have a role in the regulation of the brain regions that are involved in the anxiety onset, mainly the basolateral and central nucleus subregions of the amygdala [98].

TFR2 IN INTRACELLULAR IRON TRAFFICKING

It is still under debate if Tfr2 α contributes to iron introduction inside the cells. When the protein was characterized, it was reported that *in vitro* was able to introduce iron inside cells [2], but its contribution to intracellular iron amount *in vivo* seems to be quite negligible, since Tfr1 deficient mice present severe iron deficiency not compensated by the presence of Tfr2 [1].

Conversely, Tfr2 α seems to have a role in intracellular iron trafficking, at least in specific cell types. The first evidence about it was found in dopaminergic neurons, where a novel Tf/Tfr2 α -mediated iron transport pathway to the mitochondria has been reported [3]. Disruption of this

Tf/TfR2 α -dependent system has been associated with PD, this finding **highlights** the role of iron accumulation in this movement disorder [3]. In this regard, protective association between some Tf and TfR2 α genetic haplotypes and PD was reported, suggesting that Tf or a **Tf/TfR2 α** complex may play a role in the etiology of this disorders [99].

More recently, a similar TfR2 α function in iron delivery to mitochondria has been convincingly demonstrated in erythroid cells. In intermediate stage of human erythroid cells maturation, TfR2 α results to be present in cytoplasmic multi-organelles complexes formed by lysosomes surrounded by mitochondria and to be co-regulated with several proteins among which ionic channels and proteins involved in lysosomal modification and in mitochondrial membrane contacts to other intracellular organelles [100]. Therefore, TfR2 α evidenced in lysosomes **has been proposed** to be involved in iron delivery from these organelles to mitochondria for Hb synthesis. Considering the above mentioned evidence about a specific role of TfR2 α in erythropoiesis, this data might represent one of the molecular processes at the basis of TfR2 α erythropoietic function/s. (**Figure 4**).




Figure 4. Schematic model illustrating the putative intracellular Tfr2 α role in erythroid cells. Tfr2 α together with ATPVOC, aV-ATPase that contributes to vesicular acidification and lysosomal activity, Mcoln1, a lysosomal divalent cation channel, lysosomal cathepsin B (CTSB) and iron importer DMT1 could be involved in iron delivery from lysosome to mitochondria, with the collaboration of Mitofusin-2 (MFN-2), a mitochondrial outer membrane protein involved in mitochondria-endoplasmic reticulum contacts.

On the light of the above data, Tfr2 α involvement in iron delivery to the mitochondria notably seems to work at least in the two compartments in which this protein is significantly produced, brain and bone marrow. Although a similar mechanism has been evidenced also in cell lines derived from other organs (HeLa and hepatoma cell lines) [52,101], it remains an open question if this Tfr2 α function is present also in other cell types

TFR2 IN OTHER DISEASES

Being a “regulator of the iron regulator” hepcidin, Tfr2 transcriptional analysis was attempted to unravel if the Tfr2 isoforms could be reliable markers for some disorders in which iron perturbations occur.

Tfr2 in cancer

Cancer cells need increased amount of iron for their growth and iron importers up modulation confer to these cells a selective advantage.

Since its characterization, the significant transcription of *TFR2* gene appeared evident in BM cancer cells, in particular erythroid leukemic cells [5], but also in myeloid malignant cells [2], while Tfr2 β seemed much more prevalent than the Tfr2 α isoform in chronic B cells lymphocytic leukemia (B-CLL) [102].

Due to high *TFR2* expression in erythroid lineage and to its functional relationship with the erythropoietin receptor (EpoR) [68], *TFR2* transcription was evaluated in patients with myelodysplastic syndromes (MDS), a hematopoietic disorder with a variable risk to evolve in acute leukemia (AML) and in which chronic anemia can be corrected by Epo injection [103]. It has been found that Tfr2 α and Tfr2 β isoforms, as well as EPOR transcript, have lower level of transcription in BM from high risk MDS patients, such as RAEB2, compared to controls and low risk MDS cells [104]. Likewise, AML patients with high level of Tfr2 α and Tfr2 β present an increased survival [105]. Therefore, Tfr2 isoforms might represent good predictive markers for MDS/AML prognosis. Calzolari et al. 2007 [106] demonstrated *TFR2* expression in colon and ovarian cancer cell lines as well as in lymphoma and glioblastoma (GBM). Moreover, in glioblastoma TB10 cell line under hypoxic condition, a marked increase of *TFR2* transcription was observed. In these cell lines, *TFR2* high expression is probably correlated with cells proliferation, since Tfr2 silencing inhibited GBM cells growth. Surprisingly, tumor cells from GBM patients with high *TFR2* transcriptional levels present a better prognosis compared to patient with low transcripts. Although, this is probably due to the fact that the highly increased Tfr2 expressing cells proliferate so are more sensitive to temozolomide, the anti-proliferative drug used in GBM therapy, more than to a direct involvement of Tfr2 in the disease course [107].

Tfr2 in Alzheimer disease (AD)

Alzheimer disease (AD) is another common degenerative disorder in which iron perturbation has been demonstrated [108].

A recent genetic study from 116 AD patients has found that a Tfr2 single nucleotide polymorphism (rs 7385804) and a Tfr2 haplotype, composed by two SNPs (rs 7385804 and rs 4434553) are associated with a decreased AD susceptibility [109].

In the same study, a significant decrease of Tfr2 transcription was found in peripheral blood mononuclear cells (PBMC) from AD patients compared to healthy controls ($p < 0.001$) [110].

TFR2 AS A THERAPEUTIC TARGET

In our opinion, three might be the major application of Tfr2 α as a therapeutic target: a) as Hepc regulator it could be a target in disorders in which Hepc amount is in some way inadequate to body iron availability; b) since Tfr2 α production is selective in specific organs and cell types, it could represent a selective target to correct iron perturbation in these organs; c) being Tfr2 a membrane protein that is able to link **plasma** Tf and to be internalized with it, this property could be utilized as a vector for drugs.

As **mentioned** above, Tfr2 α is involved in regulation of Hepc and, **in** consequence, in iron modulation according to body/organs needs. Among the Hepc related disorders there are all the hereditary hemochromatosis forms (**Table 1**) and the secondary iron overload disorders like hemoglobinopathy, where abnormally low Hepc amount leads to iron overload.

In this regard, just published data demonstrate that Tfr2 KO BM transplantation in a Beta-thalassemia intermedia (β -TI) mouse models significantly improved these animals erythropoiesis, opening a new way to the therapy of this very common disorder [111].

In IRIDA and anemia of chronic disease (ACD), where abnormally high Hepc causes the onset of an iron deficiency condition [111], Tfr2 down-modulation might be beneficial to decrease Hepc hyper-production. A preliminary study on mice revealed that Tfr2 silencing through small interfering RNAs (Tfr2-siRNA) in a single dose led to a significant Hepc down-modulation and increase transferrin saturation within 24 hours post-administration, persisting for more than two weeks, and to recovery from anemia in animal models of ACD [112].

Beside the liver, in brain Tfr2 regulates the production of local Hepc and iron amount in the CNS since Tfr2 KO mice brain have a blunted Hepc response to brain iron overload [97]. In recent time, it has emerged that brain Hepc production is altered in several neurodegenerative disorders: down-modulated in Alzheimer and Parkinson disease [113,114] and up-modulated in restless legs syndrome (RLS) [115]. It might be worthwhile to further investigate if Tfr2 is involved in these Hepc variations and consider using anti Tfr2 antibodies or siRNA based therapy to rescue Hepc physiologic values in RLS. Nowadays siRNA delivery to brain is quite difficult due to the presence of **BBB**, but the ongoing studies on nanoparticles use to target siRNA in specific sites could allow to increase the efficacy of this therapy [116].

An alternative therapeutic approach aims to deliver blocking antibody in the brain exploiting the BBB physiologic activities. It is based on the use of anti-Tfr1 antibodies, since Tfr1 well expressed in BBB endothelial cells and is involved in receptor-mediated transcytosis. Indeed, it has been demonstrated that the bispecific Tfr1/BACE1 (β -amyloid cleaving enzyme-1) antibody resulted to be effective in decreasing β -amyloid concentration in brain [117].

Due to Tfr1/Tfr2 high homology and the common cellular internalization through a receptor mediated endocytosis pathway, one could hypothesize a parallel Tfr2 based approach for CNS blocking antibody delivery. Nevertheless, it should be further confirmed Tfr2 presence on BBB endothelial cells and its involvement in transcytosis.

Recent studies **have aimed at exploiting** the Tfr1 as a vehicle for drug delivery inside the cells through endocytosis, often utilizing Tfr1 natural ligand, Tf, conjugated with different synthetic molecules [1]. Moreover, since Tfr1 is able to bind and internalize FtH also [118], FtH nanocages conjugated with a PARP inhibitor, olaparib, was developed for breast cancer therapy [119].

Due to its strict homology to Tfr1, one might hypothesize that Tfr2 α could also be utilized to deliver drugs inside cells. This approach would be particularly useful for two main Tfr2 α features; Tfr2 α selective expression in particular tissues (hepatic, erythroid and in CNS) and Tfr2 α high expression in several tumor cells, sometimes with Tfr1 and sometimes without it. High expression of Tfr1 and Tfr2 α were **in fact** detected in tumor and para-cancerous normal liver tissues collected from 41 patients with **hepato cellular carcinoma (HCC)** [120]. Tfr2 α is also **highly** expressed in brain tumor cells in several cases of anaplastic astrocytoma and glioblastoma, but not in normal brain or endothelial brain cells [121].

Furthermore, on the light of the new data on Tfr2 α function in delivery iron to mitochondria, Tfr2 α could possibly represent a good vehicle for drug delivery in these organelles [122], paving the way to tailored therapies for mitochondrial iron unbalanced disorders, notably Friederich Ataxia [123].

Unfortunately, too few functional data are available at the moment on Tfr2 β isoform to foresee possible therapeutic applications.

CONCLUSIONS

The process of body iron homeostasis is complex, and since there is no apparent significant excretory pathway, the amount of iron absorbed from mature duodenal enterocytes and that recycled by macrophages need to be tightly regulated. **From** the last decades, iron pathways have been enriched of new regulatory proteins whose functions **remains to** be elucidated from the molecular point of view. Each of them could represent a potential target for a focused pharmacological therapy of disorders with iron unbalance, that still represent the most **prevalent diseases** all around the world.

ACKNOWLEDGEMENT

We would like to acknowledge Marco De Gobbi, Paolo Porporato, Myriam Hsu and Delia Galeno for their help in manuscript editing.

REFERENCES

1. Kawabata H. Transferrin and transferrin receptors update. *Free Radic. Biol. Med.* **2018**, pii: S0891-5849(18)31160-2. doi: 10.1016/j.freeradbiomed.2018.06.037.
2. Kawabata H.; Yang R.; Hirama T.; Vuong P.T.; Kawano S.; Gombart A.F.; Koeffler H.P. Molecular cloning of transferrin receptor 2. A new member of the transferrin receptor-like family. *J. Biol. Chem.* **1999**, 274, 20826-20832.
3. Mastroberardino P.G.; Hoffman E.K.; Horowitz M.P.; Betarbet R.; Taylor G.; Cheng D.; Na H.M.; Gutekunst C.A.; Gearing M.; Trojanowski J.Q.; Anderson M.; Chu C.T.; Peng J.; Greenamyre J.T. A novel transferrin/TfR2-mediated mitochondrial iron transport system is disrupted in Parkinson's disease. *Neurobiol. Dis.* **2009**, 34, 417-431. doi: 10.1016/j.nbd.2009.02.009.
4. Pagani A.; Vieillevoye M.; Nai A.; Rausa M.; Ladli M.; Lacombe C.; Mayeux P.; Verdier F.; Camaschella C.; Silvestri L. Regulation of cell surface transferrin receptor-2 by iron-dependent cleavage and release of a soluble form. *Haematologica* **2015**,100, 458-465. doi: 10.3324/haematol.2014.118521.
5. Kawabata H.; Germain R.S.; Ikezoe T.; Tong X.; Green E.M.; Gombart A.F.; Koeffler H.P. Regulation of expression of murine transferrin receptor 2. *Blood* **2001**, 98, 1949-1954.
6. Matsuo S.; Ogawa M.; Muckenthaler M.U.; Mizui Y.; Sasaki S.; Fujimura T.; Takizawa M.; Ariga N.; Ozaki H.; Sakaguchi M.; Gonzalez F.J.; Inoue Y. Hepatocyte Nuclear Factor 4 α controls iron metabolism and regulates transferrin receptor 2 in mouse liver. *J. Biol. Chem.* **2015**, 290, 30855-30865. doi: 10.1074/jbc.M115.694414.
7. Muckenthaler M.U.; Galy B.; Hentze M.W. Systemic iron homeostasis and the iron-responsive element/iron-regulatory protein (IRE/IRP) regulatory network. *Annu. Rev. Nutr.* **2008**, 28, 197-213. doi: 10.1146/annurev.nutr.28.061807.155521.
8. Chen J.; Enns CA. CD81 promotes both the degradation of transferrin receptor 2 (TfR2) and the TfR2-mediated maintenance of hepcidin expression. *J Biol Chem.* **2015**, 290, 7841-7850. doi: 10.1074/jbc.M114.632778.
9. Camaschella C.; Roetto A.; Calì A.; De Gobbi M.; Garozzo G.; Carella M.; Majorano N.; Totaro A.; Gasparini P. The gene TFR2 is mutated in a new type of haemochromatosis mapping to 7q22. *Nat. Genet.* **2000**, 25, 14-15.
10. Brissot P.; Cavey T.; Ropert M.; Guggenbuhl P.; Loréal O. Genetic hemochromatosis: Pathophysiology; diagnostic and therapeutic management. *Presse Med.* **2017**, 46, e288-e295. doi: 10.1016/j.lpm.2017.05.037.
11. De Gobbi M.; Roetto A. TFR2-Related Hereditary Hemochromatosis. In: GeneReviews® Adam M.P., Ardinger H.H., Pagon R.A., Wallace S.E., Bean L.J.H., Stephens K., Amemiya A., Eds., [Internet]. Seattle (WA): University of Washington, Seattle, USA, 2005 Aug 29 [updated **2018** Feb 15], pp.1993-2018. ISSN: 2372-0697
12. Biasiotto G.; Belloli S.; Ruggeri G.; Zanella I.; Gerardi G.; Corrado M.; Gobbi E.; Albertini A.; Arosio P. Identification of new mutations of the HFE, hepcidin, and transferrin receptor 2 genes by denaturing HPLC analysis of individuals with biochemical indications of iron overload. *Clin Chem.* **2003**, 49, 1981-1988.

13. Roetto A.; Totaro A.; Piperno A.; Piga A.; Longo F.; Garozzo G.; Cali A.; De Gobbi M.; Gasparini P.; Camaschella C. New mutations inactivating transferrin receptor 2 in hemochromatosis type 3. *Blood*, **2001**, 97, 2555-2260.
14. Bardou-Jacquet E.; Cunat S.; Beaumont-Epinette M.P.; Kannengiesser C.; Causse X.; Sauvion S.; Pouliquen B.; Deugnier Y.; David V.; Loréal O.; Aguilar-Martinez P.; Brissot P.; Jouanolle A.M. Variable age of onset and clinical severity in transferrin receptor 2 related haemochromatosis: novel observations. *Br J Haematol.* **2013**, 162, 278-281. doi: 10.1111/bjh.12350.
15. Majore S.; Milano F.; Binni F.; Stuppia L.; Cerrone A.; Tafuri A.; De Bernardo C.; Palka G.; Grammatico P. Homozygous p.M172K mutation of the TFR2 gene in an Italian family with type 3 hereditary hemochromatosis and early onset iron overload. *Haematologica* **2006**, 91(8 Suppl), ECR33.
16. Finberg K.E. Regulation of systemic iron homeostasis. *Curr. Opin. Hematol.* **2013**, 20, 208-214. doi: 10.1097/MOH.0b013e32835f5a47.
17. Hunter H.N.; Fulton D.B.; Ganz T.; Vogel H.J. The solution structure of human hepcidin; a peptide hormone with antimicrobial activity that is involved in iron uptake and hereditary hemochromatosis. *J. Biol. Chem.* **2002**, 277, 37597–37603.
18. Ganz T.; Nemeth E. Hepcidin and iron homeostasis. *Biochim. Biophys. Acta* **2012**, 1823, 1434-1443. doi: 10.1016/j.bbamcr.2012.01.014
19. Courselaud B.; Pigeon C.; Inoue Y.; Inoue J.; Gonzalez F.J.; Leroyer P.; Gilot D.; Boudjema K.; Guguen-Guillouzo C.; Brissot P.; Loreal O.; Ilyin G. C/EBPalpha regulates hepatic transcription of hepcidin;an antimicrobial peptide and regulator of iron metabolism. Cross-talk between C/EBP pathway and iron metabolism. *J. Biol. Chem.* **2002**, 277: 41163–41170.
20. Canali S.; Zumbrennen-Bullough K.B.; Core A.B.; Wang C.Y.; Nairz M.; Bouley R.; Swirski F.K.; Babitt JL. Endothelial cells produce bone morphogenetic protein 6 required for iron homeostasis in mice. *Blood*, **2017**, 129, 405-414. doi: 10.1182/blood-2016-06-721571.
21. Steinbicker A.U.; Bartnikas T.B.; Lohmeyer L.K.; Leyton P.; Mayeur C.; Kao S.M.; Pappas A.E.; Peterson R.T.; Bloch D.B.; Yu P.B.; Fleming M.D.; Bloch K.D. Perturbation of hepcidin expression by BMP type I receptor deletion induces iron overload in mice. *Blood*, **2011**, 118, 4224-4230. doi: 10.1182/blood-2011-03-339952.
22. Mayeur C.; Leyton P.A.; Kolodziej S.A.; Yu B.; Bloch K.D. BMP type II receptors have redundant roles in the regulation of hepatic hepcidin gene expression and iron metabolism. *Blood*, **2014**, 124, 2116-2123. doi: 10.1182/blood-2014-04-572644.
23. Enns C.A.; Ahmed R.; Zhang AS. Neogenin interacts with matriptase-2 to facilitate hemojuvelin cleavage. *J Biol. Chem.* **2012**, 287, 35104-35117. doi: 10.1074/jbc.M112.363937.
24. Canali S; Wang CY; Zumbrennen-Bullough KB; Bayer A; Babitt JL. Bone morphogenetic protein 2 controls iron homeostasis in mice independent of Bmp6. *Am. J. Hematol.* **2017**, 92, 1204-1213. doi: 10.1002/ajh.24888.
25. Schmidt P.J.; Toran P.T.; Giannetti A.M.; Bjorkman P.J.; Andrews N.C. The transferrin receptor modulates Hfe-dependent regulation of hepcidin expression. *Cell Metab.* **2008**, 7, 205-214. doi: 10.1016/j.cmet.2007.11.016.
26. Gao J.; Chen J.; Kramer M.; Tsukamoto H.; Zhang A.S.; Enns C.A. Interaction of the hereditary hemochromatosis protein HFE with transferrin receptor 2 is required for transferrin-induced hepcidin expression. *Cell Metab.* **2009**, 9, 217-227. doi: 10.1016/j.cmet.2009.01.010.
27. D'Alessio F.; Hentze M.; Muckenthaler M.U. The hemochromatosis proteins HFE; Tfr2; and HJV form a membrane-associated protein complex for hepcidin regulation. *J. Hepatol.* **2012**, 57, 1052-1060. doi:10.1016/j.jhep.2012.06.015.

28. Wallace D.F.; Summerville L.; Crampton E.M.; Frazer D.M.; Anderson G.J.; Subramaniam VN. Combined deletion of Hfe and transferrin receptor 2 in mice leads to marked dysregulation of hepcidin and iron overload. *Hepatology* **2009**, 50,1992-2000. doi: 10.1002/hep.23198.
29. Poli M.; Luscieti S.; Gandini V.; Maccarinelli F.; Finazzi D.; Silvestri L.; Roetto A.; Arosio P. Transferrin receptor 2 and HFE regulate furin expression via mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/Erk) signaling. Implications for transferrin- dependent hepcidin regulation. *Haematologica* **2010**, 95, 1832-1840.
30. Corradini E.; Rozier M.; Meynard D.; Odhiambo A.; Lin H.Y.; Feng Q.; Migas M.C.; Britton R.S.; Babitt J.L.; Fleming RE. Iron regulation of hepcidin despite attenuated Smad1,5,8 signaling in mice without transferrin receptor 2 or Hfe. *Gastroenterology* **2011**, 141, 1907-1914. doi: 10.1053/j.gastro.2011.06.077.
31. Silvestri L.; Pagani A.; Nai A.; De Domenico I.; Kaplan J.; Camaschella C. The serine protease matriptase-2 (TMPRSS6) inhibits hepcidin activation by cleaving membrane hemojuvelin. *Cell Metab.* **2008**, 8, 502-511. doi: 10.1016/j.cmet.2008.09.012.
32. Meynard D.; Vaja V.; Sun C.C.; Corradini E.; Chen S.; López-Otín C.; Grgurevic L.; Hong C.C.; Stirnberg M.; Gütschow M; Vukicevic S.; Babitt J.L., Lin H.Y. Regulation of TMPRSS6 by BMP6 and iron in human cells and mice. *Blood*, **2011**, 118, 747-756. doi: 10.1182/blood-2011-04-348698.
33. Nicolas G.; Chauvet C.; Viatte L.; Danan J.L.; Bigard X.; Devaux I.; Beaumont C.; Kahn A.; Vaulont S. The gene encoding the iron regulatory peptide hepcidin is regulated by anemia; hypoxia; and inflammation. *J. Clin. Invest.* **2002**, 110, 1037–1044.
34. Donovan A.; Lima C.A.; Pinkus J.L.; Pinkus G.S.; Zon L.I.; Robine S.; Andrews N.C. The iron exporter ferroportin/Slc40a1 is essential for iron homeostasis. *Cell Metab.* **2005**, 1, 191–200.
35. Nemeth E.; Tuttle M.S.; Powelson J.; Vaughn M.B.; Donovan A.; Ward D.M.; Ganz T.; Kaplan J. Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science* **2004**, 306, 2090–2093.
36. Papanikolaou G., Pantopoulos K. Systemic iron homeostasis and erythropoiesis. *IUBMB Life.* **2017**, 69, 399-413. doi: 10.1002/iub.1629.
37. Wrighting D.M.; Andrews N.C. Interleukin-6 induces hepcidin expression through STAT3. *Blood* **2006**, 108: 3204–3209.
38. Armitage A.E.; Eddowes L.A.; Gileadi U.; Cole S.; Spottiswoode N.; Selvakumar T.A.; Ho L.P.; Townsend A.R.; Drakesmith H. Hepcidin regulation by innate immune and infectious stimuli. *Blood*, **2011**, 118, 4129-4139. doi: 10.1182/blood-2011-04-351957.
39. Piperno A.; Galimberti S.; Mariani R.; Pelucchi S.; Ravasi G.; Lombardi C.; Bilo G.; Revera M.; Giuliano A.; Faini A.; Mainini V.; Westerman M.; Ganz T; Valsecchi M.G.; Mancina G.; Parati G.; HIGHCARE investigators. Modulation of hepcidin production during hypoxia-induced erythropoiesis in humans in vivo: data from the HIGHCARE project. *Blood* **2011**, 117, 2953-2959. doi: 10.1182/blood-2010-08-299859.
40. Talbot N.P.; Lakhal S.; Smith T.G.; Privat C.; Nickol A.H.; Rivera-Ch. M.; León-Velarde F.; Dorrington K.L.; Mole D.R.; Robbins P.A. Regulation of hepcidin expression at high altitude. *Blood*, **2012**, 119, 857-860. doi: 10.1182/blood-2011-03-341776.
41. Maurer E.; Gütschow M.; Stirnberg M. Matriptase-2 (TMPRSS6) is directly up-regulated by hypoxia inducible factor-1: identification of a hypoxia-responsive element in the TMPRSS6 promoter region. *Biol. Chem.* **2012**, 393, 535-540. doi: 10.1515/hsz-2011-0221.
42. Kawabata H.; Fleming R.E.; Gui D.; Moon S.Y.; Saitoh T.; O’Kelly J.; Umehara Y.; Wano Y.; Said J.W.; Koefler H.P. Expression of hepcidin is down-regulated in Tfr2 mutant mice manifesting a phenotype of hereditary hemochromatosis. *Blood* **2005**, 105, 376–381.
43. Nemeth E.; Roetto A.; Garozzo G.; Ganz T.; Camaschella C. Hepcidin is decreased in

- TFR2 hemochromatosis. *Blood* **2005**, 105, 1803–1806.
44. Du X.; She E.; Gelbart T.; Truksa J.; Lee P.; Xia Y.; Khovananth K.; Mudd S.; Mann N.; Moresco E.M.; Beutler E.; Beutler B. The serine protease TMPRSS6 is required to sense iron deficiency. *Science* **2008**, 320, 1088-1092. doi: 10.1126/science.1157121.
 45. Finberg K.E.; Heeney M.M.; Campagna D.R.; Aydinok Y.; Pearson H.A.; Hartman K.R.; Mayo M.M.; Samuel S.M.; Strouse J.J.; Markianos K.; Andrews N.C.; Fleming M.D. Mutations in TMPRSS6 cause iron-refractory iron deficiency anemia (IRIDA). *Nat. Genet.* **2008**, 40, 569-571. doi: 10.1038/ng.130.
 46. Folgueras A.R.; de Lara F.M.; Pendás A.M.; Garabaya C.; Rodríguez F.; Astudillo A.; Bernal T.; Cabanillas R.; López-Otín C.; Velasco G. Membrane-bound serine protease matriptase-2 (Tmprss6) is an essential regulator of iron homeostasis. *Blood* **2008**, 112, 2539-2545.
 47. Calzolari A.; Raggi C.; Deaglio S.; Sposi N.M.; Stafnes M.; Fecchi K.; Parolini I.; Malavasi F.; Peschle C.; Sargiacomo M.; Testa U. Tfr2 localizes in lipid raft domains and is released in exosomes to activate signal transduction along the MAPK pathway. *J. Cell Sci.* **2006**, 119, 4486-4498.
 48. Simons K.; Toomre D. Lipid rafts and signal transduction. *Nat. Rev. Mol. Cell. Biol.* **2000**, 1, 31-39.
 49. Chen J.; Wang J.; Meyers K.R.; Enns C.A. Transferrin-directed internalization and cycling of transferrin receptor 2. *Traffic* **2009**, 10, 1488-1501. doi: 10.1111/j.1600-0854.2009.00961.x.
 50. Johnson M.B.; Enns C.A. Diferric transferrin regulates transferrin receptor 2 protein stability. *Blood* **2004**, 104, 4287-4293.
 51. Chen J.; Enns C.A. The cytoplasmic domain of transferrin receptor 2 dictates its stability and response to holo-transferrin in Hep3B cells. *J Biol. Chem.* **2007**, 282, 6201-6209.
 52. Johnson M.B.; Chen J.; Murchison N.; Green F.A.; Enns C.A. Transferrin receptor 2: evidence for ligand-induced stabilization and redirection to a recycling pathway. *Mol. Biol. Cell.* **2007**, 18, 743-754.
 53. Goswami T; Andrews NC. Hereditary hemochromatosis protein; HFE; interaction with transferrin receptor 2 suggests a molecular mechanism for mammalian iron sensing *J. Biol. Chem.* **2006**, 281, 28494-28498.
 54. Valore E.V.; Ganz T. Posttranslational processing of hepcidin in human hepatocytes is mediated by the prohormone convertase furin. *Blood Cells Mol. Dis.* **2008**, 40, 132-138.
 55. Fleming R.E.; Ahmann J.R.; Migas M.C.; Waheed A.; Koeffler H.P.; Kawabata H.; Britton R.S.; Bacon B.R.; Sly W.S. Targeted mutagenesis of the murine transferrin receptor-2 gene produces hemochromatosis. *Proc. Natl. Acad. Sci. USA*, **2002**, 99, 10653-10658.
 56. Wallace D.F.; Summerville L.; Subramaniam V.N. Targeted disruption of the hepatic transferrin receptor 2 gene in mice leads to iron overload. *Gastroenterology* **2007**, 132, 301-310.
 57. Roetto A.; Di Cunto F.; Pellegrino R.M.; Hirsch E.; Azzolino O.; Bondi A.; Defilippi I.; Carturan S.; Miniscalco B.; Riondato F.; Cilloni D.; Silengo L.; Altruda F.; Camaschella C.; Saglio G. Comparison of 3 Tfr2-deficient murine models suggests distinct functions for Tfr2-alpha and Tfr2-beta isoforms in different tissues. *Blood* **2010**, 115, 3382-3389. doi: 10.1182/blood-2009-09-240960.
 58. Fleming R.E.; Feng Q.; Britton RS. Knockout mouse models of iron homeostasis. *Annu. Rev. Nutr.* **2011**, 31, 117-137. doi: 10.1146/annurev-nutr-072610-145117.
 59. Gutschow P.; Schmidt P.J.; Han H.; Ostland V.; Bartnikas T.B.; Pettiglio M.A.; Herrera C.; Butler J.S.; Nemeth E.; Ganz T.; Fleming M.D.; Westerman M. A competitive enzyme-linked immunosorbent assay specific for murine hepcidin-1: correlation with hepatic

- mRNA expression in established and novel models of dysregulated iron homeostasis. *Haematologica* **2015**, 100, 167-177. doi: 10.3324/haematol.2014.116723.
60. Latour C.; Besson-Fournier C.; Meynard D.; Silvestri L.; Gourbeyre O.; Aguilar-Martinez P.; Schmidt P.J.; Fleming M.D.; Roth M.P.; Coppin H. Differing impact of the deletion of hemochromatosis-associated molecules HFE and transferrin receptor-2 on the iron phenotype of mice lacking bone morphogenetic protein 6 or hemojuvelin. *Hepatology* **2016**, 63, 126-137. doi: 10.1002/hep.28254.
61. Nai A.; Pellegrino R.M.; Rausa M.; Pagani A.; Boero M.; Silvestri L.; Saglio G.; Roetto A.; Camaschella C. The erythroid function of transferrin receptor 2 revealed by *Tmprss6* inactivation in different models of transferrin receptor 2 knockout mice. *Haematologica* **2014**, 99, 1016-1021. doi: 10.3324/haematol.2013.103143.
62. Rishi G.; Secondes E.S.; Wallace D.F.; Subramaniam V.N. Hematopoietic deletion of transferrin receptor 2 in mice leads to a block in erythroid differentiation during iron-deficient anemia. *Am. J. Hematol.* **2016**, 91, 812-818. doi: 10.1002/ajh.24417.
63. Marro S.; Chiabrando D.; Messana E.; Stolte J.; Turco E.; Tolosano E.; Muckenthaler M.U. Heme controls ferroportin1 (FPN1) transcription involving *Bach1*; *Nrf2* and a MARE/ARE sequence motif at position -7007 of the FPN1 promoter. *Haematologica* **2010**, 95, 1261-1268. doi: 10.3324/haematol.2009.020123.
64. Chiabrando D.; Fiorito V.; Marro S.; Silengo L.; Altruda F.; Tolosano E. Cell-specific regulation of Ferroportin transcription following experimentally-induced acute anemia in mice. *Blood Cells Mol. Dis.* **2013**, 50, 25-30. doi: 10.1016/j.bcmd.2012.08.002.
65. Zhang D.L.; Hughes R.M.; Ollivierre-Wilson H.; Ghosh M.C.; Rouault T.A. A ferroportin transcript that lacks an iron-responsive element enables duodenal and erythroid precursor cells to evade translational repression. *Cell Metab.* **2009**, 9, 461-473. doi: 10.1016/j.cmet.2009.03.006.
66. Soranzo N.; Spector T.D.; Mangino M.; Kühnel B.; Rendon A.; Teumer A.; Willenborg C.; Wright B.; Chen L.; Li M.; Salo P.; Voight B.F.; Burns P.; Laskowski R.A.; Xue Y.; Menzel S.; Altshuler D.; Bradley J.R.; Bumpstead S.; Burnett M.S.; Devaney J.; Döring A.; Elosua R.; Epstein S.E.; Erber W.; Falchi M.; Garner S.F.; Ghorji M.J.; Goodall A.H.; Gwilliam R.; Hakonarson H.H.; Hall A.S.; Hammond N.; Hengstenberg C.; Illig T.; König I.R.; Knouff C.W.; McPherson R.; Melander O.; Mooser V.; Nauck M.; Nieminen M.S.; O'Donnell C.J.; Peltonen L.; Potter S.C.; Prokisch H.; Rader D.J.; Rice C.M.; Roberts R.; Salomaa V.; Sambrook J.; Schreiber S.; Schunkert H.; Schwartz S.M.; Serbanovic-Canic J.; Sinisalo J.; Siscovick D.S.; Stark K.; Surakka I.; Stephens J.; Thompson J.R.; Völker U.; Völzke H.; Watkins N.A.; Wells G.A.; Wichmann H.E.; Van Heel D.A.; Tyler-Smith C.; Thein S.L.; Kathiresan S.; Perola M.; Reilly M.P.; Stewart A.F.; Erdmann J.; Samani N.J.; Meisinger C.; Greinacher A.; Deloukas P.; Ouwehand W.H.; Gieger C. A genome-wide meta-analysis identifies 22 loci associated with eight hematological parameters in the HaemGen consortium. *Nat. Genet.* **2009**, 41, 1182-1190. doi: 10.1038/ng.467.
67. Auer P.L.; Teumer A.; Schick U.; O'Shaughnessy A.; Lo K.S.; Chami N.; Carlson C.; de Denu S.; Dubé M.P.; Haessler J.; Jackson R.D.; Kooperberg C.; Perreault L.P.; Nauck M.; Peters U.; Rioux J.D.; Schmidt F.; Turcot V.; Völker U.; Völzke H.; Greinacher A.; Hsu L.; Tardif J.C.; Diaz G.A.; Reiner A.P.; Lettre G. Rare and low frequency coding variants in *CXCR2* and other genes are associated with hematological traits. *Nat. Genet.* **2014**, 46, 629-634. doi: 10.1038/ng.2962.
68. Forejtņikovà H.; Vieillevoye M.; Zermati Y.; Lambert M.; Pellegrino R.M.; Guihard S.; Gaudry M.; Camaschella C.; Lacombe C.; Roetto A.; Mayeux P.; Verdier F. Transferrin receptor 2 is a component of the erythropoietin receptor complex and is required for efficient erythropoiesis. *Blood* **2010**, 116, 5357-5367. doi: 10.1182/blood-2010-04-281360.
69. Nai A.; Lidonnici M.R.; Rausa M.; Mandelli G.; Pagani A.; Silvestri L.; Ferrari G.;

- Camaschella C. The second transferrin receptor regulates red blood cell production in mice. *Blood* **2015**, 125, 1170-1179. doi: 10.1182/blood-2014-08-596254.
70. Kautz L., Jung G., Valore E.V., Rivella S., Nemeth E., Ganz T. Identification of erythroid regulator of iron metabolism. *Nat Genet.* **2014**, 46, 678-684. doi: 10.1038/ng.2996.
71. Pellegrino R.M.; Riondato F.; Ferbo L.; Boero M.; Palmieri A.; Osella L.; Pollicino P.; Miniscalco B.; Saglio G.; Roetto A. Altered Erythropoiesis in Mouse Models of Type 3 Hemochromatosis. *Biomed. Res. Int.* **2017**, 2408941. doi: 10.1155/2017/2408941.
72. Xu W.; Barrientos T.; Mao L.; Rockman H.A.; Sauve A.A.; Andrews NC. Lethal cardiomyopathy in mice lacking transferrin receptor in the heart. *Cell Reports* **2015**, 13, 533–545. doi: 10.1016/j.celrep.2015.09.023.
73. Lakhal-Littleton S.; Wolna M.; Carr C.A.; Miller J.J.; Christian H.C.; Ball V.; Santos A.; Diaz R.; Biggs D.; Stillion R.; Holdship P.; Larner F.; Tyler D.J.; Clarke K.; Davies B.; Robbins P.A. Cardiac ferroportin regulates cellular iron homeostasis and is important for cardiac function. *Proc. Natl. Acad. Sci. USA* **2015**, 112: 3164–3169. doi: 10.1073/pnas.1422373112.
74. Lakhal-Littleton S.; Wolna M.; Chung Y.J.; Christian H.C.; Heather L.C.; Brescia M.; Ball V.; Diaz R.; Santos A.; Biggs D.; Clarke K.; Davies B.; Robbins P.A. An essential cell-autonomous role for hepcidin in cardiac iron homeostasis. *Elife* **2016**, 5, e19804. doi: 10.7554/eLife.19804.
75. Gammella E.; Recalcati S.; Rybinska I.; Buratti P.; Cairo G. Iron-induced damage in cardiomyopathy: oxidative-dependent and independent mechanisms. *Oxid. Med. Cell. Longev.* **2015**, 230182. doi: 10.1155/2015/230182. Epub 2015 Mar 24.
76. Boero M.; Pagliaro P.; Tullio F.; Pellegrino R.M.; Palmieri A.; Ferbo L.; Saglio G.; De Gobbi M.; Penna C.; Roetto A. A comparative study of myocardial molecular phenotypes of two Tfr β null mice: role in ischemia/reperfusion. *Biofactors* **2015**; 41(5):360-371. doi: 10.1002/biof.1237.
77. Chevion M.; Leibowitz S.; Aye N.N.; Novogrodsky O.; Singer A.; Avizemer O.; Bulvik B.; Konijn A.M.; Berenshtein E. Heart protection by ischemic preconditioning: a novel pathway initiated by iron and mediated by ferritin. *J. Mol. Cell. Cardiol.* **2008**, 45, 839-845. doi: 10.1016/j.yjmcc.2008.08.011.
78. Aquino D.; Bizzi A.; Grisoli M.; Garavaglia B.; Bruzzone M.G.; Nardocci N.; Savoiaro M.; Chiapparini L. Age-related iron deposition in the basal ganglia: quantitative analysis in healthy subjects. *Radiology* **2009**, 252, 165–172. doi: 10.1148/radiol.2522081399.
79. Pfefferbaum A.; Adalsteinsson E.; Rohlfing T.; Sullivan E.V. (2009) MRI estimates of brain iron concentration in normal aging: comparison of field-dependent (FDRI) and phase (SWI) methods. *Neuroimage* **2009**, 47, 493–500. doi: 10.1016/j.neuroimage.2009.05.006.
80. Carlson E.S.; Fretham S.J.; Unger E.; O'Connor M.; Petryk A.; Schallert T.; Rao R.; Tkac I.; Georgieff M.K. Hippocampus specific iron deficiency alters competition and cooperation between developing memory systems. *J. Neurodev. Disord.* **2010**, 2, 133–143. doi: 10.1007/s11689-010-9049-0.
81. Millichap J.G. Etiologic classification of attention-deficit/hyperactivity disorder. *Pediatrics* **2008**, 12, e358–e365.
82. Benton D. The influence of dietary status on the cognitive performance of children. *Mol. Nutr. Food Res.* **2010**, 54, 457–470. doi: 10.1002/mnfr.200900158.
83. Xu H.; Wang Y.; Song N.; Wang J.; Jiang H.; Xie J. New Progress on the Role of Glia in Iron Metabolism and Iron-Induced Degeneration of Dopamine Neurons in Parkinson's Disease. *Front Mol Neurosci.* **2018**, 10, 455. doi: 10.3389/fnmol.2017.00455.
84. McCarthy R.C.; Kosman D.J. Mechanistic analysis of iron accumulation by endothelial cells of the BBB. *Biometals* **2012**, 25, 665–675. doi: 10.1007/s10534-012-9538-6.
85. Benarroch E.E. Brain iron homeostasis and neurodegenerative disease. *Neurology* **2009**,

- 72:1436–1440. doi: 10.1212/WNL.0b013e3181a26b30
86. Mills E.; Dong X.P.; Wang F.; Xu H. Mechanisms of brain iron transport: insight into neurodegeneration and CNS disorders. *Future Med Chem.* **2010** Jan;2(1):51-64.
 87. Wang J.; Jiang H.; Xie J.X. Ferroportin1 and hephaestin are involved in the nigral iron accumulation of 6-OHDA-lesioned rats. *Eur. J. Neurosci.* **2007**, 25, 2766–2772.
 88. Morris C.M.; Candy J.M.; Keith A.B.; Oakley A.E.; Taylor G.A.; Pullen R.G.; Bloxham C.A.; Gocht A.; Edwardson J.A. Brain iron homeostasis. *J. Inorg. Biochem.* **1992**, 47, 257–265.
 89. Connor J.R.; Boeshore K.L.; Benkovic S.A.; Menzies S.L. Isoforms of ferritin have a specific cellular distribution in the brain. *J. Neurosci. Res.* **1994**, 37, 461–465.
 90. Rouault T.A. Iron metabolism in the CNS: implications for neurodegenerative diseases. *Nat. Rev. Neurosci.* **2013**, 14, 551-564. doi: 10.1038/nrn3453.
 91. Wang S.M.; Fu L.J.; Duan X.L.; Crooks D.R.; Yu P.; Qian Z.M.; Di X.J.; Li J.; Rouault T.A.; Chang Y.Z. Role of hepcidin in murine brain iron metabolism. *Cell Mol. Life Sci.* **2010**, 67, 123-133. doi: 10.1007/s00018-009-0167-3.
 92. Du F.; Qian Z.M.; Luo Q.; Yung W.H.; Ke Y. Hepcidin suppresses brain iron accumulation by downregulating iron transport proteins in iron-overloaded rats. *Mol. Neurobiol.* **2015**, 52, 101-114. doi: 10.1007/s12035-014-8847-x.
 93. Vela D. Hepcidin, an emerging and important player in brain iron homeostasis. *J. Transl. Med.* **2018**, 16, 25. doi: 10.1186/s12967-018-1399-5.
 94. Moos T.; Rosengren Nielsen T.; Skjørringe T.; Morgan E.H. Iron trafficking inside the brain. *J. Neurochem.* **2007**, 103, 1730-1740.
 95. Hänninen M.M.; Haapasalo J.; Haapasalo H.; Fleming R.E.; Britton R.S.; Bacon B.R.; Parkkila S. Expression of iron-related genes in human brain and brain tumors. *BMC Neurosci.* **2009**, 10, 36. doi: 10.1186/1471-2202-10-36.
 96. Acikyoğlu B.; Graham R.M.; Trinder D.; House M.J.; Olynyk J.K.; Scott R.J.; Milward E.A.; Johnstone D.M. Brain transcriptome perturbations in the transferrin receptor 2 mutant mouse support the case for brain changes in iron loading disorders, including effects relating to long-term depression and long-term potentiation. *Neuroscience* **2013**, 235,119-128. doi: 10.1016/j.neuroscience.2013.01.014.
 97. Pellegrino R.M.; Boda E.; Montarolo F.; Boero M.; Mezzanotte M.; Saglio G.; Buffo A.; Roetto A. Transferrin Receptor 2 Dependent Alterations of Brain Iron Metabolism Affect Anxiety Circuits in the Mouse. *Sci Rep.* **2016**, 6, 30725. doi: 10.1038/srep30725.
 98. Adhikari A. Distributed circuits underlying anxiety. *Front Behav Neurosci.* **2014**, 8, 112. doi: 10.3389/fnbeh.2014.00112.
 99. Rhodes S.L.; Buchanan D.D.; Ahmed I.; Taylor K.D.; Lorient M.A.; Sinsheimer J.S.; Bronstein J.M.; Elbaz A.; Mellick G.D.; Rotter J.I.; Ritz B. Pooled analysis of iron-related genes in Parkinson’s disease: association with transferrin. *Neurobiol. Dis.* **2014**, 62,172–178. doi: 10.1016/j.nbd.2013.09.019.
 100. Khalil S.; Holy M.; Grado S.; Fleming R.; Kurita R.; Nakamura Y.; Goldfarb A. A specialized pathway for erythroid iron delivery through lysosomal trafficking of transferrin receptor 2. *Blood Adv.* **2017**, 1, 1181-1194. doi: 10.1182/bloodadvances.2016003772
 101. Robb A.D.; Ericsson M.; Wessling-Resnick M. Transferrin receptor 2 mediates a biphasic pattern of transferrin uptake associated with ligand delivery to multivesicular bodies. *Am J Physiol Cell Physiol.* **2004**, 287, C1769-1775.
 102. Smilevska T.; Stamatopoulos K.; Samara M.; Belessi C.; Tsompanakou A; Paterakis G.; Stavroyianni N.; Athanasiadou I.; Chiotoglou I.; Hadzidimitriou A.; Athanasiadou A.; Douka V.; Saloum R.; Laoutaris N; Anagnostopoulos A.; Fassas A.; Stathakis N.; Kollia P. Transferrin receptor-1 and 2 expression in chronic lymphocytic leukemia. *Leuk Res.* **2006**, 30, 183-189.
 103. Gangat N.; Patnaik M.M.; Tefferi A. Myelodysplastic syndromes: Contemporary review

- and how we treat. *Am. J. Hematol.* **2016**, 91, 76-89. doi: 10.1002/ajh.24253.
104. Di Savino A., Gaidano V., Palmieri A., Crasto F., Volpengo A., Lorenzatti R., Scaravaglio P., Manello A., Nicoli P, Gottardi E., Saglio G., Cilloni D., De Gobbi M. Clinical significance of TFR2 and EPOR expression in bone marrow cells in myelodysplastic syndromes. *Br. J. Haematol.* **2017**, 176, 491-495. doi: 10.1111/bjh.13968.
 105. Nakamaki T.; Kawabata H.; Saito B.; Matsunawa M.; Suzuki J.; Adachi D.; Tomoyasu S.; Phillip Koeffler H. Elevated levels of transferrin receptor 2 mRNA, not transferrin receptor 1 mRNA, are associated with increased survival in acute myeloid leukaemia. *Br. J. Haematol.* **2004**, 125, 42-49.
 106. Calzolari A.; Oliviero I.; Deaglio S.; Mariani G.; Biffoni M.; Sposi N.M.; Malavasi F.; Peschle C.; Testa U. Transferrin receptor 2 is frequently expressed in human cancer cell lines. *Blood Cells Mol Dis.* **2007**, 39, 82-91.
 107. Calzolari A.; Larocca L.M.; Deaglio S.; Finisguerra V.; Boe A.; Raggi C; Ricci-Vitani L.; Pierconti F.; Malavasi F.; De Maria R.; Testa U.; Pallini R. Transferrin receptor 2 is frequently and highly expressed in glioblastomas. *Transl Oncol.* **2010**, 3,123-134.
 108. Crichton R.R.; Dexter D.T.; Ward R.J. Brain iron metabolism and its perturbation in neurological diseases. *J. Neural. Transm. (Vienna).* **2011**, 118, 301-314. doi: 10.1007/s00702-010-0470-z.
 109. Crespo Â.C.; Silva B.; Marques L.; Marcelino E.; Maruta C.; Costa S.; Timóteo A.; Vilares A.; Couto F.S.; Faustino P.; Correia A.P.; Verdelho A.; Porto G; Guerreiro M.; Herrero A.; Costa C.; de Mendonça A.; Costa L.; Martins M. Genetic and biochemical markers in patients with Alzheimer's disease support a concerted systemic iron homeostasis dysregulation. *Neurobiol Aging.* **2014**, 35, 777-785. doi: 10.1016/j.neurobiolaging.2013.10.078.
 110. Artuso I.; Lidonnici M.R.; Altamura S.; Mandelli G.; Pettinato M.; Muckenthaler M.U.; Silvestri L.; Ferrari G.; Camaschella C.; Nai A. Transferrin Receptor 2 is a potential novel therapeutic target for beta-thalassemia: evidence from a murine model. *Blood*, **2018**, pii: blood-2018-05-852277. doi: 10.1182/blood-2018-05-852277.
 111. Girelli D.; Nemeth E.; Swinkels D.W. Heparin in the diagnosis of iron disorders. *Blood* **2016**, 127, 2809-2013. doi: 10.1182/blood-2015-12-639112.
 112. Akinc A., Chan-Daniels A., Sehgal A., Foster D., Bettencourt B.R., Hettinger J., Racie T., Aubin J., Kuchimanchi S., Epstein-Barashand H., Nakayama T. Targeting the hepcidin pathway with RNAi therapeutics for the treatment of anemia. *Blood*, **2011**, 21, 688.
 113. Raha A.A.; Vaishnav R.A.; Friedland R.P.; Bomford A.; Raha-Chowdhury R. The systemic iron-regulatory proteins hepcidin and ferroportin are reduced in the brain in Alzheimer's disease. *Acta Neuropathol. Commun.* **2013**, 1, 55. doi: 10.1186/2051-5960-1-55.
 114. Chen D.; Kanthasamy A.G.; Reddy M.B. EGCG protects against 6-OHDA-induced neurotoxicity in a cell culture model. *Parkinsons Dis.* **2015**, 2015:843906. doi: 10.1155/2015/843906.
 115. Clardy S.L., Wang X., Boyer P.J.; Earley C.J.; Allen R.P.; Connor J.R. Is ferroportin-hepcidin signaling altered in restless legs syndrome? *J Neurol Sci.* **2006**, 247:173–179.
 116. Zheng M.; Tao W.; Zou Y.; Farokhzad O.C.; Shi B. Nanotechnology-Based Strategies for siRNA Brain Delivery for Disease Therapy. *Trends Biotechnol.* **2018**, 36, 562-575. doi: 10.1016/j.tibtech.2018.01.006.
 117. Couch J.A.; Yu Y.J.; Zhang Y.; Tarrant J.M.; Fuji R.N.; Meilandt W.J.; Solanoy H.; Tong R.K.; Hoyte K.; Luk W.; Lu Y.; Gadkar K.; Prabhu S.; Ordonia BA.; Nguyen Q.; Lin Y.; Lin Z.; Balazs M.; Scarce-Lavie K.; Ernst J.A.; Dennis M.S.; Watts R.J. Addressing safety liabilities of TfR bispecific antibodies that cross the blood-brain barrier. *Sci Transl Med.* **2013**, 5, 1-12. doi: 10.1126/scitranslmed.3005338.

- 118.Li L.; Fang C.J.; Ryan J.C.; Niemi E.C.; Lebrón J.A.; Björkman P.J.; Arase H.; Torti F.M.; Torti S.V.; Nakamura M.C.; Seaman W.E. Binding and uptake of H-ferritin are mediated by human transferrin receptor-1. *Proc. Natl. Acad. Sci. U S A* **2010**; 107; 3505-3510. doi: 10.1073/pnas.0913192107.
- 119.Mazzucchelli S.; Truffi M.; Baccarini F.; Beretta M.; Sorrentino L.; Bellini M.; Rizzuto M.A.; Ottria R.; Ravelli A.; Ciuffreda P.; Prosperi D.; Corsi F. H-Ferritin-nanocaged olaparib: a promising choice for both BRCA-mutated and sporadic triple negative breast cancer. *Sci. Rep.* **2017**; 7; 7505. doi: 10.1038/s41598-017-07617-7.
- 120.Sakurai K.; Sohda T.; Ueda S.; Tanaka T.; Hirano G.; Yokoyama K.; Morihara D.; Aanan A.; Takeyama Y.; Irie M.; Iwata K.; Syakado S.; Noritomiz T.; Yamashita Y.; Sakisaka S. Immunohistochemical demonstration of transferrin receptor 1 and 2 in human hepatocellular carcinoma tissue. *Hepatogastroenterology* **2014**; 61; 426-430.
- 121.Voth B.; Nagasawa D.T.; Pelargos P.E.; Chung L.K.; Ung N.; Gopen Q.; Tenn S.; Kamei D.T.; Yang I. Transferrin receptors and glioblastoma multiforme: Current findings and potential for treatment. *J. Clin. Neurosci.* **2015**; 22; 1071-1076. doi: 10.1016/j.jocn.2015.02.002.
- 122.Murphy M.P. Smith R.A. Drug delivery to mitochondria: the key to mitochondrial medicine. *Adv. Drug Deliv. Rev.* **2000**; 41; 235-250.
- 123.Bürk K. Friedreich Ataxia: current status and future prospects. *Cerebellum Ataxias* **2017**; 4. doi: 10.1186/s40673-017-0062-x.