

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Natural iron ligands promote a metal-based oxidation mechanism for the Fenton reaction in water environments

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1768966> since 2021-02-04T17:53:54Z

Published version:

DOI:10.1016/j.jhazmat.2020.122413

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

SUPPLEMENTARY MATERIAL

Natural Iron Ligands Promote a Metal-Based Oxidation Mechanism for the Fenton Reaction in Water Environments

Giulio Farinelli,[†] Marco Minella,[‡] Marco Pazzi,[‡] Stefanos Giannakis,[§] Cesar
Pulgarin,^{||} Davide Vione,^{*,‡} Alberto Tiraferri^{*,†}

[†]Department of Environment, Land and Infrastructure Engineering (DIATI), Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129, Turin, Italy

[‡]Department of Chemistry, University of Turin, Via Pietro Giuria 7, 10125 Turin, Italy

[§]Universidad Politécnica de Madrid, E.T.S. Ingenieros de Caminos, Canales y Puertos, Departamento de Ingeniería Civil: Hidráulica, Energía y Medio Ambiente, Unidad docente Ingeniería Sanitaria, c/ Profesor Aranguren, s/n, ES-28040 Madrid, España

^{||}SB, ISIC, Group of Advanced Oxidation Processes, Ecole Polytechnique Fédérale de Lausanne (EPFL), Station 6, 1015 Lausanne, Switzerland

Corresponding Authors

Email: alberto.tiraferri@polito.it; davide.vione@unito.it




Figure S1. Fe-TAML molecular structure

Table S1. Oxidation of cyclohexane at different pH values with citric acid as iron ligand

pH	A (M)	K (M)	A/K
3	2.77×10^{-9}	2.75×10^{-10}	10.07
5	3.94×10^{-9}	5.37×10^{-10}	7.35
6	3.24×10^{-9}	3.82×10^{-10}	8.48
7	5.31×10^{-9}	6.16×10^{-10}	8.62

Reactants: cyclohexane (0.1 mM), hydrogen peroxide (0.001 mM), iron(II) (0.1 μ M).

Table S2. Oxidation of cyclohexane at different pH values with malic acid as iron ligand

pH	A (M)	K (M)	A/K
3	1.82×10^{-9}	1.02×10^{-10}	17.79
5	1.65×10^{-9}	1.14×10^{-10}	14.48
6	4.56×10^{-9}	2.89×10^{-10}	15.81
7	3.47×10^{-9}	2.26×10^{-10}	15.38

Reactants: cyclohexane (0.1 mM), hydrogen peroxide (0.001 mM), iron(II) (0.1 μ M).

Table S3. Oxidation of cyclohexane at different pH values with quinic acid as iron ligand

pH	A (M)	K (M)	A/K
3	5.54×10^{-9}	1.97×10^{-10}	28.14
5	5.90×10^{-9}	1.88×10^{-10}	31.29
6	5.02×10^{-9}	2.17×10^{-10}	23.15
7	2.90×10^{-9}	1.05×10^{-10}	27.70

Reactants: cyclohexane (0.1 mM), hydrogen peroxide (0.001 mM), iron(II) (0.1 μ M).

Degradation experiments were carried out in magnetically stirred beakers (the total volume of the aqueous phase was 50 mL). The reaction mixture contained phenol, Fe^{2+} (as FeSO_4) or FeIII-TAML, H_2O_2 , a reagent for pH adjustment (HClO_4 for pH 3, NaOH for pH 10) and, where relevant, t-butanol (TBA). At scheduled time intervals, a measured 2 mL sample aliquot was withdrawn from the reaction mixture and diluted 1:1 with the same volume of methanol to stop the Fenton reaction. Phenol was quantified by high performance liquid chromatography (HPLC), using a VWR-Hitachi Elite LaChrom instrument equipped with L2455 diode array detector (DAD), L2130 quaternary pump module, L2300 column oven (set at 40 °C), L2200 autosampler (sample injection volume 60 μL), Duratec vacuum degasser and reverse-phase column Merck LiChroCART, packed with LiChrospher 100 RP18 (125 mm \times 4 mm \times 5 μm). The chromatographic elution conditions were A:B = 70:30, where A is an aqueous solution of H_3PO_4 (pH 2.8) and B is methanol, at 1.0 mL/min flow rate. These conditions gave a column dead time of 1.2 min and a phenol retention time of 5.2 min. The detection wavelength was 210 nm.

Considering that the Fenton reaction can produce both $\bullet\text{OH}$ and ferryl (FeIV) species, and because TBA is a selective $\bullet\text{OH}$ scavenger, the goal of this series of experiments was to get insight into the relative roles of $\bullet\text{OH}$ vs. ferryl in the Fenton degradation of phenol. The experimental results are reported in Figure S2 for $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ at pH 3 (S2a) and Fe-TAML + H_2O_2 at pH 10 (S2b). In the latter case, it is interesting to observe that TBA had practically no effect on phenol degradation, which is consistent with the almost exclusive involvement of ferryl in the process. The opposite result was obtained with $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ at pH 3, where the large inhibition effect carried out by TBA suggests that $\bullet\text{OH}$ plays a key role in the traditional Fenton process. Therefore, one can conclude that $\bullet\text{OH}$ is largely involved in phenol degradation by $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ at pH 3, while ferryl operates in the presence of Fe-TAML + H_2O_2 at pH 10.




Figure S2. (a) Time trends of 0.1 mM phenol in the presence of 0.1 mM Fe²⁺ and 1 mM H₂O₂, at pH 3 by HClO₄, with and without 4 mM TBA. (b) Time trends of 0.1 mM phenol in the presence of 0.01 mM Fe-TAML and 0.1 mM H₂O₂, at pH 10 by NaOH, with and without 4 mM TBA.




Figure S3. Conditional binding constant computed at pH 7 and pH 3 for a variety of ligands.

The conditional binding constants for the Fe(II) complexes $[\text{FeL}]^{(n-2)-}$ were computed as

$$K'_f = \alpha_n K_f$$

where:

K_f = binding constant for the ligand to metal complex with a 1 to 1 stoichiometry

α_n = molar fraction of the totally deprotonated ligand species computed as

$$\alpha_n = \frac{\prod_1^n K_i}{[H^+]^n + K_1[H^+]^{n-1} + K_1K_2[H^+]^{n-2} + \dots + \prod_1^n K_i}$$

where

K_i = acid dissociation constant for the H_nL ligand

All the equilibrium constants were extracted from:

Arthur E. Martell and Robert M. Smith, NIST Standard Reference Database 46 Version 4.0, NIST critically selected constants of metal complexes, NIST Standard Reference Data, Gaithersburg, MD 20899 USA, 1997.

Table S4. Oxidation of cyclohexane with EDTA as iron ligand added at different concentrations, performed at pH 3 unless otherwise stated

[EDTA]/[Fe(II)]	A (M)	K (M)	A/K
0	4.86×10^{-10}	4.25×10^{-10}	1.14
0.25	7.32×10^{-10}	3.48×10^{-10}	2.10
0.375	1.24×10^{-9}	3.45×10^{-10}	3.60
0.5	1.29×10^{-8}	5.41×10^{-10}	23.80
0.75	9.87×10^{-9}	3.57×10^{-10}	27.61
1	9.10×10^{-9}	3.93×10^{-10}	23.13
1 (pH 7)	4.60×10^{-9}	1.20×10^{-10}	38.32

Reactants: cyclohexane (0.1 mM), hydrogen peroxide (0.001 mM), iron(II) (0.1 μ M)

Table S5. Oxidation of cyclohexane with EDDS as iron ligand added at different concentrations, performed at pH 3 unless otherwise stated

[EDDS]/[Fe(II)]	A (M)	K (M)	A/K
0	4.86×10^{-10}	4.25×10^{-10}	1.14
0.25	7.88×10^{-10}	8.19×10^{-10}	0.96
0.375	8.76×10^{-10}	1.20×10^{-9}	0.73
0.5	1.14×10^{-9}	6.68×10^{-10}	1.71
0.75	5.81×10^{-10}	2.79×10^{-10}	2.08
1	6.41×10^{-10}	2.53×10^{-10}	2.53
1 (pH 7)	2.94×10^{-9}	6.41×10^{-11}	45.78

Reactants: cyclohexane (0.1 mM), hydrogen peroxide (0.001 mM), iron(II) (0.1 μ M)