

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Redox-driven changes in organic C stabilization and Fe mineral transformations in temperate hydromorphic soils**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1814252> since 2022-03-18T16:27:27Z

*Published version:*

DOI:10.1016/j.geoderma.2021.115532

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Redox-driven changes in organic C stabilization and Fe**  
2 **mineral transformations in temperate hydromorphic soils**

3

4 Beatrice Giannetta<sup>a,b</sup>, Danilo Oliveira de Souza<sup>c</sup>, Giuliana Aquilanti<sup>c</sup>, Luisella Celi<sup>a</sup>,  
5 Daniel Said-Pullicino<sup>a</sup>

6

7 *<sup>a</sup>Department of Agricultural, Forest and Food Sciences, University of Torino, Largo*  
8 *Paolo Braccini 2, 10095 Grugliasco, Italy*

9 *<sup>b</sup>Department of Biotechnology, University of Verona, Strada Le Grazie 15, 37134*  
10 *Verona, Italy (Present address)*

11 *<sup>c</sup>ELETTRA Sincrotrone Trieste S.C.p.A., S.S. 14 Km 163.5, 34149 Basovizza, Trieste, Italy*

12

13

14

15 Corresponding author. E-mail address: [beatrice.giannetta@univr.it](mailto:beatrice.giannetta@univr.it)

16

17 **Abstract**

18 Paddy soils experience long-term redox alternations affecting the interactions between  
19 the biogeochemical cycling of iron (Fe) and carbon (C). Differences in particle  
20 aggregation and soil organic matter (SOM) turnover are likely to both affect and be  
21 affected by the trajectory of Fe mineral evolution/crystallinity with redox fluctuations.  
22 We hypothesized that the legacy effects of redox cycling under paddy management  
23 affects particle aggregation, the distribution and mineralogy of Fe (hydr)oxides between  
24 particle-size fractions, and the interaction with SOM stabilization. Moreover, we  
25 expected underlying processes to be different in paddy eluvial and illuvial horizons,  
26 particularly due to the different inputs and redox conditions these horizons experience.  
27 To test these hypotheses, we evaluated the distribution of Fe species and organic C  
28 between different aggregate and particle-size fractions in topsoil (eluvial) and subsoil  
29 (illuvial) horizons of soils under long-term paddy and non-paddy management in NW  
30 Italy, as well as mineralogical changes in Fe phases by Fe K-edge Extended X-ray  
31 Adsorption Fine Structure (EXAFS) and X-ray Absorption Near Edge Structure  
32 (XANES) spectroscopy. Our findings indicate that although paddy topsoils are depleted  
33 in hydrous Fe oxides with respect to non-paddy soils, they can stabilize important  
34 amounts of C through mineral associations, particularly with finer particle-size fractions  
35 rich in less crystalline Fe phases. We also show that redox cycling can influence  
36 microaggregate stability and consequently the distribution of Fe phases and OC between  
37 intra and inter-microaggregate fractions. On the other hand, illuvial horizons under paddy  
38 management were enriched in short-range ordered hydrous Fe oxides and this contributed  
39 to enhanced microaggregate formation and C stabilization with respect to non-paddy  
40 subsoils.

41

42 **Keywords:** paddy soils, eluvial and illuvial horizons, physical fractionation, EXAFS,  
43 XANES, linear combination fitting.

## 44 1. INTRODUCTION

45 Paddy soils make up the largest anthropogenic wetlands on earth and store a large  
46 proportion of the global terrestrial carbon (C) stocks, representing a major terrestrial C  
47 pool, (ca.10 Pg) (Kirk, 2004). The greater soil organic C (SOC) accumulation rates in  
48 paddy topsoils with respect to non-paddy ones are generally assumed to be due to limited  
49 mineralization under anoxic soil conditions resulting from frequent field flooding (Cheng  
50 et al., 2009; Kalbitz et al., 2013; Kögel-Knabner et al., 2010; Pan et al., 2004). Although  
51 numerous studies attribute organic matter (OM) accumulation in paddy soils to this  
52 mechanism, there is growing evidence questioning this assumption (Huang and Hall,  
53 2017).

54 Organic matter turnover and stabilization in paddy soils are strongly linked to the  
55 dynamic interactions with redox-active minerals, in particular iron (Fe) oxyhydroxides  
56 (Huang and Hall, 2017; Hall et al., 2018; Huang et al., 2020). Short-range-ordered (SRO)  
57 minerals (e.g., ferrihydrite) are able to stabilize more OC than well-ordered minerals (e.g.,  
58 goethite, hematite and lepidocrocite) due to their larger surface area, and thus play a more  
59 important role in soil C storage mechanisms (Kramer and Chadwick, 2018; Lalonde et  
60 al., 2012; Torn et al., 1997). The role of SRO-Fe oxy-hydroxides in the preservation of  
61 SOC in paddy soils and OC persistence in deeper mineral horizons have been well-  
62 documented (Chorover et al., 2004; Huang and Hall, 2017; Kramer et al., 2012; Lalonde  
63 et al., 2012; Said-Pullicino et al., 2021; Wissing et al., 2014, 2013). However, further  
64 insights into the effects of redox alternations on Fe phase composition, crystal order and  
65 depth distribution, as well as the link between microaggregate formation and  
66 (de)stabilization, and OC stabilization mechanisms are necessary. Overall, depending on  
67 soil hydrology, paddy cultivation generally leads to a depletion in pedogenic Fe oxides in  
68 the topsoil due to their reductive dissolution and Fe leaching under reducing conditions,  
69 that is often accompanied by a relative increase in the proportion of SRO Fe oxides (Kölbl  
70 et al., 2014, Winkler et al., 2016). On the other hand, paddy subsoils are generally  
71 characterized by Fe oxide accumulation (Kölbl et al., 2014). In addition, redox  
72 fluctuations can potentially increase or decrease Fe crystallinity, depending on the  
73 environmental context (Coby et al., 2011; Ginn et al., 2017; Riedel et al., 2013; Thompson  
74 et al., 2006).

75 In upland soils SRO-Fe oxides can also promote aggregation (Oades and Waters,  
76 1991, Xue et al., 2019), which indirectly enhances OM stability (Balesdent et al., 2000;  
77 Totsche et al., 2018). However, redox fluctuations are expected to influence aggregate

78 formation and stability with implications on the physical and chemical processes  
79 regulating OM stabilization. Selective microbial reduction and reprecipitation of Fe  
80 phases due to repeated redox cycles could therefore lead to a change in their distribution  
81 between different aggregate size classes. Recent studies have shown that a portion of Fe  
82 phases in soil can be protected from reductive dissolution (e.g., dithionite extraction)  
83 presumably due to their physical protection within microaggregates or co-precipitation  
84 with SRO aluminosilicates (Coward et al., 2018; Filimonova et al., 2016), suggesting the  
85 importance of aggregation/precipitation reactions. On the other hand, several studies  
86 showed that the reductive dissolution of Fe oxides involved in aggregate binding can  
87 disrupt aggregate stability (Huang et al., 2018; Henderson et al., 2012; De-Campos et al.,  
88 2009). Detailed chemical and size related characterization of soil aggregates and their  
89 depth distribution in rice paddies are scarce, even though such information is fundamental  
90 for further understanding the mechanisms controlling the biogeochemical cycling of  
91 elements in these agro-ecosystems.

92 The objectives of this work were to compare Fe mineral evolution and OC cycling in  
93 the topsoil and subsoil of a temperate paddy soil and an adjacent upland soil. We  
94 hypothesized that (1) in paddy topsoils, Fe redox cycling can lead to a depletion of  
95 pedogenetic Fe (hydr)oxides, promote microaggregate breakdown, and cause a  
96 redistribution of Fe phases with a relative increase in the proportion of SRO oxides and  
97 OC associated with the finest “free” particle-size fractions; and (2) the enhanced inputs  
98 of dissolved/colloidal Fe and OC entering the paddy subsoil through percolation can  
99 favour the accumulation of SRO Fe phases that contribute to the formation of stable  
100 microaggregates and C retention in illuvial horizons. To test these hypotheses, we  
101 evaluated the effects of redox cycling on the mineralogical changes and  
102 distribution/redistribution of Fe species (i.e., Fe in clays, SRO Fe(III) oxides such as  
103 ferrihydrite and organic Fe(III) complexes or more well-ordered Fe oxides), and OC  
104 between different intra- and inter-microaggregate particle-size fractions in topsoil  
105 (eluvial) and subsoil (illuvial) horizons of soils under long-term paddy (P) and non-paddy  
106 (NP) management in NW Italy. In order to study Fe transformations a combination of Fe  
107 K-edge Extended X-ray Adsorption Fine Structure (EXAFS) and X-ray Absorption Near  
108 Edge Structure (XANES) spectroscopy have been applied.

109

## 110 **2. MATERIALS AND METHODS**

### 111 **2.1 Site description and sample collection**

112 The study site is located in Zeme (Province of Pavia, NW Italy) (45°11'31.8"N;  
113 8°40'3.7"E), in the plains of the River Po, between the Sesia and Agogna rivers and has a  
114 temperate climate, with a mean annual temperature of 12.5°C and mean annual  
115 precipitation of 950 mm. Soils develop on alluvial sediment with a maximum elevation  
116 of 80 m a.s.l., characterized by a high groundwater level during spring, up to 30-50 cm  
117 below surface. Two adjacent plots were identified for this study, one under maize (*Zea*  
118 *mays*) monocropping (NP, non-paddy) (45°11'33.6"N 8°40'06.1"E; 1.35 ha; Endogleyic  
119 Fluvisol) and another under paddy rice (*Oryza sativa*) monocropping (P, paddy  
120 management) established about 30 years ago (45°11'29.4"N 8°40'04.5"E; 4.43 ha; Haplic  
121 Gleysol). These non-calcareous soils generally had an acidic pH, low contents of OC and  
122 pedogenic Fe, and a sandy loam texture. A full description of the soil profiles and basic  
123 properties of the soil horizons under non-paddy and paddy management, as well as  
124 agricultural practices adopted are reported in Said-Pullicino et al. (2021). In March 2018,  
125 towards the end of the winter fallow period, topsoil and subsoil samples were collected  
126 from the Ap1, Ap2 and Bgw horizons in the non-paddy field and from the Arp1, Arp2,  
127 Arpd, Brd1 and Brd2 horizons in the paddy field from three replicated soil pits in each  
128 plot, and subsequently air dried and homogenized by passing through a 2-mm sieve. In  
129 this context, differences in analyzed soil properties under these contrasting cropping  
130 systems were attributed to the legacy of past management (non-paddy vs. paddy), and as  
131 such, were considered to be influenced by the long-term effects of different water  
132 management during the cropping season (irrigated vs. flooded) on soil redox conditions  
133 (rather than the specific redox conditions at the time of sampling). Moreover, this  
134 approach also integrated other effects, such as the slightly different quantity and quality  
135 of annual OM inputs with the incorporation of maize or rice crop residues, that were  
136 considered secondary and therefore not discussed in the present work.

137

## 138 **2.2 Microaggregate separation and particle-size fractionation**

139 The fractionation scheme adopted was intended to separate particle-size fractions  
140 distinguishing between those particles that constituted part of the stable microaggregates  
141 (intra-microaggregate particles size fractions) and particles of similar dimensions not  
142 involved in microaggregation (inter-microaggregate particle size fractions). This  
143 fractionation was carried out in three principal steps as illustrated in Figure S1:

144 (i) Soil microaggregates (53-200 µm) were first separated from all soil samples by  
145 means of a custom-built, mechanical wet sieving microaggregate isolator similar to

146 the Yoder sieve apparatus used for measuring aggregate stability, as follows.  
147 Aliquots of bulk soil (< 2 mm; 24 g) were transferred into 200  $\mu\text{m}$  bucket sieves  
148 and immersed in large beakers filled with deionized water. Rotation of the buckets  
149 for 60 min in the presence of six metal beads was sufficient to break all  
150 macroaggregates. The coarse sand and particulate OM fraction (CSa+POM)  
151 remaining in the buckets was collected while the microaggregates and fine sand  
152 (53-200  $\mu\text{m}$ ) were separated from the suspension by passing through a 53  $\mu\text{m}$  sieve.  
153 (ii) The microaggregates and fine sand were resuspended in 200 ml water to which 15  
154 drops of 1 M NaCl were added, rinsed until the supernatant had an  $\text{EC} \leq 100 \mu\text{S}$   
155  $\text{cm}^{-1}$ , and subsequently sonicated with 400  $\text{J ml}^{-1}$  to break microaggregates and  
156 ensure complete clay dispersion. Particle-size fractionation of the intra-  
157 microaggregate soil particles was carried out by sedimentation based on Stokes law  
158 to obtain coarse silt (20-53  $\mu\text{m}$ ; mCSi), fine silt (2-20  $\mu\text{m}$ ; mFSi) and clay (<2  $\mu\text{m}$ ;  
159 mCl) fractions within microaggregates. Finally, the fine sand fraction (FSa, 53-200  
160  $\mu\text{m}$ ) was separated by passing through a 53  $\mu\text{m}$  sieve.  
161 (iii) The suspension remaining from (i) above, containing the inter-microaggregate and  
162 free soil particles (< 53  $\mu\text{m}$ ) was concentrated by repeated centrifugation (10,000  
163 rpm) and supernatant siphoning, after addition of 2 ml of 1 M NaCl, taking care not  
164 to lose any fine mineral particles. The sediment was thereafter rinsed (until the  
165 supernatant had an  $\text{EC} \leq 100 \mu\text{S cm}^{-1}$ ), resuspended in 200 ml deionized water, and  
166 subsequently sonicated with 400  $\text{J ml}^{-1}$  to ensure complete clay dispersion. Particle-  
167 size fractionation was carried out by sedimentation as described in (ii) above to  
168 obtain coarse silt (CSi), fine silt (FSi) and clay (Cl) fractions between  
169 microaggregates.  
170 Each fraction was transferred into a pre-weighted container, freeze-dried, weighed and  
171 subsequently ground to < 0.5 mm for subsequent analysis.

172

### 173 **2.3 OC and Fe contents characterization**

174 Total OC content of the eight size fractions was determined by high temperature  
175 combustion using an elemental analyzer (NA 2100, Carlo Erba, Milan, Italy). Poorly  
176 crystalline Fe ( $\text{Fe}_o$ ) was determined by extraction with 0.2 M ammonium oxalate at pH 3  
177 as described by Schwertmann (1964), while more ordered Fe oxides ( $\text{Fe}_d$ ) were  
178 determined by extraction with 0.3 M citrate-bicarbonate solution in the presence of  
179 sodium dithionite as described by Mehra and Jackson (1960). Metal concentrations in the

180 extracts were determined by atomic absorption spectroscopy (AAAnalyst 400, Perkin  
181 Elmer). The ratio of the  $Fe_o/Fe_d$ , indicative of the contribution of the short-range ordered  
182 phases to total pedogenic oxides, or as the metric of Fe phase crystallinity, as well as their  
183 difference to estimate the content of more crystalline oxides ( $Fe_c$ ), were calculated.

184

#### 185 **2.4 Fe K-edge XANES and EXAFS spectroscopy**

186 X-ray absorption spectroscopic (XAS) analyses of FSa, CSi, mCSi, FSi, mFSi, Cl  
187 and mCl fractions were carried out at the XAFS beamline at Elettra Sincrotrone (Trieste,  
188 Italy) (Aquilanti et al., 2017; Di Cicco et al., 2009). The samples were prepared as pressed  
189 powders (*ca.* 15 mg) between Kapton® tape. Fe spectra were recorded in transmission  
190 mode using Si (111) monochromator, which was calibrated to the first-derivative  
191 maximum of the K-edge absorption spectrum of a metallic Fe foil (7112 eV). Spectra  
192 were acquired over an energy range between 6812-7660 eV with a variable step size (5  
193 eV on the pre-edge region, 0.2 eV on the XANES region and a k constant step after  
194 XANES equal to  $0.03 \text{ \AA}^{-1}$ ) and integration time of 2 sec per point. The monochromator  
195 was detuned to exclude higher order harmonics. Two to six scans were collected and  
196 averaged to increase the signal-to-noise ratio. Prior to sample analysis, radiation damage  
197 was evaluated by recording a series of scans for a test sample over a duration similar to  
198 that required for actual spectra acquisition, and then comparing all XANES and EXAFS  
199 features of this sequence. We found that for the analytical conditions applied no photo-  
200 reduction due to the beam was observed. It is worth to note that a remarkable feature of  
201 XAFS beamline at Elettra is the unfocused beam, which makes the photon flux evenly  
202 spread over a large area of the sample minimizing (and in our case, preventing) any beam  
203 damage.

204 As previously described in Giannetta et al. (2020a, b), a suite of linear combination fitting  
205 (LCF) techniques designed to identify the major Fe components in the studied samples  
206 by comparison with the spectra of well characterized standards, was employed to analyze  
207 both the XANES and EXAFS data. Additional methods of analysis, such as Principal  
208 Component Analysis (PCA) (Table S1) and Target transformation, were used as a  
209 complementary tool during linear combination analysis.

210 XANES pre-edge peak analysis at the Fe K-edge was used to identify differences in the  
211 relative oxidation state between samples. Although redox conditions during sample  
212 collection and handling are known to affect Fe speciation (Prietz et al., 2009), we  
213 attributed differences in Fe oxidation states to the legacy effects of long-term soil



214 management, rather than to the specific influence of reducing soil conditions during rice  
215 cropping. The energy position, intensity, and shape of the XANES are governed by the  
216 oxidation state and the specific bonding environment of the irradiated Fe atoms  
217 (coordination type, bonding symmetry, length to neighboring atoms (Westre et al., 1997;  
218 Wilke et al., 2001). Particularly, the pre-edge peak centroid energy position (PCE) allows  
219 a comparative evaluation of the oxidation state of Fe with values ranging between  
220 7113.20 eV for Fe(II) and 7114.55 eV for Fe(III), and can be used for the quantification  
221 of Fe(II)/Fe(III) ratios of different Fe bearing minerals in soils with reasonable accuracy.  
222 According to Wilke et al. (2001) a linear relationship between the PCE and Fe(II)/Fe(III)  
223 ratio can be assumed, if the symmetry around Fe atoms remains the same for all samples.  
224 The pre-edge peak model and centroid determination was based on the considerations  
225 found on Wilke et al. (2001). Two pseudo-Voigt functions were considered as a  
226 component and their energy position, intensity and width were determined after the  
227 baseline extraction, which serves to remove the main absorption edge. Accordingly, it  
228 was assessed considering positions of individual components weighted by its respective  
229 intensity after the fit. A set of standard samples were also analyzed in the same way in  
230 order to get a reference value for Fe(II) and Fe(III) energy positions, respectively, 7113.20  
231 and 7114.55 eV. Siderite was used for Fe(II) whereas the average values of ferrihydrite,  
232 goethite, hematite were used as Fe(III) standards. The choice of these particular minerals  
233 was based on their probable presence in our samples according LCF results. The relative  
234 contribution of specific Fe compounds or compound classes to soils was also assessed by  
235 LCF performed on the entire XANES spectrum. Quantitative LCF was used for the  
236 speciation of different organic and inorganic Fe-bearing compounds. LCF on all  
237 normalized spectra was performed with the spectra of illite, smectite, nontronite,  
238 ferrihydrite, goethite, hematite, lepidocrocite, magnetite, maghemite, smectite, pyrite,  
239 purpurite, and Fe(III) citrate. The energy range for the fitting was 7105-7145 eV and the  
240 number of components to be include in a fit was determined after PCA on each group of  
241 samples (from the same horizon). Eventually, target transformations were used as a  
242 further parameter in the choice of the LCF result.

243 LCF analyses of  $k^3$ -weighted Fe and EXAFS spectra were performed over a  $k$  range  
244 of 2–10  $\text{\AA}^{-1}$ . EXAFS data processing (e.g., background removal, normalization and  
245 deglitching) and spectra generation were carried out with the software Athena. The  
246 qualitative speciation of Fe in the bulk soils and size-fractions by Fe K-edge EXAFS was  
247 carried out by comparison with several standards. Fe(III)-citrate is generally used as an

248 analog model compound for Fe(III)-OM complexes because it consists of Fe bound  
249 directly to OC (O'Day et al., 2004). We assumed that Fe EXAFS can be more suitable to  
250 quantify specific Fe oxyhydroxides (e.g. ferrihydrite, goethite, and hematite) and distinct  
251 Fe-organic compounds. Fe K-edge XANES can better estimate the relative contribution  
252 of different mineral classes and groups of organic compounds with different oxidation  
253 states of the Fe atom (e.g. Fe sulphides, Fe oxyhydroxides, Fe(II) and Fe(III)-organic  
254 complexes in mineral mixtures and soil samples.

255 The qualitative analysis of the nature of backscattering atoms in higher coordination  
256 shells was conducted using the Morlet wavelet transform (WT) method as implemented  
257 in the Igor Pro developed by (Funke et al., 2005). This method complements the  
258 conventional Fourier transform (FT) analysis and reveals the energies where back  
259 scattering takes place that give rise to the FT peaks. The WT modulus was analyzed and  
260 compared with the WT modulus for reference samples with contribution from C and/or  
261 Fe backscattering in the second coordination shell.

262

### 263 **3. RESULTS**

#### 264 **3.1 OC distribution in the different aggregate and particle-size fractions**

265 Across all soil samples 40-60 % of total OC was associated with the finest particle-size  
266 fractions (<2  $\mu\text{m}$ ; Cl + mCl), while 20-40 % of total OC was associated with the coarser  
267 fractions and particulate OM (>53  $\mu\text{m}$ ; FSa + CSa+POM). Arp1 and Arp2 topsoil  
268 horizons of the P soil generally showed higher OC contents across most particle-size  
269 fractions with respect to the corresponding Ap1 and Ap2 horizons of the NP soil (Figure  
270 1), in line with the higher total SOC contents of topsoils under paddy management.  
271 However, the most significant differences were observed for the 2-20  $\mu\text{m}$  (FSi + mFSi)  
272 and >53  $\mu\text{m}$  (FSa + CSa+POM) fractions in the more superficial horizon, that were on  
273 average 2.1 and 1.4-fold greater under P compared to NP management, respectively. OC  
274 associated with the finest particle fractions (Cl + mCl) that represented the greatest  
275 proportion of SOC, were only marginally though significantly higher in the P with respect  
276 to the NP soil in the first 15 cm of soil depth.

277 Paddy management also influenced the distribution of OC between intra- and inter-  
278 microaggregate particle-size fractions in the topsoil. Whereas P management resulted in  
279 a relatively higher intra:inter-microaggregate OC ratio for the fine silt-sized fraction (2-  
280 20  $\mu\text{m}$ ), lower ratios were generally observed for the clay-sized fraction with respect to  
281 NP soils (Figure 2a). In fact, on average 82 % of silt-associated OC in the P topsoils was

282 retained within microaggregates (c.f. 70 % in NP topsoils), while only 58 % of the total  
283 OC associated with clay-sized particles was intra-microaggregate OC with respect to 72  
284 % in the NP soil.

285 Subsoil horizons under P management were particularly enriched in OC associated  
286 with the finest soil fractions (mCl + Cl; ~2.7-fold), with respect to the corresponding  
287 horizons in the NP soil (Figure 1). Moreover, in the paddy illuvial Brd1 horizon, up to 60  
288 % of the clay-sized OC fraction was associated with stable microaggregates, compared to  
289 only 40 % in the Bgw horizon of the NP soil (mCl/Cl OC = 2.6 and 0.8, respectively).  
290 Although significantly less OC was associated with the silt-sized compared to the clay-  
291 sized particles in P subsoils (Figure 1), this former fraction showed the highest intra:inter-  
292 microaggregate OC ratio (Figure 2a).

293

### 294 **3.2 Fe distribution in the different aggregate and particle-size fractions**

295 Most of the pedogenic Fe in NP topsoils (53-58 %) was associated with the finest soil  
296 particles (<2  $\mu\text{m}$ ), primarily with intra-microaggregate clay particles (Figure 3a). Only  
297 around 25% of this Fe was oxalate-extractable as most Fe was in the form of more  
298 crystalline phases (Figures 3c and e). A substantial proportion of pedogenic Fe in these  
299 topsoils (around 28 %) was also associated with the coarser sand-sized fractions  
300 (CSa+POM and FSa) with an average  $\text{Fe}_o/\text{Fe}_d$  ratio of 0.5 (Table 1). In contrast, P topsoils  
301 were greatly depleted in pedogenic Fe showing around 70% less Fe associated with finer  
302 clay-sized and coarser sand-sized fractions with respect to the corresponding horizons in  
303 the NP soils, leading to a more even distribution of pedogenic Fe across particle-size  
304 fractions (Figure 3b). However, these observed differences in total pedogenic Fe contents  
305 did not seem to greatly influence the distribution of Fe between inter and intra-  
306 microaggregate Cl fractions, while a relative increase in intra-microaggregate  $\text{Fe}_d$  was  
307 observed for the FSi fraction (Figure 2b). On the other hand, the distribution between  
308 more and less crystalline phases was influenced by management as clay and sand-sized  
309 fractions showed a greater proportion of oxalate-extractable Fe and higher  $\text{Fe}_o/\text{Fe}_d$  ratios  
310 (up to 0.8) with respect to NP topsoils (Table 1).

311 Subsoil horizons in P and NP profiles showed evident management-related differences in  
312 the distribution of pedogenic Fe between aggregate and particle-size fractions. Whereas,  
313 NP subsoil samples from the Bgw horizon showed lower total pedogenic Fe contents with  
314 respect to the overlying Ap soil horizons (Fig. 3a) albeit very similar intra:inter-  
315 microaggregate  $\text{Fe}_d$  (Figure 2b) and  $\text{Fe}_o/\text{Fe}_d$  ratios (Table 1) across fractions, P subsoils

316 showed higher total pedogenic Fe contents and oxalate-extractable Fe with respect to both  
317 P topsoils (Arp1 and Arp2 horizons) and the corresponding subsoil horizon in the NP soil  
318 (Bgw). The most enriched Brd1 illuvial horizon in the P soil had pedogenic Fe contents  
319 associated with the fine silt and clay-sized fractions that were respectively 3 and 6-fold  
320 higher compared to the corresponding subsoil Bwg horizon in the NP soil. The substantial  
321 enrichment in pedogenic Fe associated with the finest soil fraction ( $< 2 \mu\text{m}$ ) of the Brd1  
322 horizon was mainly due to an increase in  $\text{Fe}_c$  (Figure 3d) that was also responsible for the  
323 lower  $\text{Fe}_o/\text{Fe}_d$  ratio observed with respect to the topsoils (Table 1). Although only limited  
324 changes in the distribution of pedogenic Fe between intra and inter-microaggregate  
325 fractions across different particles sizes was observed with soil depth in the P soil, the  
326 fine silt and clay-sized fractions of the Brd1 showed slightly higher intra:inter-  
327 microaggregate  $\text{Fe}_d$  ratios with respect to the corresponding subsoil horizon in the NP soil  
328 (Figure 2b).

329

### 330 **3.3 Fe speciation in the different aggregate and particle-size fractions**

331 NP soils showed minor variations in centroid position values with soil depth although  
332 clear differences were observed between particle-size fractions (Figure 4). PCE values  
333 tended to increase with decreasing particle size with a maximum difference of around 0.7  
334 eV between FSa and CI fractions, albeit no differences between intra- and inter-  
335 microaggregate fractions were observed (Figures 4 and S2; Table S2). Mean PCE values  
336 for the clay-sized fractions around 7114.59 eV suggest that Fe-containing mineral phases  
337 in this fraction were mostly oxidized. On the other hand, an increasing contribution of  
338 Fe(II) was observed for the FSa and CSi fractions with PCE values around 7113.93 eV.  
339 Similar differences in PCE values between particle-size fractions were also observed in  
340 the P soils, although in contrast to NP soils, clear trends with soil depth were noted (Figure  
341 4). Across all particle-size fractions, lower PCE values were generally obtained for  
342 topsoils with respect to subsoil horizons suggesting a greater contribution of Fe(II) to total  
343 Fe in the superficial horizons. The difference in PCE values between topsoils and subsoils  
344 was generally less pronounced for the finest clay-sized fractions (0.15 eV) with respect  
345 to the other fractions (0.18-0.34 eV), suggesting a greater contribution of reduced Fe in  
346 the coarse-sized fractions separated from P topsoils.

347 The deconvolution of the XANES spectra in the energy range between 7105 eV and 7145  
348 eV of both NP and P soils by LCF (Table 2) showed that Fe phases were adequately  
349 described by three or four components. The major differences in the relative contribution

350 of different Fe phases were linked to the different particle-size fractions rather than soil  
351 management and depth. According to the LCF results (Table 2), in the NP soil the FSa  
352 fractions contained relatively high proportions of chlorite (32-34%) that tended to  
353 decrease with decreasing particle-size, and siderite (26-33%) that was absent in the other  
354 fractions. Other minor phases in this fraction included illite (13-21%) and hematite (20-  
355 22%). Similarly, Fe phases in the coarse silt-sized fractions (CSi and mCSi) were also  
356 represented by chlorite (22-27%), illite (34-66%) and hematite (13-27%). Minor  
357 proportions of magnetite were also detected in this fraction (5-14%). Fine silt-sized  
358 fractions (FSi and mFSi) generally showed smaller proportions of chlorite (12-27%) and  
359 illite, though other Fe-containing minerals like smectite and nontronite were detected, as  
360 well as minor contributions (<10%) from organically complexed Fe(III). In contrast to  
361 the other fractions, the finest clay-sized fractions (Cl and mCl) were characterized by the  
362 absence of illite and the presence of relatively high amounts of nontronite that represented  
363 up to 79-83% of the Fe phases. This particle-size fraction also contained variable  
364 contributions from Fe oxide phases (e.g., lepidocrocite and ferrihydrite) as well as  
365 organically complexed Fe(III). Overall, in NP, the major differences in the relative  
366 contribution of different Fe phases were linked to the different particle-size fractions  
367 rather than depth.

368 In the P soils, the FSa fraction contained similar contents of chlorite (32-34%), and  
369 siderite (27-32%) as observed for the same fraction in NP. Other minor phases in this  
370 fraction included illite and hematite. Fe phases in the coarse silt-sized fractions were also  
371 represented by chlorite (18-27%), illite (39-75%) and hematite (9-23%) though with  
372 slightly different relative contributions. As for NP soils, fine silt-sized fractions (FSi and  
373 mFSi) generally contained chlorite (23-29%) and illite, smectite (16-57%) and nontronite  
374 (7-32%), as well as minor contributions (<10%) from organically complexed Fe(III). The  
375 clay-sized fractions were characterized by the absence of illite and the presence of  
376 relatively high amounts of nontronite that represented up to 79-84% of the Fe phases.  
377 This particle-size fraction also contained variable contributions from Fe oxide phases  
378 (e.g. lepidocrocite and ferrihydrite) as well as organically complexed Fe(III). These were  
379 most abundant in P subsoils.

380

### 381 **3.4 Fe EXAFS linear combination fitting and WT results**

382 EXAFS spectra of all size-fractions were accurately reconstructed by three or four  
383 components, and quantitative results of the LCF evidenced that Fe phases in all NP and

384 P soils were mainly characterized by the presence of chlorite, ferrihydrite, lepidocrocite,  
385 and Fe(III)-citrate (Figure 5, Tables 3 and S3).

386 LCF results for the size fractions, with the only exception of FSa, using three components  
387 are reported in Figure 6 and the sample fit (dotted lines) is labeled with the R-factor. Fe  
388 K-edge EXAFS spectra showed features at  $\sim 4.0 \text{ \AA}^{-1}$  (maximum 1),  $\sim 6.5 \text{ \AA}^{-1}$  (maximum  
389 2),  $\sim 7.5 \text{ \AA}^{-1}$  (maximum 3),  $\sim 8.5 \text{ \AA}^{-1}$  (maximum 4). The maximum at  $4.0 \text{ \AA}^{-1}$  has different  
390 features in ferrihydrite and lepidocrocite (insert in Figure 5). The maximum at  $7.5 \text{ \AA}^{-1}$  is  
391 characteristic of ferrihydrite, whereas the small maximum at  $8 \text{ \AA}^{-1}$  is typical of  
392 lepidocrocite and magnetite, and a broad one describes Fe(III)-citrate spectrum. The  
393 maximum at  $8.5 \text{ \AA}^{-1}$  is also a characteristic feature of ferrihydrite. The maximum at  $10$   
394  $\text{ \AA}^{-1}$  is only characteristic of lepidocrocite (insert in Figure 5). The maxima at  $6.5$  and  $7.5$   
395  $\text{ \AA}^{-1}$  (as also at  $8.5 \text{ \AA}^{-1}$ ) are characteristic of ferrihydrite (Figure 5).

396 Quantitatively, NP soils evidenced an increase in the relative abundance of ferrihydrite  
397 and Fe(III)-citrate phases with decreasing particle size, together constituting around 20-  
398 24 % and 9-21%, respectively in the coarse silt (CSi, mCSi) and both 31-43% in the clay-  
399 sized (Cl, mCl) fractions, with no evident trends with soil depth (Table 3). On the other  
400 hand, the relative abundance of chlorite was lower in the clay-sized fractions (4-15 %)   
401 with respect to the other fractions (30-49%), while lepidocrocite accounted for around  
402 11-19% of Fe phases across size fractions. Only the coarser FSa fraction was  
403 characterized by the presence of more crystalline Fe phases like hematite and goethite.  
404 Similarly, P soils showed an increasing abundance of ferrihydrite and Fe(III)-citrate  
405 phases with decreasing particle size, from 8-31 % and 9-29 % in the coarse-silt fractions  
406 (CSi, mCSi) to 9-41 % and 22-46 % in the clay-sized (Cl, mCl) fractions, respectively.  
407 Chlorite ranged from 9 to 66 % and lepidocrocite from 11 to 19 % across size fractions.  
408 However, in contrast to NP soils, P soils evidenced lower contents of ferrihydrite in the  
409 topsoils and more pronounced variations in the relative abundance of ferrihydrite, more  
410 crystalline Fe phases and Fe(III)-citrate with soil depth (Table 3). Considering exclusively  
411 the finest clay-sized fraction, the relative content of ferrihydrite increased from around  
412 14-20 % in the Arp1 and Arp2 horizons to 40% in the Brd1 horizon, while Fe(III)-citrate  
413 decreased from 41-46 % in the topsoils to 30% in the subsoils. Magnetite (up to 15 %)   
414 was also detected associated with the clay-sized fractions in the Brd2 horizon.

415 A key in the study of Fe associated with OM by EXAFS is a proper analysis of the  
416 second coordination shell. In traditional analysis of EXAFS spectra it is difficult to  
417 separate the contributions from different backscattering atoms in higher coordination

418 shells present at similar bonding distances from the central Fe atom. Complementary to  
419 LCF, WT provides useful information simultaneously resolving data in both wave space  
420 ( $k$ ) and interatomic distance ( $R$ ) space, and the resulting  $k$ - $R$  maps are helpful for  
421 improving the fitting model and understanding elemental speciation in the sample. In  
422 addition, WT analysis was used to qualitatively test the presence of Fe back-scatterers in  
423 the second coordination shell of Fe. The Morlet wavelet transforms of  $k^3$ -weighted Fe K-  
424 edge EXAFS spectra of the samples over a  $R + \Delta R$ -range of 2.2-4.0 Å was calculated and  
425 the resulting wavelet plots compared with those of Fe(III) reference compounds  
426 (ferrihydrite, Fe(III)-citrate, chlorite). Figure S3 shows the WT plots of the standards. In  
427 the ferrihydrite plot, the Fe shell contributes a strong feature at 7-8 Å<sup>-1</sup> and 2.75 Å  
428 (Daugherty et al., 2017). In the Fe(III)-citrate standard, the absence of Fe backscattering  
429 is accompanied by new features at distances of 2.0–2.5 Å and indicates back-scattering  
430 from lighter atoms, in agreement with single and multiple backscattering from C/O in the  
431 second and third coordination shells of tris(oxalato)Fe(III) (Daugherty et al., 2017;  
432 Karlsson and Persson, 2010). Moreover, these features appear at lower energies (3–4 Å<sup>-1</sup>)  
433 than those of Fe.

434 The WT of the EXAFS data ( $\eta=8$ ,  $\sigma=1$ ) of the CSi, FSi, Cl and relative aggregated  
435 fractions showed that both the intensity caused by light back-scatterers at low  $k$ -values  
436 and by heavier atoms at higher  $k$  varied among the samples (Figure 6), compared to the  
437 WT plots of standards (Figure S3). A general trend was observed with decreasing particle-  
438 size from the CSi to the Cl fraction in the NP topsoil horizons (Ap1 and Ap2; Figure 6a).  
439 The clay samples generally have more intense maxima near 8 Å<sup>-1</sup>, typical of ferrihydrite.  
440 A slight increase in the Fe-SOM signal contribution at above 2.2 Å and 2–4 Å<sup>-1</sup> is also  
441 present, as reported in the EXAFS LCF results, while the CSi fractions clearly match with  
442 the high chlorite content. In the Bgw fraction the increasing content of ferrihydrite (up to  
443 50% considering 3 components) is evident in the intense peak at 8 Å<sup>-1</sup> in the Cl and mCl.  
444 Although an increase in Fe oxides and organically-complexed forms occurred from the  
445 coarse to the fine fractions, significant differences were not found ranging from the  
446 topsoil to the subsoil in NP soil.

447 Considering the equivalent topsoil horizons under P management, spectra of the Arp1  
448 and Arp2 horizons are characterized by features indicating an increase in Fe-SOM and  
449 ferrihydrite with decreasing particle-size from the CSi to the Cl fraction (Figure 6b). In  
450 addition, the mCl fraction, as also reported in the LCF performed using three components,  
451 clearly matches lepidocrocite features. In the Arpd horizon of the P soil the increasing

452 contribution of ferrihydrite (from 21 to up to 28% from the CSi to the Cl fraction) and  
453 Fe-SOM (from ca. 18 to ca. 38% from the CSi to the Cl fraction) are also reported in the  
454 WT of the EXAFS data. In the Brd1 horizon the higher Fe(III)-SOM content in the Cl  
455 with respect to the mCl fraction is evident. As also evidenced by the LCF, CSi contains  
456 less ferrihydrite than mCSi. Compared to the Brd1 horizon, in the Brd2 horizon the  
457 ferrihydrite content decreased, whereas the increased Fe(III)-SOM contents, in all the  
458 fractions, appear as a clear feature.

459

## 460 **4. DISCUSSION**

### 461 **4.1 Influence of paddy management on OC distribution in the aggregate and** 462 **particle-size fractions in topsoils and subsoils**

463 The larger accumulation of OM in soils under paddy management with respect to other  
464 arable ecosystems has been widely recognized to contribute to the C sink functions of rice  
465 agro-ecosystems (Kalbitz et al., 2013; Kögel-Knabner et al., 2010; Wissing et al., 2011).  
466 This has often been attributed to the retarded decomposition of OC under waterlogged  
467 conditions during several months of paddy field flooding every year (Sahrawat, 2004),  
468 although enhanced soil microaggregation (Wissing et al., 2014) and specific organo-  
469 mineral associations have also been shown to drive OC accrual in these soils (Winkler et  
470 al., 2019). Apart from confirming the greater OC contents in P with respect to NP topsoils,  
471 this study also suggests that retarded decomposition could be partly responsible for the  
472 higher contents of particulate OM associated with the coarser size fractions under paddy  
473 management. This is in line with the findings of Chen et al. (2021) who showed that SOC  
474 sequestration in paddy soils is partly linked to the accumulation of crop residue-derived  
475 labile C. Although paddy management resulted in a greater amount of particulate OM  
476 responsible for around 25 % of total SOC, mineral-associated OM still represented the  
477 greatest proportion of SOC. The interaction of OC with soil minerals and its subsequent  
478 stabilization against microbial mineralization largely depend on soil redox conditions  
479 (Eusterhues et al., 2014). In fact, long-term paddy management favored OC stabilization  
480 through this mechanism as P soils showed higher amounts of OC associated with the finer  
481 soil fractions with respect to NP soils, not only in the topsoil but also in the subsoil.  
482 Moreover, our results evidenced that whereas the clay-sized fraction is the main OC sink  
483 in both paddy topsoil and illuvial horizons (40 and 47-62 %, respectively), OC associated  
484 with the fine silt-sized fraction can also represent a substantial proportion of total SOC in  
485 P topsoils (on average 23 % of total SOC). This was assumed to be linked to the presence



486 of a substantial proportion of pedogenetic Fe oxides (26 %) in this fraction probably due  
487 to their redistribution from the finer to coarser fractions with repeated cycles of Fe(III)  
488 reduction and re-oxidation/precipitation that characterize these topsoils.

489 OC stabilization in the paddy soils was not only affected by particle size but also  
490 related to redox-driven changes in microaggregate formation and stability. Paddy  
491 management resulted in relatively lower contents of OC associated with the intra-  
492 microaggregate clay-sized particles in the topsoil with respect to the NP soil, but favored  
493 the presence of OC associated with the intra-microaggregate fine silt-sized particles.  
494 Aggregate stability may be somewhat compromised in soils subjected to alternating redox  
495 conditions possibly due to the disruptive energy occurring upon slaking (Six et al., 2000),  
496 and reductive dissolution of Fe oxides that are known to serve as binding agents holding  
497 aluminosilicate particles together (Blankinship et al., 2017; De-Campos et al., 2009). Our  
498 results suggest that clay-sized particles are more prone to dispersion during  
499 microaggregate breakdown under alternating redox conditions probably due to their more  
500 negative surface charge and greater susceptibility of the SRO-Fe oxides holding  
501 aluminosilicate particles together to microbial reduction. In contrast, the higher amount  
502 of OC associated with silt-sized particles within microaggregates under paddy  
503 management suggests that this size fraction could be less susceptible to aggregate  
504 breakdown under anoxic conditions and therefore represent an important pool of stable  
505 OC in these soils. This is also corroborated by the higher ratio of intra:inter-  
506 microaggregate pedogenic Fe and lower  $Fe_o/Fe_d$  ratio observed for the silt-sized with  
507 respect to the clay-sized fractions in the P topsoils. Our findings therefore partially  
508 confirm our first hypothesis that paddy management may affect the distribution of OC  
509 between different particle-size fractions due to the influence of redox cycling on  
510 aggregate stability. Huang et al. (2018) previously reported how management-induced  
511 soil aggregate turnover and OC stabilization was strongly linked to Fe cycling in a rice-  
512 wheat crop rotation, particularly when exogenous OM was applied. Similarly, De-  
513 Campos et al. (2009) showed that the reductive dissolution of poorly ordered Fe oxides  
514 involved in aggregate binding may lead to changes in the morphology of soil particles,  
515 favour colloid dispersion and contribute to a decrease in aggregate stability under anoxic  
516 conditions. However, notwithstanding the depletion of total pedogenetic Fe in the topsoil  
517 (about 50% less  $Fe_d$ ), more OC is stabilized by association with the finest soil particles  
518 both within and outside microaggregates (about 25% more OC) with respect to non-paddy  
519 soils. In paddy topsoils coprecipitation, known to retain more OC with respect to surface

520 adsorption on Fe (hydr)oxides and leading to the formation of Fe-OM associations with  
521 a high C/Fe ratio (Sodano et al., 2017), may in fact represent an important mechanism for  
522 protected OM from microbial degradation (Ginn et al., 2017) and thus contribute to OC  
523 accumulation. This was also confirmed by EXAFS LCF that evidenced a substantial  
524 proportion of Fe-OM phases (up to 45%) and ferrihydrite (up to 20%) in the clay-sized  
525 fractions (Table 3).

526 The significant enrichment in OC associated with intra-microaggregate clay-sized  
527 particles in the P subsoil horizons when compared to the corresponding horizons in the  
528 NP soil suggest that microaggregate formation and stabilization in the subsoil may also  
529 contribute to the preservation of OC and could be linked to the mobilization of dissolved  
530 OC and Fe from the topsoil during the rice cropping season when the paddy fields are  
531 flooded (Said-Pullicino et al., 2016), confirming our second hypothesis. In a previous  
532 study (Said-Pullicino et al., 2021) we provided evidence that colloid transport, together  
533 with the percolation of  $\text{Fe}^{2+}$  and subsequent precipitation of SRO oxides in the oxic  
534 subsoil horizons, could contribute to the accumulation of Fe and OC in the illuvial  
535 horizons. In the present study, we are further showing that these redox-driven dynamics  
536 may also contribute to microaggregate formation and soil OC stabilization in the deeper  
537 soil horizons of temperate paddy soils.

538

#### 539 **4.2 Fe mineral transformations and distribution in the different aggregate and** 540 **particle-size fractions**

541 As expected, paddy management resulted in a general depletion in both SRO and  
542 crystalline Fe oxides in the topsoil with respect to the non-paddy soil, probably due to the  
543 reductive dissolution of Fe oxides and subsequent leaching into the subsoil with repeated  
544 redox cycles, as suggested by Said-Pullicino et al. (2021). Nonetheless, the higher  $\text{Fe}_o/\text{Fe}_d$   
545 ratio of the clay-sized fractions (Table 1) indicate a relative increase in SRO Fe oxides  
546 associated with this fraction primarily in the form of ferrihydrite and Fe-OM complexes  
547 (Table 3). In redox-active paddy topsoils, repeated reductive dissolution of both  
548 crystalline and less ordered Fe oxides under anoxic conditions with subsequent release of  
549 high concentrations of dissolved  $\text{Fe}^{2+}$  and DOC (Bertora et al., 2018; Said-Pullicino et  
550 al., 2016), followed by the oxidative re-precipitation with the re-establishment of oxic  
551 conditions or in the  $\text{O}_2$ -rich rice rhizosphere, can lead to the formation of new mineral  
552 phases, mainly SRO oxides more or less associated with phyllosilicate particles (Sodano  
553 et al., 2016) or Fe-OC coprecipitates (Sodano et al., 2017). The presence of minor

554 amounts of lepidocrocite (around 12-15 %) also suggests that the Fe<sup>2+</sup>-catalyzed  
555 recrystallization or solid state transformations of ferrihydrite to more thermodynamically  
556 stable phases can also occur to some extent. Chen and Thompson (2021) report that the  
557 nature of the new Fe minerals formed as a result of Fe(II) oxidation may be directed by  
558 the relative presence of OM and Fe phases in the soil, that can either hinder or promote  
559 crystal growth and the formation of more crystalline phases, respectively. It is thus highly  
560 likely that in the studied soils that are relatively poor in pedogenic Fe but that hold  
561 important amounts of OM (Said-Pullicino et al., 2021), the trajectory of Fe minerals under  
562 paddy management tends towards the relative accumulation of SRO Fe(III) phases. Our  
563 results further show that redox-driven Fe phase transformations may also involve changes  
564 in microaggregate stability that consequently influences the association of new mineral  
565 phases with more aggregated (for fine silt-sized) or dispersed (for clay-sized) mineral  
566 particles (Figure 2b). EXAFS LCF suggests that hydrous Fe minerals present in the clay  
567 and silt-sized fraction of paddy topsoils are mainly represented by ferrihydrite (about 18  
568 and 14 %, respectively) and Fe-OM phases (about 45 and 30 %, respectively) confirming  
569 the role these size-fractions may play in OC retention.

570 The lower PCE values generally observed across particle-size fractions in paddy topsoils  
571 with respect to both paddy subsoils and non-paddy topsoils (Figure 4) suggest that in this  
572 redox-active horizon a proportion of Fe in the mineral phase may remain in its reduced  
573 form even after field drainage. XANES LCF suggests that this may be due to the reduction  
574 of structural Fe present in phyllosilicate structures (nontronite and smectite in the finer  
575 fractions, and chlorite in the coarser fractions) under anaerobic conditions (Table 2).  
576 Various studies have shown that the biological reduction of Fe(III) in phyllosilicates like  
577 nontronite (Zhao et al., 2013) and illite-smectite mixed layers (Shelobolina et al., 2012)  
578 is possible in soils subject to reducing conditions (Liao et al., 2019; Neumann et al., 2015;  
579 Ribeiro et al., 2009; Dong et al., 2009 and references therein) to an extent that depends  
580 on the location of Fe(III) within the clay structure (Shi et al. 2021). However, the similar  
581 proportions of nontronite in P and NP topsoils suggest that this redox-cycling under paddy  
582 management did not lead to *in situ* loss of structural Fe. Similar results were obtained by  
583 Vogelsang et al. (2016) after one year of incubation of nontronite in a paddy soil. Akter  
584 et al. (2018) have also shown that reduction of octahedral Fe<sup>3+</sup> in clay minerals and their  
585 interstratified forms could be responsible for accepting electrons during OM  
586 mineralization under anaerobic conditions. Vertical redox gradients and OM contents in  
587 hydromorphic soils can therefore exert a major control on the oxidation state of structural

588 Fe in phyllosilicates and other Fe containing minerals (Chen et al. 2017). However, with  
589 the re-establishment of oxic conditions, structural Fe in clay minerals can be rapidly  
590 reoxidized (within days) to Fe(III) through abiotic or biotic oxidation, as reported for  
591 chlorite and nontronite-associated Fe(II) by Zhang et al. (2011) and Shi et al. (2021). It is  
592 however possible that in under oxic conditions reduced Fe was preferentially reoxidized  
593 in smaller particles having higher contact surfaces with O<sub>2</sub> while reoxidation was slower  
594 in larger size fractions and within microaggregates. This would partially explain the  
595 increasing presence of reduced Fe(II) with increasing particle size in the redox-dynamic  
596 topsoils, even if the soils were already oxic during sampling. The lower PCE values in  
597 the coarser soil fractions may also be due to the precipitation of Fe(II) carbonates.  
598 XANES LCF evidenced that the precipitation of siderite is evidently not associated with  
599 the fine particle-sizes like most other secondary minerals, but eventually other  
600 mechanisms (e.g., linked with high rates of CO<sub>2</sub> production during the mineralization of  
601 particulate OM) may drive their precipitation on coarser fractions (Table 2). However,  
602 due to the presence of similar proportions of siderite in the coarse fractions of both paddy  
603 and non-paddy topsoils and subsoils we cannot exclude a lithogenic origin of this mineral.  
604 Siderite oxidation kinetics are known to be relatively slow as structurally bound Fe(II) in  
605 carbonates are not particularly electro-active due to the low Fe(II) surface density, and  
606 rates of oxidation are generally slower with increasing nominal particle size (Hoving et  
607 al. 2017) or due to surface passivation of the mineral by organic ligands (Rothwell and  
608 Kretzschmar, 2020). Nonetheless, the different trend in PCE values for this fraction in  
609 paddy and non-paddy soils suggests that redox-cycling in the former can affect the  
610 oxidation state of Fe phases along the soil profile and that Fe K-edge XANES can  
611 adequately evaluate these changes.

612 Our results also evidenced that paddy illuvial horizons just beneath the plough pan can  
613 accumulate remarkable amounts of Fe (hydr)oxides particularly associated with the finest  
614 particle fractions and also contribute to microaggregate formation and stabilization  
615 (Figure 3). This is consistent with the inputs of colloidal and/or dissolved Fe from the  
616 topsoil and the larger Fe oxide stocks in the subsoil (about 40 % higher) with respect to  
617 adjacent non-paddy soils reported for temperate paddy soils by Said-Pullicino et al.  
618 (2021). EXAFS LCF suggested that most of the Fe phases associated with the finest  
619 particle fractions in the Brd1 horizon were represented by ferrihydrite (40%) and Fe-OM  
620 forms (30%) confirming that these subsoil horizons can play an important role in OC  
621 accumulation in rice paddy soils. However, we did not observe substantial differences in

622 the mineral composition of pedogenetic Fe phases in paddy and non-paddy subsoils  
623 (Table 3). Nor was the transformation of ferrihydrite into more crystalline phases  
624 particularly expressed because interaction with dissolved OM could have inhibited the  
625 Fe<sup>2+</sup>-catalyzed transformation into more crystalline phases even in the subsoil.  
626 Nonetheless, the increasing trend in PCE with soil depth under paddy but not under non-  
627 paddy management (Figure 4) clearly suggests that redox cycling in paddy topsoils  
628 contribute reduced Fe forms, while Fe phases in the subsoil are mostly oxidized, except  
629 for the coarser-size fractions where siderite was detected (Table 2). In these paddy fields,  
630 subsoil horizons just beneath the plough pan generally remain prevalently oxic even when  
631 the topsoils are flooded, as the rising ground water table during spring and summer (up  
632 to around 40 cm below the soil surface) does not limit oxygen diffusion until late in the  
633 cropping season (mid-July).

#### 634 **4.3 Potential and limitation of Fe K-edge XANES and EXAFS for Fe speciation in soil fractions**

635 This work together with the results of previous studies carried out on well-defined binary  
636 and ternary mixtures of different Fe-bearing minerals (e.g., Wilke et al., 2001; O'Day et  
637 al., 2004; Prietzel et al., 2007) showed that Fe K-edge XANES can provide an estimate  
638 of the relative contribution of different mineral classes and groups of organic compounds  
639 with different oxidation state of the Fe atom (e.g., Fe(II) sulphides versus Fe(III)  
640 oxyhydroxides; Fe(II)-organic complexes versus Fe(III)-organic complexes) by  
641 calculating the PCE and applying LCF to the entire XANES spectrum. In soils this  
642 evaluation is complicated by the presence of mixed Fe phases that form during  
643 pedogenesis. Therefore, accurate quantification of different Fe phases in soils by Fe K-  
644 edge XANES will probably always remain difficult to achieve particularly in  
645 environments experiencing dynamic redox conditions. At present, LCF conducted on the  
646 XANES spectrum in an energy range between 7105 and 7145 eV allows the quantification  
647 of specific minerals in soils approximately, but not accurately. The accuracy of the results  
648 of the LCF procedure is strongly affected by the correctness of the applied set of predictor  
649 variables. LCF on the XANES spectrum revealed a systematic influence of increasing  
650 oxidizing conditions on the Fe speciation in the topsoil and the subsoil of paddy soil: the  
651 contribution of Fe(III) compounds increased with soil depth and decreasing particle size  
652 fractions, especially in the aggregated ones. This study highlights the potential of Fe K-  
653 edge XANES for the determination of Fe speciation in redoximorphic soils and soil  
654 fractions, despite its present problems and limitations. With Fe K-edge XANES, the  
655 proportion of Fe(II) and Fe(III) and the relative contribution of different Fe compounds

656 in soil samples and soil aggregates can be evaluated. However, although Fe K-edge  
657 XANES can estimate fairly well the relative contribution of different mineral classes and  
658 groups of organic compounds with different oxidation states of the Fe atom in mineral  
659 mixtures and soil samples, it fails to quantify specific Fe oxyhydroxides (e.g., ferrihydrite,  
660 goethite, and hematite) or distinct Fe-organic compounds. Thus, a combination with Fe  
661 EXAFS, as also WT, allows the speciation of Fe in different fractions and aggregates in  
662 soils subjected to dynamic redox conditions.

663

## 664 **5. CONCLUSIONS**

665 Evaluating the mechanisms for OC stabilization in paddy soils still deserves attention  
666 particularly due to the complex interactions between C cycling and changes in the  
667 contents and mineralogy of hydrous Fe oxides in redox-active systems. In this study, we  
668 linked the enhanced stabilization of OC in temperate paddy soils to the redox-driven  
669 changes in Fe phases (evaluated by Fe K-edge XANES and EXAFS spectroscopy) and  
670 their distribution between aggregate and particle-size fractions. Our findings have shown  
671 that although paddy topsoils show higher contents of particulate OC in the coarser size-  
672 fractions with respect to non-paddy soils, most of the SOC is associated with the finer  
673 soil fractions, not only in the topsoil but also in the subsoil. This suggests that the  
674 management-induced differences in C stocks is not primarily due to the limited  
675 decomposition of crop residues under paddy management, but rather to the redox-driven  
676 changes in the association of OC with soil minerals. We show that OC stabilization  
677 through interaction with minerals is also affected by microaggregate stability. Paddy  
678 management may lead to microaggregate breakdown with a preferential release of clay-  
679 sized particles rich in both SRO Fe oxides and OC, while intra-microaggregate fine silt-  
680 sized particles may also serve as an important OC sink. Long-term alternating redox  
681 conditions generally resulted in paddy topsoil horizons that were depleted in pedogenic  
682 Fe with respect to non-paddy soils, and led to a redistribution of Fe phases across different  
683 particle size fraction often as less crystalline phases (primarily ferrihydrite and Fe-OM  
684 associations). Nonetheless, the higher C contents indicate that higher C loadings can be  
685 achieved under these redox-dynamic environments. Moreover, illuvial horizons under  
686 paddy management were enriched in SRO hydrous Fe oxides and this contributed to  
687 enhanced microaggregate formation and C stabilization with respect to non-paddy  
688 subsoils.

689

690 **Acknowledgements**

691 This study was partly funded by the University of Torino as part of the research project  
692 "Colloid mobility and organic matter stabilization in hydromorphic soils". We also like  
693 to express our gratitude to Marco Romani for his assistance with sample collection.

## References

- Akter, M., Deroo, H., De Grave, E., Van Alboom, T., Kader, M.A., Pierreux, S., Begum, M.A., Boeckx, P., Sleutel, S., 2018. Link between paddy soil mineral nitrogen release and iron and manganese reduction examined in a rice pot growth experiment. *Geoderma* 326, 9–21. <https://doi.org/10.1016/j.geoderma.2018.04.002>
- Aquilanti, G., Giorgetti, M., Dominko, R., Stievano, L., Arčon, I., Novello, N., Olivi, L., 2017. *Operando* characterization of batteries using x-ray absorption spectroscopy: advances at the beamline XAFS at synchrotron Elettra. *J. Phys. D. Appl. Phys.* 50, 074001. <https://doi.org/10.1088/1361-6463/aa519a>
- Balesdent, J., Chenu, C., Balabane, M., 2000. Relationship of soil organic matter dynamics to physical protection and tillage. *Soil Tillage Res.* 53, 215–230. [https://doi.org/10.1016/S0167-1987\(99\)00107-5](https://doi.org/10.1016/S0167-1987(99)00107-5)
- Bertora, C., Cucu, M.A., Lerda, C., Peyron, M., Bardi, L., Gorra, R., Sacco, D., Celi, L., Said-Pullicino, D., 2018. Dissolved organic carbon cycling, methane emissions and related microbial populations in temperate rice paddies with contrasting straw and water management. *Agric. Ecosyst. Environ.* 265, 292–306.
- Blankinship, J.C., Fonte, S.J., Six J, Schimel, J.P., 2016. Plant versus microbial controls on soil aggregate stability in a seasonally dry ecosystem. *Geoderma* 272, 39–50. <https://doi.org/10.1016/j.geoderma.2016.03.008>
- Chen, X., Hu, Y., Xia, Y., Zheng, S., Ma, C., Rui, Y., He, H., Huang, D., Zhang, Z., Ge, T., Wu, J., Guggenberger, G., Kuzyakov, Y., Su, Y., 2021. Contrasting pathways of carbon sequestration in paddy and upland soils. *Glob. Chang. Biol.* 27, 2478–2490. <https://doi.org/10.1111/gcb.15595>
- Chen, C., Kukkadapu, R.K., Lazareva, O., Sparks, D.L., 2017. Solid-phase Fe speciation along the vertical redox gradients in floodplains using XAS and Mössbauer spectroscopies. *Environ. Sci. Technol.* 2017, 51, 7903–7912. <https://doi.org/10.1021/acs.est.7b00700>
- Chen, C., Thompson, A., 2021. The influence of native soil organic matter and minerals on ferrous iron oxidation. *Geochim. Cosmochim. Acta* 292, 254–270. <https://doi.org/10.1016/j.gca.2020.10.002>
- Cheng, Y.Q., Yang, L.Z., Cao, Z.H., Ci, E., Yin, S., 2009. Chronosequential changes of selected pedogenic properties in paddy soils as compared with non-paddy soils. *Geoderma* 151, 31–41. <https://doi.org/10.1016/j.geoderma.2009.03.016>
- Chorover, J., Amistadi, M.K., Chadwick, O.A., 2004. Surface charge evolution of mineral-organic complexes during pedogenesis in Hawaiian basalt. *Geochim. Cosmochim. Acta* 68, 4859–4876. <https://doi.org/10.1016/j.gca.2004.06.005>
- Coby, A.J., Picardal, F., Shelobolina, E., Xu, H., Roden, E.E., 2011. Repeated anaerobic microbial redox cycling of iron. *Appl. Environ. Microbiol.* 77, 6036–6042. <https://doi.org/10.1128/AEM.00276-11>
- Coward, E.K., Thompson, A., Plante, A.F., 2018. Contrasting Fe speciation in two humid forest soils: insight into organomineral associations in redox-active environments. *Geochim. Cosmochim. Acta* 238, 68–84. <https://doi.org/10.1016/j.gca.2018.07.007>
- Daugherty, E.E., Gilbert, B., Nico, P.S., Borch, T., 2017. Complexation and redox buffering of iron(II) by dissolved organic matter. *Environ. Sci. Technol.* 51, 11096–11104. <https://doi.org/10.1021/acs.est.7b03152>



- De-Campos, A.B., Mamedov, A.I., Huang, C.-H., 2009. Short-term reducing conditions decrease soil aggregation. *Soil Sci. Soc. Am. J.* 73, 550-559. <https://doi.org/10.2136/sssaj2007.0425>
- Di Cicco, A., Aquilanti, G., Minicucci, M., Principi, E., Novello, N., Cognigni, A., Olivi, L., 2009. Novel XAFS capabilities at ELETTRA synchrotron light source. *J. Phys. Conf. Ser.* 190. <https://doi.org/10.1088/1742-6596/190/1/012043>
- Dong, H.L., Jaisi, D.P., Kim, J., Zhang, G.X., 2009. Microbe-clay mineral interactions. *Am. Mineral.* 94, 1505–1519. <https://doi.org/10.2138/am.2009.3246>
- Eusterhues, K., Hädrich, A., Neidhardt, J., Küsel, K., Keller, T.F., Jandt, K.D., Totsche, K.U., 2014. Reduction of ferrihydrite with adsorbed and coprecipitated organic matter: Microbial reduction by *Geobacter bremensis* vs. abiotic reduction by Na-dithionite. *Biogeosciences* 11, 4953–4966. <https://doi.org/10.5194/bg-11-4953-2014>
- Filimonova, S., Kaufhold, S., Wagner, F.E., Häusler, W., Kögel-Knabner, I., 2016. The role of allophane nano-structure and Fe oxide speciation for hosting soil organic matter in an allophanic Andosol. *Geochim. Cosmochim. Acta* 180, 284–302. <https://doi.org/10.1016/j.gca.2016.02.033>
- Funke, H., Scheinost, A.C., Chukalina, M., 2005. Wavelet analysis of extended x-ray absorption fine structure data. *Phys. Rev. B* 71, 094110. <https://doi.org/10.1103/PhysRevB.71.094110>
- Giannetta, B., Plaza, C., Siebecker, M.G., Aquilanti, G., Vischetti, C., Plaisier, J.R., Juanco, M., Sparks, D.L., Zacccone, C., 2020a. Iron speciation in organic matter fractions isolated from soils amended with biochar and organic fertilizers. *Environ. Sci. Technol.* 54, 5093–5101. <https://doi.org/10.1021/acs.est.0c00042>
- Giannetta, B., Siebecker, M.G., Zacccone, C., Plaza, C., Rovira, P., Vischetti, C., Sparks, D.L., 2020b. Iron(III) fate after complexation with soil organic matter in fine silt and clay fractions: An EXAFS spectroscopic approach. *Soil Tillage Res.* 200, 104617. <https://doi.org/10.1016/j.still.2020.104617>
- Ginn, B., Meile, C., Wilmoth, J., Tang, Y., Thompson, A., 2017. Rapid iron reduction rates are stimulated by high-amplitude redox fluctuations in a tropical forest soil. *Environ. Sci. Technol.* 51, 3250–3259. <https://doi.org/10.1021/acs.est.6b05709>
- Hall, S.J., Berhe, A.A., Thompson, A., 2018. Order from disorder: do soil organic matter composition and turnover co-vary with iron phase crystallinity? *Biogeochemistry* 140, 93–110. <https://doi.org/10.1007/s10533-018-0476-4>
- Henderson, R., Kabengi, N., Mantripragada, N., Cabrera, M., Hassan, S., Thompson, A., 2012. Anoxia-induced release of colloid- and nanoparticle-bound phosphorus in grassland soils. *Environ. Sci. Technol.* 46, 11727–11734. <https://doi.org/10.1021/es302395r>
- Hoving, A.L., Sander, M., Bruggeman, C., Behrends, T., 2017. Redox properties of clay-rich sediments as assessed by mediated electrochemical analysis: Separating pyrite, siderite and structural Fe in clay minerals. *Chemical Geology*, 457, 149-161. <https://doi.org/10.1016/j.chemgeo.2017.03.022>
- Huang, W., Hall, S.J., 2017. Elevated moisture stimulates carbon loss from mineral soils by releasing protected organic matter. *Nat. Commun.* 8, 1774. <https://doi.org/10.1038/s41467-017-01998-z>
- Huang, X., Tang, H., Kang, W., Yu, G., Ran, W., Jianping Hong, J., Shen, Q., 2018. Redox interface-associated organo-mineral interactions: A mechanism for C sequestration under a rice-wheat cropping system. *Soil Biol. Biochem.* 120, 12-23. <https://doi.org/10.1016/j.soilbio.2018.01.031>
- Huang, X., Kang, W., Guo, J., Wang, L., Tang, H., Li, T., Yu, G., Ran, W., Hong, J., Shen, Q., 2020. Highly reactive nanomineral assembly in soil colloids: implications for paddy soil carbon

- storage. *Sci. Total Environ.* 703, 134728. <https://doi.org/10.1016/j.scitotenv.2019.134728>
- Kalbitz, K., Kaiser, K., Fiedler, S., Kölbl, A., Amelung, W., Bräuer, T., Cao, Z., Don, A., Grootes, P., Jahn, R., Schwark, L., Vogelsang, V., Wissing, L., Kögel-Knabner, I., 2013. The carbon count of 2000 years of rice cultivation. *Glob. Chang. Biol.* 19, 1107–1113. <https://doi.org/10.1111/gcb.12080>
- Karlsson, T., Persson, P., 2010. Coordination chemistry and hydrolysis of Fe(III) in a peat humic acid studied by X-ray absorption spectroscopy. *Geochim. Cosmochim. Acta* 74, 30–40. <https://doi.org/10.1016/j.gca.2009.09.023>
- Kirk, G. 2004. *The Biogeochemistry of Submerged Soils*. John Wiley & Sons, Chichester, U.K. 291 pp. ISBN 0-470-86301-3.
- Kögel-Knabner, I., Amelung, W., Cao, Z., Fiedler, S., Frenzel, P., Jahn, R., Kalbitz, K., Kölbl, A., Schloter, M., 2010. Biogeochemistry of paddy soils. *Geoderma* 157, 1–14. <https://doi.org/10.1016/j.geoderma.2010.03.009>
- Kölbl, A., Schad, P., Jahn, R., Amelung, W., Bannert, A., Cao, Z.H., Fiedler, S., Kalbitz, K., Lehdorff, E., Müller-Niggemann, C., Schloter, M., Schwark, L., Vogelsang, V., Wissing, L., Kögel-Knabner, I., 2014. Accelerated soil formation due to paddy management on marshlands (Zhejiang Province, China). *Geoderma* 228–229, 67–89. <https://doi.org/10.1016/j.geoderma.2013.09.005>
- Kramer, M.G., Chadwick, O.A., 2018. Climate-driven thresholds in reactive mineral retention of soil carbon at the global scale. *Nat. Clim. Chang.* 8, 1104–1108. <https://doi.org/10.1038/s41558-018-0341-4>
- Kramer, M.G., Sanderman, J., Chadwick, O.A., Chorover, J., Vitousek, P.M., 2012. Long-term carbon storage through retention of dissolved aromatic acids by reactive particles in soil. *Glob. Chang. Biol.* 18, 2594–2605. <https://doi.org/10.1111/j.1365-2486.2012.02681.x>
- Lalonde, K., Mucci, A., Ouellet, A., Gélinas, Y., 2012. Preservation of organic matter in sediments promoted by iron. *Nature* 483, 198–200. <https://doi.org/10.1038/nature10855>
- Liao, W., Yuan, S., Liu, X., Tong, M., 2019. Anoxic storage regenerates reactive Fe(II) in reduced nontronite with short-term oxidation. *Geochim. Cosmochim. Acta* 257, 96–109. <https://doi.org/10.1016/j.gca.2019.04.027>
- Mehra, O.P., Jackson, M.L., 1960. Iron oxide removal from soils and clays by dithionite–citrate systems buffered with sodium bicarbonate. *Clay Clay Miner.* 7, 317–327. <https://doi.org/10.1016/B978-0-08-009235-5.50026-7>
- Neumann, A., Wu, L., Li, W., Beard, B.L., Johnson, C.M., Rosso, K.M., Friedrich, A.J., Scherer, M.M., 2015. Atom exchange between aqueous Fe(II) and structural Fe in clay minerals. *Environ. Sci. Technol.* 49, 2786–2795. <https://doi.org/10.1021/es504984q>
- O’Day, P.A., Rivera, N., Root, R., Carroll, S.A., 2004. X-ray absorption spectroscopic study of Fe reference compounds for the analysis of natural sediments. *Am. Mineral.* 89, 572–585. <https://doi.org/10.2138/am-2004-0412>
- Oades, J.M., Waters, A.G., 1991. Aggregate hierarchy in soils. *Aust. J. Soil Res.* 29, 815–825. <https://doi.org/10.1071/SR9910815>
- Pan, G., Li, L., Wu, L., Zhang, X., 2004. Storage and sequestration potential of topsoil organic carbon in China’s paddy soils. *Glob. Chang. Biol.* 10, 79–92. <https://doi.org/10.1111/j.1365-2486.2003.00717.x>
- Prietzl, J., Thieme, J., Eusterhues, K., Eichert, D., 2007. Iron speciation in soils and soil aggregates

- by synchrotron-based X-ray microspectroscopy (XANES,  $\mu$ -XANES). *Eur. J. Soil Sci.* 58, 1027–1041. <https://doi.org/10.1111/j.1365-2389.2006.00882.x>
- Prietzl, J., Tyufekchieva, N., Eusterhues, K., Kögel-Knabner, I., Thienme, J., Paterson, D., McNulty, I., de Jonge, M., Eichert, D., Salomé, M., 2009. Anoxic versus oxic sample pretreatment: effects on the speciation of sulfur and iron in well-aerated and wetland soils as assessed by X-ray absorption near-edge spectroscopy (XANES). *Geoderma* 153, 318–330. <https://doi.org/10.1016/j.geoderma.2009.08.015>
- Ribeiro, F.R., Fabris, J.D., Kostka, J.E., Komadel, P., Stucki, J.W., 2009. Comparisons of structural iron reduction in smectites by bacteria and dithionite: II. A variable-temperature Mössbauer spectroscopic study of Garfield nontronite. *Pure Appl. Chem.* 81, 1499–1509. <https://doi.org/10.1351/PAC-CON-08-11-16>
- Riedel, T., Zak, D., Biester, H., Dittmar, T., 2013. Iron traps terrestrially derived dissolved organic matter at redox interfaces. *Proc. Natl. Acad. Sci. U. S. A.* 110, 10101–10105. <https://doi.org/10.1073/pnas.1221487110>
- Rothwell, K., Kretzschmar, R., 2020. Siderite oxidation in the presence of organic ligands. EGU General Assembly 2020-16894. <https://doi.org/10.5194/egusphere-egu2020-16894>
- Said-Pullicino, D., Giannetta, B., Demeglio, B., Missong, A., Gottselig, N., Romani, M., Bol, R., Klumpp, E., Celi, L., 2021. Redox-driven changes in water-dispersible colloids and their role in carbon cycling in hydromorphic soils. *Geoderma* 385, 114894. <https://doi.org/10.1016/j.geoderma.2020.114894>
- Said-Pullicino, D., Miniotti, E.F., Sodano, M., Bertora, C., Lerda, C., Chiaradia, E.A., Romani, M., Cesari de Maria, S., Sacco, D., Celi, L., 2016. Linking dissolved organic carbon cycling to organic carbon fluxes in rice paddies under different water management practices. *Plant Soil* 401, 273–290. <https://doi.org/10.1007/s11104-015-2751-7>
- Sawrat K.L. (2004) Organic matter accumulation in submerged soils. *Adv. Agron.* 81, 169–201.
- Schwertmann, U., 1964. Differenzierung der Eisenoxide des Bodens durch Extraktion mit saurer Ammonium-oxalat Lösung. *Z Pflanzenernaehr Dueng Bodenkd* 105, 194–202.
- Shelobolina, E., Konishi, H., Xu, H., Benzine, J., Xiong, M.Y., Wu, T., Blöthe, M., Roden, E., 2012. Isolation of phyllosilicate-iron redox cycling microorganisms from an illite-smectite rich hydromorphic soil. *Front. Microbiol.* 3, 1–10. <https://doi.org/10.3389/fmicb.2012.00134>
- Shi, B., Smeaton, C., Roden, E., Lee, S., Liu, K., Xu, H., Kendall, B., Johnson, C., Parsons, C.T., Van Cappellen, P., 2021. Consecutive Fe redox cycles decrease bio-reducible Fe(III) and Fe isotope fractions by eliminating small clay particles. *Geochim. Cosmochim. Acta* 308, 118–135. <https://doi.org/10.1016/j.gca.2021.05.040>
- Six, J., Elliott, E.T., Paustian, K., 2000. Soil macroaggregate turnover and microaggregate formation: A mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.* 32, 2099–2103. [https://doi.org/10.1016/S0038-0717\(00\)00179-6](https://doi.org/10.1016/S0038-0717(00)00179-6)
- Sodano, M., Lerda, C., Nisticò, R., Martin, M., Magnacca, G., Celi, L., Said-Pullicino, D., 2017. Dissolved organic carbon retention by coprecipitation during the oxidation of ferrous iron. *Geoderma* 307, 19–29. <https://doi.org/10.1016/j.geoderma.2017.07.022>
- Sodano, M., Said-Pullicino, D., Fiori, A.F., Catoni, M., Martin, M., Celi, L., 2016. Sorption of paddy soil-derived dissolved organic matter on hydrous iron oxide-vermiculite mineral phases. *Geoderma* 261, 169–177. <https://doi.org/10.1016/j.geoderma.2015.07.014>
- Thompson, A., Chadwick, O.A., Boman, S., Chorover, J., 2006. Colloid mobilization during soil iron

- redox oscillations. *Environ. Sci. Technol.* 40, 5743–5749. <https://doi.org/10.1021/es061203b>
- Torn, M.S., Trumbore, S.E., Chadwick, O.A., Vitousek, P.M., Hendricks, D.M., 1997. Mineral control of soil organic carbon storage and turnover content were measured by horizon down to the depth at which. *Nature* 389, 3601–3603. <https://doi.org/10.1038/38260>
- Totsche, K.U., Amelung, W., Gerzabek, M.H., Guggenberger, G., Klumpp, E., Knief, C., Lehndorff, E., Mikutta, R., Peth, S., Prechtel, A., Ray, N., Kögel-Knabner, I., 2018. Microaggregates in soils. *J. Plant Nutr. Soil Sci.* 181, 104–136. <https://doi.org/10.1002/jpln.201600451>
- Vogelsang, V., Fiedler, S., Jahn, R., Kaiser, K., 2016. In-situ transformation of iron-bearing minerals in marshland-derived paddy subsoil. *Eur. J. Soil Sci.* 67, 676–685. <https://doi.org/10.1111/ejss.12374>
- Westre, T.E., Kennepohl, P., DeWitt, J.G., Hedman, B., Hodgson, K.O., Solomon, E.I., 1997. A multiplet analysis of Fe K-edge 1s → 3d pre-Edge features of iron complexes. *J. Am. Chem. Soc.* 119, 6297–6314. <https://doi.org/10.1021/ja964352a>
- Wilke, M., Farges, F., Petit, P.E., Brown, G.E., Martin, F., 2001. Oxidation state and coordination of Fe in minerals: An Fe K-XANES spectroscopic study. *Am. Mineral.* 86, 714–730. <https://doi.org/10.2138/am-2001-5-612>
- Winkler, P., Kaiser, K., Kölbl, A., Kühn, T., Schad, P., Urbanski, L., Fiedler, S., Lehndorff, E., Kalbitz, K., Utami, S.R., Cao, Z., Zhang, G., Jahn, R., Kögel-Knabner, I., 2016. Response of Vertisols, Andosols, and Alisols to paddy management. *Geoderma* 261, 23–35. <https://doi.org/10.1016/j.geoderma.2015.06.017>
- Winkler, P., Kaiser, K., Jahn, R., Mikutta, R., Fiedler, S., Cerli, C., Kölbl, A., Schulz, S., Jankowska, M., Schloter, M., Müller-Niggemann, C., Schwark, L., Woche, S.K., Kümmel, S., Utami, S.R., Kalbitz, K., 2019. Tracing organic carbon and microbial community structure in mineralogically different soils exposed to redox fluctuations. *Biogeochemistry* 143, 31–54. <https://doi.org/10.1007/s10533-019-00548-7>
- Wissing, L., Kölbl, A., Häusler, W., Schad, P., Cao, Z.H., Kögel-Knabner, I., 2013. Management-induced organic carbon accumulation in paddy soils: The role of organo-mineral associations. *Soil Tillage Res.* 126, 60–71. <https://doi.org/10.1016/j.still.2012.08.004>
- Wissing, L., Kölbl, A., Schad, P., Bräuer, T., Cao, Z.H., Kögel-Knabner, I., 2014. Organic carbon accumulation on soil mineral surfaces in paddy soils derived from tidal wetlands. *Geoderma* 228–229, 90–103. <https://doi.org/10.1016/j.geoderma.2013.12.012>
- Wissing, L., Kölbl, A., Vogelsang, V., Fu, J.R., Cao, Z.H., Kögel-Knabner, I., 2011. Organic carbon accumulation in a 2000-year chronosequence of paddy soil evolution. *Catena* 87, 376–385. <https://doi.org/10.1016/j.catena.2011.07.007>
- Xue, B., Huang, L., Huang, Y., Zhou, F., Li, F., Kubar, K.A., Li, X., Lu, J., Zhu, J., 2019. Roles of soil organic carbon and iron oxides on aggregate formation and stability in two paddy soils. *Soil Tillage Res.* 187, 161–171. <https://doi.org/10.1016/j.still.2018.12.010>
- Zhang, G., Burgos, W.D., Senko, J.M., Bishop, M.E., Dong, H., Boyanov, M.I., Kemner, K.M., 2011. Microbial reduction of chlorite and uranium followed by air oxidation. *Chemical Geology* 283, 242–250. <https://doi.org/10.1016/j.chemgeo.2011.01.021>
- Zhao, L., Dong, H., Kukkadapu, R., Agrawal, A., Liu, D., Zhang, J., Edelman, R.E., 2013. Biological oxidation of Fe(II) in reduced nontronite coupled with nitrate reduction by *Pseudogulbenkiania* sp. Strain 2002. *Geochim. Cosmochim. Acta* 119, 231–247. <https://doi.org/10.1016/j.gca.2013.05.033>



### Figure captions

**Figure 1.** Soil OC distribution between different aggregate and particle-size fractions in the (a) non-paddy and (b) paddy soils as a function of soil depth. CSa+POM, coarse sand and particulate OM; FSa, fine sand; CSi, coarse silt; FSi, fine silt; Cl, clay. Black bars represent the proportion of fraction OC within microaggregates for the coarse silt (mCSi), fine silt (mFSi), and clay (mCl) fractions. Error bars represent the standard error.

**Figure 2.** Intra:inter-microaggregate ratio of (a) OC and (b) pedogenic Fe ( $Fe_d$ ) associated with coarse silt (CSi), fine silt (FSi), and clay (Cl) particle-size fractions as a function of depth for non-paddy and paddy soils.

**Figure 3.** Distribution of dithionite-citrate-bicarbonate-extractable Fe ( $Fe_d$ ; a, b), oxalate-extractable Fe ( $Fe_o$ ; c, d), and crystalline Fe (hydr)oxides ( $Fe_c$ ; e, f) between different aggregate and particle-size fractions in the non-paddy (a, c, e) and paddy (b, d, f) soils as a function of soil depth. CSa+POM, coarse sand and particulate OM; FSa, fine sand; CSi, coarse silt; FSi, fine silt; Cl, clay. Black bars represent the proportion of fraction pedogenic Fe within microaggregates for the coarse silt (mCSi), fine silt (mFSi), and clay (mCl) fractions.

**Figure 4.** Changes in the Fe-K edge XANES pre-edge centroid position for (a) fine sand (FSa), inter and intra-microaggregate coarse silt (CSi and mCSi), (b) inter and intra-microaggregate fine silt (FSi and mFSi), and (c) inter and intra-microaggregate clay (Cl and mCl) fractions in non-paddy and paddy soils as a function of soil depth.

**Figure 5.**  $k^3$ -weighted EXAFS spectra of the coarse silt (CSi), fine silt (FSi), clay (Cl), coarse silt in aggregates (mCSi), fine silt in aggregates (mFSi), clay in aggregates (mCl) fractions of Ap1, Ap2, Bgw horizons of non-paddy soil and Arp1, Arp2, Arpd, Brd1 and Brd2 horizons of paddy soil. The red dotted lines represent the linear combination fits of the sample spectra ( $n=3$ ). R-factors for the fits are displayed.

**Figure 6.** High-resolution WT plots of of the coarse silt (CSi), fine silt (FSi), clay (Cl), coarse silt in aggregates (mCSi), fine silt in aggregates (mFSi), clay in aggregates (mCl) fractions of non- paddy (a) and paddy (b) soils. Data are plotted as a function of  $k$  ( $\text{\AA}^{-1}$ )

on the x axis and  $R$  (Å) on the y axis in the range 2.2–4.0 (Å).

Figure 1






Figure 2




Figure 3




Figure 4




Figure 5





Figure 6

a)





**Table 1.** Ratio of oxalate-extractable Fe (Fe<sub>o</sub>) to dithionite-citrate-bicarbonate-extractable Fe (Fe<sub>d</sub>) in different particle-size fractions for each horizon in the non-paddy and paddy soils.

Soil/Horizon	CSa+POM	FSa	CSi + mCSi	FSi + mFSi	Cl + mCl
<b>Non-paddy</b>					
Ap1	0.5	0.7	0.9	0.4	0.3
Ap2	0.4	0.4	0.6	0.3	0.2
Bgw	0.5	0.2	0.3	0.1	0.3
<b>Paddy</b>					
Arp1	0.8	0.7	0.2	0.3	0.5
Arp2	0.8	1.0	0.3	0.2	0.8
Arpd	0.7	0.8	0.5	0.5	0.2
Brd1	0.5	0.7	0.4	0.3	0.2
Brd2	0.7	1.0	0.6	0.2	0.2

CSa+POM, coarse sand and particulate OM; FSa, fine sand; CSi, inter-microaggregate coarse silt; mCSi, intra-microaggregate coarse silt; FSi, inter-microaggregate fine silt; mFSi, intra-microaggregate fine silt; Cl, inter-microaggregate clay; mCl, intra-microaggregate clay fraction.

**Table 2.** Results for the LCF performed on the Fe K-edge XANES data of the fine sand (FSa), coarse silt (CSi), fine silt (FSi), clay (Cl), coarse silt in aggregates (mCSi), fine silt in aggregates (mFSi), clay in aggregates (mCl) fractions of Ap1, Ap2, Bgw horizons of non-paddy soil and Arp1, Arp2, Arpd, Brd1, and Brd2 horizons of paddy soil.

	Chi2	Component 1	%	Component 2	%	Component 3	%	Component 4	%	Sum
Ap1 FSa	0.033087096	Chlorite	32	Siderite	26	Illite	21	Hematite	20	100
Ap1 CSi	0.019310413	Chlorite	26	Illite	48	Hematite	21	Magnetite	5	100
Ap1 mCSi	0.008858912	Chlorite	22	Illite	66	Hematite	13			100
Ap1 FSi	0.008196354	Chlorite	27	Smectite	42	Nontronite	19	Lepidocrocite	11	100
Ap1 mFSi	0.006696096	Chlorite	26	Smectite	47	Illite	23	FeIII citrate	4	100
Ap1 Cl	0.010688951	Chlorite	8	Jacobsite	13	Nontronite	79			100
Ap1 mCl	0.010184488	Chlorite	8	Jacobsite	13	Nontronite	79			100
Ap2 FSa	0.039485435	Chlorite	33	Siderite	33	Illite	13	Hematite	20	100
Ap2 CSi	0.022535003	Chlorite	27	Illite	46	Hematite	27			100
Ap2 mCSi	0.027156644	Chlorite	26	Illite	37	Hematite	23	Magnetite	14	100
Ap2 FSi	0.006497925	Chlorite	25	Illite	26	Smectite	42	FeIII citrate	7	100
Ap2 mFSi	0.427969307	Chlorite	12	Jacobsite	38	Nontronite	50			100
Ap2 Cl	0.009034114	Chlorite	11	Smectite	65	Lepidocrocite	24			100
Ap2 mCl	0.013101914	Chlorite	11	Nontronite	84	Lepidocrocite	5			100
Bgw FSa	0.038458154	Chlorite	34	Siderite	29	Illite	14	Hematite	22	100
Bgw CSi	0.032032481	Chlorite	26	Illite	36	Hematite	27	Magnetite	11	100
Bgw mCSi	0.026492885	Chlorite	25	Illite	34	Hematite	26	Magnetite	14	100
Bgw FSi	0.007402408	Chlorite	26	Smectite	61	Goethite	9	FeIII citrate	4	100
Bgw mFSi	0.007050961	Chlorite	23	Smectite	25	Illite	42	FeIII citrate	9	100
Bgw Cl	0.013186125	Chlorite	8	Nontronite	77	Ferrihydrite	15			100



Bgw mCl	0.014873833	Chlorite	10	Nontronite	83	FeIII citrate	7	100
Arp1 FSa	0.027628912	Chlorite	33	Siderite	29	Illite	19	100
Arp1 CSi	0.01276442	Chlorite	27	Illite	64	Magnetite	9	100
Arp1 mCSi	0.009723367	Chlorite	24	Illite	68	Hematite	9	100
Arp1 FSi	0.009919808	Chlorite	29	Illite	26	Smectite	22	100
Arp1 mFSi	0.008831991	Chlorite	29	Illite	42	Smectite	22	100
Arp1 Cl	0.010773327	Chlorite	19	Smectite	80	FeIII citrate	1	100
Arp1 mCl	0.010542052	Chlorite	16	Jacobsite	5	Nontronite	79	100
Arp2 FSa	0.034934128	Chlorite	33	Siderite	32	Hematite	22	100
Arp2 CSi	0.01158174	Chlorite	27	Illite	60	Hematite	11	100
Arp2 mCSi	0.047753368	Chlorite	26	Illite	40	Magnetite	34	100
Arp2 FSi	0.00624999	Chlorite	26	Illite	52	Smectite	22	100
Arp2 mFSi	0.006762771	Chlorite	26	Illite	58	Smectite	16	100
Arp2 Cl	0.008425091	Chlorite	19	Smectite	74	Lepidocrocite	7	100
Arp2 mCl	0.007167858	Chlorite	18	Smectite	65	Ferrihydrite	17	100
Arpd FSa	0.032190435	Chlorite	34	Siderite	32	Illite	14	100
Arpd CSi	0.017333956	Chlorite	27	Illite	47	Hematite	18	100
Arpd mCSi	0.005592672	Chlorite	19	Illite	75	Fe III citrate	6	100
Arpd FSi	0.007043644	Chlorite	28	Smectite	33	Nontronite	25	100
Arpd mFSi	0.008959662	Chlorite	28	Illite	44	Smectite	21	100
Arpd Cl	0.008911101	Chlorite	12	Jacobsite	11	Nontronite	78	100
Arpd mCl	0.009600908	Chlorite	11	Jacobsite	12	Nontronite	77	100
Brd1 FSa	0.029485302	Chlorite	32	Siderite	27	Illite	19	100
Brd1 CSi	0.008067183	Chlorite	20	Illite	63	Hematite	17	100
Brd1 mCSi	0.004323186	Chlorite	19	Illite	53	Smectite	14	100
Brd1 FSi	0.005927233	Chlorite	23	Smectite	35	Nontronite	32	100
Brd1 mFSi	0.066203605	Chlorite		Smectite		Nontronite		
						FeIII citrate	9	
						FeIII citrate		

Brd1 Cl	0.015676096	Chlorite	10	Nontronite	84	FeIII citrate	6		100
Brd1 mCl	0.014728171	Chlorite	9	Nontronite	83	Fe(III)-citrate	8		100
Brd2 FSa	0.031482016	Chlorite	32	Siderite	31	Illite	16	Hematite	20
Brd2 CSi	0.00581651	Chlorite	18	Illite	75	Maghemite	7		100
Brd2 mCSi	0.022010179	Chlorite	25	Illite	39	Hematite	23	Magnetite	12
Brd2 FSi	0.008325141	Chlorite	26	Smectite	57	Goethite	17		100
Brd2 mFSi	0.007011355	Chlorite	25	Illite	28	Smectite	40	FeIII citrate	6
Brd2 Cl	0.012284473	Chlorite	9	Nontronite	81	Ferrihydrite	11		100
Brd2 mCl	0.010851395	Chlorite	10	Smectite	66	Lepidocrocite	24		100

**Table 3.** Results for the four-components LCF performed on the  $k^3$ - weighted Fe K-edge EXAFS data of the fine sand (FSa) coarse silt (CSi), fine silt (FSi), clay (Cl), coarse silt in aggregates (mCSi), fine silt in aggregates (mFSi), clay in aggregates (mCl) fractions of Ap1, Ap2, Bgw horizons of non-paddy soil and Arp1, Arp2, Arpd, Brd1, and Brd2 horizons of paddy soil. Results are expressed in percentage of the fitting components corresponding to experimental spectra of model compounds reported by O'Day et al. (2004) chlorite, ferrihydrite, lepidocrocite, and Fe(III)-citrate.

	<b>R-factor</b>	<b>Component 1</b>	<b>%</b>	<b>Component 2</b>	<b>%</b>	<b>Component 3</b>	<b>%</b>	<b>Component 4</b>	<b>%</b>	<b>Sum</b>
Ap1 FSa	0.079749	Chlorite	44	Magnetite	8	Goethite	33	Lepidocrocite	15	100
Ap1 CSi	0.0429419	Chlorite	46	Ferrihydrite	21	Lepidocrocite	17	FeIII citrate	17	100
Ap1 mCSi	0.0343152	Chlorite	44	Ferrihydrite	20	Lepidocrocite	15	FeIII citrate	21	100
Ap1 FSi	0.0339569	Chlorite	35	Ferrihydrite	19	Lepidocrocite	13	FeIII citrate	33	100
Ap1 mFSi	0.0289672	Chlorite	34	Ferrihydrite	22	Lepidocrocite	13	FeIII citrate	31	100
Ap1 Cl	0.0350046	Chlorite	13	Ferrihydrite	31	Lepidocrocite	14	FeIII citrate	43	100
Ap1 mCl	0.0339516	Chlorite	13	Ferrihydrite	31	Lepidocrocite	13	FeIII citrate	42	100
Ap2 FSa	0.0945382	Chlorite	49	Illite	19	Hematite	13	Lepidocrocite	19	100
Ap2 CSi	0.0425902	Chlorite	49	Ferrihydrite	20	Lepidocrocite	16	FeIII citrate	15	100
Ap2 mCSi	0.0529809	Chlorite	49	Ferrihydrite	24	Lepidocrocite	17	FeIII citrate	11	100
Ap2 FSi	0.0316848	Chlorite	35	Ferrihydrite	22	Lepidocrocite	13	FeIII citrate	30	100
Ap2 mFSi	0.0324117	Chlorite	36	Ferrihydrite	21	Lepidocrocite	12	FeIII citrate	31	100
Ap2 Cl	0.0406223	Chlorite	15	Ferrihydrite	36	Lepidocrocite	11	FeIII citrate	38	100
Ap2 mCl	0.0405023	Chlorite	13	Ferrihydrite	35	Lepidocrocite	12	FeIII citrate	40	100
Bgw FSa	0.0561351	Chlorite	47	Illite	14	Hematite	14	Lepidocrocite	25	100
Bgw CSi	0.0293375	Chlorite	48	Ferrihydrite	24	Lepidocrocite	19	Fe III citrate	9	100
Bgw mCSi	0.0565707	Chlorite	48	Ferrihydrite	24	Lepidocrocite	17	Fe III citrate	11	100
Bgw FSi	0.0119554	Chlorite	30	Ferrihydrite	25	Lepidocrocite	15	Fe III citrate	30	100
Bgw mFSi	0.0302615	Chlorite	32	Ferrihydrite	25	Lepidocrocite	14	Fe III citrate	29	100
Bgw Cl	0.0484722	Chlorite	4	Ferrihydrite	42	Lepidocrocite	17	FeIII citrate	37	100

Bgw mCl	0.02219	Chlorite	12	Ferrihydrite	43	Lepidocrocite	15	Fe III citrate	31	100
Arp1 FSa	0.1303528	Chlorite	45	Illite	16	Ferrihydrite	28	Lepidocrocite	12	100
Arp1 CSi	0.0431159	Chlorite	54	Ferrihydrite	8	Lepidocrocite	14	FeIII citrate	23	100
Arp1 mCSi	0.0394779	Chlorite	46	Ferrihydrite	15	Lepidocrocite	16	FeIII citrate	23	100
Arp1 FSi	0.0128322	Chlorite	43	Ferrihydrite	9	Lepidocrocite	13	FeIII citrate	34	100
Arp1 mFSi	0.0117425	Chlorite	45	Ferrihydrite	13	Lepidocrocite	13	FeIII citrate	29	100
Arp1 Cl	0.0513891	Chlorite	26	Ferrihydrite	14	Lepidocrocite	14	FeIII citrate	46	100
Arp1 mCl	0.0461159	Chlorite	25	Ferrihydrite	16	Lepidocrocite	14	FeIII citrate	45	100
Arp2 FSa	0.1116709	Chlorite	50	Magnetite	10	Hematite	13	Lepidocrocite	27	100
Arp2 CSi	0.0604463	Chlorite	51	Ferrihydrite	13	Lepidocrocite	17	FeIII citrate	18	100
Arp2 mCSi	0.0320848	Chlorite	56	Ferrihydrite	16	Lepidocrocite	19	FeIII citrate	9	100
Arp2 FSi	0.0339822	Chlorite	46	Ferrihydrite	19	Lepidocrocite	11	FeIII citrate	24	100
Arp2 mFSi	0.0130281	Chlorite	46	Ferrihydrite	15	Lepidocrocite	14	FeIII citrate	26	100
Arp2 Cl	0.0395401	Chlorite	25	Ferrihydrite	20	Lepidocrocite	14	FeIII citrate	41	100
Arp2 mCl	0.0545088	Chlorite	23	Ferrihydrite	21	Lepidocrocite	15	FeIII citrate	42	100
Arpd FSa	0.1353997	Chlorite	44	Magnetite	11	Ferrihydrite	35	FeIII citrate	10	100
Arpd CSi	0.0491993	Chlorite	41	Ferrihydrite	23	Lepidocrocite	19	FeIII citrate	18	100
Arpd mCSi	0.0141161	Chlorite	41	Ferrihydrite	21	Lepidocrocite	18	FeIII citrate	21	100
Arpd FSi	0.0246098	Chlorite	36	Ferrihydrite	21	Lepidocrocite	16	FeIII citrate	27	100
Arpd mFSi	0.0442138	Chlorite	41	Ferrihydrite	13	Lepidocrocite	16	FeIII citrate	31	100
Arpd Cl	0.035265	Chlorite	18	Ferrihydrite	26	Lepidocrocite	18	FeIII citrate	38	100
Arpd mCl	0.0274329	Chlorite	17	Ferrihydrite	28	Lepidocrocite	20	FeIII citrate	35	100
Brd1 FSa	0.0858789	Chlorite	49	Nontronite	27	Hematite	13	Magnetite	11	100
Brd1 CSi	0.0526621	Chlorite	35	Ferrihydrite	29	Lepidocrocite	16	FeIII citrate	20	100
Brd1 mCSi	0.0123727	Chlorite	31	Ferrihydrite	31	Lepidocrocite	16	FeIII citrate	22	100
Brd1 FSi	0.2231373	Chlorite	66	Goethite	21	Lepidocrocite		FeIII citrate	13	100
Brd1 mFSi	0.011673	Chlorite	27	Ferrihydrite	30	Lepidocrocite	13	FeIII citrate	30	100

Brd1 Cl	0.0242448	Chlorite	10	Ferrihydrite	40	Lepidocrocite	19	FeIII citrate	31	100
Brd1 mCl	0.0230121	Chlorite	9	Ferrihydrite	41	Lepidocrocite	19	FeIII citrate	31	100
Brd2 FSa	0.0843658	Chlorite	51	Nontronite	31	Hematite	10	Magnetite	8	100
Brd2 CSi	0.0699578	Chlorite	36	Ferrihydrite	24	Lepidocrocite	12	Fe III citrate	29	100
Brd2 mCSi	0.0539459	Chlorite	47	Ferrihydrite	24	Lepidocrocite	14	Lepidocrocite	15	100
Brd2 FSi	0.0467505	Chlorite	29	Ferrihydrite	24	Lepidocrocite	11	Fe III citrate	36	100
Brd2 mFSi	0.0382681	Chlorite	36	Ferrihydrite	23	Lepidocrocite	11	Fe III citrate	30	100
Brd2 Cl	0.1897291	Chlorite	41	Goethite	22	Magnetite	15	FeIII citrate	22	100
Brd2 mCl	0.0416766	Chlorite	11	Ferrihydrite	40	Magnetite	6	Fe III citrate	43	100

