

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Bioprotection strategies in winemaking

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1851245> since 2022-06-22T15:24:09Z

Published version:

DOI:10.1016/j.ijfoodmicro.2022.109532

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Bioprotection strategies in winemaking**

2

3 Paola Di Gianvito, Vasileios Englezos, Kalliopi Rantsiou, Luca Cocolin

4

5 Università degli Studi di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Largo

6 Braccini 2, 10095 Grugliasco, Italy.

7

8 *Corresponding author: Luca Cocolin, Fax: +39-011-6708553, e-mail: lucasimone.cocolin@unito.it.

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27 **Abstract**

28

29 Worldwide the interest for biological control of food spoilage microorganisms has significantly
30 increased over the last decade. Wine makes no exception to this trend, as consumer demands for
31 wines free of preservatives that are considered negative for human health, increase. Biological control
32 during wine fermentation aims at producing high quality wines, while minimizing, or even
33 eliminating, the use of chemical additives. Its success lies in the inoculation of microorganisms to
34 prevent, inhibit or kill undesired microbes, therefore maintaining wine spoilage at the lowest level.
35 The food industry already makes use of this practice, with dedicated commercial microbes already
36 on the market. In winemaking, there are commercial microbes currently under investigation,
37 particularly with the aim to reduce or replace the use of sulfur dioxide. In this review, the potential
38 of wine yeasts and lactic acid bacteria as bioprotection agents and their mechanisms of action during
39 wine fermentation are presented.

40

41 **Keywords:** Bioprotection mechanisms; Wine fermentation; Antagonism strategies; Competition
42 strategies; Yeasts; Lactic Acid bacteria

43

44

45

46

47

48

49

50

51

52

53 1. Introduction

54

55 There is a growing concern about the negative effects on human health of chemical additives
56 used in foods and drinks to improve their stability and shelf life. For this reason, several research
57 lines are focusing on the development of alternative strategies to avoid spoilage. Bioprotection (or
58 biopreservation or biological control) fits well in this new concept of spoilage prevention, and several
59 microorganisms or their antimicrobial products have already been identified as bioprotection agents
60 (Dokka et al., 2018; Mewa-Ngongang et al., 2019; Mukherjee et al., 2020; Pu et al., 2014; Singh et
61 al., 2018;). This practice is widely used, mainly in agriculture and food industries for the protection
62 of fruits from postharvest spoilage microorganisms, and the extension of the shelf life of food
63 products (Ferraz et al., 2019; Keswani et al., 2019). For example, a commercial product based on
64 *Aureobasidium pullulans* was developed to protect grapes, strawberry and tomato against
65 *Penicillium*, *Botrytis* and *Monilinia*, while a product based on *Metschnikowia fructicola* is used to
66 protect strawberry, blueberry, grape, stone fruit and pome against *Botrytis*, *Penicillium*, *Rhizopus*,
67 *Aspergillus* and *Monilinia* (Zhang et al., 2020). To date, there is a patent (WO 2015/110484) related
68 to the exploitation of a *Lactiplantibacillus plantarum* strain as bioprotective agent against undesired
69 microorganisms in wine (Krieger-Weber et al., 2020). Different genera of bacteria (e.g.
70 *Lactiplantibacillus*, *Bacillus*, *Pseudomonas*, *Streptomyces*), fungi (*Trichoderma*) and yeasts
71 (belonging mainly to the genera, *Candida*, *Cryptococcus* and *Aureobasidium*) are commercially
72 available for these purposes (Droby et al., 2016; López-Seijas et al., 2020; Mukherjee et al., 2020).

73 The bioprotection strategy consists of the inoculation of living microorganisms (Bio
74 Protective Cultures, BPCs), or the addition of their metabolites (Bio Protective Metabolites, BPMs),
75 in purified form, during the production of the food or there after. These microorganisms prevent food
76 microbial spoilage through different bioprotection mechanisms that can be divided in passive
77 (competition for space, nutrient and oxygen) and active antagonistic strategies (production of
78 antimicrobial molecules) (Pandin et al., 2017). The addition of BPCs during the early stages of the

79 production process could also enhance the characteristics of fermented foods, such as flavour, texture
80 and nutritional value (Gaggia et al., 2011).

81 In the wine industry, the most common chemical additive is sulfur dioxide (SO₂). This
82 compound is considered an essential tool for winemakers, due to its low-cost and its antioxidant and
83 antimicrobial properties against a wide spectrum of microorganisms (Roullier-Gall et al., 2017).
84 Therefore, its presence during both the fermentation and the storage of the wine is desired to avoid
85 spoilage. In spite of the advantages, high doses of SO₂ may cause consumer health problems,
86 particularly in sensitive individuals (Guerrero and Cantos-Villar, 2015). For this reason, the World
87 Health organization (WHO) encourages alternative methods to reduce, or even eliminate, its use in
88 wine production. Consequently, several alternative technologies able to control microorganisms have
89 been investigated. These include ultrasound, ultraviolet radiation, pulsed electric field, electrolyzed
90 water, high hydrostatic pressure treatments, or the addition of lysozyme, sorbic acid, dimethyl
91 dicarbonate and chitosan (Guerrero and Cantos-Villar, 2015). However, a definitive substitute for
92 SO₂ has not yet been found, especially for wines undergoing long-term storage (Giacosa et al., 2019).

93 The application of bioprotection strategies in viticulture has increased significantly in the last
94 years, with studies investigating its applicability against grapevine trunk diseases (Mondello and
95 Songy, 2018), the fungal pathogen *Botrytis cinerea* causing bunch (Carbó et al., 2018; Garrido et al.,
96 2017; Jacometti et al., 2010) and sour rot (Carbó et al., 2018) in the vineyard (Nardi et al., 2020).
97 However, little is known about the application of bioprotection strategies during or after the
98 completion of the fermentation process (Simonin et al., 2018). Bioprotective strategies found in wine
99 yeasts and Lactic Acid Bacteria (LAB) are represented in Fig. 1. In this review, the current knowledge
100 about bioprotection opportunities and mechanisms exerted by wine yeasts and LAB, during
101 fermentation to control spoilage microorganisms, will be discussed.

102

103 **2. Bioprotection mechanisms**

104

105 Wine microorganisms form complex ecological interactive webs, which result in the
106 dominance of a specific species or strains able to determine the final quality of wine. In inoculated
107 wine fermentations, the use of a starter culture aims to establish dominance of the inoculated culture
108 and determine wine characteristics. Conversely, the BPCs aim to prevent the growth of the spoilage
109 microorganisms, without affecting the performances of useful microorganisms responsible for the
110 alcoholic/malolactic fermentation (Nardi, 2020). Furthermore, BPCs may also be directly responsible
111 for wine fermentation (Table 1). Understanding the antagonistic mechanisms that confer
112 bioprotection activity to the inoculated microorganisms is fundamental for their successful
113 application as BPCs. These competition strategies can be either passive or active (Bauer et al., 2018)
114 as shown in Fig. 2. The difference is that passive competition strategies are exerted by all
115 microorganisms, while active competition strategies are performed only by some microorganisms
116 that, consequently, result stronger than the other and able to dominate the fermentation process.

117

118 *2.1. Passive competition strategies*

119

120 Passive competition strategies comprise all the mechanisms that are beneficial including
121 strategies to take away resources from competitors (exploitative competition), or to achieve a superior
122 positioning within the niche (Parijs and Steenackers, 2018). During wine fermentation, the factors
123 that mainly modulate the fitness advantage of some species above others are the ability to rapidly
124 uptake nutrients (carbohydrates, nitrogen, vitamins, sterols and microelements) and oxygen or to
125 colonize the space (Curiel et al., 2017; Petigonnnet et al., 2019; Prior et al., 2019; Rollero et al. 2018;
126 Siedler et al., 2019; Taillandier et al. 2014).

127

128 *2.1.2. Nutrients competition*

129

130 In a specific environment all microorganisms compete with each other for nutrients necessary
131 for their growth (Liu et al., 2015). Passive competitive microorganisms are able to consume the
132 substrates faster and in a more efficient way than the competitors, producing higher levels of biomass
133 (Boynton et al., 2019). Proteomic and transcriptomic studies have revealed that, during mixed wine
134 fermentation, an increased resource uptake (glucose, nitrogen, oxygen, sterols, copper, iron and
135 vitamins) occurring in several yeast strains, compared to the respective control pure culture
136 fermentations. These yeasts belong to *Saccharomyces cerevisiae* (Alonso-del-Real et al., 2019;
137 Barbosa et al., 2015; Curiel et al., 2017; Kosel et al., 2017; Peng et al., 2019; Rossouw et al., 2012;
138 Shekhawat et al., 2019; Tronchoni et al., 2017;), *Torulaspora delbrueckii* (Prior et al., 2019; Su et al.,
139 2020), even in presence of different nitrogen sources (Roca-Mesa et al., 2020), and *Lachancea*
140 *thermotolerans* (Shekhawat et al. 2019). Furthermore, nutrient competition was identified as the
141 major antagonistic strategy exhibited by *Lpb. plantarum* (Sielder et al., 2020). A metagenomic
142 analysis revealed that during wine fermentation, this species has the largest number of
143 phosphotransferase systems (the major mechanism used by bacteria for carbohydrates uptake) among
144 the bacteria observed in wine (Melkonian et al., 2019). The ability of the inoculated microorganisms
145 to subtract high levels of nutrients from the medium, could increase the microbial stability in the
146 wines at the end of the fermentation process by inhibiting the growth of unwanted microorganisms
147 and reduce or even avoid the use of SO₂. Since, residual nitrogen levels could favor the growth of
148 undesirable microorganisms in unwanted steps of wine production chain and could act as precursors
149 for the formation of biogenic amines (Restuccia et al., 2018). However, particular attention should
150 be paid in order to find the most appropriate strain selection that fits with the fermentation conditions,
151 since low nitrogen concentration may lead to stuck or sluggish fermentations (Bisson et al., 1999).

152 Competition for nutrients can also be achieved by the secretion of extracellular molecules
153 such as digestive enzymes, or siderophores (nutrient exclusion) (Ghoul and Mitri, 2016). In this
154 context, antibacterial and antifungal activity of *Metschnikovia pulcherrima* and *Metschnikovia*
155 *fructicola* was mainly attributed to the iron depletion caused by the insoluble maroon-red

156 pulcherrimin pigment (Lachance, 2016; Sipiczki, 2006; 2020). In particular, this antimicrobial feature
157 is the result of the reduction of the iron in the medium, due to the precipitation of iron (III) ions,
158 caused the precursor of the pulcherrimin pigment called pulcherriminic acid. As a result, low iron
159 levels in the medium could limit the growth of the microorganisms that require this compound for
160 their growth. This mechanism was found to have an inhibitory effect against the following
161 microorganisms: *Candida tropicalis*, *Candida albicans*, *Brettanomyces/Dekkera*, *Hanseniaspora* and
162 *Pichia* genera, *B. cinerea* and *Penicillium* (Morata et al., 2019a). While, contradictory results were
163 observed against *S. cerevisiae* (Kantor et al., 2015, Melvyda et al., 2020, Oro et al., 2019). This
164 behaviour let us to hypothesize a strain specific inhibition. Therefore, careful selection of *S. cerevisiae*
165 strains should be performed to find the most suitable partner of *M. pulcherrima* in mixed culture
166 fermentations to reduce ethanol content and enhance the aroma volatile fraction of the wines (Jolly
167 et al., 2014). However, there was not a clear correlation between the antagonistic activity and the iron
168 deprivation by pulcherrimin (Gore-Lloyd et al., 2019; Saravanakumar et al., 2008). In a recent study,
169 a genome comparison between a *M. pulcherrima* wild type and three pigmentless mutants, allowed
170 the identification of a point mutation in the *MPUL0C08850* gene, which encodes for an ortholog of
171 *S. cerevisiae SNF2* gene. The resultant truncated form of this transcriptional regulator caused a down-
172 regulation of the transcription of *PUL* genes (pulcherrimin biosynthesis and utilisation) explaining
173 why the colourless strains showed an antimicrobial behaviour even if it was less than the wild type
174 (Gore-Lloyd et al., 2019). These authors speculated that the antagonistic activity of *M. pulcherrima*
175 was due to the combination of the effect of pulcherrimin and diverse proteins regulated by *SNF2* such
176 as genes encoding different types of transporters or secreted proteases (Gore-Lloyd et al., 2019).

177 A similar system was also identified in many Lactobacilli, including *Lpb. plantarum*. In
178 particular, Sielder et al. (2020) demonstrated that the Lactobacilli showed an overexpression of a
179 manganese transporter (*mntH1* gene) during milk fermentation. The depletion of this microelement
180 in the media resulted in the reduction of the growth of several yeasts (also present in wine) such as
181 *Debariomyces hansenii* and *T. delbrueckii*. These authors found that for *T. delbrueckii* the addition

182 of manganese restored the growth only partially and hypothesized that other mechanisms
183 (antimicrobial compounds production) were involved in the antagonistic interactions.

184

185 *2.1.3. Space competition*

186

187 Another passive competition phenotype is the ability to colonize the space where competitors are
188 established (Ghool and Mitri, 2016) or grow faster than other microorganisms. With this purpose,
189 microorganisms are able of change their lifestyle forming multicellular structures like biofilms
190 (Váchová and Palková, 2018). These structures may exhibit different phenotypic characteristics,
191 compared to free-floating cells. Usually, after the adhesion of single cells to the host surface, the
192 biofilm formation determines the modification of the cell wall properties due to the secretion of
193 extracellular matrix and often the formation of hyphae and pseudohyphae (Freimoser et al., 2019).
194 The biofilm formation is considered a desired characteristic in biocontrol yeasts and has been widely
195 investigated. For example, *M. pulcherrima* and *Pichia kluyveri* are able to form biofilms on grape
196 berries surface (Cordero-Bueso et al., 2017; Parafati et al., 2015; Pawlikowska et al., 2019). This
197 ability confers a great competitive advantage to these yeasts to colonize both intact and damaged
198 grapes as well as establishing themselves in grape juice and starting the fermentation process even in
199 harsh conditions such as those found in desiccated grapes (Cordero-Bueso et al., 2017).
200 Microorganisms with ability to form biofilms are generally characterised by a higher competitiveness
201 against pathogens due to the higher environment persistence and higher biocontrol activity (Freimoser
202 et al., 2019). Furthermore, it has been demonstrated that *Oenococcus oeni* is able to produce
203 exopolysaccharides and form biofilms on cellar surfaces (Dimopoulou et al., 2015). This organization
204 is formed by stress-tolerant cells able for efficient malolactic fermentation under winemaking
205 conditions (Bastard et al., 2016).

206

207 *2.1.4. Competition for oxygen*

208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233

The presence of oxygen during the fermentation process is considered as one of the key factors that regulate the presence and population of yeast species. This molecule could positively affect ethanol tolerance and the fermentation performance of yeast cells, through its key role in the production of sterols and unsaturated fatty acids (Albergaria and Arneborg, 2016). Grape must contains relatively low dissolved oxygen levels and becomes rapidly anaerobic due to the fast consumption by the microbial consortium and the grape-derived polyphenol oxidase (Romano et al., 2019). Therefore, the concentration of oxygen may have an important role in the survival and dominance of a specific yeast during the fermentation process. Indeed, the low levels could promote the growth of species that are able to grow in anaerobic conditions, like *S. cerevisiae* (Hansen et al., 2001). While, on the other hand low oxygen levels can influence the growth and death rate of several non-*Saccharomyces* yeast species like *T. delbrueckii*, *L. thermotolerans*, *Starmerella bacillaris* and *Hanseniaspora vinai* and reduce the competitiveness against *S. cerevisiae* (Englezos et al., 2018a, Hansen et al., 2001, Yan et al., 2020). Oxygen addition is also believed to enhance the impact of non-*Saccharomyces* yeasts on the chemical and aroma composition of the wines produced by mixed culture fermentations (Englezos et al., 2018a, Hansen et al., 2001, Yan et al., 2020). While the limited addition of oxygen at the beginning of mixed fermentations with *M. pulcherrima* or *Starm. bacillaris* and *S. cerevisiae* could help to reduce the ethanol content in wines due to the higher sugar consumption of the first yeast species (Varela et al., 2017). However, a fine tuning of oxygen addition is required, since excessive oxygenation might lead to stuck fermentation due to increased antagonism between non-*Saccharomyces* and *S. cerevisiae* yeasts, as previously observed for *H. vinai* and *S. cerevisiae* (Yan et al., 2020). Interestingly, the use of selected yeasts species or strains within species could potentially protect the wine against oxidation due to the rapid consumption of oxygen, as previously demonstrated for *T. delbrueckii* (Simonin et al., 2018). The effect of oxygen on yeast interactions during mixed fermentations and as consequence on wine characteristics needs further investigation.

234 2.2. Active competition strategies

235

236 Generally, active competition includes the strategies exploited to confer a disadvantage to the
237 unwanted species through a direct and active interference (Ghoul and Mitri, 2016) due to the
238 production of antimicrobial compounds or through a contact dependent inhibition (Parijs and
239 Steenackers, 2018). During fermentation, yeasts release ethanol that is the most studied antimicrobial
240 compound. Its toxicity on cells is well understood (Gao and Fleet, 1988). Yeasts are also able to
241 release SO₂ as result of their metabolism (Andorrà et al., 2018). Some *S. cerevisiae* strains can
242 produce more than 100 mg/L of this compound (Rauhut, 2017), while information regarding its
243 production by non-*Saccharomyces* yeasts are scarce, probably because these yeasts are generally
244 more sensitive to SO₂ than *S. cerevisiae* (Jolly et al., 2014). However, some spoilage yeasts namely
245 *Zygosaccharomyces bailii* and *Brettanomyces* spp. are more tolerant to these conditions than *S.*
246 *cerevisiae* and therefore are still detected in wines with relatively high SO₂ levels.

247 Several studies highlighted that an array of other antimicrobial compounds are produced in
248 wine environment by yeasts (Albergaria and Arneborg, 2016; Balmaseda et al., 2018; Liu et al., 2017)
249 and bacteria (Bartle et al, 2019; Siedler et al., 2019) of oenological interest. These molecules differ
250 and possess diverse inhibition mechanisms. Antimicrobial compounds are actually subjected to an
251 extensive research with the purpose to exploit their action during wine fermentation. However, it is
252 important to consider that these compounds are often produced after the logarithmic stage and
253 therefore their competitive ability is strictly correlated with the growth performance of the inoculated
254 strain (Mazzucco et al., 2019; Singh, 2018; Villaba et al., 2016). However, the use of these
255 compounds for commercial purpose cannot be authorised without the approval by the International
256 Office of Vine and Wine (OIV) and/or the national regulators (Mehlomakulu et al., 2015). On the
257 other hand, the yeasts producing these compounds can be used if their origin is oenological. In the
258 next sections, the active antagonistic strategies used by yeasts and LAB that confer a possible
259 bioprotection in wine will be discussed. These molecules can be grouped according to the chemical

260 groups they possess. In particular, nitrogen antimicrobial compounds are antimicrobial peptides,
261 killer toxins, bacteriocins and lytic enzymes as listed in the next section.

262

263 2.2.2. Antimicrobial peptides and proteins

264

265 2.2.1.1 Antimicrobial peptides

266

267 Antimicrobial peptides (AMPs) are low molecular weight oligopeptides with a varying
268 number (from five to over a hundred) of amino acids and high sequence variability. Furthermore, the
269 majority of AMPs present charged and hydrophobic amino acids at physiological pH, defining the
270 amphipathic nature of these molecules (Zhang et al., 2019). AMPs are evolutionary conserved in the
271 genome and have a broad antimicrobial spectrum of action from viruses to parasites (Bahar and Ren,
272 2013). Up to date, more than 5,000 AMPs have been discovered from prokaryotes (e.g., bacteria) and
273 eukaryotes (e.g., protozoa, fungi, plants, insects, and animals) or synthesized (Zhao et al., 2013).

274 Several studies reported that *S. cerevisiae* is able to secrete a proteinaceous antimicrobial
275 compound able to inhibit *O. oeni* growth and as a consequence malolactic fermentation (Comitini et
276 al., 2005; Osborne and Edwards, 2007; Mendoza et al., 2010; Nehme et al., 2010; Rizk et al., 2016).
277 Furthermore, the AMP “saccharomycin” produced by *S. cerevisiae* was found to act as a natural
278 biocide (2-10kDa) against several wine-related non-*S. cerevisiae* yeasts such as *Hanseniaspora*
279 *uvarum* (Pérez-Nevado et al., 2006), *Brettanomyces/D. bruxellensis* and LABs (Branco et al., 2014).
280 These authors demonstrated that saccharomycin is comprises by two anionic (isoelectric point of
281 4.37) peptides (AMP1 and AMP2/3) able to alter intracellular pH, membrane permeability and
282 cultivability of non-*Saccharomyces* strains. These two peptides are fragments of the glycolytic
283 enzyme glyceraldehyde-3-phosphate dehydrogenase (GAPDH); which is considered an energy
284 metabolism-related enzyme, able to generate NADH in glycolysis (Branco et al., 2017).

285 The production of these molecules is widespread among wine related yeasts and bacteria and
286 the use of yeasts able to release AMPs was recently proposed for the biocontrol against undesired
287 microorganisms. However, this strategy has been poorly applied to wine-related microorganisms
288 (Peña and Ganga, 2019). A recent study reported the production of an AMP by the non-
289 *Saccharomyces* strain *Candida intermedia* LAMAP1790. The application of this wine strain affected
290 the growth of several strains of *Brettanomyces bruxellensis*, without influencing *S. cerevisiae*
291 performance during fermentation (Peña and Ganga, 2019). These findings opens new prospects
292 regarding the use of these AMPs as alternatives to SO₂ in wine production industrial process
293 (Albergaria and Arneborg, 2016). Since, some spoilage microorganisms like *Pichia* spp., *Dekkera*
294 spp. and *Z. bailii* could tolerate medium-high SO₂ levels and the AMPs might have a key role in
295 controlling unwanted microorganisms without affecting the beneficial microorganisms.

296

297 2.2.1.2 Killer toxins

298

299 Killer toxins play an important role in the defence mechanisms of the yeast starter culture
300 against undesired microorganisms. This character consists in the production of proteins or
301 glycoproteins (molecular weight between 30 and 70 kDa) that are lethal for sensitive cells
302 taxonomically related or not. It was demonstrated that the action of these compounds is mediated by
303 the presence of two kinds of specific receptors in the sensitive microorganism cell wall (Liu et al.,
304 2015). Even if this characteristic was first observed in *S. cerevisiae*, it is well distributed in several
305 yeast genera like *Candida*, *Ogataea*, *Pichia*, *Williopsis*, *Tetrapisispora*, *Schwanniomyces*,
306 *Debaryomyces*, *Ustilago*, *Cryptococcus*, *Metschnikowia*, *Kluyveromyces* and *Zygosaccharomyces*
307 (de Ullivarri et al., 2014; Liu et al., 2015). Killer toxins are encoded chromosomally, or their
308 production has been related with the presence of dsRNA viruses, or linear dsDNA plasmids (Belda
309 et al., 2017), however this is the case of some of the non-*Saccharomyces* species listed above. Killer
310 yeasts have been used to control pathogenic fungi in plants (Schmitt & Breinig, 2002) and to develop

311 antimycotics for the treatment of human and animal infections (Liu et al., 2015). Furthermore, they
312 were purposed as an alternative strategy of biocontrol to combat contaminating wild yeasts in food
313 industries and in particular in winemaking (Çorbacı, and Uçar, 2018; de Ullivarri et al., 2014;
314 Mazzucco, et al., 2019; Santos et al., 2011; Villaba et al., 2016;).

315 The possibility of using *S. cerevisiae* as BPCs (active antagonistic phenotype) in wine was
316 first introduced in the 1960s, when it was found that *S. cerevisiae* strains are able to release killer
317 toxins and inhibit other *S. cerevisiae* strains as well as spoilage yeast species using reduced SO₂
318 concentrations (de Ullivarri et al., 2014). These authors proposed the use of a sequential inoculation
319 in wine of two *S. cerevisiae* killer strains (Cf8 and M12) to improve the biocontrol against undesired
320 wine yeasts *Dekkera anomala* and *Z. bailii*. This new strategy showed a reduction in the growth of
321 the undesired yeasts to about 78% and 50%, respectively. Subsequently, Oro et al. (2016) found that
322 Kwkt and Pikt, two killer toxins produced by *Kluyveromyces wickerhamii* and *Wickerhamomyces*
323 *anomalus*, respectively, showed an antimicrobial activity against *B. bruxellensis*. These authors
324 demonstrated that this spoilage wine yeast, after a treatment of 0.7 mg/l of SO₂ at pH 3.2, entered in
325 a viable but non culturable state. Divergently, after treatment with Kwkt or Pikt, *B. bruxellensis* D46
326 strain completely lost its viability. Furthermore, Mehlomakulu et al. (2017) exposed this spoilage
327 yeast to the killer toxin CpKT1 produced by *Candida pyralidae* and demonstrated that the loss of
328 viability was due to the induction of cell membrane and cell wall damages. Mazzucco et al. (2019)
329 also observed an inhibition against *B. bruxellensis*. These authors, demonstrated that the killer toxin
330 SeKT produced by *Saccharomyces eubayanus* in wine, could be used for the biocontrol of four
331 common spoilage wine yeasts, namely *B. bruxellensis*, *Pichia membranifaciens*, *Pichia*
332 *guilliermondii* and *Pichia manshurica*.

333 Villaba and colleagues (2016) investigated the ability of *T. delbrueckii* to release a killer toxin
334 (TdKT). In particular, the NPCC 1033 strain was tested for its killer activity against some undesired
335 wine yeasts, namely *H. uvarum*, *B. bruxellensis*, *P. guilliermondii* and *P. membranifaciens* (Lopes
336 and Sangorrín, 2010). It was also verified that the TdKT toxin was active in oenological conditions

337 (at different pH and temperatures values, ethanol, glucose and SO₂ concentrations simulating several
338 wine environmental conditions). It was also found that some *L. thermotolerans* strains secreted killer
339 toxins against *S. cerevisiae* (Aponte and Blaiotta 2016), while Nally et al. (2018) demonstrated that
340 two strains were able to inhibit *Aspergillus* without affecting *S. cerevisiae* in a mixed fermentation.
341 Some *P. kluyveri* strains were found able to release killer toxins in the medium inhibiting the growth
342 of some food and beverage spoilage yeast genera like *Dekkera*, *Kluyveromyces*, *Pichia*,
343 *Saccharomyces* and *Zygosaccharomyces* (with the highest activity against *D. bruxellensis*) (Labrani
344 et al., 2015; Zagorc et al., 2001). Finally, the use of killer strains in sparkling wine production was
345 proposed in order to accelerate the maturation time (Todd et al., 2000, Velázquez *et al.*, 2016, 2019).
346 Since, the production of this category of wines using the traditional Methode Champenoise is
347 characterized by a secondary fermentation in a bottle of a base wine, followed by a long period of
348 aging in which the yeast cells are in contact with the fermented wine. During this period, which may
349 last several months, yeast cells undergo autolysis and release cell components into the wine with
350 positive repercussions on the aroma, mouthfeel and foaming properties of wines (Di Gianvito et al.,
351 2019). Such investigations, suggested that killer *T. delbrueckii* strains and sensitive yeasts in mixed
352 cultures could accelerate the onset of yeast autolysis, improving the organoleptic quality and foam
353 properties of sparkling wines, without affecting fermentation kinetics (Velázquez et al., 2016, 2019).
354 Another study demonstrated that the use of a combination of a killer *S. cerevisiae* strain and a sensitive
355 *Saccharomyces bayanus* induced the autolysis of the sensitive yeast in a shorter time, compared to
356 the control wine (without the addition of these yeasts), under oenological conditions (Lombardi et al.,
357 2016). In particular, in a pilot scale production, the selected strains influenced the concentrations of
358 free amino acids, total proteins and polysaccharides reaching, after 3 months of aging, the levels
359 showed by control wines after nine months of aging. These findings, raise the possibility of using a
360 biotechnological approach based on specific combinations of killer/sensitive strains or different types
361 of killer strains to reduce the time needed to release of cell components in the medium and

362 consequently the time for sparkling wine production with economic benefits for the producers
363 (Mannazzu et al., 2019).

364
365 2.2.1.3 Bacteriocins and bacteriocin-like inhibitory substance (BLIS)

366

367 Many wine related LAB are known for their ability to produce proteinaceous antimicrobial
368 molecules called bacteriocins, active against other bacteria that are closely related to the producing
369 organism growing in the same medium (Ndlovu et al., 2015). Several studies highlighted the ability
370 of wine LAB to produce bacteriocins, however, bacteriocin producing activity has not yet been
371 demonstrated under wine conditions (Krieger-Weber et al., 2020) and their efficacy in wine is only
372 demonstrated in combination with SO₂ (Royo-Bezares et al., 2007). In fact, the production and
373 application of bacteriocins in food is influenced by the physical and chemical conditions of food and
374 the presence of competitive microorganisms (Singh et al., 2018).

375 Knoll et al. (2008) found that both indigenous and commercial *O. oeni* strains showed an
376 antagonistic activity against several wine-related LAB strains *in vitro*. These authors identified some
377 putative bacteriocin-encoding genes, but they did not demonstrate the release of these bacteriocins in
378 wine. Recently, Lasik-Kurdyś and Sip (2019) reported the ability of one *O. oeni* strain to synthesize
379 a peptidic BLIS active against *Leuconostoc mesenteroides* and *Pediococcus pentosaceus*. In
380 particular, a synergistic effect between the presence of *O. oeni* (competition for nutrients and space)
381 the release of BLIS and other inhibition compounds (organic acids and H₂O₂) was highlighted. It was
382 further found that the removal of viable cells, as well as the neutralization of the supernatant or the
383 treatment with the catalase enzyme, significantly decreased or removed the inhibition activity against
384 some LAB indicator strains. Although several studies have indicated the use of bacteriocins in food
385 and beverages production, the application of these molecules in wine has a great potential, but it's
386 not yet approved. Interestingly, studies have proposed the use of bacteriocins to control biofilm
387 formation on stainless steel tanks and surfaces (Nel et al., 2002), suggesting that their application

388 could be a valid response to the chemical cleaning detergents with a negative impact to the
389 environment.

390

391 2.2.1.4 Lytic enzymes

392

393 The production of cell wall lytic enzymes by wine related yeasts and bacteria is a less known
394 form of antagonism. However, there are growing evidences that suggest the involvement of cell-wall
395 degrading enzymes in the mechanism of action of BPCs. Lytic enzymes such as β -glucanase
396 determine the leakage of cell contents by cleaving the β -glucan polysaccharide of fungi and bacteria,
397 such as *Penicillium expansum*, *Fusarium oxysporum* and *B. cinerea* (Edison et al., 2018; Wisniewski
398 et al., 1991). Cordero-Bueso et al., (2017), observed the release of lytic enzymes by *P. kluyveri* that
399 could result in the degradation of the fungal cell wall (β -1, 3-glucanase, proteolytic and pectinolytic
400 activities). Moreover, on grapes *A. pullulans* releases extracellular β -1, 3-glucanase which could play
401 a role in the biocontrol activity towards grape pathogens (Bozoudi and Tsaltas, 2018). Lastly, several
402 strains of the *M. pulcherrima* clade secrete extracellular chitinase and β -1, 3-glucanases able to
403 degrade the cell wall of *B. cinerea*, however their impact is strain and matrix-dependent (Sipiczki,
404 2020).

405

406 2.2.2. Volatile Organic Compounds (VOCs)

407

408 Among mechanisms of antagonism, VOCs are considered superior over the nonvolatile
409 compounds ones because direct contact and closeness among microorganisms are not required. In
410 fact, these small organic molecules ($<C_{20}$) can diffuse to greater distances in heterogeneous
411 environment, since are characterized by a low molecular mass (100 to 500 Da), a lipophilic moiety,
412 low solubility in the water, low boiling point and a high vapor pressure that permit a readily
413 volatilization at ambient temperatures (Zhang et al., 2021). VOCs are molecules potentially produced

414 by all living microorganisms and can be ascribed to several molecular classes, including
415 hydrocarbons, alcohols, thioalcohols, aldehydes, ketones, thioesters, cyclohexanes, heterocyclic
416 compounds, phenols, and benzene derivatives and can be produced by the primary metabolism or via
417 specialized secondary metabolic pathways.

418 It is widely known that wines produced by mixed fermentations have a complex aroma profile,
419 but the role of these volatile molecules in microorganisms' interactions is still poorly understood.
420 This is the case of the higher alcohols tryptophol, phenylethanol, and tyrosol which are produced at
421 relative high levels by *S. cerevisiae*. Recently, interactions studies highlighted their role as
422 antagonistic VOCs. In fact, through *in vitro* experiments, it was demonstrated that these molecules
423 caused the reduction of the growth of non-*Saccharomyces* like *H uvarum*, *Starm. bacillaris*, *T.*
424 *delbrueckii* and *M. pulcherrima*, even at low concentration (Gonzalez et al., 2018).

425 In another study, Farbo et al. (2018) attributed the bioprotective activity of *L. thermotolerans*
426 against toxinogenic fungi (*Aspergillus* spp.) to the production of VOCs, particularly 2-phenylethanol.
427 These authors demonstrated that this volatile compound was able to downregulate key genes
428 implicated in Ochratoxin A biosynthesis in a species-dependent way. The implication of this higher
429 alcohol in the antagonistic role during alcoholic fermentation was also suggested by the
430 transcriptomic profile of the yeast during fermentation. In fact, it was highlighted that in a mixed
431 fermentation with *S. cerevisiae*, *L. thermotolerans* showed an up regulation of all genes involved in
432 the conversion of phenylalanine to phenylethanol (Shekhawat et al., 2019).

433 Furthermore, through *in vitro* experiments, it was demonstrated that VOCs such as isoamyl
434 acetate, isoamyl alcohol, 2-phenyl ethylacetate and 2-phenyl ethanol were responsible of the
435 bioprotective activity of *P. kluyveri* on grapes (Mewa-Ngongang et al., 2019b). These authors have
436 also observed that this yeast inhibited the growth of wine spoilage yeasts (*D. bruxellensis* and *D.*
437 *anomala*) and fruit fungi (*B. cinerea*, *Colletotrichum acutatum* and *Rhizopus stolonifer* (Mewa-
438 Ngongang et al., 2019a, b). A biocontrol ability on grape berries was also observed for another non-
439 *Saccharomyces* yeast, namely *Starm. bacillaris*. This species is able to survive until the last stages of

440 the fermentation process, contributing significantly to wine quality (Englezos et al., 2017). Its
441 antagonistic activity was investigated against fungal infections such as *Alternaria alternata* (Prendes
442 et al., 2018) and *B. cinerea* (Lemos Junior et al., 2016). In the latter study, through *in vivo* tests, it
443 was found that a co-inoculation of 14 *Starm. bacillaris* strains in artificially wounded grape berries
444 lead to a decrease of *B. cinerea* infection. A recent study attributed this bioprotection ability to the
445 production of VOCs and in particular to benzyl alcohol (Lemos Junior et al., 2020). In line with this
446 finding, several authors observed an increased amount of this alcohol in mixed fermentations with *S.*
447 *cerevisiae* (Englezos et al., 2019a; Binati et al., 2020). Antimicrobial activity against *B. cinerea* was
448 also observed for *M. pulcherrima* even if in a strain dependent way, due to the production of ethyl
449 alcohol and ethyl acetate (Parafati et al., 2015; Contarino et al., 2019). The inhibitory effect of yeast-
450 derived VOCs against postharvest food pathogens was greatly investigated and proposed as an
451 effective biocontrol strategy to inhibit the growth of *B. cinerea*, *Colletotrichum acutatum*, *P.*
452 *expansum*, *Penicillium digitatum* and *Penicillium italicum* (Contarino et al., 2019). VOCs could be
453 considered as ideal antimicrobials, since contact between biocontrol agent and food pathogen is not
454 required, to inhibit the growth of the last. Such approach could be useful on postharvest table grapes
455 to improve shelf life and therefore reduce food loss during storage. Further studies are necessary to
456 understand the type and the concentration of VOCs that could be considered crucial in the biocontrol
457 mechanism.

458

459 2.2.3. Organic acids

460

461 Organic acids production by the BPCs could play an important role on their dominance over the
462 indigenous microorganisms, increasing the microbial stability of the wine. Bagheri et al. (2018) found
463 that in a consortium of six non-*Saccharomyces* and a *S. cerevisiae* strains, a higher concentration of
464 *Starm. bacillaris* resulted in the disappearance of *Candida parapsilosis* and *M. pulcherrima*. This
465 antagonistic ability was attributed to the capacity to produce high levels of organic acids, increasing

466 the total acidity and reducing pH of the wines. In wine conditions, *L. thermotolerans* bioprotective
467 ability was investigated in combination with a *Lpb. plantarum* strain (Rubio-Bretón et al., 2018). The
468 results showed the potential of this inoculation strategy as alternatives to SO₂ addition. The final
469 wines had a higher titratable acidity and esters concentration than control wines inoculated with *S.*
470 *cerevisiae*. The antimicrobial activity was attributed to the production of high lactic acid amounts by
471 both microorganisms and consequently to the decrease of the pH (Gobbi et al., 2013; Rubio-Bretón
472 et al., 2018; Vilela, 2018). This peculiar feature is of great interest in wine starter cultures, since
473 current climate change in many wine-growing regions may cause a shift in grapes composition,
474 increasing sugar content and decreasing must acidity (Mira de Orduña, 2010). In a recent study,
475 Tremonte et al., (2017) demonstrated that the highest antimicrobial activity of 106 *Lpb. plantarum*
476 strains was exerted when growing cells were present. It was also highlighted that the use of a cell free
477 supernatant reduced the bioprotective ability of all strains and in particular that the main antagonistic
478 activity was due to the production of organic acids. Interestingly, in this study a relationship between
479 the antagonistic activity and the origin of the strains was underlined, with the wine strains showing
480 the strongest growth inhibition capacity. Generally, biocontrol activity, by the decrease of pH due to
481 the organic acid production could have an impact on wine color stability, due to the enhance of SO₂
482 properties and the shift of the equilibrium of anthocyanins from the colourless to colored form
483 (Giacosa et al., 2019).

484

485 2.2.4. Cell-to-cell contact

486

487 Among the interaction mechanisms, different researches underlined the ability of yeasts to
488 form cellular aggregates called flocs in different environments, wine included. These flocs can be
489 formed by cells of the same strain, by different strains of the same species or by microorganisms of
490 different species (Rossouw et al., 2015). This physical association increases the metabolic exchange
491 among cells and, consequently, determines the ecological interactions like inhibition or stimulation.

492 It was demonstrated that in a wine environment this cell-to-cell contact mechanism is involved in
493 antagonistic interactions among microorganisms. This phenomenon also called co-aggregation was
494 firstly observed in *S. cerevisiae* against several wine yeasts namely *L. thermotolerans*, *T. delbrueckii*,
495 *H. uvarum* and *Starm. bacillaris* (Alonso-del-Real et al., 2019; Englezos et al., 2019b; Kemsawasd
496 et al., 2015; Nissen and Arneborg, 2003; Petigonnet et al., 2019; Pietrafesa et al., 2020; Renault et
497 al., 2013; Rossouw et al., 2018; Shekhawat et al., 2019). It was also observed between different strains
498 of *S. cerevisiae* and is involved in dominance (Pérez-Torrado et al., 2018).

499 Co-aggregation is mainly studied using a double compartment bioreactor where two species
500 are separated by a membrane that permits the exchange of metabolites (Nissen and Arneborg, 2003),
501 however this system limits the nature and extent of the interactions investigated because it does not
502 permit the immediate and complete transfer of all relevant metabolites and macromolecules (Rossouw
503 et al., 2018).

504 Recently, in *S. cerevisiae* the genetic basis of this ability was mainly attributed to the
505 expression of *FLO* family genes, which encode for adhesive proteins on cell wall surface (Rossouw
506 et al., 2018). In fact, transcriptomic analyses highlighted that, during wine and sparkling wine
507 fermentation, *S. cerevisiae* is achieved in a general cell wall remodelling with the up-regulation of
508 genes involved in asexual cell aggregation (*FLO* genes) (Shekhawat et al., 2019; Di Gianvito et al.,
509 2018). Genetic analyses suggested that non-*Saccharomyces* yeasts like *H. uvarum* contain adhesion-
510 related domains in their genome (Pu et al., 2014). However, the antagonistic role in these
511 microorganisms is not yet investigated.

512

513 **3. Conclusions and future perspectives**

514

515 Success in biological control of unwanted microorganisms requires consideration of the wine
516 fermentation ecology as a balanced ecosystem, where the resident microbiota contribute to enhance
517 wine quality, while maintaining spoilage microorganisms at the lowest levels. Implementing

518 fermentation conditions that favour the population of microorganisms with antimicrobial activity is
519 essential to this purpose. However, the use of chemical additives such as SO₂ may still be needed
520 when the concentration of microbial spoilage metabolites is high, especially in wines with low or
521 moderate ethanol levels. Despite the limitations observed in the different stages, the use of viable
522 cells of antagonist microorganisms well adapted in specific conditions or their derived antimicrobial
523 metabolites are of relevance in winemaking industry, since they could help to reduce or even replace
524 chemical additives. In the same context, the selection of yeasts and bacteria of oenological interest
525 with the double role of starter culture and BPCs represents an opportunity for winemakers to enhance
526 wine quality and reduce the production costs, as the demand for sustainable wines will continue to
527 grow. A better knowledge of the fermentation conditions and oenological practices which modulate
528 microorganism's performance, will allow a greater management of specific unwanted
529 microorganisms during the alcoholic and/or malolactic fermentation. Therefore, studies aiming to
530 further investigate the interaction mechanisms among the starter culture and BPCs and the spoilage
531 microorganisms will be of great value in the near future for this field of research.

532

533 **Funding**

534

535 This research did not receive any specific grant from funding agencies in the public, commercial, or
536 not-for-profit sectors.

537

538 **Declaration of competing interest**

539

540 The authors declare no conflict of interest.

541

542 **References**

543

544 Albergaria, H., Arneborg, N., 2016. Dominance of *Saccharomyces cerevisiae* in alcoholic
545 fermentation processes: role of physiological fitness and microbial interactions. Appl. Microbiol.
546 Biotechnol. 100(5), 2035-2046. <http://dx.doi.org/10.1007/s00253-015-7255-0>.

547

548 Alonso-del-Real, J., Pérez-Torrado, R., Querol, A., Barrio, E., 2019. Dominance of wine
549 *Saccharomyces cerevisiae* strains over *Saccharomyces kudriavzevii* in industrial fermentation
550 competitions is related to an acceleration of nutrient uptake and utilization. Environ. Microbiol. 21(5),
551 1627-1644. <http://dx.doi.org/10.1111/1462-2920.14536>.

552

553 Andorrà, I., Martín, L., Nart, E., Puxeu, M., Hidalgo, C., Ferrer-Gallego, R., 2018. Effect of grape
554 juice composition and nutrient supplementation on the production of sulfur dioxide and carboxylic
555 compounds by *Saccharomyces cerevisiae*. Aust. J. Grape Wine Res. 24(2), 260-266.
556 <http://dx.doi.org/10.1111/ajgw.12325>.

557

558 Aponte, M., Blaiotta, G., 2016. Potential role of yeast strains isolated from grapes in the production
559 of Taurasi DOCG. Front. Microbiol. 7, 809. <https://doi.org/10.3389/fmicb.2016.00809>.

560

561 Bagheri, B., Zambelli, P., Vigentini, I., Bauer, F. F., Setati, M. E., 2018. Investigating the effect of
562 selected non-*Saccharomyces* species on wine ecosystem function and major volatiles. Front. Bioeng.
563 Biotechnol. 6, 169. <https://doi.org/10.3389/fbioe.2018.00169>.

564

565 Bahar, A. A., Ren, D., 2013. Antimicrobial peptides. Pharmaceuticals. 6(12), 1543-1575.
566 <https://doi.org/10.3390/ph6121543>.

567

568 Balmaseda, A., Bordons, A., Reguant, C., Bautista-Gallego, J., 2018. Non-*Saccharomyces* in wine:
569 effect upon *Oenococcus oeni* and malolactic fermentation. *Front. Microbiol.* 9, 534.
570 <https://doi.org/10.3389/fmicb.2018.00534>.

571

572 Barbosa, C., Mendes-Faia, A., Lage, P., Mira, N. P., Mendes-Ferreira, A., 2015. Genomic expression
573 program of *Saccharomyces cerevisiae* along a mixed-culture wine fermentation with *Hanseniaspora*
574 *guilliermondii*. *Microb. Cell Factories.* 14(1), 124. <https://doi.org/10.1186/s12934-015-0318-1>.

575

576 Bartle, L., Sumbly, K., Sundstrom, J., Jiranek, V., 2019. The microbial challenge of winemaking:
577 yeast-bacteria compatibility. *FEMS Yeast Res.* 19(4), foz040. <https://doi.org/10.1093/femsyr/foz040>.

578

579 Bastard, A., Coelho, C., Briandet, R., Canette, A., Gougeon, R., Alexandre, H., Guzzo, J., Weidmann,
580 S., 2016. Effect of biofilm formation by *Oenococcus oeni* on malolactic fermentation and the release
581 of aromatic compounds in wine. *Front. Microbiol.* 7, 613. <https://doi.org/10.3389/fmicb.2016.00613>.

582

583 Bauer, M. A., Kainz, K., Carmona-Gutierrez, D., Madeo, F., 2018. Microbial wars: competition in
584 ecological niches and within the microbiome. *Microb. Cell.* 5(5), 215.
585 <https://doi.org/10.15698/mic2018.05.628>.

586

587 Belda, I., Ruiz, J., Alonso, A., Marquina, D., Santos, A., 2017. The biology of *Pichia*
588 *membranifaciens* killer toxins. *Toxins.* 9(4), 112. <https://doi.org/10.3390/toxins9040112>.

589

590 Benito, S., 2018a. The impact of *Torulasporea delbrueckii* yeast in winemaking. *Appl. Microbiol.*
591 *Biotechnol.* 102(7), 3081-3094. <https://doi.org/10.1007/s00253-018-8849-0>.

592

593 Benito, S., 2018b. The impacts of *Lachancea thermotolerans* yeast strains on winemaking Appl.
594 Microbiol. Biotechnol. 102 (16), 6775-6790. <https://doi.org/10.1007/s00253-018-9117-z>.
595

596 Binati, R. L., Junior, W. J. L., Luzzini, G., Slaghenaufi, D., Ugliano, M., Torriani, S., 2020.
597 Contribution of non-*Saccharomyces* yeasts to wine volatile and sensory diversity: A study on
598 *Lachancea thermotolerans*, *Metschnikowia* spp. and *Starmerella bacillaris* strains isolated in Italy.
599 Int. J. Food Microbiol. 318, 108470. <https://doi.org/10.1016/j.ijfoodmicro.2019.108470>.
600

601 Bisson, L. F., 1999. Stuck and sluggish fermentations. Am. J. Enol. Vitic. 50, 107-119.
602

603 Boynton, P. J., Peterson, C. N., Pringle, A., 2019. Superior dispersal ability can lead to persistent
604 ecological dominance throughout succession. Appl. Environ. Microbiol. 85(6), e02421-18.
605 <https://doi.org/10.1128/AEM.02421-18>.
606

607 Bozoudi, D., Tsaltas, D., 2018. The multiple and versatile roles of *Aureobasidium pullulans* in the
608 vitivicultural sector. Fermentation. 4(4), 85. <https://doi.org/10.3390/fermentation4040085>.
609

610 Branco, P., Francisco, D., Chambon, C., Hébraud, M., Arneborg, N., Almeida, M. G., Caldeira, J.,
611 Albergaria, H., 2014. Identification of novel GAPDH-derived antimicrobial peptides secreted by
612 *Saccharomyces cerevisiae* and involved in wine microbial interactions. Appl. Microbiol. Biotechnol.
613 98, 843-853. <https://doi.org/10.1007/s00253-013-5411-y>.
614

615 Branco, P., Francisco, D., Monteiro, M., Almeida, M. G., Caldeira, J., Arneborg, N., Prista, C.,
616 Albergaria, H., 2017. Antimicrobial properties and death-inducing mechanisms of saccharomycin, a
617 biocide secreted by *Saccharomyces cerevisiae*. Appl. Microbiol. Biotechnol. 101, 159-171.
618 <https://doi.org/10.1007/s00253-016-7755-6>.

619

620 Brizuela, N., Tymczynszyn, E. E., Semorile, L. C., La Hens, D. V., Delfederico, L., Hollmann, A.,
621 Bravo-Ferrada, B., 2019. *Lactobacillus plantarum* as a malolactic starter culture in winemaking: A
622 new (old) player?. *Electron. J. Biotechnol.* 38, 10-18. <https://doi.org/10.1016/j.ejbt.2018.12.002>.

623

624 Carbó, A., Torres, R., Teixidó, N., Usall, J., Magan, N., Medina, A., 2018. Predicted ecological niches
625 and environmental resilience of different formulations of the biocontrol yeast *Candida sake* CPA-1
626 using the Bioscreen C. *BioControl.* 63(6), 855-866. <https://doi.org/10.1007/s10526-018-09910-4>.

627

628 Comitini, F., Ferretti, R., Clementi, F., Mannuzzu, I., Ciani, M., 2005. Interactions between
629 *Saccharomyces cerevisiae* and malolactic bacteria: preliminary characterization of a yeast
630 proteinaceous compound(s) active against *Oenococcus oeni*. *J. Appl. Microbiol.* 99, 105-111.
631 doi:10.1111/j.1365-2672.2005.02579.x.

632

633 Contarino, R., Brighina, S., Fallico, B., Cirvilleri, G., Parafati, L., Restuccia, C., 2019. Volatile
634 organic compounds (VOCs) produced by biocontrol yeasts. *Food Microbiol.* 82, 70-74.
635 <https://doi.org/10.1016/j.fm.2019.01.008>.

636

637 Çorbacı, C., Uçar, F. B., 2018. Purification, characterization and *in vivo* biocontrol efficiency of killer
638 toxins from *Debaryomyces hansenii* strains. *Int. J. Biol. Macromol.* 119, 1077-1082.
639 <https://doi.org/10.1016/j.ijbiomac.2018.07.121>.

640

641 Cordero-Bueso, G., Mangieri, N., Maghradze, D., Foschino, R., Valdetara, F., Cantoral, J. M.,
642 Vigentini, I., 2017. Wild grape-associated yeasts as promising biocontrol agents against *Vitis vinifera*
643 fungal pathogens. *Front. Microbiol.* 8, 2025. <https://doi.org/10.3389/fmicb.2017.02025>.

644

645 Curiel, J. A., Morales, P., Gonzalez, R., Tronchoni, J., 2017. Different non-*Saccharomyces* yeast
646 species stimulate nutrient consumption in *S. cerevisiae* mixed cultures. *Front. Microbiol.* 8, 2121.
647 <https://doi.org/10.3389/fmicb.2017.02121>.

648

649 de Ullivarri, M. F., Mendoza, L. M., Raya, R. R., 2014. Killer activity of *Saccharomyces cerevisiae*
650 strains: partial characterization and strategies to improve the biocontrol efficacy in winemaking.
651 *Antonie Leeuwenhoek.* 106(5), 865-878. <https://doi.org/10.1007/s10482-014-0256-7>.

652

653 Di Gianvito, P., Arfelli, G., Suzzi, G., Tofalo, R., 2019. New trends in sparkling wine production:
654 yeast rational selection. In *Alcoholic Beverages*, Woodhead Publishing., pp. 347-386.

655

656 Di Gianvito, P., Tesnière, C., Suzzi, G., Blondin, B., Tofalo, R., 2018. Different genetic responses to
657 oenological conditions between a flocculent wine yeast and its *FLO5* deleted strain: insights from the
658 transcriptome. *Food Res. Int.* 114, 178-186. <https://doi.org/10.1016/j.foodres.2018.07.061>.

659

660 Dimopoulou, M., Bardeau, T., Ramonet, P.Y., Miot-Certier, C., Claisse, O., Doco, T., Petrel, M.,
661 Lucas, P., Dols-Lafargue, M., 2015. Exopolysaccharides produced by *Oenococcus oeni*: from
662 genomic and phenotypic analysis to technological valorization. *Food Microbiol.* 53, 10-17.
663 <https://doi.org/10.1016/j.fm.2015.07.011>.

664

665 Dimopoulou, M., Raffenne, J., Claisse, O., Miot-Sertier, C., Iturmendi, N., Moine, V., Coulon, J.,
666 Dols-Lafargue, M., 2018. *Oenococcus oeni* exopolysaccharide biosynthesis, a tool to improve
667 malolactic starter performance. *Front. Microbiol.* 9, 1276. <https://doi.org/10.3389/fmicb.2018.01276>.

668

669 Dokka, A., Kotilinga Reddy, Y., Spoorthy, G. S., Reddy, I. S., 2018. Protective cultures-a review. *Int.*
670 *J. Curr. Microbiol. App. Sci.* 7(6), 228-238. <https://doi.org/10.20546/ijemas.2018.706.028>.

671

672 Droby, S., Wisniewski, M., Teixidó, N., Spadaro, D., Jijakli, M. H., 2016. The science, development,
673 and commercialization of postharvest biocontrol products. *Postharvest Biol. Technol.* 122, 22-29.
674 <https://doi.org/10.1016/j.postharvbio.2016.04.006>.

675

676 Edison, L. K., Shiburaj, S., Pradeep, N. S., 2018. Microbial beta glucanase in agriculture. In: Kumar,
677 P., Patra, K. J., Chandra, P. (Ed.), *Advances in Microbial Biotechnology*. Apple Academic Press. P.,
678 New York, pp. 53-72.

679

680 Englezos, V., Cravero, F., Torchio, F., Rantsiou, K., Ortiz-Julien, A., Lambri, M., Gerbi, V., Rolle,
681 L., Cocolin, L., 2018a. Oxygen availability and strain combination modulate yeast growth dynamics
682 in mixed culture fermentations of grape must with *Starmerella bacillaris* and *Saccharomyces*
683 *cerevisiae*. *Food Microbiol.* 69, 179-188. <https://doi.org/10.1016/j.fm.2017.08.007>.

684

685 Englezos, V., Giacosa, S., Rantsiou, K., Rolle, L., Cocolin, L., 2017. *Starmerella bacillaris* in
686 winemaking: opportunities and risks. *Curr. Opin. Food Sci.* 17, 30-35.
687 <https://doi.org/10.1016/j.cofs.2017.08.007>.

688

689 Englezos, V., Pollon, M., Rantsiou, K., Ortiz-Julien, A., Botto, R., Rio Segade, S., Giacosa S., Rolle
690 L., Cocolin, L., 2019a. *Saccharomyces cerevisiae*-*Starmerella bacillaris* strains interaction
691 modulates chemical and volatile profile in red wine mixed fermentations. *Food Res. Int.* 122, 392-
692 401. <https://doi.org/10.1016/j.foodres.2019.03.072>.

693

694 Englezos, V., Rantsiou, K., Giacosa, S., Rio Segade, S., Rolle, L., Cocolin, L., 2019b. Cell-to-cell
695 contact mechanism modulates *Starmerella bacillaris* death in mixed culture fermentations with

696 *Saccharomyces cerevisiae*. Int. J. Food Microbiol. 289, 106-114.
697 <https://doi.org/10.1016/j.ijfoodmicro.2018.09.009>.
698
699 Farbo, M. G., Urgeghe, P. P., Fiori, S., Marcello, A., Oggiano, S., Balmas, V., Hassan, Z. U. I., Jaoua,
700 S., Migheli, Q., 2018. Effect of yeast volatile organic compounds on ochratoxin A-producing
701 *Aspergillus carbonarius* and *Aspergillus ochraceus*. Int. J. Food Microbiol. 284, 1-10.
702 <https://doi.org/10.1016/j.ijfoodmicro.2018.06.023>.
703
704 Ferraz, P., Cássio, F., Lucas, C., 2019. Potential of yeasts as biocontrol agents of the phytopathogen
705 causing cacao Witches' Broom Disease: is microbial warfare a solution?. Front. Microbiol. 10, 1766.
706 <https://doi.org/10.3389/fmicb.2019.01766>.
707
708 Fiori, S., Urgeghe, P. P., Hammami, W., Razzu, S., Jaoua, S., Migheli, Q., 2014. Biocontrol activity
709 of four non-and low-fermenting yeast strains against *Aspergillus carbonarius* and their ability to
710 remove ochratoxin A from grape juice. Int. J. Food Microbiol. 189, 45-50.
711 <https://doi.org/10.1016/j.ijfoodmicro.2014.07.020>.
712
713 Gaggia, F., Di Gioia, D., Baffoni, L., Biavati, B., 2011. The role of protective and probiotic cultures
714 in food and feed and their impact in food safety. Trends Food Sci. Technol. 22, S58-S66.
715 <https://doi.org/10.1016/j.tifs.2011.03.003>.
716
717 Gao, C., Fleet, G. H., 1988. The effects of temperature and pH on the ethanol tolerance of the wine
718 yeasts, *Saccharomyces cerevisiae*, *Candida stellate* and *Kloeckera apiculata*. J. Appl. Bacteriol. 65,
719 405-410. <https://doi.org/10.1111/j.1365-2672.1988.tb01909.x>.
720

721 Garrido, C. C., Usall, J., Torres, R., Teixidó, N., 2017. Effective control of *Botrytis* bunch rot in
722 commercial vineyards by large-scale application of *Candida sake* CPA-1. *BioControl*. 62, 161-173.
723 <https://doi.org/10.1007/s10526-017-9789-9>.

724

725 Ghoul, M., Mitri, S., 2016. The ecology and evolution of microbial competition. *Trends Microbiol.*
726 24(10), 833-845. <https://doi.org/10.1016/j.tim.2016.06.011>.

727

728 Giacosa, S., Río Segade, S., Cagnasso, E., Caudana, A., Rolle, L., Gerbi, V., 2019. SO₂ in wines:
729 rational use and possible alternatives. In: Morata, A. (Ed.), *Red Wine Technology*. Academic Press,
730 Cambridge, MA, USA, pp. 309-321.

731

732 Gobbi, M., Comitini, F., Domizio, P., Romani, C., Lencioni, L., Mannazzu, I., Ciani, M., 2013.
733 *Lachancea thermotolerans* and *Saccharomyces cerevisiae* in simultaneous and sequential co-
734 fermentation: a strategy to enhance acidity and improve the overall quality of wine. *Food Microbiol.*
735 33(2), 271-281. <https://doi.org/10.1016/j.fm.2012.10.004>.

736

737 González, B., Vázquez, J., Morcillo-Parra, M. Á., Mas, A., Torija, M. J., Beltran, G., 2018. The
738 production of aromatic alcohols in non-*Saccharomyces* wine yeast is modulated by nutrient
739 availability. *Food Microbiol.* 74, 64-74. <https://doi.org/10.1016/j.fm.2018.03.003>.

740

741 Gore-Lloyd, D., Sumann, I., Brachmann, A. O., Schneeberger, K., Ortiz-Merino, R. A., Moreno-
742 Beltrán, M., Schläfli, M., Kirner, P., Santos Kron, A., Rueda-Mejia, M. P., Somerville, V., Wolfe, K.
743 H., Piel, J., Ahrens, C. H., Henk, D., Freimoser, F. M., 2019. Snf2 controls pulcherriminic acid
744 biosynthesis and antifungal activity of the biocontrol yeast *Metschnikowia pulcherrima*. *Molecular*
745 *Microbiol.* 112(1), 317-332. <https://doi.org/10.1111/mmi.14272>.

746

747 Guerrero, R. F., Cantos-Villar, E., 2015. Demonstrating the efficiency of sulphur dioxide
748 replacements in wine: A parameter review. Trends Food Sci Technol. 42(1), 27-43.
749 <https://doi.org/10.1016/j.tifs.2014.11.004>.

750

751 Jacometti, M. A., Wratten, S. D., Walter, M., 2010. Alternatives to synthetic fungicides for *Botrytis*
752 *cinerea* management in vineyards. Aust. J. Grape Wine Res. 16(1), 154-172.
753 <https://doi.org/10.1111/j.1755-0238.2009.0067.x>.

754

755 Jolly, N. P., Varela, C., Pretorius, I. S., 2014. Not your ordinary yeast: non-*Saccharomyces* yeasts in
756 wine production uncovered. FEMS Yeast Res. 14(2), 215-237. [https://doi.org/10.1111/1567-
757 1364.12111](https://doi.org/10.1111/1567-1364.12111).

758

759 Kántor, A., Hutková, J., Petrová, J., Hleba, L., Kacňániová, M., 2015. Antimicrobial activity of
760 pulcherrimin pigment produced by *Metschnikowia pulcherrima* against various yeast species. J.
761 Microbiol. Biotechnol. Food Sci. 5, 282-285.

762

763 Kemsawasd, V., Branco, P., Almeida, M. G., Caldeira, J., Albergaria, H., Arneborg, N., 2015. Cell-
764 to-cell contact and antimicrobial peptides play a combined role in the death of *Lachancea*
765 *thermotolerans* during mixed-culture alcoholic fermentation with *Saccharomyces cerevisiae*. FEMS
766 Microbiol. Lett. 362(14), fnv103. <https://doi.org/10.1093/femsle/fnv103>.

767

768 Keswani, C., Singh, H. B., Hermosa, R., García-Estrada, C., Caradus, J., He, Y. W., Mezaache-
769 Aichour, S., Glare, T. R., Borriss, R., Vinale, F., Sansinenea, E., 2019. Antimicrobial secondary
770 metabolites from agriculturally important fungi as next biocontrol agents. Appl. Microbiol.
771 Biotechnol. 103(23-24), 9287-9303. <https://doi.org/10.1007/s00253-019-10209-2103>.

772

773 Knoll, C., Divol, B., Du Toit, M., 2008. Genetic screening of lactic acid bacteria of oenological origin
774 for bacteriocin-encoding genes. Food Microbiol. 25(8), 983-991.
775 <https://doi.org/10.1016/j.fm.2008.06.010>.

776

777 Kosel, J., Čadež, N., Schuller, D., Carreto, L., Franco-Duarte, R., Raspor, P., 2017. The influence of
778 *Dekkera bruxellensis* on the transcriptome of *Saccharomyces cerevisiae* and on the aromatic profile
779 of synthetic wine must. FEMS Yeast Res. 17(4), fox018. <https://doi.org/10.1093/femsyr/fox018>.

780

781 Krieger-Weber, S., Heras, J. M., Suarez, C., 2020. *Lactobacillus plantarum*, a new biological tool to
782 control malolactic fermentation: a review and an outlook. Beverages. 6(2), 23.
783 <https://doi.org/10.3390/beverages6020023>.

784

785 Kuchen, B., Maturano, Y. P., Mestre, M. V., Combina, M., Toro, M. E., Vazquez, F., 2019. Selection
786 of native non-*Saccharomyces* yeasts with biocontrol activity against spoilage yeasts in order to
787 produce healthy regional wines. Fermentation, 5(3), 60.
788 <https://doi.org/10.3390/fermentation5030060>.

789

790 Labbani, F. Z. K., Turchetti, B., Bennamoun, L., Dakhmouche, S., Roberti, R., Corazzi, L., Meraihi,
791 Z., Buzzini, P., 2015. A novel killer protein from *Pichia kluyveri* isolated from an Algerian soil:
792 purification and characterization of its *in vitro* activity against food and beverage spoilage yeasts.
793 Antonie Leeuwenhoek. 107(4), 961-970. <https://doi.org/10.1007/s10482-015-0388-4>.

794

795 Lachance, M. A., 2016. *Metschnikowia*: half tetrads, a regicide and the fountain of youth. Yeast.
796 33(11), 563-574. <https://doi.org/10.1002/yea.3208>.

797

798 Lasik-Kurdyś, M., Sip, A., 2019. Evaluation of the antimicrobial activity of bacteriocin-like
799 inhibitory substances of enological importance produced by *Oenococcus oeni* isolated from wine.
800 Eur. Food Res. Technol. 245(2), 375-382. <https://doi.org/10.1007/s00217-018-3169-2>.
801

802 Lemos Junior, W. J., Fernandes, W. J., Bovo, B., Nadai, C., Crosato, G., Carlot, M., Favaron, F.,
803 Giacomini A., Corich, V., 2016. Biocontrol ability and action mechanism of *Starmerella bacillaris*
804 (synonym *Candida zemplinina*) isolated from wine musts against gray mold disease agent *Botrytis*
805 *cinerea* on grape and their effects on alcoholic fermentation. Front Microbiol. 7, 1249.
806 <https://doi.org/10.3389/fmicb.2016.01499>.
807

808 Lemos Junior, W. J., Binati, R. L., Felis, G. E., Slaghenaufi, D., Ugliano, M., Torriani, S., 2020.
809 Volatile organic compounds from *Starmerella bacillaris* to control gray mold on apples and modulate
810 cider aroma profile. Food Microbiol. 89, 103446. <https://doi.org/10.1016/j.fm.2020.103446>.
811

812 Liu, G. L., Chi, Z., Wang, G. Y., Wang, Z. P., Li, Y., Chi, Z. M., 2015. Yeast killer toxins, molecular
813 mechanisms of their action and their applications. Crit. Rev. Biotechnol. 35(2), 222-234.
814 <https://doi.org/10.3109/07388551.2013.833582>.
815

816 Liu, Y., Rousseaux, S., Tourdot-Maréchal, R., Sadoudi, M., Gougeon, R., Schmitt-Kopplin, P.,
817 Alexandre, H., 2017. Wine microbiome: a dynamic world of microbial interactions. Crit. Rev. Food
818 Sci. Nutr. 57(4), 856-873. <https://doi.org/10.1080/10408398.2014.983591>.
819

820 Lombardi, S. J., De Leonardis, A., Lustrato, G., Testa, B., Iorizzo, M., 2016. Yeast autolysis in
821 sparkling wine aging: use of killer and sensitive *Saccharomyces cerevisiae* strains in co-culture.
822 Recent Pat. Biotechnol. 9, 223-230.
823

824 Lopes, C.A., Sangorrín, M.P., 2010. Optimization of killer assays for yeast selection protocols. Rev.
825 Argent. Microbiol. 42, 298-306. <https://doi.org/10.1590/S0325-75412010000400011>.

826

827 López-Seijas, J., García-Fraga, B., da Silva, A. F., Sieiro, C., 2020. Wine Lactic Acid Bacteria with
828 antimicrobial activity as potential biocontrol agents against *Fusarium oxysporum* f. sp. *lycopersici*.
829 Agronomy. 10(1), 31. <https://doi.org/10.3390/agronomy10010031>.

830

831 Mazzucco, M. B., Ganga, M. A., Sangorrín, M. P., 2019. Production of a novel killer toxin from
832 *Saccharomyces eubayanus* using agro-industrial waste and its application against wine spoilage
833 yeasts. Antonie Leeuwenhoek. 112(7), 965-973. <https://doi.org/10.1007/s10482-019-01231-5>.

834

835 Mannazzu, I., Domizio, P., Carboni, G., Zara, S., Zara, G., Comitini, F., Budroni, M., Ciani, M., 2019.
836 Yeast killer toxins: from ecological significance to application, Crit. Rev. Biotechnol. 35 (5), 603-
837 617. <https://doi.org/10.1080/07388551.2019.1601679>.

838

839 Mehlomakulu, N. N., Prior, K. J., Setati, M. E., Divol, B., 2017. *Candida pyralidae* killer toxin
840 disrupts the cell wall of *Brettanomyces bruxellensis* in red grape juice. J. Appl. Microbiol. 122(3),
841 747-758. <https://doi.org/10.1111/jam.13383>.

842

843 Mehlomakulu, N. N., Setati, M. E., Divol, B., 2015. Non-*Saccharomyces* killer toxins: possible
844 biocontrol agents against *Brettanomyces* in wine?. S. Afr. J. Enol. Vitic. 36(1), 94-104.

845

846 Melkonian, C., Gottstein, W., Blasche, S., Kim, Y., Abel-Kistrup, M., Swiegers, H., Saerens, S.,
847 Edwards, N., Patil, K. R., Teusink, B., Molenaar, D., 2019. Finding functional differences between
848 species in a microbial community: case studies in wine fermentation and Kefir culture. Front.
849 Microbiol. 10, 1347. <https://doi.org/10.3389/fmicb.2019.01347>.

850

851 Melvydas, V., Svediene, J., Skridlaite, G., Vaiciuniene, J., Garjonyte, R., 2020. *In vitro* inhibition of
852 *Saccharomyces cerevisiae* growth by *Metschnikowia* spp. triggered by fast removal of iron via two
853 ways. Braz. J. Microbiol., 51(4), 1953-1964. <https://doi.org/10.1007/s42770-020-00357-3>.

854

855 Mendoza, L. M., de Nadra, M. C. M., Farías, M. E., 2010. Antagonistic interaction between yeasts
856 and lactic acid bacteria of oenological relevance: partial characterization of inhibitory compounds
857 produced by yeasts. Food Res. Int. 43(8), 1990-1998. <https://doi.org/10.1016/j.foodres.2010.05.017>.

858

859 Mewa-Ngongang, M., du Plessis, H. W., Ntwampe, S. K. O., Chidi, B. S., Hutchinson, U. F., Mekuto,
860 L., Jolly, N. P., 2019a. The use of *Candida pyralidae* and *Pichia kluyveri* to control spoilage
861 microorganisms of raw fruits used for beverage production. Foods. 8(10), 454.
862 <https://doi.org/10.3390/foods8100454>.

863

864 Mewa-Ngongang, M., du Plessis, H. W., Ntwampe, S. K., Chidi, B. S., Hutchinson, U. F., Mekuto,
865 L., Jolly, N. P., 2019b. Grape pomace extracts as fermentation medium for the production of potential
866 biopreservation compounds. Foods. 8(2), 51. <https://doi.org/10.3390/foods8020051>.

867

868 Mira de Orduña R., 2010. Climate change associated effects on grape and wine quality and
869 production. Food Res. Int. 43:1844-1855. doi:10. 1016/j.foodres.2010.05.001.

870

871 Mondello, V., Songy, A., Battiston, E., Pinto, C., Coppin, C., Trotel-Aziz, P., Clément, C., Mugnai,
872 L., Fontaine, F., 2018. Grapevine trunk diseases: a review of fifteen years of trials for their control
873 with chemicals and biocontrol agents. Plant Dis. 102(7), 1189-1217. [https://doi.org/10.1094/PDIS-](https://doi.org/10.1094/PDIS-08-17-1181-FE)
874 08-17-1181-FE.

875

876 Morata, A., Loira, I., Tesfaye, W., Bañuelos, M. A., González, C., Suárez Lepe, J. A., 2018.
877 *Lachancea thermotolerans* applications in wine technology. *Fermentation*. 4(3), 53.
878 <https://doi.org/10.3390/fermentation4030053>.
879
880 Morata, A., Loira, I., Escott, C., del Fresno, J. M., Bañuelos, M. A., Suárez-Lepe, J. A., 2019a.
881 Applications of *Metschnikowia pulcherrima* in Wine Biotechnology. *Fermentation*. 5(3), 63.
882 <https://doi.org/10.3390/fermentation5030063>.
883
884 Morata, A., 2019b. Enological Repercussions of Non-*Saccharomyces* Species in Wine
885 Biotechnology. *Fermentation*. 5, 72. <https://doi.org/10.3390/fermentation5030072>.
886
887 Mukherjee, A., Verma, J. P., Gaurav, A. K., Chouhan, G. K., Patel, J. S., Hesham, A. E. L., 2020.
888 Yeast a potential bio-agent: future for plant growth and postharvest disease management for
889 sustainable agriculture. *Appl. Microbiol. Biotechnol.* 104, 1497-1510.
890 <https://doi.org/10.1007/s00253-019-10321-3>.
891
892
893 Nally, M. C., Ponsone, M. L., Pesce, V. M., Toro, M. E., Vazquez, F., Chulze, S., 2018. Evaluation
894 of behaviour of *Lachancea thermotolerans* biocontrol agents on grape fermentations. *Lett. Appl.*
895 *Microbiol.* 67(1), 89-96. <https://doi.org/10.1111/lam.13001>.
896
897 Nardi, T., 2020. Microbial resources as a tool for enhancing sustainability in winemaking.
898 *Microorganisms*. 8(4), 507. <https://doi.org/10.3390/microorganisms8040507>.
899

900 Ndlovu, B., Schoeman, H., Franz, C. M. A. P., Du Toit, M., 2015. Screening, identification and
901 characterization of bacteriocins produced by wine-isolated LAB strains. J. Appl. Microbiol. 118(4),
902 1007-1022. <https://doi.org/10.1111/jam.12752>.

903

904 Nehme, N., Mathieu, F., Taillandier, P., 2010. Impact of the co-culture of *Saccharomyces cerevisiae*
905 – *Oenococcus oeni* on malolactic fermentation and partial characterization of a yeast-derived
906 inhibitory peptidic fraction. Food Microbiol. 27, 150-57. <https://doi.org/10.1016/j.fm.2009.09.008>.

907

908 Nel, H. A., Bauer, R., Wolfaardt, G. M., Dicks, L. M. T., 2002. Effect of bacteriocins pediocin PD-
909 1, plantaricin 423, and nisin on biofilms of *Oenococcus oeni* on a stainless steel surface. Am J Enol
910 Vitic. 53, 191-196.

911

912 Nissen, P., Arneborg, N., 2003. Characterization of early deaths of non-*Saccharomyces* yeasts in
913 mixed cultures with *Saccharomyces cerevisiae*. Arch. Microbiol. 180(4), 257-263.
914 <https://doi.org/10.1007/s00203-003-0585-9>.

915

916 Oro, L., Ciani, M., Comitini, F., 2014. Antimicrobial activity of *Metschnikowia pulcherrima* on wine
917 yeasts. J. Appl. Microbiol. 116, 1209-1217. <https://doi.org/10.1111/jam.12446>.

918

919 Oro, L., Ciani, M., Bizzaro, D., Comitini, F., 2016. Evaluation of damage induced by Kwkt and Pikt
920 zymocins against *Brettanomyces/Dekkera* spoilage yeast, as compared to sulphur dioxide. J. Appl.
921 Microbiol. 121(1), 207-214. <https://doi.org/10.1111/jam.13121>.

922

923 Osborne, J. P., Edwards, C. G., 2007. Inhibition of malolactic fermentation by a peptide produced by
924 *Saccharomyces cerevisiae* during alcoholic fermentation. Int. J. Food Microbiol. 118, 27-34.
925 <https://doi.org/10.1016/j.ijfoodmicro.2007.05.007>.

926

927 Pandin, C., Le Coq, D., Canette, A., Aymerich, S., Briandet, R., 2017. Should the biofilm mode of
928 life be taken into consideration for microbial biocontrol agents?. *Microb. Biotechnol.* 10(4), 719-734.
929 <https://doi.org/10.1111/1751-7915.12693>.

930

931 Parafati, L., Vitale, A., Restuccia, C., Cirvillieri, G., 2015. Biocontrol ability and action mechanism
932 of food-isolated yeast strains against *Botrytis cinerea* causing post-harvest bunch rot of table grape.
933 *Food Microbiol.* 47, 85-92. <https://doi.org/10.1016/j.fm.2014.11.013>.

934

935 Parapouli, M., Vasileiadis, A., Afendra, A. S., Hatziloukas, E., 2020. *Saccharomyces cerevisiae* and
936 its industrial applications. *AIMS microbial.* 6(1), 1-31. <https://doi.org/10.3934/microbiol.2020001>.

937

938 Parijs, I., Steenackers, H. P., 2018. Competitive inter-species interactions underlie the increased
939 antimicrobial tolerance in multispecies brewery biofilms. *ISME J.* 12(8), 2061-2075.
940 <https://doi.org/10.1038/s41396-018-0146-5>.

941

942 Pawlikowska, E., James, S. A., Breierova, E., Antolak, H., Kregiel, D., 2019. Biocontrol capability
943 of local *Metschnikowia* sp. isolates. *Antonie Leeuwenhoek.* 112(10), 1425-1445.
944 <https://doi.org/10.1007/s10482-019-01272-w>.

945

946 Pawlowska, A. M., Zannini, E., Coffey, A., Arendt, E. K., 2012. "Green preservatives": combating
947 fungi in the food and feed industry by applying antifungal lactic acid bacteria. In Jeyakumar H. (Ed.),
948 *Advances in food and nutrition research.* Academic Press, New York, Vol. 66, pp. 217-238.

949

950 Peña, R., Ganga, M. A., 2019. Novel antimicrobial peptides produced by *Candida intermedia*
951 LAMAP1790 active against the wine-spoilage yeast *Brettanomyces bruxellensis*. *Antonie*
952 *Leeuwenhoek*. 112(2), 297-304. <https://doi.org/10.1007/s10482-018-1159-9>.
953
954 Peng, C., Andersen, B., Arshid, S., Larsen, M. R., Albergaria, H., Lametsch, R., Arneborg, N., 2019.
955 Proteomics insights into the responses of *Saccharomyces cerevisiae* during mixed-culture alcoholic
956 fermentation with *Lachancea thermotolerans*. *FEMS Microbiol. Ecol.* 95(9), fiz126.
957 <https://doi.org/10.1093/femsec/fiz126>.
958
959 Pérez-Navado, F., Albergaria, H., Hogg, T., Gírio, F., 2006. Cellular death of two non-
960 *Saccharomyces* wine-related yeasts during mixed fermentations with *Saccharomyces cerevisiae*. *Int.*
961 *J. Food Microbiol.* 108, 336-345. <https://doi.org/10.1016/j.ijfoodmicro.2005.12.012>.
962
963 Pérez-Torrado, R., Rantsiou, K., Perrone, B., Navarro-Tapia, E., Querol, A., Cocolin, L., 2017.
964 Ecological interactions among *Saccharomyces cerevisiae* strains: insight into the dominance
965 phenomenon. *Sci. Rep.* 7, 43603. <https://doi.org/10.1038/srep43603>.
966
967 Petitgonnet, C., Klein, G. L., Roullier-Gall, C., Schmitt-Kopplin, P., Quintanilla-Casas, B., Vichi, S.,
968 Julien-Davi, D., Alexandre, H., 2019. Influence of cell-cell contact between *L. thermotolerans* and *S.*
969 *cerevisiae* on yeast interactions and the exo-metabolome. *Food Microbiol.* 83, 122-133.
970 <https://doi.org/10.1016/j.fm.2019.05.005>.
971
972 Pietrafesa, A., Capece, A., Pietrafesa, R., Bely, M., Romano, P., 2020. *Saccharomyces cerevisiae* and
973 *Hanseniaspora uvarum* mixed starter cultures: influence of microbial/physical interactions on wine
974 characteristics. *Yeast.* 37(11), 609-621. <https://doi.org/10.1002/yea.3506>.
975

976 Porter, T. J., Divol, B., Setati, M. E., 2019. *Lachancea* yeast species: Origin, biochemical
977 characteristics and oenological significance. *Food Res. Int.* 119, 378-389.
978 <https://doi.org/10.1016/j.foodres.2019.02.003>.
979

980 Prendes, L. P., Merín, M. G., Fontana, A. R., Bottini, R. A., Ramirez, M. L., de Ambrosini, V. I. M.,
981 2018. Isolation, identification and selection of antagonistic yeast against *Alternaria alternata*
982 infection and tenuazonic acid production in wine grapes from Argentina. *Int. J. Food Microbiol.* 266,
983 14-20. <https://doi.org/10.1016/j.ijfoodmicro.2017.10.033>.
984

985 Prior, K. J., Bauer, F. F., Divol, B., 2019. The utilisation of nitrogenous compound by commercial
986 non-*Saccharomyces* yeasts associated with wine. *Food Microbiol.* 79, 75-84.
987 <https://doi.org/10.1016/j.fm.2018.12.002>.
988

989 Pu, L., Jingfan, F., Kai, C., Chao-an, L., Yunjiang, C., 2014. Phenylethanol promotes adhesion and
990 biofilm formation of the antagonistic yeast *Kloeckera apiculata* for the control of blue mold on citrus.
991 *FEMS Yeast Res.* 14(4), 536-546. <https://doi.org/10.1111/1567-1364.12139>.
992

993 Rauhut, D., 2017. Usage and formation of sulphur compounds. In *Biology of Microorganisms on*
994 *Grapes, in Must and in Wine*, Springer, Cham., pp. 255-291..
995

996 Renault, P. E., Albertin, W., Bely, M., 2013. An innovative tool reveals interaction mechanisms
997 among yeast populations under oenological conditions. *Appl. Microbiol. Biotechnol.* 97, 4105-4119.
998 <https://doi.org/10.1007/s00253-012-4660-5>.
999

1000 Restuccia, D., Loizzo, M. R., Spizzirri, U. G., 2018. Accumulation of biogenic amines in wine: role
1001 of alcoholic and malolactic fermentation. *Ferment.* 4(1), 6.
1002 <https://doi.org/10.3390/fermentation4010006>.
1003

1004 Rizk, Z., El Rayess, Y., Ghanem, C., Mathieu, F., Taillandier, P., Nehme, N., 2016. Impact of
1005 inhibitory peptides released by *Saccharomyces cerevisiae* BDX on the malolactic fermentation
1006 performed by *Oenococcus oeni* Vitilactic F. *Int. J. Food Microbiol.* 233, 90-96.
1007 <https://doi.org/10.1016/j.ijfoodmicro.2016.06.018>.
1008

1009 Roca-Mesa, H., Sendra, S., Mas, A., Beltran, G., Torija, M. J., 2020. Nitrogen preferences during
1010 alcoholic fermentation of different non-*Saccharomyces* yeasts of oenological interest.
1011 *Microorganisms*, 8(2), 157. <https://doi.org/10.3390/microorganisms8020157>.
1012

1013 Rojo-Bezares, B., Saenz, Y., Zarazaga, M., Torres, C., Ruiz-Larrea, F., 2007. Antimicrobial activity
1014 of nisin against *Oenococcus oeni* and other wine bacteria. *Int. J. Food Microbiol.* 116(1):32-36.
1015 <https://doi.org/10.1016/j.ijfoodmicro.2006.12.020>.
1016

1017 Rollero, S., Bloem, A., Ortiz-Julien, A., Camarasa, C., Divol, B., 2018. Altered fermentation
1018 performances, growth, and metabolic footprints reveal competition for nutrients between yeast
1019 species inoculated in synthetic grape juice-like medium. *Front. Microbiol.* 9, 196.
1020 <https://doi.org/10.3389/fmicb.2018.00196>.
1021

1022 Rossouw, D., Bagheri, B., Setati, M. E., Bauer, F. F., 2015. Co-flocculation of yeast species, a new
1023 mechanism to govern population dynamics in microbial ecosystems. *PloS one*, 10(8).
1024 <https://doi.org/10.1371/journal.pone.0136249>.
1025

1026 Rossouw, D., Du Toit, M., Bauer, F. F., 2012. The impact of co-inoculation with *Oenococcus oeni*
1027 on the transcriptome of *Saccharomyces cerevisiae* and on the flavour-active metabolite profiles during
1028 fermentation in synthetic must. *Food Microbiol.* 29(1), 121-131.
1029 <https://doi.org/10.1016/j.fm.2011.09.006>.
1030
1031 Rossouw, D., Meiring, S. P., Bauer, F. F., 2018. Modifying *Saccharomyces cerevisiae* adhesion
1032 properties regulates yeast ecosystem dynamics. *mSphere.* 3(5), e00383-18.
1033 <https://doi.org/10.1128/mSphere.00383-18>.
1034
1035 Roullier-Gall, C., Hemmler, D., Gonsior, M., Li, Y., Nikolantonaki, M., Aron, A., Coelho, C.,
1036 Gougeon, R. D., Schmitt-Kopplin, P., 2017. Sulfites and the wine metabolome. *Food Chem.* 237,
1037 106-113. <https://doi.org/10.1016/j.foodchem.2017.05.039>.
1038
1039 Rubio-Bretón, P., Gonzalo-Diago, A., Iribarren, M., Garde-Cerdán, T., Pérez-Álvarez, E. P., 2018.
1040 Bioprotection as a tool to free additives winemaking: Effect on sensorial, anthocyanic and aromatic
1041 profile of young red wines. *LWT-Food Sci. Technol.* 98, 458-464.
1042 <https://doi.org/10.1016/j.lwt.2018.08.050>.
1043
1044 Santos, A., Navascués, E., Bravo, E., Marquina, D., 2011. *Ustilago maydis* killer toxin as a new tool
1045 for the biocontrol of the wine spoilage yeast *Brettanomyces bruxellensis*. *Int. J. Food Microbiol.*
1046 145(1), 147-154. <https://doi.org/10.1016/j.ijfoodmicro.2010.12.005>.
1047
1048 Saravanakumar, D., Spadaro, D., Garibaldi, A., Gullino, M. L., 2009. Detection of enzymatic activity
1049 and partial sequence of a chitinase gene in *Metschnikowia pulcherrima* strain MACH1 used as post-
1050 harvest biocontrol agent. *Eur. J. Plant Pathol.* 123(2), 183-193. [https://doi.org/10.1007/s10658-008-](https://doi.org/10.1007/s10658-008-9355-5)
1051 9355-5.

1052

1053 Saravanakumar, D., Ciavorella, A., Spadaro, D., Garibaldi, A., Gullino, M. L., 2008. *Metschnikowia*
1054 *pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and *Penicillium*
1055 *expansum* in apples through iron depletion. *Postharvest Biol. Technol.* 49, 121-128.
1056 <https://doi.org/10.1016/j.postharvbio.2007.11.006>.

1057

1058 Schmitt, M. J., Breinig, F., 2002. The viral killer system in yeast: from molecular biology to
1059 application. *FEMS Microbiol. Rev.* 26(3), 257-276. [https://doi.org/10.1111/j.1574-](https://doi.org/10.1111/j.1574-6976.2002.tb00614.x)
1060 [6976.2002.tb00614.x](https://doi.org/10.1111/j.1574-6976.2002.tb00614.x).

1061

1062 Shekhawat, K., Patterson, H., Bauer, F. F., Setati, M. E., 2019. RNA-seq based transcriptional analysis
1063 of *Saccharomyces cerevisiae* and *Lachancea thermotolerans* in mixed-culture fermentations under
1064 anaerobic conditions. *BMC Genomics.* 20(1), 145. <https://doi.org/10.1186/s12864-019-5511-x>.

1065

1066 Siedler, S., Balti, R., Neves, A. R., 2019. Bioprotective mechanisms of lactic acid bacteria against
1067 fungal spoilage of food. *Curr. Opin. Biotechnol.* 56, 138-146.
1068 <https://doi.org/10.1016/j.copbio.2018.11.015>.

1069

1070 Siedler, S., Rau, M. H., Bidstrup, S., Vento, J. M., Aunbjerg, S. D., Bosma, E. F., McNair, L. M.,
1071 Beisel, C. L., Neves, A. R., 2020. Competitive exclusion is a major bioprotective mechanism of
1072 lactobacilli against fungal spoilage in fermented milk products. *Appl. Environ. Microbiol.* 86(7),
1073 e02312-19. <https://doi.org/10.1128/AEM.02312-19>

1074

1075 Simonin, S., Alexandre, H., Nikolantonaki, M., Coelho, C., Tourdot-Maréchal, R., 2018. Inoculation
1076 of *Torulaspora delbrueckii* as a bio-protection agent in winemaking. *Food Res. Int.* 107, 451-461.
1077 <https://doi.org/10.1016/j.foodres.2018.02.034>

1078

1079 Singh, V. P., 2018. Recent approaches in food bio-preservation-a review. *Open. Vet. J.* 8(1), 104-
1080 111. <https://doi.org/10.4314/ovj.v8i1.16>.

1081

1082 Sipiczki, M., 2006. *Metschnikowia* strains isolated from botrytized grapes antagonize fungal and
1083 bacterial growth by iron depletion. *Appl. Environ. Microbiol.* 72(10), 6716-6724.
1084 <https://doi.org/10.1128/AEM.01275-06>.

1085

1086 Sipiczki, M., 2016. Overwintering of vineyard yeasts: survival of interacting yeast communities in
1087 grapes mummified on vines. *Front. Microbiol.* 7, 212. <https://doi.org/10.3389/fmicb.2016.00212>

1088

1089 Sipiczki, M., 2020. *Metschnikowia pulcherrima* and related pulcherrimin-producing yeasts: fuzzy
1090 species boundaries and complex antimicrobial antagonism. *Microorganisms.* 8(7), 1029.
1091 <https://doi.org/10.3390/microorganisms8071029>.

1092

1093 Su, Y., Seguinot, P., Sanchez, I., Ortiz-Julien, A., Heras, J. M., Querol, A., Camarasa, C., Guillamón,
1094 J. M., 2020. Nitrogen sources preferences of non-*Saccharomyces* yeasts to sustain growth and
1095 fermentation under winemaking conditions. *Food Microbiol.* 85, 103287.
1096 <https://doi.org/10.1016/j.fm.2019.103287>.

1097

1098 Taillandier, P., Lai, Q. P., Julien-Ortiz, A., Brandam, C., 2014. Interactions between *Torulaspota*
1099 *delbrueckii* and *Saccharomyces cerevisiae* in wine fermentation: influence of inoculation and
1100 nitrogen content. *World J. Microbiol. Biotechnol.* 30, 1959-1967. [https://doi.org/10.1007/s11274-](https://doi.org/10.1007/s11274-014-1618-z)
1101 [014-1618-z](https://doi.org/10.1007/s11274-014-1618-z).

1102

1103 Todd, B. E. N., Fleet, G. H., Henscke, P. A., 2000. Promotion of autolysis through the interaction of
1104 killer and sensitive yeasts: potential application in sparkling wine production. *Am. J. Enol. Vitic.* 51,
1105 65-72.
1106

1107 Tremonte, P., Pannella, G., Succi, M., Tipaldi, L., Sturchio, M., Coppola, R., Luongo, D., Sorrentino,
1108 E., 2017. Antimicrobial activity of *Lactobacillus plantarum* strains isolated from different
1109 environments: a preliminary study. *Int. Food Res. J.* 24(2), 852-859.
1110

1111 Tronchoni, J., Curiel, J. A., Morales, P., Torres-Pérez, R., Gonzalez, R., 2017. Early transcriptional
1112 response to biotic stress in mixed starter fermentations involving *Saccharomyces cerevisiae* and
1113 *Torulaspora delbrueckii*. *Int. J. Food Microbiol.* 241, 60-68. [https://doi.org/](https://doi.org/10.1016/j.ijfoodmicro.2016.10.017)
1114 [10.1016/j.ijfoodmicro.2016.10.017](https://doi.org/10.1016/j.ijfoodmicro.2016.10.017).
1115

1116 Váchová, L., Palková, Z., 2018. How structured yeast multicellular communities live, age and die?.
1117 *FEMS Yeast Res.* 18(4), foy033. <https://doi.org/10.1093/femsyr/foy033>.
1118

1119 Varela, C., Borneman, A. R., 2017. Yeasts found in vineyards and wineries. *Yeast* 34(3), 111-128.
1120 <https://doi.org/10.1002/yea.3219>.
1121

1122 Velázquez, R., Zamora, E., Álvarez, M., Álvarez, M. L., Ramírez, M., 2016. Using mixed inocula of
1123 *Saccharomyces cerevisiae* killer strains to improve the quality of traditional sparkling-wine. *Food*
1124 *Microbiol.* 59, 150-160. <https://doi.org/10.1016/j.fm.2016.06.006>.
1125

1126 Velázquez, R., Zamora, E., Álvarez, M. L., Ramírez, M., 2019. Using *Torulaspora delbrueckii* killer
1127 yeasts in the elaboration of base wine and traditional sparkling wine. *Int. J. Food Microbiol.* 289, 134-
1128 144. <https://doi.org/10.1016/j.ijfoodmicro.2018.09.010>.

1129

1130 Vilela, A., 2018. *Lachancea thermotolerans*, the non-*Saccharomyces* yeast that reduces the volatile
1131 acidity of wines. *Fermentation*. 4(3), 56. <https://doi.org/10.3390/fermentation4030056>.

1132

1133 Vilela, A., 2019. Use of nonconventional yeasts for modulating wine acidity. *Fermentation*. 5(1), 27.
1134 <https://doi.org/10.3390/fermentation5010027>.

1135

1136 Villalba, M. L., Sáez, J. S., Del Monaco, S., Lopes, C. A., Sangorrín, M. P., 2016. TdKT, a new killer
1137 toxin produced by *Torulaspota delbrueckii* effective against wine spoilage yeasts. *Int. J. Food*
1138 *Microbiol.* 217, 94-100. <https://doi.org/10.1016/j.ijfoodmicro.2015.10.006>.

1139

1140 Wang, C., Mas, A., Esteve-Zarzoso, B., 2015. Interaction between *Hanseniaspora uvarum* and
1141 *Saccharomyces cerevisiae* during alcoholic fermentation. *Int. J. Food Microbiol.* 206, 67-74.
1142 <https://doi.org/10.1016/j.ijfoodmicro.2015.04.022>.

1143

1144 Wang, C., Mas, A., Esteve-Zarzoso, B., 2016. The interaction between *Saccharomyces cerevisiae*
1145 and non-*Saccharomyces* yeast during alcoholic fermentation is species and strain specific. *Front.*
1146 *Microbiol.* 7, 502. <https://doi.org/10.3389/fmicb.2016.00502>.

1147

1148 Wisniewski, M., Biles, C., Droby, S., McLaughlin, R., Wilson, C., Chalutz, E., 1991. Mode of action
1149 of the postharvest biocontrol yeast *Pichia guillermondii*. I. Characterization of attachment to *Botrytis*
1150 *cinerea*. *Physiol. Mol. Plant Pathol.* 39(4), 245-258. [https://doi.org/10.1016/0885-5765\(91\)90033-E](https://doi.org/10.1016/0885-5765(91)90033-E).

1151

1152 Zagorc, T., Maráz, A., Cadez, N., Jemec, K. P., Péter, G., Resnik, M., Nemanič, J., Raspor, P., 2001.
1153 Indigenous wine killer yeasts and their application as a starter culture in wine fermentation. *Food*
1154 *Microbiol.* 18(4), 441-451. <https://doi.org/10.1006/fmic.2001.0422>.

1155

1156 Zhang, H., Du, H., Xu, Y., 2021. Volatile organic compound-mediated antifungal activity of *Pichia*
1157 spp. and its effect on the metabolic profiles of fermentation communities. *Appl. Environ. Microbiol.*
1158 87(9), e02992-20. <https://doi.org/10.1128/AEM.02992-20>.

1159

1160 Zhang, C., Yang, M., Ericsson, A. C., 2019. Antimicrobial peptides: potential application in liver
1161 cancer. *Front. Microbiol.* 10, 1257. <https://doi.org/10.3389/fmicb.2019.01257>.

1162

1163 Zhao, X., Wu, H., Lu, H., Li, G., Huang, Q., 2013. LAMP: a database linking antimicrobial peptides.
1164 *PloS one.* 8(6), e66557. <https://doi.org/10.1371/journal.pone.0066557>.

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181 **Figure captions**

1182 **Fig. 1 Antagonistic behaviour demonstrated in wine microorganisms. Antagonistic behaviour**
1183 **was investigated *in vitro* (A), on grapes (B) and in wine (C).**

1184 **Fig. 2 Antagonistic strategies of wine microorganisms**

1185

1186

1187 **Table 1**

1188 Role of starter cultures in wine and their bioprotective strategies

Yeasts	Starter ability	References	Bioprotective mechanisms		References
			Passive strategies	Active strategies	
<i>Saccharomyces cerevisiae</i>	- Alcoholic fermentation conduction - Positive impact on the organoleptic quality of wines	Parapouli et al., 2020	- Fast nutrient uptake - Multicellular consortia formation	- Toxic compounds production - Ethanol - SO ₂ - Antimicrobial peptides (saccharomycin; killer toxins) - VOCs (higher alcohols) - Cell-to-cell contact	Alonso-del-Real et al., 2019 Tronchoni et al., 2017 Rossouw et al., 2015; 2018 Albergaria and Arneborg, 2016 González et al., 2018 Englezos et al., 2019b
<i>Torulaspota delbrueckii</i>	- Low acetic acid production - Reduction of ethanol content - Increase of glycerol content - Release of mannoproteins and polysaccharides - Production of high levels of fruity esters, thiols, and terpenes and lower amounts of higher alcohols	Benito, 2018a	- Fast nutrient uptake	- Toxic compounds production - Ethanol - Antimicrobial peptides (Tdk Killer toxin)	Prior et al., 2019 Su et al., 2020 Simonin et al., 2018 Villaba et al., 2016

<i>Metschnikowia pulcherrima</i>	<ul style="list-style-type: none"> - Low acetic acid production - Ethanol content reduction - Increase of glycerol content - Production of high levels of esters and higher alcohols (isobutanol and phenylethanol) - Enhances varietal and fruity aromas 	Morata et al., 2019a	<ul style="list-style-type: none"> - Iron immobilization (pulcherrimic acid production) - high growth rate and a short lag phase - Biofilm formation 	<ul style="list-style-type: none"> - Toxic compounds production - Ethanol - Lytic enzymes - VOCs (ethyl alcohol and ethyl acetate) 	<ul style="list-style-type: none"> Sipiczki, 2006, Sipiczki, 2020 Gore-Lloyd et al., 2019 Kunchen et al., 2019 Cordero-Bueso et al., 2017 Pawlikowska et al., 2019 Contarino et al., 2019 Saravanakumar et al., 2009
<i>Lachancea thermotolerans</i>	<ul style="list-style-type: none"> - Increase of titratable acidity - Ethanol content reduction - Increase of glycerol content - Production of high levels of ethyl lactate, ethyl hexanoate and 2-phenylethanol - Reduction of phenylethyl acetate 	Benito, 2018b Morata et al., 2019b Porter et al., 2019	<ul style="list-style-type: none"> - Fast nutrient uptake - Short lag phase and ability to survive until the end of fermentation - Filamentous growth/ flocculation 	<ul style="list-style-type: none"> - Toxic compounds production - Ethanol - Lactic acid - Killer toxins - VOCs (2- phenylethanol) 	<ul style="list-style-type: none"> Fiori et al., 2014 Farbo et al., 2018 Shekhawat et al., 2019 Rubio-Bretón et al., 2018 Vilela, 2018 Nally et al., 2018
<i>Pichia kluyveri</i>	<ul style="list-style-type: none"> - Ethanol content reduction - Increase of glycerol content - Production of thiols (hotrienol and linalool oxide) - enhance varietal aromas 	Varela and Borneman, 2017	<ul style="list-style-type: none"> - Biofilm formation 	<ul style="list-style-type: none"> - Toxic compounds production - Ethanol - Killer toxins - Lytic enzymes - VOCs (isoamyl acetate, isoamyl alcohol, 2-phenyl 	<ul style="list-style-type: none"> Cordero-Bueso et al., 2017 Sipiczki, 2016 Prior et al., 2019 Labbani et al., 2015 Mewa-Ngongang et al., 2019a,b Englezos et al. (under preparation)

ethylacetate and 2-phenylethanol)

<i>Starmerella bacillaris</i>	- Ethanol content reduction - Increase of glycerol content - Production of high levels of linalool, organic acids	Englezos et al., 2017	- Relative fast ammonium uptake	- Toxic compounds production - Ethanol - Killer toxins - Lytic enzymes - VOCs (benzyl alcohol)	Prendes et al., 2018 Lemos Junior et al., 2016, 2020 Englezos et al., 2018, 2019b Binati et al., 2020 Roca Mesa et al., 2020 Bagheri et al., 2018
-------------------------------	---	-----------------------	---------------------------------	--	--

Lactic Acid


Bacteria

<i>Oenococcus oeni</i>	- Decrease of titratable acidity - Diacetyl production – ‘buttery’ flavour - Release of secondary metabolites that impact on wine aroma and flavor		- Better adaptation to wine conditions - Biofilm formation	- Toxic compounds production - Lactic acid - BLIS	Dimopoulou et al., 2015, 2018 Bastard et al., 2016 Knoll et al., 2008 Lasik-Kurdyś and Sip, 2019
<i>Lactiplantibacillus plantarum</i>	- Titratable acidity decrease - Diacetyl production – ‘buttery’ flavour	Krieger-Weber et al., 2020 Brizuela et al., 2019	- Competitive exclusion - Nutrient competition - Biofilm formation	- Toxic compounds production - Lactic acid - Bacteriocins	Rubio-Breton et al., 2018 Melkonian et al., 2019 Sielder et al., 2020 Tremonte et al., 2017 Bartle et al., 2019

- Positive effects on the organoleptic properties of wine due to a diverse array of enzymes
- Improvement of color in red wines

1190 **Figure captions**


1191 **Fig. 1**



1192

1193 **Fig. 2**

1194



1195

1196

1197

1198

1199