

RESEARCH ARTICLE

Unintentional injuries and potential determinants of falls in young children: Results from the Piccolipiù Italian birth cohort

Martina Culasso^{1*}, Daniela Porta¹, Sonia Brescianini², Luigi Gagliardi³, Paola Michelozzi¹, Costanza Pizzi⁴, Luca Ronfani⁵, Franca Rusconi⁶, Liza Vecchi Brumatti⁵, Federica Asta¹

1 Department of Epidemiology, Lazio Regional Health Service—ASL ROMA1, Rome, Italy, **2** Center for Behavioral Science and Mental Health, Istituto Superiore di Sanità, Rome, Italy, **3** Division of Neonatology and Pediatrics, Ospedale Versilia, Viareggio, AUSL Toscana Nord Ovest, Pisa, Italy, **4** Department of Medical Sciences, University of Turin, Turin, Italy, **5** Institute for Maternal and Child Health—IRCCS "Burlo Garofolo", Trieste, Italy, **6** AUSL Toscana Nord Ovest, Pisa, Italy

* m.culasso@deplazio.it



OPEN ACCESS

Citation: Culasso M, Porta D, Brescianini S, Gagliardi L, Michelozzi P, Pizzi C, et al. (2022) Unintentional injuries and potential determinants of falls in young children: Results from the Piccolipiù Italian birth cohort. *PLoS ONE* 17(10): e0275521. <https://doi.org/10.1371/journal.pone.0275521>

Editor: Angela Lupattelli, University of Oslo, NORWAY

Received: November 26, 2021

Accepted: September 19, 2022

Published: October 3, 2022

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0275521>

Copyright: © 2022 Culasso et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Despite the Piccolipiù data cannot be publicly shared because of privacy concerns, we will be pleased to share de-identified data upon request. Data requests may be sent to

Abstract

Objectives

Unintentional injuries such as falls, are particularly frequent in early childhood. To date, epidemiological studies in this field have been carried out using routine data sources or registries and many studies were observational studies with a cross-sectional design. The aims of the study are to describe unintentional injuries in the first two years of life in the Piccolipiù birth cohort, and to investigate the association between mother and children characteristics and the First Event of Raised surface Fall (FERF).

Methods

This longitudinal observational study included 3038 children from an Italian birth cohort. Data on socio-demographic factors, socio-economic indicators, maternal health and lifestyle characteristics and child's sleeping behavior, obtained from questionnaires completed at birth, 12 and 24 months of age, were considered in the analyses as potential risk factors of FERF. Time of occurrence of FERF was analyzed using the Kaplan-Meier method. The multivariable analysis for time to event was carried out using a Cox proportional hazards model.

Results

Falls from raised surfaces are the leading cause of unintentional injuries in the cohort with 610 (21.1%) and 577 (20.0%) cases among children during the first and second year of life, respectively. An increased risk of FERF was associated with several risk factors: maternal psychological distress (HR 1.41, 95%CI 1.10–1.81), maternal alcohol intake (HR 1.26, 95% CI 1.10–1.45), and child's sleeping problems (HR 1.28, 95%CI 1.09–1.51). Children with older aged mothers (HR 0.98, 95%CI 0.96–0.99) and living in northern Italy (HR 0.64, 95% CI 0.55–0.75) had a lower risk of FERF.

the following health institution: ASL Roma1 Borgo Santo Spirito 3, 00193 Rome Italy email: dipepi@deplazio.it.

Funding: The Piccolipiù study was approved and initially funded by the Italian National Centre for Disease Prevention and Control (CCM grant 2010) and by the Italian Ministry of Health (art 12 and 12bis D.lgs 502/92). This work was supported by the Italian Ministry of Health, through the contribution given to the Institute for Maternal and Child Health IRCCS Burlo Garofolo, Trieste, Italy (RC 12/12). Funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Conclusion

The results of the study suggest that a higher risk of FERF is associated with socio-demographic factors, maternal characteristics and child sleeping behavior that could hinder parent empowerment.

Introduction

Unintentional injuries are a leading cause of death among children, an important health threat and a public health issue [1]. The pattern and etiology of injuries and their outcome vary substantially within populations and across countries but worldwide, approximately 950,000 children die every year due to unintentional injury [2]. Previous studies investigating the circumstances leading to child mortality following injury have found that most injuries could be prevented [3–5].

Unintentional injuries due to falls, near-drowning and burns are particularly frequent in early childhood, even in high-income countries [6]. A population-based cohort study conducted in Japan estimated that more than 60% of children aged 1.5 years were affected by unintentional injuries in their first year of life [6]. Individual and family-related risk factors associated to child injury from previous studies include: gender (being a male), having a large number of siblings (3 or more children), having mothers with psychological, and/or behavioral problems and having a young mother (aged ≤ 22 years) [7–13].

So far, epidemiological studies on unintentional injuries in children have been mainly carried out using routine data sources and registries [14–17], which included only severe injuries reported to secondary health care service, or have often used the cross-sectional design [8, 12, 18], which is generally considered of low-quality. Moreover, the contribution from birth and child cohorts in this field was focused on school-children and adolescents [19–22]. European studies including preschoolers have been mainly conducted in Northern countries [23–25], and the evidence on Mediterranean countries is limited.

Piccolipiù is a prospective cohort of newborns enrolled in Italy. Since fetal and infant life are periods of rapid development, characterized by high susceptibility to exposures, this prospective cohort was set up to investigate the effects of environmental exposures, parental conditions and social factors acting during pre-natal and early post-natal life on infant and child health and development [26].

The aims of the present study are to describe unintentional injuries in early life, with a particular focus on those due to falls from a raised surface, and to identify potential risk factors associated with the First Event of Raised surface Fall (FERF) during the first two years of life in the Piccolipiù cohort.

Methods

Study population

Piccolipiù is the name of a birth cohort of 3,358 children enrolled in six maternity wards located in five Italian cities (Florence, Rome, Trieste, Turin, and Viareggio) between October 2011 and March 2015 [26]. Women were contacted during pregnancy or at delivery and a written informed consent form for participation was signed by both parents at enrollment. Enrolled mothers were asked to complete a baseline questionnaire, with questions on demographics, environmental exposures, and mother's health. Additional information was obtained

either from medical records or directly from the mother within the 48 hours after delivery. Parents were then contacted at 6, 12, 24 months, 4 and 6 years after delivery and asked to fill in self-administrated questionnaires which included information on demographics, environmental exposures, lifestyle, and mother's and child health. The Piccolipiù study was approved by the Ethics Committee of the Local Health Unit Roma E, national coordinator of the project (Prot. CE/82 09/06/2011), and of each local center [27].

For the purpose of this study, only data obtained at birth ($n = 3,038$), at 12 months ($n = 2,897$) and 24 months ($n = 2,751$) of age were considered. The response rates were 99.4%, 87.2% and 83.8%, respectively.

Study outcomes

Data on unintentional injuries were provided in both 12 and 24 month's questionnaires, where parents reported if their children had experienced at least one of the following injuries in the previous twelve months (multiple answers were allowed): falls from a raised surface (e.g. bed, furniture, table, chair), burns/scalds, poisoning, foreign body ingestion, road traffic injuries and other injuries. Moreover, details about circumstances of each fall from raised surface were collected; in particular where and when the fall happened, who was with the child, type/site of injury (multiple choice answer) and the treatment required.

Potential risk factors

A series of potential risk factors were considered in the analyses, namely: socio-demographic factors, socio-economic indicators, maternal health and lifestyle characteristics and child's sleeping behavior.

The enrollment area (center/north of Italy), child gender, maternal age when the child was born (<30 , $30-34$, $> = 35$), maternal education level, paternal occupation (employed, not employed), maternal smoking (no/yes) during pregnancy and alcohol consumption (no/yes) during pregnancy were obtained from the baseline questionnaire. Maternal educational level was classified in three categories: low (primary school), medium (secondary school) and high (university degree).

All others variables, including maternal occupation (employed, not employed), siblings (no/yes), day care attendance (no/ yes), maternal psychological distress (high distress, moderate distress, feel good), and variables concerning child sleeping patterns, were assessed at 12 months.

Moreover, the Equivalised Household Income Indicator (EHII), was used as indicator of the total disposable monthly household income at birth, standardized by household size and composition in our cohort [28]. Piccolipiù cohort data (maternal age, cohabitation status, country of birth, educational level, occupational status and occupational code; paternal/partner age, country of birth, educational level, occupational status and occupational code; and household size and tenure status) and external data from the Italian 2011 "European Union Statistics on Income and Living Conditions" (EU- SILC) survey [29, 30] were considered. The EHII was reported as a three-level categorical variable by using the 25th (933 Euro) and 75th (1,810 Euro) percentiles of total monthly disposable household income in the Italy-EUSILC survey as cut-off thresholds.

To measure maternal psychological distress, the Italian version of the 12-items General Health Questionnaire (GHQ-12) was administrated at the same time as the 12 months questionnaire. This is a self-administered questionnaire aimed at detecting current levels of general (not psychotic) psychiatric morbidity, mainly in the anxiety/depression spectrum over the past two weeks. It has been used extensively in many community and hospital settings in different

countries, including Italy and showed high validity and reproducibility [31, 32]. Mothers were asked to rate the degree to which they had experienced several symptoms and/or mood states. Answers are reported according to a four-level Likert-type scale (from “not at all” to “much more than usual”). The GHQ-12 was reported on a two-level scale by collapsing the four Likert categories into two (coded 0–1) [31]. The total score was computed by summing up the single item scores. Thus, the theoretical range was 0–12, with higher values indicating more severe distress. We considered scores between 2 and 4 to identify moderate psychological distress, and ≥ 5 to identify severe distress. These cut-offs have previously been adopted in the literature to screen mental problems [31, 33–35].

Children’s sleeping behavior was assessed considering data provided in the questionnaire: (i) time needed to fall asleep (a categorical variable was created: ≤ 30 minutes, > 30 minutes); (ii) where the child sleeps (a categorical variable was created: in parents’ bed (co-sleeping), in a room with others, in a room alone); (iii) the use of a comfort object while sleeping such as a pacifier, a cuddly toy, sucking the thumb (no/yes); (iv) parental perception of child’s sleeping behavior. The latter, was recoded into a dichotomous variable (no/yes), in which “yes” comprised both “somewhat of a problem” and “quite a problem” responses.

Statistical analysis

Firstly, unintentional injuries from birth to 12 months and from 13 to 24 months were described as absolute and relative frequencies.

Time to occurrence of FERF was defined as the time from birth to the first raised surface fall event. In case of multiple falls, the first one was considered. Children without a documented fall from raised surface at the end of the study were censored at the date of the last available questionnaire. For example, for a child who didn’t have any fall event and had the 24 months’ questionnaire filled in (the end of study period or follow-up), the time to event considered was 24. The time of occurrence of FERF was analyzed using the Kaplan-Meier method.

The Log-rank test was used to compare time of occurrence of FERF and all predictive factors.

All variables described in the previous section (“Potential risk factors”) were included in the univariable analysis based on Log-rank test (S1 Table) and those significant (p -value < 0.05), at this first stage, were analyzed with the Chi-Square test to evaluate their independence. Risk factors significant in the univariate analysis based on Log-rank test and not correlated among each other were included in the multivariable analysis. Time to event analysis was performed using Cox proportional hazard models, without testing interaction between covariates. We ran Cox proportional hazard models considering mother-child pairs with complete information of outcome and potential risk factors ($N = 2,886$). We evaluated multicollinearity in the multivariable model using the variance inflation factor (VIF), considering the presence of collinearity when VIF was higher than 10. The proportional hazards assumption was assessed through scaled Schoenfeld residuals and no relevant departure was detected. Since the missing rate was lower than 10%, we handled missing as missing at random (the pattern of missingness is not related to other variables in the dataset). All analyses were conducted using STATA 12 software (StataCorp).

Results

In Table 1 unintentional injuries in children occurring in the first and second year of life are described by leading cause.

During the first year of life, 746 out of 2,896 children (25.8%) who filled in the questionnaire at 12 months had at least one unintentional injury, for a total number of 793 injuries.

Table 1. Leading cause of unintentional injuries during the first and second year of life.

	0–12 months (children injured = 746)		13–24 months (children injured = 994)	
	N	%	N	%
Fall from a raised surface	610	76.9	577	50.3
Burn/scald	39	4.9	81	7.1
Poisoning	3	0.4	13	1.1
Foreign body ingestion	24	3.0	25	2.2
Road accident	10	1.3	15	1.3
Other injury	107	13.5	435	38.0
Total	793	100	1146	100

<https://doi.org/10.1371/journal.pone.0275521.t001>

During the second year of life, 994 out of 2,751 children (36.1%) who filled in the questionnaire at 24 months had at least one unintentional injury, for a total number of 1,146 injuries. Falls from raised surface were the most frequent leading cause of unintentional injury, both in the first and in the second year of life (respectively 76.9% and 50.3%).

Table 2 summarizes details of falls from a raised surface in the cohort.

During the first year of life, most falls occurred after eight months (55.4%), under parental supervision (92%) and mainly from the bed (57%). During the second year of life, falls from a raised surface occurred more frequently after twenty months (40.6%) again under parents' supervision (84.7%) and mainly from the bed (34.8%). Interestingly, the percentage of children fallen in the nursery was very low (0.8% and 3.8% in the first and in the second year respectively). Moreover, the percentage of children taken to an Emergency Department or hospitalized as a result of the fall was similar in the first two years (around 22%).

Potential risk factors in the 2,886 mother-child pairs considered in the present study are reported in **Table 3**.

Overall, during the first two years of life, 996 out of 2886 children (35.9%) reported a FERF.

Fig 1 displays the Kaplan-Meier survival function for the probability of FERF overall. At 12 months of age 79% of children had never experienced a fall (95% CI 77.5%-80.4%), while at 24 months of age the percentage declined to 64% (95% CI 62%-65.7%).

The results of the adjusted Cox model, including the predictive variables significant in the previous univariable analysis based on Log-rank test (**S1 Table**), are shown in **Table 4**. No collinearity was observed between predictors included in our model (VIF = 1.46).

A lower risk of FERF was observed in children living in Northern Italy (HR 0.66, 95% CI 0.57–0.77), compared to central Italy. With regards to maternal age, a lower risk was found among children with mothers aged 30–34 years (HR 0.80, 95% CI 0.66–0.96) and >35 years (HR 0.71, 95% CI 0.60–0.85) compared to those with younger mothers (<30 years old). Moreover, an increased risk of FERF was observed in children with mothers with a moderate (HR 1.41, 95% CI 1.18–1.69) and high (HR 1.50, 95% CI 1.12–2.01) psychological distress. Maternal alcohol intake during pregnancy was also found to be associated with an increasing risk of FERF among children (HR 1.23, 95% CI 1.07–1.41). Furthermore, an increased risk of FERF was found among children having sleeping problems (HR 1.33, 95% CI 1.13–1.56).

Discussion

Our study identified maternal age, maternal distress, maternal alcohol consumption, child sleeping problems and enrollment area as predictive factors of FERF.

Children whose mothers were relatively older (> = 30 years old), had a lower risk of FERF. This finding is coherent with the evidence in the literature, in particular a study conducted in

Table 2. Description of every raised surface fall reported during the first and the second year of life.

	0–12 months (raised surface falls reported = 766)			13–24 months (raised surface falls reported = 706)	
	N	%		N	%
Age at fall			Age at fall		
1–4 months	56	7.4	13–16 months	150	21.6
5–8 months	282	37.2	17–20 months	262	37.8
9–12 months	420	55.4	21–24 months	282	40.6
Place where fall happened			Place where fall happened		
Bedroom	479	62.8	Bedroom	268	38.1
Living room	147	19.3	Living room	218	31.0
Nursery	6	0.8	Nursery	27	3.8
Playground/garden	31	4.1	Playground/garden	75	10.7
Other	100	13.1	Other	115	16.4
From where the child fell			From where the child fell		
Table/chair/high chair	89	11.6	Table/chair/high chair	194	27.5
Bed	437	57.0	Bed	246	34.8
Changing table	46	6.0	Changing table	16	2.3
Arms	15	2.0	Arms	20	2.8
Other (e.g. stroller, stairs, etc)	179	23.4	Other (e.g. stroller, stairs, etc)	230	32.6
Who was with the child			Who was with the child		
Parents	690	92.0	Parents	594	84.7
Granparents/relatives	42	5.6	Granparents	58	8.3
Siblings	8	1.1	Siblings	13	1.9
Baby sitter	10	1.3	Baby sitter	17	2.4
Other	0	0.0	Other	19	2.7
Injury type/site of injury*			Injury type/site of injury*		
Bruise	419	59.8	Bruise	319	40.9
Bleeding	63	9.0	Bleeding	73	9.4
Wound	18	2.6	Wound	61	7.8
Fracture	8	1.1	Fracture	14	1.8
Blow	48	6.8	Blow	54	6.9
Head injury	14	2.0	Head injury	15	1.9
Other	131	18.7	Nothing	219	28.1
			Other	25	3.2
Treatment			Treatment		
Medical examination	45	5.9	Medical examination	18	2.6
Emergency Hospital Department	152	20.1	Emergency Hospital Department	153	22.0
Hospitalization	16	2.1	Hospitalization	1	0.1
Nothing	540	71.3	Nothing	517	74.2
Other	4	0.5	Other	8	1.1

Total number may vary across variables due to missing values

* question with multiple choice

<https://doi.org/10.1371/journal.pone.0275521.t002>

the UK reported that younger maternal age is an important risk factor for unintentional injuries among preschoolers [23]. As Mitton and colleagues hypothesized, younger mothers may be less aware of the risks that children encounter and could be more prone to injuries growing up [36].

Table 3. Potential risk factors of the 2,886 mother-child pairs.

	N	%
Enrollment area		
Central Italy	1929	66.8
Northern Italy	957	33.2
Child gender		
Male	1471	51.0
Female	1415	49.0
Maternal age at delivery		
<30	556	19.3
30–34	992	34.4
> = 35	1,338	46.3
Maternal education		
Primary school	301	10.4
Secondary school	1,242	43.1
University degree or above	1,343	46.5
Maternal employment		
No	739	25.6
Yes	2,091	72.5
Not respondant	56	1.9
Paternal employment		
No	84	2.9
Yes	2,784	96.5
Non-responders	18	0.6
EHII*		
Low income	126	4.4
Medium income	1,607	55.7
High income	965	33.4
Non-responders	188	6.5
Number of siblings		
No	1,707	59.2
Yes	1,175	40.7
Non-responders	4	0.1
Day care attendance		
No	2,000	69.3
Yes	708	24.5
Non-responders	178	6.2
Maternal distress		
Feel good	2,146	74
Moderate distress	335	13,0
High distress	106	4.1
Non-responders	299	10.4
Maternal smoking during pregnancy		
No	2,261	78.3
Yes	623	21.6
Non-responders	2	0.1
Maternal alcohol intake during pregnancy		
No	1,161	40.2
Yes	1,667	57.8

(Continued)

Table 3. (Continued)

	N	%
Non-responders	58	2.0
Time needed to fall asleep		
< = 30 min	2,576	89.3
>30 min	221	7.7
Non-responders	89	3.0
Sleeping problems (reported by parents)		
No	2,341	81.1
Yes	480	16.6
Non-responders	65	2.3
Where the child sleeps		
Own bed	636	22
Bed in a room with others	1,554	53.9
Parents bed (Cosleeping)	584	20.2
Non-responders	112	3.9
The child uses a comfort object for sleeping		
No	1,005	34.8
Yes	1,820	63.1
Non-responders	61	2.1

Non-responders indicates an unanswered question

* low income = <933 euro, medium income = 934–1809, high income = > = 1810 euro

<https://doi.org/10.1371/journal.pone.0275521.t003>

Regarding maternal psychological distress, our study found that children with mothers with the highest GHQ-12 score are at higher risk of FERF. Factors affecting the association between maternal psychological health or depression and unintentional injury in children of pediatric age (0–3 years) have already been investigated in two longitudinal studies conducted in the UK and US [12, 25]. One possible explanation linking maternal depression and children's risk of injury is that chronically depressed mothers do not appropriately safeguard the physical environments children engage in [37, 38]. Furthermore, Phelan and colleagues reported that child supervision behavior on children aged 0–36 months differed between mothers suffering from depression and those who do not [39]. In contrast to these findings, a study conducted in UK in a deprived setting, found that maternal depressive symptoms, stress and a lack of social support do not influence the adoption of safety practices (such as smoke alarms, fireguards, safe storage of sharp objects/medicines, stair gates and window locks) [40].

Maternal alcohol consumption was associated with an increased risk of their child having a FERF occurring in the first two years of life in our study. This finding is coherent with previous studies where parental alcohol consumption has been found to increase the risk of unintentional injuries [18, 41].

Children with sleeping problems were found at higher risk of FERF compared to children with no sleeping problems. Sleeping difficulties in children might be correlated to sleep deprivation in parents. Considering that parental supervision plays a critical role in maintaining child safety, the consequence of parental sleep deprivation might be indicative of a less attentive or effective supervision of their children [42, 43].

The geographical differences in the risk of FERF observed in our study is somewhat difficult to explain with the data at hand and requires further analysis to evaluate if safety behaviors and risk perceptions differ by regional, social and cultural setting in Italy.




Fig 1. Kaplan Meier survival function for the probability of first fall from a raised surface overall during the first two years of life. FERF = First Event of Raised surface Fall.

<https://doi.org/10.1371/journal.pone.0275521.g001>

Table 4. Estimated hazard ratios (HRs) for FERF from the multivariable Cox model.

		FERF 0–24 months	
		HR	CI 95%
Enrollment site	Central Italy	1,00	
	Northern Italy	0.66	0.57–0.77
Maternal age	<30	1,00	
	30–34	0.80	0.66–0.96
	> = 35	0.71	0.60–0.85
Maternal distress	Feel good	1,00	
	Moderate distress	1.41	1.18–1.69
	High distress	1.50	1.12–2.01
Maternal alcohol intake	No	1,00	
	Yes	1.23	1.07–1.41
Child sleeping problems	No	1,00	
	Yes	1.33	1.13–1.56

<https://doi.org/10.1371/journal.pone.0275521.t004>

No association was found between FERF and factors such as maternal education, parental employment (a proxy of availability for supervision of the child at home), having siblings, EHII, maternal smoking during pregnancy (which could indicate a lower attention to the needs and health of the child), nursery attendance, thus we were not able to give strength to the role of “moderation” of welfare factors in injuries, as previously shown by other authors [17].

Although it is well known that boys experience injuries and falls more often than girls, this study did not show any difference by gender [9–11, 44]. Although the mechanisms through which this disparity may arise are not entirely clear, it has been suggested that boys are generally more active than girls thus are more likely to incur in injuries at an earlier age [44].

Furthermore, it is important to recall that our analysis included FERF occurring during the first year of life in which child-related factors may be less important compared to mother-related factors, potentially having a role in explaining the null association found between gender and FERF.

Several limitations of this study should be mentioned. First of all, since all the information about unintentional injuries were collected from self-reported questionnaires, parents can intentionally or unintentionally underreport injuries that children have experienced; a questionnaire administrated by a properly trained interviewer, might have reduced the risk of bias and ensure a higher quality of data, reducing definitely the amount of missing data. A social desirability bias may exist so that fall injuries can be underreported; moreover, a recall bias due to a relatively long time from injury to questionnaires may occur [45, 46]. Second, collected data included maternal and child factors, but not environmental factors (such as baby gates, window guards, restraining strap use, etc.) or other factors related to the kinetic energy on impact (such as fall height and cushioning capacity of the surface of impact).

Finally, since this study considered only the first event of fall from raised surface, future studies could be focused to assess multiple and repeated falls and evaluated if risk factors are confirmed or differ.

Conclusion

Despite the limitations mentioned above, the present study represents a crucial investigation regarding the predictive factors of FERF in children, simultaneously explored in early life, and adds evidence in this field of research, where the role of birth cohorts is limited.

Results of this study suggest that unintentional injuries in early life can be addressed by interventions and policies that target supervision of the child, especially during pregnancy and toddlers' early life, when parental role is critical to prevent childhood injury.

Further investigations will be essential to strengthen these findings, by means of which policy makers and health professionals could design prevention strategies to empower parents and significantly reduce unintentional injuries in early life.

Supporting information

S1 Table. Results of the Log-rank test.

(PDF)

Acknowledgments

Our thanks go to all the families who took part in this study and to the whole Piccolipiù team which includes the following research scientists and computer/laboratory technicians: Paola Lorusso, Maria Gabellieri, Valentina Ziroli, Valentina Colelli, Chelo Salatino, Silvia Narduzzi,

Sara Fioravanti, Giulia Poggesi, Veronica Montelatici, Antonella Ranieli, Maura Bin, Veronica Tognin, Assunta Rasulo, Laura Fiorini.

We also thank Annarita Vestri and Alessandra Spagnoli for the support provided on the statistical analyses, and Francesca de'Donato and Ursula Kirchmayer for the revision of English language. Lastly, special thanks to Manuela De Sario who helped us in revising the manuscript.

Author Contributions

Conceptualization: Martina Culasso.

Data curation: Daniela Porta, Sonia Brescianini, Luigi Gagliardi, Paola Michelozzi, Luca Ronfani, Franca Rusconi.

Formal analysis: Martina Culasso, Federica Asta.

Investigation: Martina Culasso.

Methodology: Martina Culasso, Federica Asta.

Project administration: Daniela Porta.

Software: Martina Culasso.

Supervision: Martina Culasso, Daniela Porta, Sonia Brescianini, Luigi Gagliardi, Paola Michelozzi, Costanza Pizzi, Luca Ronfani, Franca Rusconi, Liza Vecchi Brumatti, Federica Asta.

Writing – original draft: Martina Culasso.

Writing – review & editing: Martina Culasso, Daniela Porta, Sonia Brescianini, Luigi Gagliardi, Paola Michelozzi, Costanza Pizzi, Luca Ronfani, Franca Rusconi, Liza Vecchi Brumatti, Federica Asta.

References

1. Sethi D, Towner E, Vincenten J, Segui-Gomez M, Racioppi F. European report on child injury prevention 2008. World Health Organization. Regional Office for Europe. <https://apps.who.int/iris/handle/10665/326500> (accessed on 18 Jul 2022).
2. Rimsza ME, Schackner RA, Bowen KA, Marshall W. Can child deaths be prevented? The Arizona Child Fatality Review Program experience. *Pediatrics* 2002; 110 (1 Pt 1): e11. <https://doi.org/10.1542/peds.110.1.e11> PMID: 12093992
3. Kendrick D, Barlow J, Hampshire A, Stewart-Brown S, Polnay L. Parenting interventions and the prevention of unintentional injuries in childhood: systematic review and meta-analysis. *Child Care Health Dev* 2008; 34: 682–95. <https://doi.org/10.1111/j.1365-2214.2008.00849.x> PMID: 18796060
4. Fergusson DM, Grant H, Horwood LJ, Ridder EM. Randomized trial of the Early Start program of home visitation. *Pediatrics* 2005; 116: e803–9. <https://doi.org/10.1542/peds.2005-0948> PMID: 16322138
5. Kitzman H, Olds DL, Henderson CR Jr, Hanks C, Cole R, Tatelbaum R, et al. Effect of prenatal and infancy home visitation by nurses on pregnancy outcomes, childhood injuries, and repeated childbearing. A randomized controlled trial. *JAMA* 1997; 278: 644–52. PMID: 9272896
6. Fujiwara T, Okuyama M, Takahashi K. Paternal involvement in childcare and unintentional injury of young children: A population-based cohort study in Japan. *International Journal of Epidemiology* 2010; 39(2): 588–97. <https://doi.org/10.1093/ije/dyp340> PMID: 19923202
7. Cooper C, Dennison EM, Leufkens HG, Bishop N, Van Staa TP. Epidemiology of childhood fractures in Britain: a study using the general practice research database. *J Bone Miner Res* 2004; 19(12): 1976–81. <https://doi.org/10.1359/JBMR.040902> PMID: 15537440
8. Lalloo R, Sheiham A. Risk factors for childhood major and minor head and other injuries in a nationally representative sample. *Injury* 2003; 34(4): 261–6. [https://doi.org/10.1016/s0020-1383\(02\)00277-2](https://doi.org/10.1016/s0020-1383(02)00277-2) PMID: 12667777
9. Park SH, Cho BM, Oh SM. Head injuries from falls in preschool children. *Yonsei Med J* 2004; 45(2): 229–32. <https://doi.org/10.3349/ymj.2004.45.2.229> PMID: 15118993

10. Pittone ML, Attia MW. Patterns of injury associated with routine childhood falls. *Pediatr Emerg Care* 2006; 22(7): 470–474. <https://doi.org/10.1097/01.pec.0000226869.41803.50> PMID: 16871104
11. Love PF, Tepas JJ 3rd, Wludyka PS, Masnita-lusan C. Fall-related pediatric brain injuries: the role of race, age, and sex. *J Trauma* 2009; 67(Suppl 1): S12–5.
12. Schwebel DC, Brezausek CM. Chronic maternal depression and children's injury risk. *Journal of Pediatric Psychology* 2008; 33(10): 1108–16. <https://doi.org/10.1093/jpepsy/jsn046> PMID: 18474518
13. Ekéus C, Christensson K, Hjern A. Unintentional and violent injuries among pre-school children of teenage mothers in Sweden: a national cohort study. *J Epidemiol Community Health* 2004; 58(8): 680–5. <https://doi.org/10.1136/jech.2003.015255> PMID: 15252071
14. Reinberg O, Lutz N, Reinberg A, Mechkouri M. Trauma does not happen at random. Predictable rhythm pattern of injury occurrence in a cohort of 15,110 children. *J Pediatr Surg* 2005; 40(5): 819–825. <https://doi.org/10.1016/j.jpedsurg.2005.01.050> PMID: 15937822
15. D'Souza AL, Smith GA, McKenzie LB. Bunk bed-related injuries among children and adolescents treated in emergency departments in the United States, 1990–2005. *Pediatrics* 2008; 121(6): 1696–702. <https://doi.org/10.1542/peds.2007-2555> PMID: 18519473
16. Kendrick D, Marsh P. How useful are sociodemographic characteristics in identifying children at risk of unintentional injury? *Public Health* 2001; 115(2): 103–107. <https://doi.org/10.1038/sj.ph.1900737> PMID: 11406774
17. Shenassa ED, Stubbendick A, Brown MJ. Social disparities in housing and related pediatric injury: a multilevel study. *Am J Public Health*. 2004; 94(4): 633–9. <https://doi.org/10.2105/ajph.94.4.633> PMID: 15054017
18. Damashek A, Kuhn J. Toddlers' unintentional injuries: the role of maternal-reported paternal and maternal supervision. *J Pediatr Psychol* 2013; 38(3): 265–75. <https://doi.org/10.1093/jpepsy/jss113> PMID: 23143608
19. Chen G, Smith GA, Deng S, Chen D, Kelleher K, Xiang H. Psychological symptoms and nonfatal unintentional injuries among Chinese adolescents: a prospective study. *J Adolesc Health* 2005; 37:460–6. <https://doi.org/10.1016/j.jadohealth.2004.09.005> PMID: 16310123
20. McKinlay A, Dalrymple-Alford JC, Horwood LJ, Ferguson DM. Long-term outcomes after mild head injury in early childhood. *J Neurol Neurosurg Psychiatr* 2002; 73: 281–8.
21. Jones IE, Williams SM, Goulding A. Associations of birth weight and length, childhood size, and smoking with bone fractures during growth: evidence from a birth cohort study. *Am J Epidemiol* 2004; 159: 343–50. <https://doi.org/10.1093/aje/kwh052> PMID: 14769637
22. Shepherd J, Farrington D, Potts J. Impact of antisocial lifestyle on health. [Erratum in *J Public Health (Oxf)* 2005;27:312–13.] *J Public Health (Oxf)* 2004; 26: 347–52. <https://doi.org/10.1093/pubmed/fdh169> PMID: 15598851
23. Orton E, Kendrick D, West J, Tata LJ. Independent risk factors for injury in preschool children: Three population-based nested case-control studies using routine primary care data. *PLoS ONE* 2012; 7(4): e35193. <https://doi.org/10.1371/journal.pone.0035193> PMID: 22496906
24. Pearce A, Li L, Abbas J, Ferguson B, Graham H, Law C. Does childcare influence socioeconomic inequalities in unintentional injury? Findings from the UK Millennium Cohort Study. *J Epidemiol Community Health* 2010; 64(2): 161–166. <https://doi.org/10.1136/jech.2009.092643> PMID: 19934167
25. O'Connor TG, Davies L, Dunn J, Golding J. Distribution of accidents, injuries, and illnesses by family type. ALSPAC Study Team. Avon Longitudinal Study of Pregnancy and Childhood. *Pediatrics* 2000; 106(5): e68. <https://doi.org/10.1542/peds.106.5.e68> PMID: 11061805
26. Farchi S, Forastiere F, Vecchi Brumatti L, Alviti S, Arnofi A, Bernardini T, et al. Piccolipiù, a multicenter birth cohort in Italy: protocol of the study. *BMC Pediatr* 2014; 7: 14–36. <https://doi.org/10.1186/1471-2431-14-36> PMID: 24506846
27. Toccaceli V, Serino L, Stazi MA. Informed consent, and an ethico-legal framework for paediatric observational research and biobanking: the experience of an Italian birth cohort study. *Cell and Tissue Bank* 2014; 15(4): 579–90.
28. Pizzi C, Richiardi R, Charles MA, Heude B, Lanoe JL, Lioret S, et al. Measuring Child Socio-Economic Position in Birth Cohort Research: The Development of a Novel Standardized Household Income Indicator. *Int J Environ Res Public Health* 2020; 17(5): 1700. <https://doi.org/10.3390/ijerph17051700> PMID: 32150940
29. European Union Statistics on Income and Living Conditions (EUSILC). Available from: <https://ec.europa.eu/eurostat/web/microdata/european-union-statistics-on-income-and-living-conditions> (accessed on 18 July 2019).

30. Arora VS, Karanikolos M, Clair M, Reeves A, Stuckler D, McKee M. Data Resource Profile: The European Union Statistics on Income and Living Conditions (EU-SILC). *Int. J. Epidemiol.* 2015; 44(2): 451–461. <https://doi.org/10.1093/ije/dyv069> PMID: 25948659
31. Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O, et al. The validity of the GHQ in the WHO study of mental illness in general health care. *Psychological Medicine* 1997; 27(1): 191–7.
32. Piccinelli M, Bisoffi G, Bon MG, Cunico L, Tansella M. Validity and test-retest reliability of the Italian version of the 12-item general health questionnaire in general practice: A comparison between three scoring methods. *Comprehensive Psychiatry* 1993; 34(3): 198–205. [https://doi.org/10.1016/0010-440x\(93\)90048-9](https://doi.org/10.1016/0010-440x(93)90048-9) PMID: 8339539
33. Hoeymans N, Garssen AA, Westert GP, Verhaak PF. Measuring mental health of the Dutch population: A comparison of the GHQ-12 and the MHI-5. *Health and Quality of Life Outcomes* 2004; 2: 23. <https://doi.org/10.1186/1477-7525-2-23> PMID: 15132745
34. Sampogna F, Picardi A, Chren MM, Melchi CF, Pasquini P, Masini C, et al. Association between poorer quality of life and psychiatric morbidity in patients with different dermatological conditions. *Psychosomatic Medicine* 2004; 66 (4): 620–4. <https://doi.org/10.1097/01.psy.0000132869.96872.b2> PMID: 15272112
35. Uner S, Ozcebe H, Telatar TG, Tezcan S. Assessment of mental health of university students with GHQ-12. *Turkish Journal of Medicine Sciences* 2008; 38(5): 437–46.
36. Mytton J, Towner E, Brussoni M, Gray S. Unintentional injuries in school-aged children and adolescents: lessons from a systematic review of cohort studies. *Inj Prev.* 2009; 15(2): 111–. <https://doi.org/10.1136/ip.2008.019471> PMID: 19346424
37. Leiferman J. The effect of maternal depressive symptomatology on maternal behaviors associated with child health. *Health Educ Behav.* 2002; 29(5): 596–607. <https://doi.org/10.1177/109019802237027> PMID: 12238703
38. McLennan JD, Kotelchuck M. Parental prevention practices for young children in the context of maternal depression. *Pediatrics* 2000; 105(5): 1090–5. <https://doi.org/10.1542/peds.105.5.1090> PMID: 10790467
39. Phelan KJ, Morrongiello BA, Khoury JC, Xu Y, Liddy S, Lanphear B. Maternal supervision of children during their first 3 years of life: The influence of maternal depression and child gender. *Journal of Pediatric Psychology*. 2014; 39(3): 349–57. <https://doi.org/10.1093/jpepsy/jst090> PMID: 24357732
40. Mulvaney C, Kendrick D. Do maternal depressive symptoms, stress and a lack of social support influence whether mothers living in deprived circumstances adopt safety practices for the prevention of childhood injury? *Child Care Health Dev.* 2006; 32: 311–9. <https://doi.org/10.1111/j.1365-2214.2006.00590.x> PMID: 16634976
41. Bijur PE, Kurzon M, Overpeck MD, Scheidt PC. Parental alcohol use, problem drinking, and children's injuries. *JAMA* 1992; 267(23): 3166–71. PMID: 1593737
42. Boergers J, Hart C, Owens JA, Streisand R, Spirito A. Child sleep disorders: Associations with parental sleep duration and daytime sleepiness. *J Fam Psychol.* 2007; 21(1):88–94. <https://doi.org/10.1037/0893-3200.21.1.88> PMID: 17371113
43. Morrongiello BA. Caregiver supervision and child-injury risk: I. Issues in defining and measuring supervision; II. Findings and directions for future research. *J Pediatric Psychology*. 2005; 30(7): 536–52. <https://doi.org/10.1093/jpepsy/jsi041> PMID: 16166243
44. Matheny AP. Children's unintentional injuries and gender: Differentiation by environmental and psychosocial aspects. *Children's Environments Quarterly* 1991; 8(3–4):51–61.
45. Cummings P, Rivara FP, Thompson RS, Reid RJ. Ability of parents to recall the injuries of their young children. *Inj Prev.* 2005; 11(1): 43–7. <https://doi.org/10.1136/ip.2004.006833> PMID: 15691989
46. Pless CE, Pless IB. How well they remember. The accuracy of parent reports. *Arch Pediatr Adolesc Med.* 1995; 149(5): 553–8. <https://doi.org/10.1001/archpedi.1995.02170180083016> PMID: 7735412