

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Selection of food cultures with protective properties for cooked ham

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1889894> since 2024-12-13T11:10:28Z

Published version:

DOI:10.1016/j.fm.2023.104218

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Selection of food cultures with protective properties for cooked ham**

2

3 Valentina Alessandria*^a, Ilario Ferrocino^a, Valerio Carta^a, Veronique Zuliani^b, Tim Martin Seibert^c, Jakob
4 Soeltoft-Jensen^d, Kalliopi Rantsiou^a, Luca Cocolin^a

5

6 ^aDipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Largo Paolo
7 Braccini n°2, 10095 Grugliasco (TO), Italy

8 ^bChr. Hansen SAS, Saint-Germain-lès-Arpajon, France

9 ^cChr. Hansen GmbH, Pohlheim, Germany

10 ^dChr. Hansen A/S, Hoersholm, Denmark

11

12

13

14

15

16 * Corresponding author: Valentina Alessandria, Phone: +39-011-6708873, Fax: +39-011-6708549

17 Email: valentina.alessandria@unito.it

18

19

20

21

22

23

24

25

26

27 **Abstract**

28 Sliced cooked ham stored in modified atmosphere packaging (MAP) can be spoiled by lactic acid bacteria
29 (LAB) which are dominating under psychrotrophic conditions. Depending on the strains, the colonization can
30 result in a premature spoilage characterized by off-flavors, gas and slime production, discoloration, and
31 acidification. The purpose of this study was the isolation, identification and characterization of potential food
32 culture with protective properties, able to prevent or delay spoilage in cooked-ham. The first step was to
33 identify by means of microbiological analysis, the microbial consortia both in unspoiled and in spoiled lots of
34 sliced cooked ham by the use of media for the detection lactic acid bacteria and total viable count. Counts
35 ranged from values lower than 1 Log CFU/g to 9 Log CFU/g in spoiled and unflawed samples.

36 The interaction between consortia was then studied in order to screen for strains able to inhibit spoilage
37 consortia. Strains showing antimicrobial activity were identified and characterized by molecular methods and
38 tested for their physiological features. Among a total of 140 strains isolated, nine were selected for their ability
39 to inhibit a large number of spoilage consortia, to grow and ferment at 4°C and to produce bacteriocins. The
40 effectiveness of the fermentation made by food culture was evaluated, through challenge tests *in situ*, analysing
41 the microbial profiles of artificially inoculated cooked-ham slices during storage by high throughput 16S rRNA
42 gene sequencing. The native population *in situ* resulted competitive against the inoculated strains and only one
43 strain was able to significantly reduce the native populations reaching about 46.7 % of the relative abundance.
44 The results obtained in this study provide information about the selection of autochthonous LAB on the base
45 of their action against spoilage consortia, in order to select protective potential cultures able to improve the
46 microbial quality of sliced cooked ham.

47

48

49

50 **Keyword:** Sliced cooked ham, Bioprotection, Lactic acid bacteria, Microbial Consortia, Bacteriocins.

51

52

53

54 1. Introduction

55 Cooked ham constitutes over 26% of the volume of the delicatessen products sold in Europe, where Spain,
56 France and Italy are the biggest consumers (Casiraghi et al., 2007). The final quality of the products depends
57 both on the raw material used and the processing, which normally includes injection of brine and tumbling and
58 cooking reaching at least 75°C core. These products are very susceptible to bacterial spoilage and several
59 preservation techniques are combined to optimize their microbial quality.

60 Thermal processing is a critical step in cooked ham production and plays an important role in the microbiota
61 selection. The process is not always effective against thermo-tolerant vegetative bacteria and has no effect on
62 the re-contamination that may occur during slicing and packaging (Franz & von Holy, 1996; Vasilopoulos et
63 al., 2010). The spoilage microorganisms that colonize cooked ham stored in modified atmosphere packaging
64 (MAP) are, amongst other, psychrotrophic lactic acid bacteria (LAB) such as *Leuconostoc*, *Latilactobacillus*
65 and *Carnobacterium* (Vasilopoulos et al., 2008). The use of gas mixture of N₂ and CO₂ in MAP products favors
66 microbiota that withstand microaerophilic conditions, usually LAB species and *Brochothrix thermosphacta*
67 (Casaburi et al., 2011; Geeraerts et al., 2017; la Stora et al., 2012; Samelis et al., 2000; Vasilopoulos et al.,
68 2015). LAB growth is also encouraged by presence of sodium chloride and sodium nitrite and a reduced water
69 activity. The overwhelming colonization may result in premature spoilage and subsequent food waste,
70 characterized by accelerated pH decrease, gas and slime production, discoloration, and/or off-flavor formation.
71 As mentioned above, the contamination can come mainly from post processing contamination. Therefore, apart
72 from cooking and good sanitation practices, the use of various post-cooking or post-slicing technologies such
73 as high pressure, antimicrobial treatments and use of food cultures can be useful for the microbial and sensory
74 quality preservation of cooked meat (Comi et al., 2016; Vermeiren et al., 2004). Biopreservation through
75 controlled fermentation can thus contribute to reduce food waste and extend the shelf life of the products. The
76 mechanism of biopreservation is driven by a multiple mode of actions such as the production of organic acids,
77 competitive exclusion (spatial volume and substrate availability), quorum sensing or by metabolites with
78 antimicrobial properties such as bacteriocins, natural peptides with the potential to inhibit the growth of
79 foodborne pathogens or spoilage bacteria in food products (Deegan et al., 2006). The production of
80 bacteriocins that leads to a better preservation of the product is a characteristic of different starter cultures

81 (Cleveland et al., 2001). Strains of different genera of LAB have been identified as bacteriocin producers and
82 their action against pathogen bacteria as *Listeria monocytogenes* in fermented meat has been deeply
83 investigated (Franciosa et al., 2018).

84 The purpose of this study was the isolation and identification of potential strains selected based on their action
85 against spoilage consortia. The potential food cultures with protective properties were tested for their
86 effectiveness against spoilage consortia, isolated from sliced cooked ham samples coming from different
87 processing plants in Europe.

88

89 **2. Material and methods**

90 *2.1 Microbiological analysis of sliced cooked ham samples*

91 A total of 67 sliced cooked ham samples, packed in modified atmosphere, were supplied by 10 producers
92 located in different areas in Europe and sent to the DISAFA laboratory. Companies were named with letters
93 from “A” to “L”. Among 67 samples, 30 were indicated as spoiled by the producers and their defects confirmed
94 at the laboratory. Criteria used from the companies to identify samples as spoiled were: package swelling,
95 presence of slime, off-odors and off-flavors and discoloration. Samples were divided in spoiled (30 samples)
96 and unspoiled (37 samples); for each sample several subsamples were supplied. The subsamples of the
97 unspoiled ones were stored at 7°C until the end of the shelf-life defined by the producers. Microbiological
98 analyses were conducted on spoiled samples the day of arrival, while the subsamples of the unspoiled were
99 analyzed also during their shelf life (at the beginning [B], middle [M] and end [E] of shelf life). The three
100 times of sampling were planned on the base of their shelf life (provided by producers), knowing the day of
101 production and the expiry day. Plate counts were adopted in order to isolate the predominant populations. Ten
102 grams of each sample were diluted in 90 mL of sterile Ringer solution (Oxoid, Milan, Italy) and homogenized
103 in a blender (Stomacher 400 Circulator, International PBI S.p.a, Italy) for 2 min at room temperature. For
104 enumeration, appropriate decimal dilutions were prepared and plated, in triplicate, on different media: lactic
105 acid bacteria (LAB) on de Man Rogosa Sharp agar in microaerophilic conditions (MRS, Oxoid), while total
106 viable count on BHI agar (Oxoid), incubated at 30°C and 37°C respectively for 48 h. In both cases samples
107 were spread on the plates.

108

109 *2.2 Selection of isolates with antimicrobial activity*

110 Colonies from spoiled hams were recovered from representative count plates of MRS and BHI by adding 1
111 mL of Ringer's solution, scraping and collecting the cell suspension. The recovered cell suspension represented
112 the microbial populations composing the consortia and they were used as indicators in assays performed to
113 detect antimicrobial capabilities. Among the 30 spoiled samples, 5 of them with the most evident defects
114 (mainly gas production/blowing) were selected and their corresponding spoilage consortia were used. To
115 screen a large quantity of LAB colonies for antimicrobial activity, the plates of the sampling of the unspoiled
116 hams, after counting, were overlaid with a layer of MRS or BHI agar containing the consortia recovered from
117 the spoiled samples (used as indicators). Specifically, 1% of the suspension was added to soft-agar media (agar
118 [10 g/l]) and plates of unspoiled samples were covered with a layer of soft agar containing the spoilage
119 consortium and incubated at 30°C for 24 h. After incubation colonies that displayed inhibition halo were
120 marked, collected from the original plates and cultivated in BHI or MRS as appropriate. In the case of unspoiled
121 samples, count plates (with a maximum of 300 colonies) of MRS and BHI were replicated by "replica plates"
122 (Dal Bello et al., 2010) in order to test the same population against all the spoilage consortia selected. All the
123 spoiling consortia were dispensed in Eppendorf tubes with glycerol and stored at -80°C.

124 A total of 140 colonies were selected for their antimicrobial activity and subjected to Gram staining and
125 catalase test. In order to confirm the antimicrobial activity, each of the 140 isolates was tested against the
126 spoilage consortia through agar-well-diffusion assay (AWDA), as described by Toba et al. (1991). Fifty µL of
127 an overnight broth culture of each isolate were placed into a well of 5 mm diameter in a soft agar plates (agar
128 [10 g/l], either MRS or BHI) containing the spoilage consortium. After incubation for 24 h at 30 and 37°C,
129 respectively for MRS and BHI, they were examined for the presence of inhibition zones. The isolates able to
130 inhibit more than the 60% of spoiled consortia tested, were selected. In order to determine the nature of the
131 inhibition, isolates were growth overnight in MRS and BHI broth and after removal of the cells, through a 0.2
132 mm filter (Millipore, Italy), and the pH neutralization with KOH (1 M), proteinase K (25 mg/mL) was added
133 to the supernatant and a new AWDA was performed. The absence of inhibition halos after the proteinase K
134 treatment confirmed the proteinaceous origin of the antimicrobial compound.

135 With the aim of finding potential starter for cooked ham, the isolates exerting antimicrobial activity were also
136 evaluated for their ability to grow at refrigeration temperature. A 1% (v/v) inoculum was added to BHI and
137 MRS broth and incubated at 4°C for 72 h. Tubes that displayed value of 5 compared to the McFarland scale
138 were considered positive.

139

140 *2.3 DNA extraction from bacteriocin producing isolates and molecular identification and* 141 *characterization*

142 Genomic DNA of the bacteriocin producing isolates was extracted as described in Cocolin et al. (2001).
143 Isolates were first grouped by their fingerprinting profile by the means of PCR 16S–23S rRNA gene spacer
144 analysis (RSA). The RSA was carried out with primers G1 (GAAGTCGTAACAAGG) and L1
145 (CAAGGCATCCACCGT) (Dolci et al., 2008). Representative isolates from each group were then identified
146 by amplified the variable V1 and V3 regions of 16S rRNA gene (Cocolin et al., 2001). PCR products were
147 sequenced and the resultant sequences were aligned with those in Gene Bank using the Blastn program
148 (Altschul et al., 1997) to determine the known relatives.

149 Rep-PCR fingerprinting was performed for molecular characterization with the single oligonucleotide primer
150 (GTG)₅, following the protocol described by Gevers et al. (2001). Products of rep-PCR were run in a 2%
151 agarose gel and the fingerprints obtained were visualized under ultraviolet light using UVI pro platinum 1.1
152 Gel Software (Eppendorf, Germany) and analyzed with BioNumerics 4.6 software package (Applied Maths,
153 Belgium). The similarity among digitalized profiles was calculated using the Pearson correlation and an
154 average linkage (UPGMA) dendrogram was derived from the profiles.

155 The methods adopted in the next steps of the analyses were chosen on the basis of the results of the
156 identification of strains.

157

158 *2.4 LAB bacteriocin activity (AU)*

159 The AWDA method was adopted to titrate the bacteriocin produced. Isolates that possessed antimicrobial
160 activity were grown overnight in M17 added either with glucose or lactose in concentration of 0.5 % p/v. M17
161 medium was chosen because of its suitable composition for the growth of *Lactococcus* spp. In order to quantify

162 the bacteriocin activity, the cell free overnight supernatants were two-fold serially diluted in PBS (pH 7)
163 solution reaching a dilution of 1: 4096. Fifty μ L of each dilution was added to BHI soft agar plate containing
164 1% inoculum of an overnight culture of *L. monocytogenes* EGDe as an indicator. The bacteriocin activity was
165 expressed in arbitrary units (AU)/ml. One arbitrary unit was defined as the reciprocal of the highest dilution
166 yielding a clear inhibition zone of the indicator strain (Ananou et al., 2005). Four *Lactococcus lactis* strains
167 (strains 41FL18, 41FLL3, 39FL2 and 41FL1) from the DISAFA collection, isolated by artisanal products from
168 North West of Italy (Dal Bello et al., 2010) were included in the work because of their ability to produce
169 bacteriocins. The values of bacteriocin activity shown in this study were the averages of three assays.
170 Data were subjected to one-way ANOVA, and Duncan test was used to determine differences in the inhibition
171 activity expressed as AU/ml at $p < 0.05$, using the statistical software, STATISTICA 7.0 for Windows (Statsoft,
172 Tulsa, USA).

173

174 *2.5 Targeting the bacteriocin encoding gene by PCR amplification*

175 On the base of the results of the isolates identification, PCR was used to amplify the following target genes:
176 nisin A (Choi et al., 2000), lactococcin A (M. I. Martínez et al., 1998), lactococcin B (van Belkum et al., 1992),
177 lactococcin 513 (Villani et al., 2001), lactococcin 972 (B. Martínez et al., 1999), lacticin RM (Villani et al.,
178 2001) and lacticin 481 (Piard et al., 1993), as performed in Dal Bello et al. (2010). The reaction was performed
179 in a final volume of 25 μ L containing 1X PCR buffer, 1.5 mmol/L $MgCl_2$, 0.2 μ mol/L dNTP, 0.2 μ mol/L of
180 each bacteriocin primer and 1 U of Taq polymerase (Sigma Aldrich). The amplification cycles used were 95°C
181 for 5 min as initial step, 95° C for 1 min, annealing at 42°C (for the primers of Nis A and Nis Z), 56°C (for
182 the primers of lactococcins and lacticins) for 1 min, and 72 °C for 1 min for the next 30 cycles, 72 °C for 5
183 min concluded the amplification as reported in (Dal Bello et al., 2010). Fragments were visualized on 2%
184 agarose gels added of ethidium bromide using the 100 bp ladder (Promega, Italy) as the molecular weight
185 marker.

186

187 *2.6 In situ challenge tests*

188 Challenge tests were carried out on cooked ham slices packaged in a white room of a Piedmont ham plant the
189 same day of analysis, inoculating the most performing bacteriocin producing isolates. Strains isolated from
190 spoiled samples were also included in the trials to simulate the interaction in the occurrence of spoilage
191 (Raimondi et al., 2019). Four spoilage consortia (SC) were prepared: consortium 1 was composed by strains
192 34, 36, 37 and 38, respectively identified as *Leuconostoc carnosum*, *Kazachstania servazzi*, *Candida sake* and
193 *Yarrowia lipolytica*, consortium 2 by strain 46 (*Enterococcus gilvus*), consortium 3 by *Leuconostoc carnosum*
194 1 and *Latilactobacillus curvatus* 11 and consortium 4 by strains 18,19 and 20 belonging to *Latilactobacillus*
195 *sakei* species.

196 The potential food cultures with protective properties (BC) selected and SC were inoculated to a final
197 concentration of 6 Log CFU/g and 4 Log CFU/g, respectively, alone and in interaction. Negative controls with
198 only Ringer solution were carried out for each trial. Two batches were carried out for each trial and biological
199 replicates were carried out for each trial.

200 Briefly, the strains were cultured for 24h and cell counts were performed by the preparation of appropriate
201 decimal dilutions and plated on the appropriate media (MRS and M17 for LAB, Malt Agar for yeasts (Oxoid))
202 and incubated at 30°C for 48 h. Cells were washed twice with Ringer solution and 10 mL of cells suspension
203 were filled in spray bottles in order to reach a final count on ham (sliced the day of the inoculum in a white
204 room) of 4 and 6 Log CFU/g for SC and BC respectively. Four slices were collected sterile and inoculated in
205 both sides using spray bottles. The inoculated hams were packed on modified atmosphere (70% N₂ and 30%
206 CO₂) and stored at 7°C for 10 weeks (abuse time).

207 About 10 g from each sample before the inoculum and after 10 weeks of storage were homogenized with
208 Ringer solution (Oxoid, Milan, Italy) and analyzed as previously described using M17 (Oxoid, Milan) and
209 MRS media in triplicate. Samples inoculated with consortium 1 were also analyzed using WLN for the
210 detection of yeast.

211 The pH was measured by immersing the pH probe of a digital pH meter (micropH2001; Crison, Barcelona,
212 Spain) in the first decimal dilution before inoculation and after the 10 weeks.

213

214 *2.8 DNA extraction from ham samples*

215 Before the inoculum and after 10 weeks of storage, 1 ml of the first 10-fold serial dilution was collected and
216 directly centrifuged at maximum speed for 5 minutes. Nucleic acid was extracted by pooling two biological
217 replicates from each sampling point from each trial by using the MasterPure™ Complete DNA and RNA
218 Purification Kit (Epicentre, Madison, WI, USA) following the manufacturer's instructions. The nucleic acids
219 were resuspended in 50 µl of sterile water. DNA was quantified by using the Nanodrop 1000 (Thermo
220 Scientific, Milano, Italy) and standardized at 100 ng/µl.

221

222 *2.9 16S rRNA amplicon target sequencing*

223 DNA sequencing was used to assess the microbiota by amplification of the V3-V4 region of the 16S rRNA
224 gene using the primers and protocols described by Klindworth et al. (2013). The PCR products were purified
225 and tagged by using the Nextera XT index kit (Illumina), according to the manufacturer's instructions.
226 Sequencing was performed with a MiSeq instrument (Illumina) with V3 chemistry and generated 250-bp
227 paired-end reads, according to the manufacturer's instructions.

228

229 *2.10 Bioinformatics and statistical analysis*

230 After sequencing raw reads were first joined through flash and analyzed with QIIME 1.9.0 software (Caporaso
231 et al., 2010) and the pipeline described by Ferrocino et al. (2017). Briefly after quality filtering OTUs were
232 clustered at 99% of similarity and centroids sequence were mapped against the green genes by means of the
233 RDP Classifier. OTU table was rarefied at the lowest number of sequence and displays the highest taxonomic
234 level. Table was used to build a principal-component analysis (PCA) as a function of the production by using
235 the made4 package of R.

236

237 **3. Results**

238 *3.1 Microbial counts*

239 Among 67 samples, 30, coming from 8 different plants, were defined spoiled on the basis of their main defects:
240 they showed off-flavors formation, gas and slime production, discoloration, and acidification. From each plant

241 different number of samples were supplied. Table 1 reports the microbial counts of the spoiled samples. Both
242 MRS and BHI counts are shown.

243 In spoiled samples, counts ranged from values lower than 1 Log CFU/g to 9 Log CFU/g (the lowest from C
244 and L plants, the highest from I) in both MRS and BHI media. Important differences between the two media
245 were observed for plant F, where LAB counts were higher than population grown on BHI while in the case of
246 plant C, counts resulted higher in BHI for two of the three samples supplied. In the case of plant L, no colonies
247 were observed in the BHI plates (< 1 Log CFU/g).

248 For the samples that were classified as unspoiled when arrived at the laboratory (37 out of 67), the microbial
249 dynamics during shelf life were monitored on MRS and BHI plates. All the samples at the beginning of the
250 shelf life showed a load under 4 Log CFU/g, except samples of plant B where counts reached values of 6 Log
251 CFU/g while samples from plant C and E showed counts under the detection limit (< 1 Log CFU/g). In Figure
252 1, LAB counts on MRS medium at the beginning (B), middle (M) and end (E) of shelf life are reported.

253 In samples from company B, the counts were the highest from the beginning to the end if compared to the
254 other samples. Regarding the other companies, the microbial counts were similar at the middle and at the end
255 of shelf life except for samples produced by company C, where more variability was observed.

256

257 3.2 Detection of antimicrobial activity and molecular identification

258 Five spoilage consortia coming from samples with the most evident defects were selected for the inhibition
259 assays. Among 37 consortia from unspoiled samples, 20 showed inhibition against all consortia. A total of 140
260 colonies, with potential antimicrobial activity, were observed. The isolates were then tested against all the
261 spoilage consortia. After this second selection, 17 isolates were chosen according to their ability to inhibit
262 more than 60% of the spoilage consortia. All 17 isolates originated only from samples of the same plant (plant
263 B). The isolates were then identified by RSA and 16S rRNA gene sequencing. Among 17, 16 resulted to be
264 *Lactococcus lactis* subsp. *lactis*, while one was identified as *Enterococcus faecium*. Due to their possible
265 exploitation to preserve sliced cooked ham, the ability to grow at 4°C was investigated for the *L. lactis* strains
266 while the strain identified as *E. faecium* was not used or tested further.

267 In Table 2 the inhibition activity (expressed as percentage of inhibited consortia tested in MRS and BHI media)
268 of the 16 *L. lactis* strains are reported along with their ability to grow at refrigeration temperatures. Only three
269 strains did not show growth expressed as turbidity in BHI broth. All the strains expressed higher inhibition
270 towards spoilage consortia in MRS compared to those in BHI medium.

271 In order to identify the nature of the antimicrobial compound, the AWDA was carried out on these 16 strains
272 after neutralization and proteinase K assay. The results highlighted that among 16 isolates, 7 produced halos
273 for acidity while 9 were putative bacteriocin producers. These nine strains were selected and subjected to Rep-
274 PCR followed by cluster analysis where a high level of similarity (90%) was observed (data not shown).
275 The 9 bacteriocinogenic *L. lactis* strains (named A16, A14, A30, A34, A35, A36, A51 and A52) were subjected
276 to the identification of the bacteriocin genes by PCR. In addition, the four *L. lactis* strains (41FL18, 41FLL3,
277 39FL2 and 41FL1) belonging to the culture collection of DISAFA were included from this point forward of
278 the study. The presence of the genes responsible for the nisin A was revealed in all the isolates. Strain 39FL2
279 resulted positive also for the production of lactococin Ga, G β , Qa and Q β while A13, A16 and 41FL1 a part
280 from the presence of *nisA* possessed the gene encoding for lactococin Ga, G β . The bacteriocin activity,
281 expressed as arbitrary units (AU)/ml was also determined (Figure 2). The activity was calculated on M17
282 medium, with the addition of glucose and lactose.

283 The results highlighted that the different sugar added to the medium was determinant for the bacteriocin
284 production: all the tested strains showed a higher bacteriocin titer with the presence of glucose (significant
285 differences were observed for strains 39FL2 and A16) and the highest bacteriocin activity was detected for
286 strain A16 with a value of 2346 AU/ml in presence of glucose and for strain 39FL2 with a titer of 7253 AU/ml.
287 The difference in the bacteriocin production was not related to different growth rates in glucose and lactose
288 (data not shown).

289

290 *3.3 In situ challenge tests*

291 Challenge tests were carried out using the four DISAFA strains as well as strain A16 that was the most
292 performing strain, considering the bacteriocin titer, among the strains isolated from unspoilied sliced cooked
293 ham samples.

294 Non inoculated sliced cooked hams at time zero showed an average of pH of about 6.3 that decreased after 10
295 weeks reaching average value of 5.3. Same trend was observed for all the inoculated samples, with final pH
296 values ranging from 5.06 to 5.3.

297 Microbial counts at time zero showed a maximum of 3 Log CFU/g on both MRS and M17 media, reaching
298 values of 8 Log CFU/g after 10 weeks in all the samples. Regarding yeast population analyzed for samples
299 inoculated with consortium 1, all the samples presented values of about 5 Log CFU/g. No significant
300 differences were observed between the counts of the different trials, including inoculated and control samples
301 (data not shown).

302 Concerning the visual observation, no intense defects were observed after 10 weeks of storage in any of the
303 trials, but discoloration and slime production were randomly observed also when food cultures were
304 inoculated. Only samples inoculated with strain 39FL2 did not show any defects also when in interaction with
305 spoilage consortia (except in the case of consortium 4 where discolorations were observed). In order to follow
306 the development of the microbial populations in the challenge tests performed, amplicon (16S rRNA encoding
307 gene) based sequencing was performed. A total of 90 samples were analyzed and results are shown in Figure
308 3. Not inoculated samples showed a simple microbiota composition across the entire datasets characterized by
309 the presence of *L. sakei*, *Leuc. carnosum* and *Weissella viridescens* that accounted for 68,39%, 26,02% and
310 5,17% of the relative abundance respectively (Figure 3). We observed that *L. sakei* was the dominant OTU
311 also in hams inoculated with the four spoilage consortia together with *Leuc. carnosum*. In samples of
312 consortium 2 *Leu. carnosum* represented above 46,5% of the total OTUs. In addition, *Enterococcus*
313 *pseudoavium* was found in samples inoculated with consortia 2 and 3.

314 Comparing the relative abundance of the samples inoculated with the food cultures, it was evident how strain
315 39FL2 was the most competitive against the autochthonous microbiota (Figure 3c) reaching about 46.7 % of
316 the relative abundance. Strains 41FL1, 41FL18 and 41FLL3, despite the inoculum of 6 Log CFU/g, reached
317 respectively 14. 8%, 8% and 14% of the relative abundance after the 10 weeks of storage. Interestingly, by the
318 application of these three food cultures, a reduction of *L. sakei* from 68.4% to about 57% was observed while
319 the % of *Leuc. carnosum* remained constant at about 26% for all the samples. In general, the five starter strains

320 were more competitive against consortium 1 composed by 4 different species (*Leuc. carnosum*, *K. servazzi*, *C.*
321 *sake* and *Y. lipolytica*).

322 The trials with the A16 strain (Figure 4) were carried out in a different day and for this reason it had another
323 control compared to the other strains. This control was also composed mainly by *L. sakei*, *Leuc. carnosum* and
324 *Weissella viridescens*. *L. lactis* was detected, with an average abundance of 15 and 23% in trials with consortia
325 1 and 2 respectively, but it was below the detection limit when co-inoculated with consortia 3 and 4. In the
326 sample with the inoculum of strain A16 alone, a decrease of *L. sakei* and *L. carnosum*, compared to the control,
327 was registered (Figure 4). When co-inoculated with consortium 2, the A16 strain showed a good ability to
328 inhibit *Enterococcus* spp.

329 Strain 39FL2 resulted the most competitive against consortium 1, 2 and 3 covering respectively 31, 24 and
330 28% of the relative abundance. Only with consortium 4 (composed by three strains of *L. sakei*), *L. lactis* 39FL2
331 only reached 3.6% of the relative abundance. By Principal Component Analysis, a significant shift in the
332 microbiota was observed only when strain 39FL2 was used (Figure 5). In this case it is possible to observe a
333 certain degree of separation in the samples inoculated with only the 39FL2 strain compared to the control and
334 to the samples inoculated with the spoilage consortia.

335 In the case of strain 41FL1, the control and the samples inoculated with food culture showed a high degree of
336 similarity. In the other cases the production with food cultures were not well separated by the ones with
337 spoilage consortia (data not shown).

338

339 **4. Discussion**

340 Sliced cooked ham is a perishable meat product having low salt content, pH around 6 and water activity higher
341 than 0.945. These physicochemical parameters represent only small hurdles to inhibit microorganisms
342 associated to postprocessing contamination (Aymerich et al., 2005). Re-pasteurization after slicing and
343 packaging is not recommended because of the consequent release of meat juices. In this context the application,
344 after slicing, of alternative hurdles, such as natural antimicrobials, are useful tools in the preservation of this
345 product.

346 Part of the LAB species are generally recognized as safe bacteria and play an important role in preservation of
347 fermented foods. Their ability to produce bacteriocins may be useful also in the preservation of products such
348 as cooked ham (Laursen et al., 2009). The shelf life of processed meat products, if stored under anaerobic
349 conditions or MAP conditions, is generally dictated by LAB growth.

350 In this study, samples of cooked ham were analyzed and microbial counts at the end of shelf life seemed not
351 to be connected to the defects: samples coming from the same plant showed similar microbial counts
352 independently from the fact that they presented or not sensorial defect. Samples from company A and D
353 showed higher counts at the end of shelf life in acceptable samples, if compared to the spoiled ones (Table 1
354 and Figure 1). This underlines that the microbial count is not the only parameter that has to be taken into
355 account to describe or predict spoilage, and it becomes of great importance to properly understand the ecology
356 of the samples when spoilage occurs. Unexpectedly, spoiled samples of plant C (on MRS) and of plant L,
357 showed microbial counts under the detection limit (Table 1). However it has to be considered that spoilage can
358 be driven also by other microorganism that can not be detected by the media and conditions used in this
359 research.

360 As found by Raimondi et al. (2019), the microbiota analysis at the end of the shelf life of unspoiled and spoiled
361 samples of cooked hams, did not unveil any specific taxon that could be surely associated to spoilage,
362 indicating that a negative evolution can be due to strain dependent characteristics (Ercolini et al., 2010).
363 Different microbial loads were observed between companies, underlining how the production plant is
364 determinant in the microbial development (Table 1). Moreover, raw materials and processing steps are of
365 primary importance in the definition of the final microbiota. The hygiene conditions throughout the processing
366 line have great impact on the initial microbial levels, on the microbiota composition (Vihavainen et al., 2007)
367 and consequently influence the onset of spoilage. LAB can contaminate the products during the post-cooking
368 processing. The control of potential spoilage microorganisms through the use of other microorganisms is of
369 great interest. Several studies have demonstrated that autochthonous microbiota can influence the reduction of
370 pathogens such as *L. monocytogenes* in cooked ham (Hwang & Sheen, 2011), while others underlined how
371 autochthonous microbiota do not influence the growth of the pathogens in samples of modified-atmosphere-

372 packaged poultry (Barakat & Harris, 1999). A probable reason for this discrepancy may be attributed to
373 difference in the microbiota investigated in the cited studies.

374 In this study, selection of potential food cultures with protective properties and active towards spoilage
375 microbial consortia, was performed. An advanced experimental design was applied to identify strains with
376 potential to inhibit entire microbial consortia rather than focusing on an isolated, putative spoilage
377 microorganism. The study was focused on the main bacterial spoilage of cooked ham and in particular on LAB
378 that are dominating under psychrotrophic conditions: for this reason, MRS medium, commonly used for LAB
379 detection, was chosen together with BHI, selected for the detection of microbial viable counts. A limitation of
380 the method could be linked to the difficulties of specific microbial groups involved in spoilage of cooked ham
381 as *Carnobacterium* spp. to grow on MRS and BHI media: future studies could adapt the methods including also
382 other specific media. Among 37 consortia coming from unspoiled sliced cooked ham, from 6 plants, 20 showed
383 an inhibition against spoilage consortia. On the basis of the inhibition observed in lab conditions, 140 isolates
384 were selected. These isolates were tested against all the consortia coming from spoiled samples and 17 isolates
385 confirmed a good inhibition against more than 60% of the spoiled consortia tested (Table 2). The selected
386 inhibitor isolates were identified as *L. lactis* (16/17) and *E. faecium* (1/17). Surprisingly the strains selected for
387 their attitude to inhibit the largest number of spoilage consortia came from the ham samples produced in one
388 plant. Only *E. faecium* was not considered further because of the controversial role of this genus *Enterococcus*
389 in food microbiology. Despite the important role in contributing in the taste and flavor of various fermented
390 foods through proteolysis, lipolysis, members of this genus have been associated with human infections
391 (Foulquié Moreno et al., 2006). Furthermore, due to its ability to resist to a wide variety of antibiotic and
392 through the skills to exchange genetic information by conjugation, it may spread antibiotic resistance among
393 non-pathogenic organisms (Coconcelli et al., 2003). In this light, among the two species, our attention was
394 focused on *L. lactis* species. Comi et al. (2016) used this species to improve the shelf life of cooked bacon
395 where it showed its ability to reduce the risk of *Leuconostoc mesenteroides* spoilage. Comi et al. (2011) also
396 evaluated the oxygen scavenging ability of *L. lactis* ssp. *lactis*, to improve the quality of sliced cooked ham
397 packaged under MAP. Suspensions were spread on sliced cooked ham packed with MAP (oxygen < 0.5%),
398 stored at 4°C for 30 days under light and samples were analyzed during the shelf life to evaluate the quality.

399 *L. lactis ssp. lactis* was able to reduce the oxygen concentration, the lipid oxidation and inhibited the
400 autochthonous LAB, extending the shelf-life of the products to 30–35 days. It was shown that this species
401 influenced the flavor and odor and the stability of the red color of the sliced cooked ham. Despite the high
402 similarity observed by rep-PCR (data not shown), the selected strains showed a different attitude to produce
403 bacteriocins. From the technological point of view, thanks to their provenience from unspoiled samples and,
404 at the same time, to their ability to inhibit a great number of spoilage consortia, plus their attitude to grow at
405 refrigerated temperature, they can be considered in the elaboration of new potential food cultures after the
406 evaluation of their sensory impact on the final product.

407 It has to be considered that the bacteriocin production is strongly dependent on the medium composition
408 (Gänzle et al., 1999) and on the temperature. The results of the present study confirmed glucose as a better
409 carbon source for nisin producing microbes (Guerra et al., 2001). Furthermore comparing the inhibition of the
410 strains against spoiled consortia higher inhibition was detected in MRS than in BHI for all the strains: these
411 phenomena can be explained by the better ability of the LAB to grow in MRS compared to the BHI. The
412 composition of the substrate represented by the food matrix and the surrounding conditions have to be
413 considered also in the understanding of the *in situ* inhibition ability of the strains. As reported by Chollet et
414 al., (2008) also sodium chloride concentration can influence the nisin concentration and its antimicrobial
415 activity.

416 In the challenge tests performed in this study, strain A16 was selected because of its ability to produce the
417 higher titer of bacteriocin compared to the others selected strains. Taking in consideration the bacteriocin titer,
418 strains present in the DISAFA collection were also included in the challenge tests because of their ability to
419 produce bacteriocins. The autochthonous microbiota resulted competitive against the strains selected for their
420 antimicrobial properties inoculated at 6 Log CFU/g. It has to be considered that the analysis of the 16S
421 amplicons were performed to have a generic overview: the other population could have covered the presence
422 of the BCs because in a higher concentration. Strain 39FL2 was able to significantly reduce the autochthonous
423 populations, representing the 46.7% of the total OTUs, while in the other cases the inoculated strains covered
424 maximum the 15% of the OTUs (Figure 3). Therefore strain 39FL2 resulted to be the better performing strain

425 in the inhibition of indigenous cooked ham populations and this could be associated to its ability to produce
426 lactococcins together with nisins.

427

428 **5. Conclusion**

429 This study provides a detailed analysis of cooked ham coming from plants located in different European
430 countries with own processing procedures. The interaction between a high number of consortia coming from
431 unspoiled and spoiled samples allowed to select strains able to inhibit consortia of different spoiled hams. The
432 power of the study was the selection connected to the inhibition against consortia and not against single or
433 indicator strains. The ability of the strains to inhibit *in vitro* of a high number of spoilage consortia remarks
434 the effectiveness of this methods in the selection of bioprotective cultures. However further investigations need
435 to be conducted on real production where different factors can affect the action of bioprotective cultures,
436 especially for the strong competition of the autochthonous population as resulted by the challenge tests
437 conducted in this study.

438

439

440 **Acknowledgments**

441 This study was supported by Christian Hansen.

442

443

444

445

446 **Reference**

447 Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997).
448 Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. In *Nucleic*
449 *Acids Research* (Vol. 25, Issue 17). <https://doi.org/10.1093/nar/25.17.3389>

450 Ananou, S., Maqueda, M., Martínez-Bueno, M., Gálvez, A., & Valdivia, E. (2005). Control of
451 *Staphylococcus aureus* in sausages by enterocin AS-48. *Meat Science*, 71(3).
452 <https://doi.org/10.1016/j.meatsci.2005.04.039>

- 453 Aymerich, T., Jofré, A., Garriga, M., & Hugas, M. (2005). Inhibition of *Listeria monocytogenes* and
454 *Salmonella* by natural antimicrobials and high hydrostatic pressure in sliced cooked ham. *Journal of*
455 *Food Protection*, 68(1). <https://doi.org/10.4315/0362-028X-68.1.173>
- 456 Barakat, R. K., & Harris, L. J. (1999). Growth of *Listeria monocytogenes* and *Yersinia enterocolitica* on
457 cooked modified-atmosphere-packaged poultry in the presence and absence of a naturally occurring
458 microbiota. *Applied and Environmental Microbiology*, 65(1). [https://doi.org/10.1128/aem.65.1.342-](https://doi.org/10.1128/aem.65.1.342-345.1999)
459 345.1999
- 460 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Pêa,
461 A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R.
462 E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., ... Knight, R. (2010). QIIME allows
463 analysis of high-throughput community sequencing data. In *Nature Methods* (Vol. 7, Issue 5).
464 <https://doi.org/10.1038/nmeth.f.303>
- 465 Casaburi, A., Nasi, A., Ferrocino, I., di Monaco, R., Mauriello, G., Villani, F., & Ercolini, D. (2011).
466 Spoilage-related activity of *Carnobacterium maltaromaticum* strains in air-stored and vacuum-packed
467 meat. *Applied and Environmental Microbiology*, 77(20). <https://doi.org/10.1128/AEM.05304-11>
- 468 Casiraghi, E., Alamprese, C., & Pompei, C. (2007). Cooked ham classification on the basis of brine injection
469 level and pork breeding country. *LWT - Food Science and Technology*, 40(1).
470 <https://doi.org/10.1016/j.lwt.2005.07.007>
- 471 Choi, H. J., Cheigh, C. I., Kim, S. B., & Pyun, Y. R. (2000). Production of a nisin-like bacteriocin by
472 *Lactococcus lactis* subsp. *lactis* A164 isolated from Kimchi. *Journal of Applied Microbiology*, 88(4).
473 <https://doi.org/10.1046/j.1365-2672.2000.00976.x>
- 474 Chollet, E., Sebti, I., Martial-Gros, A., & Degraeve, P. (2008). Nisin preliminary study as a potential
475 preservative for sliced ripened cheese: NaCl, fat and enzymes influence on nisin concentration and its
476 antimicrobial activity. *Food Control*, 19(10), 982–989. <https://doi.org/10.1016/j.foodcont.2007.10.005>
- 477 Cleveland, J., Montville, T. J., Nes, I. F., & Chikindas, M. L. (2001). Bacteriocins: safe, natural
478 antimicrobials for food preservation. In *International Journal of Food Microbiology* (Vol. 71).
479 www.elsevier.com/locate/ijfoodmicro
- 480 Cocconcelli, P. S., Cattivelli, D., & Gazzola, S. (2003). Gene transfer of vancomycin and tetracycline
481 resistances among *Enterococcus faecalis* during cheese and sausage fermentations. *International*
482 *Journal of Food Microbiology*, 88(2–3). [https://doi.org/10.1016/S0168-1605\(03\)00194-6](https://doi.org/10.1016/S0168-1605(03)00194-6)
- 483 Cocolin, L., Manzano, M., Cantoni, C., & Comi, G. (2001). Denaturing Gradient Gel Electrophoresis
484 Analysis of the 16S rRNA Gene V1 Region to Monitor Dynamic Changes in the Bacterial Population
485 during Fermentation of Italian Sausages. *Applied and Environmental Microbiology*, 67(3–12).
486 <https://doi.org/10.1128/aem.67.11.5113-5121.2001>
- 487 Comi, G., Andyanto, D., Manzano, M., & Iacumin, L. (2016). *Lactococcus lactis* and *Lactobacillus sakei* as
488 bio-protective culture to eliminate *Leuconostoc mesenteroides* spoilage and improve the shelf life and
489 sensorial characteristics of commercial cooked bacon. *Food Microbiology*, 58.
490 <https://doi.org/10.1016/j.fm.2016.03.001>

- 491 Dal Bello, B., Rantsiou, K., Bellio, A., Zeppa, G., Ambrosoli, R., Civera, T., & Cocolin, L. (2010).
492 Microbial ecology of artisanal products from North West of Italy and antimicrobial activity of the
493 autochthonous populations. *LWT - Food Science and Technology*, 43(7).
494 <https://doi.org/10.1016/j.lwt.2010.03.008>
- 495 Deegan, L. H., Cotter, P. D., Hill, C., & Ross, P. (2006). Bacteriocins: Biological tools for bio-preservation
496 and shelf-life extension. In *International Dairy Journal* (Vol. 16, Issue 9).
497 <https://doi.org/10.1016/j.idairyj.2005.10.026>
- 498 Dolci, P., Alessandria, V., Rantsiou, K., Rolle, L., Zeppa, G., & Cocolin, L. (2008). Microbial dynamics of
499 Castelmagno PDO, a traditional Italian cheese, with a focus on lactic acid bacteria ecology.
500 *International Journal of Food Microbiology*, 122(3), 302–311.
- 501 Ercolini, D., Casaburi, A., Nasi, A., Ferrocino, I., di Monaco, R., Ferranti, P., Mauriello, G., & Villani, F.
502 (2010). Different molecular types of *Pseudomonas fragi* have the same overall behaviour as meat
503 spoilers. *International Journal of Food Microbiology*, 142(1–2).
504 <https://doi.org/10.1016/j.ijfoodmicro.2010.06.012>
- 505 Ferrocino, I., Bellio, A., Romano, A., Macori, G., Rantsiou, K., Decastelli, L., & Cocolin, L. (2017). RNA-
506 based amplicon sequencing reveals microbiota development during ripening of artisanal versus
507 industrial Lard d'Arnad. *Applied and Environmental Microbiology*, 83(16).
508 <https://doi.org/10.1128/AEM.00983-17>
- 509 Foulquié Moreno, M. R., Sarantinopoulos, P., Tsakalidou, E., & de Vuyst, L. (2006). The role and
510 application of enterococci in food and health. In *International Journal of Food Microbiology* (Vol. 106,
511 Issue 1). <https://doi.org/10.1016/j.ijfoodmicro.2005.06.026>
- 512 Franciosa, I., Alessandria, V., Dolci, P., Rantsiou, K., & Cocolin, L. (2018). Sausage fermentation and starter
513 cultures in the era of molecular biology methods. *International Journal of Food Microbiology*, 279.
514 <https://doi.org/10.1016/j.ijfoodmicro.2018.04.038>
- 515 Franz, C. M. A. P., & von Holy, A. (1996). Bacterial populations associated with pasteurized vacuum-
516 packed Vienna sausages. In *Food Microbiology* (Vol. 13).
- 517 Gänzle, M. G., Weber, S., & Hammes, W. P. (1999). Effect of ecological factors on the inhibitory spectrum
518 and activity of bacteriocins. *International Journal of Food Microbiology*, 46(3).
519 [https://doi.org/10.1016/S0168-1605\(98\)00205-0](https://doi.org/10.1016/S0168-1605(98)00205-0)
- 520 Geeraerts, W., Pothakos, V., de Vuyst, L., & Leroy, F. (2017). Diversity of the dominant bacterial species on
521 sliced cooked pork products at expiration date in the Belgian retail. *Food Microbiology*, 65.
522 <https://doi.org/10.1016/j.fm.2017.03.007>
- 523 Gevers, D., Huys, G., & Swings, J. (2001). Applicability of rep-PCR fingerprinting for identification of
524 *Lactobacillus* species. *FEMS Microbiology Letters*, 205(1). [https://doi.org/10.1016/S0378-
525 1097\(01\)00439-6](https://doi.org/10.1016/S0378-1097(01)00439-6)
- 526 Guerra, N. P., Rua, M. L., & Pastrana, L. (2001). Nutritional factors affecting the production of two
527 bacteriocins from lactic acid bacteria on whey. *International Journal of Food Microbiology*, 70(3).
528 [https://doi.org/10.1016/S0168-1605\(01\)00551-7](https://doi.org/10.1016/S0168-1605(01)00551-7)

- 529 Hwang, C. A., & Sheen, S. (2011). Growth characteristics of *Listeria monocytogenes* as affected by a native
530 microflora in cooked ham under refrigerated and temperature abuse conditions. *Food Microbiology*,
531 28(3). <https://doi.org/10.1016/j.fm.2010.03.004>
- 532 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O. (2013).
533 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation
534 sequencing-based diversity studies. *Nucleic Acids Research*, 41(1). <https://doi.org/10.1093/nar/gks808>
- 535 la Storia, A., Ferrocino, I., Torrieri, E., di Monaco, R., Mauriello, G., Villani, F., & Ercolini, D. (2012). A
536 combination of modified atmosphere and antimicrobial packaging to extend the shelf-life of beefsteaks
537 stored at chill temperature. *International Journal of Food Microbiology*, 158(3).
538 <https://doi.org/10.1016/j.ijfoodmicro.2012.07.011>
- 539 Laursen, B. G., Byrne, D. v., Kirkegaard, J. B., & Leisner, J. J. (2009). Lactic acid bacteria associated with a
540 heat-processed pork product and sources of variation affecting chemical indices of spoilage and sensory
541 characteristics. *Journal of Applied Microbiology*, 106(2). <https://doi.org/10.1111/j.1365-2672.2008.04045.x>
- 543 Martínez, B., Fernández, M., Suárez, J. E., & Rodríguez, A. (1999). Synthesis of lactococcin 972, a
544 bacteriocin produced by *Lactococcus lactis* IPLA 972, depends on the expression of a plasmid-encoded
545 bicistronic operon. *Microbiology*, 145(11). <https://doi.org/10.1099/00221287-145-11-3155>
- 546 Martínez, M. I., Rodríguez, E., Medina, M., Hernández, P. E., & Rodríguez, J. M. (1998). Detection of
547 specific bacteriocin-producing lactic acid bacteria by colony hybridization. *Journal of Applied*
548 *Microbiology*, 84(6). <https://doi.org/10.1046/j.1365-2672.1998.00448.x>
- 549 Piard, J. C., Kuipers, O. P., Rollema, H. S., Desmazeaud, M. J., & de Vos, W. M. (1993). Structure,
550 organization, and expression of the *lct* gene for lactacin 481, a novel lantibiotic produced by
551 *Lactococcus lactis*. *Journal of Biological Chemistry*, 268(22). [https://doi.org/10.1016/s0021-9258\(19\)85429-7](https://doi.org/10.1016/s0021-9258(19)85429-7)
- 553 Raimondi, S., Luciani, R., Sirangelo, T. M., Amaretti, A., Leonardi, A., Ulrici, A., Foca, G., D'Auria, G.,
554 Moya, A., Zuliani, V., Seibert, T. M., Søltøft-Jensen, J., & Rossi, M. (2019). Microbiota of sliced
555 cooked ham packaged in modified atmosphere throughout the shelf life: Microbiota of sliced cooked
556 ham in MAP. *International Journal of Food Microbiology*, 289.
557 <https://doi.org/10.1016/j.ijfoodmicro.2018.09.017>
- 558 Samelis, J., Kakouri, A., & Rementzis, J. (2000). Selective effect of the product type and the packaging
559 conditions on the species of lactic acid bacteria dominating the spoilage microbial association of
560 cooked meats at 4°C. *Food Microbiology*, 17(3). <https://doi.org/10.1006/fmic.1999.0316>
- 561 Singh, V. P. (2018). Recent approaches in food bio-preservation-A review. In *Open Veterinary Journal* (Vol.
562 8, Issue 1). <https://doi.org/10.4314/ovj.v8i1.16>
- 563 Toba, T., Samant, S. K., & Itoh, T. (1991). Assay system for detecting bacteriocin in microdilution wells.
564 *Letters in Applied Microbiology*, 13(2). <https://doi.org/10.1111/j.1472-765X.1991.tb00581.x>
- 565 van Belkum, M. J., Kok, J., & Venema, G. (1992). Cloning, sequencing, and expression in *Escherichia coli*
566 of *lcnB*, a third bacteriocin determinant from the lactococcal bacteriocin plasmid p9B4-6. *Applied and*
567 *Environmental Microbiology*, 58(2). <https://doi.org/10.1128/aem.58.2.572-577.1992>

568 Vasilopoulos, C., de Maere, H., de Mey, E., Paelinck, H., de Vuyst, L., & Leroy, F. (2010). Technology-
569 induced selection towards the spoilage microbiota of artisan-type cooked ham packed under modified
570 atmosphere. *Food Microbiology*, 27(1). <https://doi.org/10.1016/j.fm.2009.08.008>

571 Vasilopoulos, C., de Vuyst, L., & Leroy, F. (2015). Shelf-life reduction as an emerging problem in cooked
572 hams underlines the need for improved preservation strategies. *Critical Reviews in Food Science and*
573 *Nutrition*, 55(10). <https://doi.org/10.1080/10408398.2012.695413>

574 Vermeiren, L., Devlieghere, F., & Debevere, J. (2004). Evaluation of meat born lactic acid bacteria as
575 protective cultures for the biopreservation of cooked meat products. *International Journal of Food*
576 *Microbiology*, 96(2). <https://doi.org/10.1016/j.ijfoodmicro.2004.03.016>

577 Vihavainen, E., Lundström, H. S., Susiluoto, T., Koort, J., Paulin, L., Auvinen, P., & Björkroth, K. J. (2007).
578 Role of broiler carcasses and processing plant air in contamination of modified-atmosphere-packaged
579 broiler products with psychrotrophic lactic acid bacteria. *Applied and Environmental Microbiology*,
580 73(4). <https://doi.org/10.1128/AEM.01644-06>

581 Villani, F., Aponte, M., Blaiotta, G., Mauriello, G., Pepe, O., & Moschetti, G. (2001). Detection and
582 characterization of a bacteriocin, garviecin L1-5, produced by *Lactococcus garvieae* isolated from raw
583 cow's milk. *Journal of Applied Microbiology*, 90(3). <https://doi.org/10.1046/j.1365-2672.2001.01261.x>

584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603

604 **Legend to figures**

605

606 **Figure 1.** Microbial evolution of the not spoiled samples during the shelf life on MRS and BHI plates at the
607 beginning (B), middle (M) and end (E) of shelf life. Results were calculated as the means of Log counts for
608 two determinations.

609

610 **Figure 2.** AU/ml of the strains on M17 with glucose or lactose in a concentration of 0.5 % p/v. *L.*
611 *monocytogenes* EGDe was used as indicator. The results are the mean of three replicates. Values with different
612 letters are significantly different, $P < 0,05$. Capital letters indicate differences among AU/ml in lactose while
613 lowercase letters among AU/ml in glucose.

614

615 **Figure 3.** Relative abundance of the major taxonomic groups. Results are average mean of the biological
616 replicates. Only OTUs with an incidence above 0.5% in at least two samples are shown. Letters a, b, c, d
617 correspond to the production with starters 41FL18, 41FLL3, 39FL2 and 41FL1 respectively. The control (first
618 bar in each histogram) is the same for all trials.

619

620 **Figure 4.** Abundance of the major taxonomic groups of production with A16 strain. Only OTUs with an
621 incidence above 0.5% in at least two samples are shown.

622

623 **Figure 5.** PCA based on the OTU abundance of the strain 39FL2 and its interaction with spoilage consortia
624 after the 10 weeks of storage.

625

626

627


628

629

630

631


632 **Figure 1**



633

634

635 **Figure 2**



636

637


638

639


640

641 **Figure 3**


41FL18




41FLL3



39FL2




41FL1



642

643

644 **Figure 4**



645


646

647

648

649

650 **Figure 5**



651

652

653

654

655

656

657

658

659

660

661

662

663

664

665 **Table 1.** Microbial count expressed in Log (CFU/g) of the spoiled samples from the different plants (indicated
 666 by letters) in MRS and BHI media. Each line corresponds to different ham samples. Results were calculated
 667 as the means of Log counts for two determinations.

668

669

Producer	BHI counts (CFU/g Log10)	MRS counts (CFU/g Log10)
F	8.12	9.22
	7.53	8.89
	7.68	8.79
	7.44	8.45
	7.05	8.39
	6.65	8.53
	5.9	8.47
	6.06	8.53
	5.58	8.69
G	7.57	7.15
	7.57	7.47
	7.73	7.6
C	5.63	<1
	6.87	6.49
	5.88	2
H	6.96	6.91
	7.1	4.68
	7.36	6.67
	7.64	7.61
D	6.35	6.37
	8.14	7.89
	5.66	7.18
	4.88	8.28
	7.16	6.45
	6.67	8.11
A	5.79	6
I	8.82	9.26
	8.3	9.19
L	<1	5.54
	<1	<1

670

671 **Table 2.** Inhibition of spoilage consortia (expressed as % of inhibited consortia of the total of 30 tested) both
 672 in BHI and MRS by the selected *L. lactis* strains. Ability of the *L. lactis* strains to grow at refrigeration
 673 temperature is also shown.

674

Selected strains	% of inhibition against spoilage consortia		*Growth at 4°C
	BHI	MRS	4°C.- 72h (BHI)
A3	63.6%	73.9%	+
A4	81.8%	100.0%	-
A5	79.2%	100.0%	-
A6	79.2%	100.0%	-
A12	70.8%	95.7%	+
A13	70.8%	100.0%	+
A14	66.7%	100.0%	+
A16	77.3%	95.7%	+
A30	70.8%	95.7%	+
A34	70.8%	95.7%	+
A35	75.0%	95.7%	+
A36	70.8%	95.7%	+
A37	75.0%	100.0%	+
A50	75.0%	100.0%	+
A51	75.0%	100.0%	+
A52	75.0%	100.0%	+

675

676

* (+/-) indicate the ability (+) and not (-) to grow at 4°C.

677