

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The Functional Profile and Antioxidant Capacity of Tomato Fruits Are Modulated by the Interaction between Microbial Biostimulants, Soil Properties, and Soil Nitrogen Status

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1944376> since 2023-11-23T09:36:37Z

Published version:

DOI:10.3390/antiox12020520

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



The functional profile and antioxidant capacity of tomato fruits are modulated by the interaction between microbial biostimulants, soil properties, and nitrogen regimen

Paola Ganugi ¹, Andrea Fiorini ², Vincenzo Tabaglio ², Federico Capra ², Gokhan Zengin ³, Paolo Bonini ⁴, Tito Caffi ², Edoardo Puglisi ¹, Marco Trevisan ¹, and Luigi Lucini ^{1,*}

¹ Department for Sustainable Food Process, Università Cattolica del Sacro Cuore, Via Emilia Parmense 84, 29122 Piacenza, Italy. E-mail: paola.ganugi@unicatt.it (P.G.); edoardo.puglisi@unicatt.it (E.P.); marco.trevisan@unicatt.it (M.T.); luigi.lucini@unicatt.it (L.L.)

² Department of Sustainable Crop Production, Università Cattolica del Sacro Cuore, Via Emilia Parmense 84, 29122 Piacenza, Italy. E-mail: andrea.fiorini@unicatt.it (A.F.); tito.caffi@unicatt.it (T.C.); vincenzo.tabaglio@unicatt.it (V.T.)

³ Department of Biology, Faculty of Science, Selcuk University, Konya Campus, Turkey. E-mail: gokhan-zengin@selcuk.edu.tr (G.Z.)

⁴ oloBion-OMICS LIFE LAB, Barcelona, Spain. E-mail: pb@olobion.ai (P.B.)

* Correspondence: luigi.lucini@unicatt.it

Abstract: The application of microbial biostimulants to plants has revealed positive effects related to nutrients uptake, stress tolerance, root development and phenological growth. However, little information is available exploiting the potential synergistic biostimulant action of microbes on the functional quality of the yields. The current research elucidated the effect of single or coupled action of biostimulants, associated with either optimal or reduced nitrogen application, on the functional quality of tomato berries. Chemical assays and untargeted metabolomics were applied to investigate *Rhizoglyphus irregularis* and *Funneliformis mosseae* administration (both being arbuscular mycorrhiza, AMF), under optimal or low N input conditions, alone or coupled to *Trichoderma atroviride* application. The coupling of AMF and *Trichoderma* fungal inoculations resulted in a synergistic biostimulant effect on tomato fruits, revealing increased concentrations of antioxidant compounds (flavonoids, lignans and small-molecular-weight phenolics) at a higher rate than the sole AMF application. However, a strong dependence of the biostimulant effect on nitrogen availability was also noticed, reflecting significant increment in antioxidant activity under sub-optimal fertility conditions and low nitrogen levels.

Keywords: phenolic compounds; carotenoids; microbial biostimulants; antioxidant activity; metabolomics.

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Last-name

Received: date

Revised: date

Accepted: date

Published: date



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The use of plant biostimulants in agriculture has increased significantly over the last 10 years, mainly due to the successful advances in research that showed beneficial effects on plants, especially in terms of nutrient use efficiency, abiotic stress tolerance, quality traits and the availability of limited nutrients in the soil and rhizosphere [1-3].

More recently, an increasing number of experimental studies have addressed research evaluating the combined use of plant biostimulants with the aim of ensuring global food security and environmental sustainability without increasing the rate of nutrient use [4,5]. Antagonistic, additive, or rather synergistic interactions among different plant biostimulants categories have been described, depending on whether the combined effect

was respectively less, equal or greater than the effect obtained by each biostimulant individually [5]. In most cases, the combined application of microbial and non-microbial biostimulants has reflected a synergistic action on plants, revealing an increase in nutrient uptake, stress tolerance, root development and phenological growth [6-9]. However, only limited scientific literature is available regarding the effect of combined applications of biostimulants on fruit quality.

Tomato plant (*Solanum lycopersicum* L.) represents one of the most grown and consumed crops worldwide, mainly due to key role in the Mediterranean human diet as fresh or processed product. Given the commercial importance of this crop, research is addressing efforts targeted to improve the biochemical composition of the fruit, including the content of potentially health-beneficial components such as antioxidants (vitamin C, lycopene etc.). For this reason, various metabolomics approaches have been extensively adopted to gain a better understanding of the biochemical determinants concerning of fruit growth and quality, both under and presence and the absence of abiotic and biotic stress factors [10].

In this context, biostimulant products, which have been abundantly investigated for their multiple benefits for plants - including nutrient uptake and use efficiency stimulation, and abiotic/biotic stress tolerance increment -, have been recently evaluated as a sustainable approach to improve food nutritional/functional values [11,12]. Concerning tomato, latest advances in metabolomics have allowed to elucidate the physiological processes involved in fruit response to biostimulant application, which seem to be linked to higher levels of antioxidants, mineral nutrients (N, P, Ca, Na, Fe, Mn and Zn), total vitamin C and phenolics [13,14]. Nevertheless, the combined action of two or more biostimulants on tomato fruit quality remains few explored.

Given this premise, this paper gets insight into the effect of single or coupled action of biostimulants, associated with a high or low nitrogen (N) application, on the functional quality of tomato berries. Specifically, the work makes use of chemical assays and untargeted metabolomics to describe and compare the impact of no-treatment and low N input, with *Rhizoglyphus irregularis* and *Funneliformis mosseae* administration, in conventional or low N input condition, either in single or coupled to *Trichoderma atroviride* application.

2. Materials and Methods

2.1 Growth conditions and experimental design

Two field experiments were conducted between May and August 2020 at Pizzacchera S.n.c. and Felletti Luca farms, respectively located near Parma (44°50'55.7"N 10°15'34.4"E) and Ferrara (44°49'49.8"N 12°07'07.6"E), Emilia-Romagna Region, Northern Italy. The two sites were selected as optimal (Pizzacchera) and sub-optimal (Felletti) soil fertility conditions. Initial soil properties in the 0-30 cm soil layer at optimal fertility were: % sand 18.2, % silt 48.5, % clay 33.3, organic matter 40.13 g kg⁻¹, pH(H₂O) 7.98, pH(CaCl₂) 7.35, % CaCO₃ 9.7, electrical conductivity (μS cm⁻¹) 295, organic carbon (g kg⁻¹) 23.28 and total soil Kjeldahl nitrogen (g kg⁻¹) 2.55. Same properties in the same soil layer at sub-optimal fertility were: % sand 93, % silt 6, % clay 1, organic matter 8.43 g kg⁻¹, pH(H₂O) 7.54, pH(CaCl₂) 7.09, % CaCO₃ 5.8, electrical conductivity (μS cm⁻¹) 346, organic carbon (g kg⁻¹) 4.89 and total soil Kjeldahl nitrogen (g kg⁻¹) 0.60.

In both farms, the experiment was performed on non-randomized plots with four pseudo-replicates, arranged to test biostimulant-based treatments on tomato berries quality under conventional and low N input conditions. The single plot size was 144 m² (30 m x 1.6 m). In details, five different treatments were compared: 1) conventional N input without microbial treatment (Control); 2) low N input (LowN); 3) mycorrhizal treatment (*R. irregularis* BEG72 and *F. mosseae* BEG234, 700 sp g⁻¹ each species) under conventional N input (AMF); 4) mycorrhizal treatment under low N input (AMF+LowN); 5) mycorrhizal

treatment coupled with *T. atroviride* application under low N input (AMF+*T. atroviride*+LowN). The microbial biostimulants were formulated commercial products supplied by Agrotecnogias Naturales (Tarragona, Spain), inoculated at transplanting according to label recommendations.

At harvest, tomato yields were determined by weighting tomato fruits from four randomly selected areas (25 m² each) from each plot. In details, commercial (red ripe berries), immature (green underripe berries), overripe (homogenously rotten berries) fruits, as well as fruits with apical rot fruits, were separated after removal of fruits from plant stems, and weighted separately. Then, yields were expressed as t/ha. The resulting values were divided by the total yield (t/ha) to respectively obtain the percentages of commercial, immature, rotten and apical rot yield. Total refractometric (Kg°Brix/ha) and optical residue (°Brix) as qualitative parameters on berries and fruit's juice were also measured. Finally, the mature yield of 5 plants per condition was collected and immediately frozen at -20 °C. Successively, the fruits were ground with liquid nitrogen using pestle and mortar for the following chemical analysis.

2.2 Carotenoids determination

Carotenoids were determined by high performance liquid chromatography with diode array detection–mass spectrometry, as previously reported [15]. Briefly, a binary elution using (A) methanol/acetonitrile/water (84:14:4, v/v/v) and (B) dichloromethane, with a 45 min gradient run at 25 °C, together with a polymeric C30 column were employed following extraction in ethanol:*n*-hexane (60:40, v/v)[16]. Detection was then set at the wavelengths 450, 348 and 286 nm and quantification done against pure reference standards.

2.3 Total phenolics and phenolic profile

Folin-Ciocalteu and AlCl₃ assays, respectively, were utilized to determine the total phenolic and flavonoid contents [17]. Results were expressed respectively as gallic acid equivalents (mg GAEs/g extract) and rutin equivalents (mg REs/g extract).

2.3.1 Determination of antioxidant and enzyme inhibitory effects

The antioxidant and enzyme inhibitory activity of the extracts was determined according to previously described methods [18]. DPPH and ABTS radical scavenging activity, cupric ion reducing antioxidant capacity (CUPRAC), and ferric ion reducing antioxidant power (FRAP) were expressed as mg Trolox equivalents (TE)/g extract. The metal chelating ability (MCA) was reported as mg EDTA equivalents (EDTAE)/g extract, whereas the total antioxidant activity (phosphomolybdenum assay, PBD) was expressed as mmol TE/g extract. AChE and BChE inhibitory activities were given as mg galanthamine equivalents (GALAE)/g extract; tyrosinase inhibitory activity was expressed as mg kojic acid equivalents (KAE)/g extract, and amylase inhibitory activities were presented as mmol acarbose equivalents (ACAE)/g extract.

2.4 UHPLC-ESI/QTOF-MS untargeted profiling of tomato berries polyphenols

Starting from the grounded samples, tomato berries were extracted as previously described by [19]. Briefly, six replicates per thesis (2 gr each) were extracted in 20 ml of 80% methanol (v/v) acidified with 0.1% formic acid (v/v), using an Ultra-turrax (Ika, T25, Staufen, Germany). Later, the extracts were centrifuged (12000 × g) and 1 ml of the resulting supernatants was transferred into vials for the analysis.

Metabolite screening of tomato berries was performed with an untargeted metabolomics approach, throughout a hybrid quadrupole-time-of-flight mass spectrometer coupled to an ultra-high performance liquid chromatographic system (UHPLC/QTOF). Specifically, A 1290 liquid chromatograph system, equipped with a binary pump and a Dual

Electrospray JetStream ionization system, coupled to a G6550 mass spectrometer detector (Agilent technologies, Santa Clara, CA, USA) was used. The mass spectrometer worked to acquire positive ions in the 100–1200 m/z rang, while a C18 column and a binary gradient consisting of 5% to 90% methanol in water (in 30 min) was used for reverse phase chromatographic separation. The volume of injection was 6 μ L for each replicate and the flow rate was 200 μ L/min, according to [20].

The Agilent Profinder B.07 software (Agilent Technologies, Santa Clara, CA, USA) was used to process the raw data mass features, according to a targeted ‘find-by-formula’ algorithm and basing on the Phenol-Explorer 3.6 database (<http://phenol-explorer.eu>). Following mass and retention time alignment, compound identification was based on both monoisotopic accurate mass and isotope pattern (accurate spacing and isotope ratio), adopting a mass tolerance of 5-ppm. Features which were not present in 100% of replications within at least one treatment were discarded.

According to the Phenol-Explorer subclass information, phenols were classified in the phenolic subclasses, whose cumulative intensities were calculated and converted in mg L⁻¹ equivalent using standard solutions, as previously described by [21].

2.5 Statistical analyses

One and Two-way analysis of variance (ANOVA) were carried out in Rstudio software in order to determine any statistically differences among tomato berries when comparing different treatments, within and between the two farms. The elaboration was performed both for qualitative, productive, and chemical parameters. In the presence of statistically difference, Duncan test was conducted for multiple comparisons between pairs of treatments.

The statistical analyses on the metabolomics dataset were performed using Mass Profiler Professional B.12.06 (Agilent technologies) software. Compound abundance was Log₂ transformed and normalized at 75th percentile and baselined against the median. Firstly, unsupervised hierarchical cluster analysis (Squared Euclidean distance, Ward’s linkage rule) was carried out using the fold-change based heat map, to highlight the relatedness among treatments and the two farms.

Successively, for both farms, one-way ANOVA and Duncan test were carried out on semi-quantitative polyphenol results to determine whether there were any statistically significant differences between all and pairs of treatments.

3. Results

3.1 Productive and qualitative parameters

Results on qualitative and quantitative parameters of tomato production at sub-optimal fertility (Table 1) didn’t show any significant difference among treatments, except for the apical rot which proved to be considerably higher in the control samples (2.1 t/ha).

On the contrary, at optimal fertility, the reduction of nitrogen doses for LowN, AMF+LowN and AMF +*T. atroviride*+LowN led to a remarkable decrease in marketable production (Table 2). Additionally, microbial treatment didn’t reveal an increase of the total yield, having obtained the highest value for the control condition (104.9 t/ha).

Table 1. One-way analysis and Duncan test for productive and qualitative parameters at sub-optimal fertility.

	Yield										Juice
	Total refractometric (Kg°brix/ha)	Marketable (t/ha)	Immature (t/ha)	Rotten (t/ha)	Apical rot (t/ha)	Total yield (t/ha)	Commer- cial (%)	Immature (%)	Rotten (%)	Apical rot (%)	Optical residue (°brix)
Treatment (T)											
LowN	3.339	68	5.4	1.4	1.2	76.1	89.3 ^a	7.1	1.9	1.5	4.91
AMF	3.321	65.8	3.8	1.1	0.8	71.5	92 ^a	5.3	1.5	1.2	5.05
AMF+LowN	3.625	70.7	4.8	1.4	1.5	78.4	90.2 ^a	6.1	1.8	1.9	5.13
AMF+ <i>T.atr.</i> +LowN	3.698	71.9	10.1	1.4	1.5	84.9	84.6 ^b	11.9	1.6	1.8	5.15
Control	3.573	70.2	7.9	0.8	2.1	81	86.7 ^b	9.8	1	2.6	5.09
Significance											
T	n.s.	n.s.	n.s.	n.s.	*	n.s.	*	n.s.	n.s.	n.s.	n.s.

*Different letters show significant difference at the 0.05 probability level.

Table 2. One-way analysis and Duncan test for productive and qualitative parameters at optimal fertility.

	Yield										Juice
	Total refractometric (Kg°brix/ha)	Marketable (t/ha)	Immature (t/ha)	Rotten (t/ha)	Apical Rot (t/ha)	Total Yield (t/ha)	Commercial (%)	Immature (%)	Rotten (%)	Apical Rot (%)	Optical residue (°brix)
Treatment (T)											
LowN	3.803	82.7 ^b	13	2.1	0.4	98.2 ^b	84.1	13.2	2.2	0.5	4.6
AMF	4.124	88.8 ^a	12.3	3.1	0.5	104.7 ^a	84.8	11.8	3	0.5	4.65
AMF+LowN	3.548	78.6 ^b	7.4	1.9	0.5	88.5 ^b	88.9	8.3	2.2	0.6	4.51
AMF+ <i>T.atr.</i> +LowN	4.215	88 ^a	10.2	2.7	1	101.9 ^a	86.4	10	2.6	1	4.79
Control	4.082	89 ^a	12.1	3.1	0.8	104.9 ^a	84.8	11.5	2.9	0.7	4.59
Significance											

T	n.s.	*	n.s.	n.s.	n.s.	*	n.s.	n.s.	n.s.	n.s.	n.s.
---	------	---	------	------	------	---	------	------	------	------	------

*Different letters show significant difference at the 0.05 probability level.



3.2 Carotenoids determination

194

Carotenoid content in tomato berries was significantly different in the two farms, highlighting –except for B-carotene- higher concentrations of each carotenoid compound in optimal fertility samples (Table S1). However, for both cultivation sites, all-trans lycopene content was the highest, reaching the maximum average concentration of 12.586 ± 1.511 and 16.781 ± 1.797 mg/100 g extract respectively in AMF+*T.atroviride*+LowN-treated sub-optimal fertility samples and LowN-treated optimal fertility samples (Tables 3 and 4). Interestingly, biostimulant application did not enhance carotenoid content in optimal fertility tomatoes, having found that, except for Z-carotene, LowN and Control treatments determined the greatest amounts of carotenoid compounds (Table 4). Contrary, in sub-optimal fertility samples, AMF+*T.atroviride*+LowN application showed the upmost concentrations of B-carotene, Z-carotene, All-E-y-carotene, 13-z-lycopene, while Cis-lycopene was mainly increased by AMF+LowN treatment (Table 3).

195
196
197
198
199
200
201
202
203
204
205
206

3.3 Total bioactive compounds determination

207

The total content of phenolics and flavonoids in tomato berries is provided in Tables 5 and 6. Focusing on sub-optimal+optimal fertility samples, total phenolic content values ranged from 10.2 mg GAE/g extract to 15 mg GAE/g extract and the greatest amount (13.1 ± 2.2 mg GAE/g extract) was observed with Low N application, followed by AMF+*T.atroviride*+LowN treatment (12.7 ± 2.3 mg GAE/g extract) (Table S2). Concerning total flavonoid content, two-way ANOVA revealed statically differences among treatments, ascribing to Control and AMF+LowN the highest concentrations, respectively of 0.9 ± 0.3 and 0.9 ± 0.2 mg GAE/g extract. Optimal fertility samples showed the topmost average value for both phenolics and flavonoids (Table 6).

208
209
210
211
212
213
214
215
216

3.4 Total antioxidant activity

217

The total antioxidant activity calculated via phosphomolybdenum [22] is provided in Tables 5 and 6. Concerning this assay, total antioxidant abilities of optimal fertility tomato samples were significantly higher (0.9 ± 0.1 mmol TE/g extract) than those of sub-optimal fertility (0.8 ± 0.1 mmol TE/g extract) (Table S2). However, keeping together the two farms, the study did not reveal any consistent difference among treatments. Curiously, looking at one-way ANOVA results, sub-optimal fertility samples enlightened a significant increment of total antioxidant activity following AMF+*T.atroviride*+Low N treatment (Table 5), while no treatment effect was pointed out for optimal fertility (Table 6).

218
219
220
221
222
223
224
225
226

3.5 Radical scavenging activity

227

The free radical scavenging activity of tomato berries was determined using DPPH and ABTS arrays and the results were presented in Tables S2, 5 & 6. Both assays are based on the quenching of these radicals through the transfer of either an electron or a hydrogen atom by antioxidant compounds. Concerning two-way ANOVA on DPPH assay data, AMF+

228
229
230
231
232

T.atroviride+LowN treatment exhibited the highest activity (2.5 ± 2.9 mg TE/g extract) and no difference was observed between sub-optimal and optimal fertility farms (Table S2). Contrary, two-way ANOVA on ABTS assay data demonstrated a remarkable higher radical scavenging activity for optimal fertility (25.7 ± 3.4 mg TE/g extract) and for LowN application (26.4 ± 4.2 mg TE/g extract), which was significantly different from AMF treatment.

233
234
235
236
237
238

3.6 Reducing power

239

The reductive ability reflects to the electron-donation ability of antioxidant compounds. The reductive ability of tomato extracts was measured with FRAP and CUPRAC assays, respectively aimed at quantifying the potential for reducing ferric to ferrous and cupric to cuprous ions [23]. Two-way ANOVA results showed a similar tendency for both assays, highlighting a superior reducing power for AMF+*T.atroviride*+LowN (21.6 ± 4.9 mg TE/g extract for FRAP and 35.1 ± 6.2 mg TE/g extract for CUPRAC) (Table 4). Regarding CUPRAC, this last treatment significantly differed from AMF+LowN, which revealed the lowest value (30.5 ± 2.4 mg TE/g extract). However, no significance was found for the farm factor, revealing a lack of effect linked due to the cultivation area.

3.7 Metal chelating activity on ferrous ions

Ferrous chelating activity, based on the measure of the ferrous ion-ferrozine complex formation, was used as an indicator of tomato berries antioxidant activity and the results were presented in Tables 4, 5 and 6. Without detaching sub-optimal and optimal fertility samples, ferrous chelating activity data ranged from 10.2 to 16.4 mg EDTAE/g extract, confirming the highest and lowest average value respectively for LowN and AMF+LowN application (Table 4). In addition, a markedly increase for sub-optimal fertility values (15.2 ± 1.4 mg EDTAE/g extract) was revealed, when compared to optimal fertility (12.8 ± 2.6 mg EDTA/g extract).

3.8 Enzyme inhibitory activity

The results for the inhibitory activity of tomato samples on α -amylase, AChE, BChE and tyrosinase were depicted in Table S2. In general, no statistical difference between enzyme inhibition treatments was observed when comparing sub-optimal and optimal fertility (Table 4). Particularly, anti- α -Amylase activity exhibited the same values for all the treatments (0.2 ± 0.1 mmol ACAE/g extract). However, AMF+*T.atroviride*+LowN and control samples values tended to be great both for BChE (2.7 ± 0.5 and 2.7 ± 0.7 mg GALAE/g extract) and tyrosinase (57.4 ± 4.9 and 56.9 ± 6.1 mg KAE/g extract), while AMF+*T. atroviride*+Low N and AMF treatments confirmed the highest AChE inhibitory activity (2.4 ± 0.1 and 2.4 ± 0.2 mg GALAE/g extract). Except for tyrosinase, farm factor was crucial to determine dissimilarity between sub-optimal and optimal fertility sample, indicating a greater average value for cholinesterase (AChE and BChE) and Tyrosinase, respectively. Nevertheless, Table 5 & 6 revealed a different trend within 2 farms: within sub-optimal fertility, AChE and tyrosinase were significantly affected by treatment, while no differences were observed for any enzyme inhibitory activity within optimal fertility.

Table 3. One-way analysis and Duncan test for carotenoid content in tomato berries of sub-optimal fertility.

273

Source of variance	Phytoene mg/100g	Phytofluene mg/100g	Z-b-carotene mg/100g	B-carotene mg/100g	Z-carotene mg/100g	All-E-y-carotene mg/100g	13-z-lycopene mg/100g	7-z-lycopene mg/100g	9-z-lycopene mg/100g	Cis lycopene mg/100g	All trans lycopene mg/100g
Thesis(T)											
LowN	3.278±0.649 ^a	1.233±0.015 ^a	0.131±0.141	0.598±0.262	0.020±0.001	0.063±0.001 ^a	0.126±0.061	0.151±0.06	0.020±0.001 ^a	0.247±0.023	11.982±4.985
AMF	1.821±1.136 ^c	0.599±0.138 ^c	0.044±0.021	0.433±0.103	0.006±0.009	0.045±0.004 ^b	0.103±0.020	0.112±0.022	0 ^b	0.175±0.043	9.451±1.763
AMF+LowN	2.521±0.475 ^b	0.839±0.141 ^b	0.033±0.010	0.543±0.088	0.008±0.014	0.063±0.010 ^a	0.142±0.037	0.175±0.036	0.008±0.012 ^b	0.256±0.044	12.415±1.648
AMF+T.atr.+LowN	2.599±0.424 ^b	0.856±0.293 ^b	0.038±0.011	0.647±0.243	0.021±0.021	0.063±0.022 ^a	0.145±0.052	0.154±0.037	0.007±0.012 ^b	0.233±0.051	12.586±1.511
Control	3.426±0.424 ^a	1.042±0.099 ^{ab}	0.082±0.079	0.486±0.189	0.014±0.021	0.071±0.015 ^a	0.135±0.053	0.191±0.074	0.008±0.012 ^b	0.215±0.017	10.435±4.551
Significance											
T	***	***	n.s.	n.s.	n.s.	*	n.s.	n.s.	*	n.s.	n.s.

Different letters show significant difference at the 0.05 (*), 0.01 (**) and 0.001 (***) probability levels

274

275

Table 4. One-way analysis and Duncan test for carotenoid content in tomato berries of optimal fertility.

276

Source of variance	Phytoene mg/100g	Phytofluene mg/100g	Z-b-carotene mg/100g	B-carotene mg/100g	Z-carotene mg/100g	All-E-y-carotene mg/100g	13-z-lycopene mg/100g	7-z-lycopene mg/100g	9-z-lycopene mg/100g	Cis lycopene mg/100g	All trans lycopene mg/100g
Thesis(T)											
LowN	3.718±0.649	1.042±0.099 ^a	0.315±0.022	0.436±0.070 ^{bc}	0 ^b	0.079±0.008	0.215±0.020	0.262±0.037	0.024±0.002	0.418±0.0766	16.781±1.797
AMF	3.310±1.182	1.056±0.280 ^{ab}	0.255±0.055	0.332±0.099 ^c	0.008±0.013 ^{ab}	0.061±0.022	0.170±0.069	0.182±0.073	0.015±0.012	0.274±0.122	12.201±3.562
AMF+LowN	2.611±0.291	0.827±0.088 ^a	0.279±0.034	0.573±0.082 ^a	0.020±0.002 ^a	0.072±0.014	0.179±0.039	0.191±0.043	0.019±0.003	0.315±0.085	13.694±2.439
AMF+T.atr.+LowN	2.763±1.997	1.051±0.087 ^{ab}	0.295±0.018	0.647±0.243 ^c	0.009±0.014 ^{ab}	0.072±0.011	0.200±0.206	0.206±0.005	0.021±0.001	0.349±0.027	14.549±0.225
Control	3.563±1.263	1.190±0.263 ^a	0.310±0.041	0.552±0.158 ^{ab}	0.009±0.014 ^b	0.079±0.021	0.220±0.060	0.278±0.113	0.023±0.006	0.468±0.235	16.072±4.187
Significance											
T	n.s.	*	n.s.	**	*	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.

Different letters show significant difference at the 0.05 (*) and 0.01 (**) probability levels

277

Table 5. One-way analysis and Duncan test for chemical assays in tomato berries of sub-optimal fertility.

278

Source of vari- ance	Total phenolic content mgGAE/g	Total fla- vonoid content mgRE/g	DPPH mgTE/g	ABTS mgTE/g	CUPRAC mgTE/g	FRAP mgTE/g	Metal chelating mgEDTAE/g	Phos- phomo- lyb- denum mmolTE/g	AChe inh. mgGALAE/g	BChe inh. mgGALAE/g	Tyrosinase mgKAE/g	Amylase mmolACAE/g
Thesis(T)												
LowN	11.9 ± 2.0	0.8 ± 0.1 ^b	1.6 ± 0.4	23.0 ± 2.9	33.3 ± 3.8 ^{ab}	20.4 ± 1.1 ^{ab}	15.4 ± 1.4	0.8 ± 0.1 ^{ab}	2.3 ± 0.1	2.4 ± 0.3	56.9 ± 4.9 ^c	0.21 ± 0.0
AMF	11.9 ± 2.2	0.9 ± 0.1 ^{ab}	1.4 v 1.3	21.6 ± 4.4	33.4 ± 5.0 ^{ab}	20.6 ± 2.5 ^{ab}	13.6 ± 1.8	0.9 ± 0.1 ^a	2.4 ± 0.1	2.2 ± 0.6	54.9 ± 4.2 ^{bc}	0.20 ± 0.0
AMF+LowN	10.6 ± 1.1	0.9 ± 0.2 ^a	1.6 ± 0.7	20.3 ± 2.5	29.5 ± 1.9 ^b	19.6 ± 0.8 ^b	14.0 ± 2.8	0.7 ± 0.1 ^b	2.2 ± 0.1	2.4 ± 0.4	56.9 ± 6.2 ^{abc}	0.21 ± 0.0
AMF+T.atr.+LowN	12.9 ± 2.9	0.8 ± 0.1 ^{ab}	3.4 ± 3.7	24.5 ± 8.3	37.4 ± 7.6 ^a	23.6 ± 6.3 ^a	15.2 ± 1.6	0.9 ± 0.2 ^a	2.4 ± 0.1	2.3 ± 0.4	62.2 ± 3.1 ^{ab}	0.21 ± 0.0
Control	11.2 ± 1.3	0.8 ± 0.1 ^b	1.2 ± 0.4	21.2 ± 0.8	31.2 ± 1.8 ^b	19.2 ± 1.0 ^b	14.4 ± 2.5	0.8 ± 0.1 ^{ab}	2.2 ± 0.1	2.7 ± 0.9	63.5 ± 4.6 ^a	0.21 ± 0.0
Significance												
T	n.s.	*	n.s.	n.s.	*	*	n.s.	*	**	n.s.	*	n.s.

Different letters show significant difference at the 0.05 (*) and 0.01 (**) probability levels

279

280

281

Table 6. One-way analysis and Duncan test for chemical assays in tomato berries of optimal fertility.

282

Source of variance	Total phenolic content mgGAE/g	Total flavonoid content mgRE/g	DPPH mgTE/g	ABTS mgTE/g	CUPRAC mgTE/g	FRAP mgTE/g	Metal chelating mgEDTAE/g	Phosphomolybdenum mmolTE/g	AChe inh. mgGALAE/g	BChe inh. mgGALAE/g	Tyrosinase mgKAE/g	Amylase mmolACAE/g
Thesis(T)												
LowN	14.4 ± 1.7 ^a	0.9 ± 0.1 ^{bc}	3.2 ± 1.6 ^a	29.8 ± 1.7 ^a	35.6 ± 5.7	21.8 ± 2.2 ^a	14.9 ± 1.4 ^a	0.9 ± 0.1	2.4 ± 0.2	2.8 ± 0.7	53.9 ± 4.7	0.19 ± 0.0
AMF	12.7 ± 0.4 ^b	0.8 ± 0.1 ^c	1.4 ± 0.5 ^b	23.2 ± 1.4 ^c	32.5 ± 1.6	19.5 ± 0.6 ^{bc}	12.5 ± 1.2 ^b	0.9 ± 0.1	2.4 ± 0.2	3.0 ± 0.7	54.7 ± 3.5	0.18 ± 0.0
AMF+LowN	12.4 ± 0.8 ^b	1.0 ± 0.2 ^{ab}	1.5 ± 0.2 ^b	23.6 ± 0.9 ^c	31.6 ± 2.5	19.2 ± 0.9 ^c	11.5 ± 1.7 ^b	0.8 ± 0.1	2.3 ± 0.1	2.9 ± 0.4	53.8 ± 3.2	0.19 ± 0.0
AMF+T.atr.+LowN	12.5 ± 1.7 ^b	0.8 ± 0.2 ^{bc}	1.6 ± 1.4 ^b	24.8 ± 3.1 ^{bc}	32.9 ± 3.6	19.7 ± 1.4 ^{bc}	12.7 ± 2.7 ^b	0.9 ± 0.1	2.4 ± 0.1	3.1 ± 0.3	53.9 ± 4.1	0.19 ± 0.0
Control	13.4 ± 0.6 ^{ab}	1.2 ± 0.3 ^a	1.8 ± 0.1 ^b	27.1 ± 3.8 ^b	34.7 ± 2.1	20.7 ± 1.2 ^{ab}	12.4 ± 1.7 ^b	0.9 ± 0.1	2.4 ± 0.1	2.7 ± 0.6	52.9 ± 2.3	0.19 ± 0.0
Significance												
T	**	**	**	***	n.s.	**	**	n.s.	n.s.	n.s.	n.s.	n.s.

Different letters show significant difference at the 0.05 (*), 0.01 (**) and 0.001 (***) probability levels

283

284

Table 7. One-way analysis and Duncan test for phenolic classes in tomato berries under sub-optimal fertility conditions.

285

Source of variance	Flavonoids mg eq. g ⁻¹ DM	Lignans mg eq. g ⁻¹ DM	Other polyphenols mg eq. g ⁻¹ DM	Phenolic acids mg eq. g ⁻¹ DM	Stilbenes mg eq. g ⁻¹ DM
Thesis (T)					
LowN	4.7 ± 1.8 ^b	13.2 ± 1.2 ^a	52.6 ± 14.9 ^a	27.7 ± 4.5	0.4 ± 0.2
AMF	5.2 ± 0.8 ^{ab}	11.7 ± 3.3 ^{ab}	37.7 ± 15.9 ^{ab}	28.2 ± 5.6	0.4 ± 0.1
AMF+LowN	6.7 ± 1.4 ^a	6.4 ± 1.8 ^c	22.1 ± 4.6 ^b	21.9 ± 8.7	0.5 ± 0.1
AMF+T.atr.+LowN	4.5 ± 1.1 ^b	8.4 ± 3.7 ^{bc}	24.8 ± 13.3 ^b	17.1 ± 9.7	0.5 ± 0.1
Control	6.1 ± 0.9 ^{ab}	11.4 ± 2.7 ^{ab}	30.9 ± 14.1 ^b	21.9 ± 8.7	0.5 ± 0.1
Significance					
T	*	**	**	n.s.	n.s.

Different letters show significant difference at the 0.05 (*) and 0.01 (**) probability levels

286

Table 8. One-way analysis and Duncan test for phenolic classes in tomato berries under optimal fertility conditions.

Source of variance	Flavonoids mg eq. g ⁻¹ DM	Lignans mg eq. g ⁻¹ DM	Other polyphenols mg eq. g ⁻¹ DM	Phenolic acids mg eq. g ⁻¹ DM	Stilbenes mg eq. g ⁻¹ DM
Thesis (T)					
LowN	3.5 ± 0.7	13.0 ± 1.9	55.0 ± 11.5	30.8 ± 7.2	0.5 ± 0.2
AMF	2.4 ± 0.3	12.8 ± 3.1	44.2 ± 16.2	32.4 ± 6.2	0.5 ± 0.1
AMF+LowN	3.1 ± 1.0	10.9 ± 4.9	31.9 ± 13.6	24.0 ± 11.7	0.5 ± 0.2
AMF+T.atr.+LowN	3.7 ± 1.0	11.2 ± 4.0	37.9 ± 19.3	27.3 ± 12.8	0.6 ± 0.1
Control	3.8 ± 1.4	12.7 ± 2.8	37.4 ± 15.7	27.7 ± 9.6	0.6 ± 0.2
Significance					
T	n.s.	n.s.	n.s.	n.s.	n.s.

287

288

289

290

291

292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310

3.9 Metabolomics untargeted analysis of the phenolic composition of tomato berries

The metabolomic analysis with the UHPLC/QTOF investigated the phenolic compositions of tomato berry sample subjected to different biostimulant treatments. Overall, when considering all the samples analyzed, 271 phenolic compounds were putatively annotated (Table S3), mostly consisting of flavonoids (115). Moreover, 21 lignans, 62 phenolic acids and 9 stilbenes were identified, while alkylmethoxyphenol, alkylphenol, curcuminoid, furanocoumarin, hydroxybenzaldehyde, hydroxybenzochetone, hydroxybenzoketone, hydroxycinnamaldehyde, hydroxycoumarin, hydroxyphenylpropene, methoxyphenol, naphthoquinone, phenolic terpene and tyrosol compounds were grouped together as “Other polyphenols” class (64).

An unsupervised multivariate approach, consisting of a fold-change-based hierarchical clustering, was carried out in order to gain insight into the biochemical processes of tomato berries which appeared to be regulated following treatments. The analysis showed a clear division between sub-optimal and optimal fertility tomatoes, describing the farm as the unique factor affecting samples arrangement, since no clusters were formed for treatment factor (Fig. 1).




Fig. 1 Unsupervised hierarchical cluster analysis (Euclidean distance; linkage rule: Ward) of tomato berries phenolic profiles amid Control, LowN, AMF, AMF+LowN and AMF+*T.atroviride*+LowN treatments at optimal (P) and sub-optimal (F) fertility. Metabolites were obtained by UHPLC-ESI/QTOF-MS untargeted analysis, and their intensities were used to create the fold-change heatmap provided here.

311
312
313
314
315
316
317
318
319
320
321
322
323
324
325

Consequently, the hierarchical clustering analysis was repeated keeping the two farm samples separated in order to better achieve similarities and distances across treatments within the same pedoclimatic conditions (Fig. 2). Regarding sub-optimal fertility, control samples were clustered together with AMF-treated tomatoes but highlighted a markedly different metabolomic profile from those samples obtained with the coupled action of AMF and *T. atroviride* under low N input (Fig. 2A). Generally, treatments under low N rates appeared to be more distant from high N-treatments, thus showing LowN

closer to AMF+LowN and AMF+*T.atroviride*+Low N, and further away from AMF and control. On the contrary, the nitrogen level within the treatment did not represent a discerning factor affecting clustering for optimal fertility, since no-separation N level-depending was achieved between samples (Fig. 2B). Here, two clusters –respectively formed by Control and AMF+*T.atroviride*+Low N, and by AMF and AMF+LowN were observed.




Fig. 2 Unsupervised hierarchical cluster analysis (Euclidean distance; linkage rule: Ward) of sub-optimal fertility (A) and optimal fertility (B) tomato berries phenolic profiles under LowN, AMF, AMF+LowN, AMF+*T.atroviride*+Low N and Control conditions. Metabolites were obtained by UHPLC-ESI/QTOF-MS untargeted analysis, and their intensities were used to create the fold-change heatmap provided here.

Finally, ANOVA and Duncan test results for phenolic compounds as equivalents per class were provided in Table 7 & 8. Regarding sub-optimal fertility, the treatment factor was statistically determinant for flavonoid class content, which varied from 2.9 to 8.1 mg eq. g⁻¹ DM and showed the highest value (6.7 ± 1.4 mg eq. g⁻¹ DM) with AMF+LowN application. Equally, a considerable treatment effect was remarked for lignan and other polyphenol classes, whose amounts were both mostly improved by AMF treatment (respectively 11.7 ± 3.3 and 37.7 ± 15.9 mg eq. g⁻¹ DM). In contrast, in optimal fertility samples, none of the phenolic classes were significantly affected by the type of treatment.

4. Discussion

Finding new sustainable technologies to improve the functional and nutraceutical values of food products while improving yield and pomological traits has become a major research challenge due to ambitious objectives of the EU “Farm to fork” strategy [24]. In this context, the present study indicated that the use of biostimulants in agriculture may lead to a general increase of fruit quality-related compounds in *S. lycopersicum* L. On the contrary, we did not find a specific pattern in terms of tomato yield response to selected biostimulants.

Notably, mycorrhizal treatment revealed remarkable accumulations of carotenoids and phenols in tomato berries, confirming the previous findings in literature [12,25,26]. Indeed, AMF have been proved to support plant accumulation of those secondary metabolites which are involved in the response to abiotic stresses and pathogens. This results in a concrete help for the plant to counteract the negative effects of the stress and, on the other hand, in the enhancement of the functional quality of edible plant parts. Specially carotenoids, implicated in plants’ defense mechanisms as antioxidants and photo-protecting molecules, play a key role against human cancer development, thus concurring to the nutraceutical quality of plant-based foods [27]. Likewise, phenolic compounds, involved in plant responses to environmental stress including wounding, pathogen attack, mineral deficiencies, and temperature stress, have been linked to a reduced risk of cardiovascular mortality for humans thanks to their high antioxidant potential [28,29].

More in details, our study highlights that a synergic biostimulant effect may be observed at the field level with the coupled inoculation of AMF and *T. atroviride*, which revealed increases in concentration for most compounds at higher rate than those due to single mycorrhizal application in our experiment. It has been reported that *T. atroviride* acts as a biocontrol agent against a many aerial and soilborne plant pathogens, by activating different mechanisms, including competition for nutrients, production of useful secondary metabolites, modification of the rhizosphere, and mycoparasitism [30]. Our results corroborate recent research, which previously showed the synergetic potential of AMF and *T. atroviride* co-inoculation, with increased plant growth, yield, nutrient uptake and stress-tolerance [8,31,32]. Trichoderma-plant associations take place following the fungus secretion of proteins which are recognized by plant receptors. Successively, the following transient suppression of plant defenses promotes the Trichoderma penetration and, in case of co-inoculation, the concomitant access to AMF [33,34]. Similarly, it has been ascribed to AMF a corresponding help in Trichoderma conidia germination [35].

In the present study, AMF+*T.atroviride*+Low N- treated samples of sub-optimal fertility showed higher values of carotenoid concentrations -including B-carotene, α -carotene, all- ϵ -y carotene, 13- β -lycopene and all trans lycopene- and total phenolic content. Similarly, the same treatment highlighted the strongest antioxidant abilities in DPPH, ABTS, phosphomolybdenum, reducing power and enzyme inhibitory assays. This suggests that the synergic biostimulant effect of AMF and *T. atroviride* is highly related to soil fertility status (as revealed here by our initial analyses on soil OM and total N), as confirmed by our results on optimal fertility showing no treatment differences in fertile soils.

Plant N uptake is greatly aided by mutualistic association with AMF, which grow and extend their hyphae in the surrounding soil. Many studies have indicated an increased inflow of N (and other nutrients) in mycorrhizal root and, consequently, associated plants have been shown higher N concentrations than non-mycorrhized plants [36-38]. Nevertheless, several previous works have reported a remarkable decrease in the colonization intensity and diversity of mycorrhizal fungi in high-quality agricultural lands, leading to a weakening of the mutually beneficial symbiosis between AMF and plant roots [39-41]. Particularly, limited benefits of mycorrhizal inoculation benefits for agricultural production have been found under high levels of N-fertilization, denoting a better promising in low-quality lands [42,43].

In the current study, results from mycorrhized berries showed evidence of improved levels of carotenoid and bioactive compound exclusively in sub-optimal fertility, confirming the plant stronger benefit from a symbiotic relationship with AMF in soil nutrient-scarce environments than in soil nutrient-rich environments [44–47].

5. Conclusions

The present field experiment suggested the single and coupled use of microbial biostimulants to improve the quality of tomato fruits. However, the results obtained highlighted a strong dependence of the biostimulant effect on soil fertilization, reflecting significant increment in antioxidant properties only under sub-optimal conditions. Further and more in-depth studies should be carried out to fully understand the molecular and biochemical processes underlying the plant-fungi associations and the resulting changes in the functional value of fruits.

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1. Table S1: Two-way analysis and Duncan test for carotenoid content in tomato berries of sub-optimal and optimal fertility. Table S2: Two-way analysis and Duncan test for chemical assays in tomato berries of sub-optimal and optimal fertility. Table S3: List of metabolites revealed with the untargeted UHPLC-ESI/QTOF-MS analysis on tomato berries.

Author Contributions: Conceptualization, L.L.; methodology, L.L., G.Z. and M.T. and A.F.; formal analysis, P.G., F.C., P.B. and G.Z.; investigation, T.C., V.T., E.P., M.T. and L.L.; resources, L.L.; data curation, P.G., M.T., E.P., A.F. and G.Z.; writing—original draft preparation, P.G., F.C. and G.Z.; writing—review and editing, M.T., A.F. and L.L.; supervision, L.L.; project administration, L.L.; funding acquisition, L.L. All authors have read and agreed to the published version of the manuscript.

Funding: The project was funded by the Emilia-Romagna region, Italy, in the framework of the Regional Development Program PSR 2014–2020, Measure 16.1.01—Focus Area 4B (project 5150325—INBIOS: Sviluppo di un approccio INtegrato a base di BIOstimolanti per la sostenibilità delle produzioni agrarie).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Raw data are enclosed as supplementary material.

Acknowledgments: The authors thank the “Romeo ed Enrica Invernizzi” Foundation (Milan, Italy) for the kind support to the metabolomic facility at Università Cattolica del Sacro Cuore.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Halpern, M., Bar-Tal, A., Ofek, M., Minz, D., Muller, T., Yermiyahu, U. The use of biostimulants for enhancing nutrient uptake. *Adv. Agron.* **2015**, *130*, 141–174.
- Van Oosten, M. J., Pepe, O., De Pascale, S., Silletti, S., Maggio, A. The role of biostimulants and bioeffectors as alleviators of abiotic stress in crop plants. *Chem. Biol. Technol. Agric.* **2017**, *4*(1), 1–12.
- Sestili, F., Roupheal, Y., Cardarelli, M., Pucci, A., Bonini, P., Canaguier, R., Colla, G. Protein hydrolysate stimulates growth in tomato coupled with N-dependent gene expression involved in N assimilation. *Front. Plant Sci.* **2018**, *9*, 1233.
- Castiglione, A. M., Mannino, G., Contartese, V., Berteà, C. M., Ertani, A. Microbial biostimulants as response to modern agriculture needs: Composition, role and application of these innovative products. *Plants* **2021**, *10*(8), 1533.
- Roupheal, Y., Colla, G. Synergistic biostimulatory action: Designing the next generation of plant biostimulants for sustainable agriculture. *Front. Plant Sci.* **2018**, *9*, 1655.
- Baldotto, L. E. B., Baldotto, M. A., Canellas, L. P., Bressan-Smith, R., Olivares, F. L. Growth promotion of pineapple 'Vitória' by humic acids and Burkholderia spp. during acclimatization. *Rev. Bras. Cienc. Solo* **2010**, *34*, 1593–1600.

7. Bettoni, M. M., Mogor, Á. F., Pauletti, V., Goicoechea, N. Growth and metabolism of onion seedlings as affected by the application of humic substances, mycorrhizal inoculation and elevated CO₂. *Sci. Hort.* **2014**, *180*, 227-235. 448
449
8. Roupael, Y., Cardarelli, M., Bonini, P., Colla, G. Synergistic action of a microbial-based biostimulant and a plant derived-protein hydrolysate enhances lettuce tolerance to alkalinity and salinity. *Front. Plant Sci.* **2017**, *8*, 131. 450
451
9. Furlan, A., Bianucci, E., Sequeira, M., Álvarez, L., Peralta, J. M., Valente, C., Guarnieri, V., Castro, S. Combined application of microbial and non-microbial biostimulants to improve growth of peanut plants exposed to abiotic stresses. In *Microbial Probiotics for Agricultural Systems*; Zúñiga-Dávila, D., González-Andrés, F., Ormeño-Orrillo, E., Eds.; Springer: Berlin, Germany, 2019; pp. 239-256. 452
453
454
455
10. De Vos, R. C., Hall, R. D., Moing, A. Metabolomics of a model fruit: tomato. *Biol. Plant Metab.* **2011**, *43*, 109-155. 456
11. Calvo, P., Nelson, L., Kloepper, J. W. Agricultural uses of plant biostimulants. *Plant and soil* **2014**, *383*(1), 3-41. 457
12. Ganugi, P., Martinelli, E., Lucini, L. Microbial biostimulants as a sustainable approach to improve the functional quality in plant-based foods: A review. *Curr. Opin. Food Sci.* **2021**, *41*, 217-223. 458
459
13. Paul, K., Sorrentino, M., Lucini, L., Roupael, Y., Cardarelli, M., Bonini, P., Colla, G. Understanding the biostimulant action of vegetal-derived protein hydrolysates by high-throughput plant phenotyping and metabolomics: A case study on tomato. *Front. Plant Sci.* **2019**, *10*, 47. 460
461
462
14. Abou Chehade, L., Al Chami, Z., De Pascali, S. A., Cavoski, I., Fanizzi, F. P. Biostimulants from food processing by-products: agronomic, quality and metabolic impacts on organic tomato (*Solanum lycopersicum* L.). *J. Sci. Food and Agric.* **2018**, *98*(4), 1426-1436. 463
464
465
15. Hsu, B.Y., Pu, Y.S., B., Inbaraj, S., Chen, B.H. An improved high performance liquid chromatography–diode array detection–mass spectrometry method for determination of carotenoids and their precursors phytoene and phytofluene in human serum. *Journal of Chromatography B.* **2012**, *899*, 36-45. 466
467
468
16. Salomon, M.V., Piccoli, P., Fontana, A. Simultaneous determination of carotenoids with different polarities in tomato products using a C30 core-shell column based approach. *Microchem. J.* **2020**, *159*, 105390. 469
470
17. Zengin, G., & Aktumsek, A. Investigation of antioxidant potentials of solvent extracts from different anatomical parts of *Asphodeline anatolica* E. Tuzlaci: An endemic plant to Turkey. *Afr. J. Trad., Compl. Alt. Med.* **2014**, *11*(2), 481-488. 471
472
18. Sengul, U., Zengin, G., Locatelli, M., Bahadori, B., Mocan, A., Bellagamba, G., De Luca, E., Mollica, A., Aktumsek, A. Cytotoxic and enzyme inhibitory potential of two *Potentilla* species (*P. speciosa* L. and *P. reptans* Willd.) and their chemical composition. *Front. Pharm.* **2017**, *8*, 290. 473
474
475
19. Rocchetti, G., Senizza, B., Putnik, P., Bursac Kovačević, D., Barba, F. J., Trevisan, M., Lucini, L. Untargeted screening of the bound/free phenolic composition in tomato cultivars for industrial transformation. *J. Sci. Food Agric.* **2019**, *99*(14), 6173-6181. 476
477
20. Zuluaga, M. Y. A., Milani, K. M. L., Miras-Moreno, M. B., Lucini, L., Valentinuzzi, F., Mimmo, T., de Oliveira, A. L. M. The adaptive metabolomic profile and functional activity of tomato rhizosphere are revealed upon PGPB inoculation under saline stress. *Env. Exp. Bot.* **2021**, *189*, 104552. 478
479
480
21. Rocchetti, G., Blasi, F., Montesano, D., Ghisoni, S., Marcotullio, M. C., Sabatini, S., Lucini, L. Impact of conventional/non-conventional extraction methods on the untargeted phenolic profile of *Moringa oleifera* leaves. *Food Res. Int.* **2019**, *115*, 319-327. 481
482
22. Prieto, P., Pineda, M., Aguilar, M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *An. Bioch.* **1999**, *269*(2), 337-341. 483
484
23. Zengin, G., Mollica, A., Aktumsek, A., Picot, C. M. N., Mahomoodally, M. F. In vitro and in silico insights of *Cupressus sempervirens*, *Artemisia absinthium* and *Lippia triphylla*: Bridging traditional knowledge and scientific validation. *EuJIM* **2017**, *12*, 135-141. 485
486
487
24. European Commission, 2020. Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions: A Farm to Fork Strategy for a fair, healthy and environmentally-friendly food system COM/2020/381 final. 488
489
490
25. Aseel, D. G., Rashad, Y. M., Hammad, S. M. Arbuscular mycorrhizal fungi trigger transcriptional expression of flavonoid and chlorogenic acid biosynthetic pathways genes in tomato against Tomato Mosaic Virus. *Sci. Rep.* **2019**, *9*(1), 1-10. 491
492
26. Rodriguez-Heredia, M., Djian-Caporalino, C., Ponchet, M., Lapeyre, L., Canaguier, R., Fazari, A., Offroy-Chave, M. Protective effects of mycorrhizal association in tomato and pepper against *Meloidogyne incognita* infection, and mycorrhizal networks for early mycorrhization of low mycotrophic plants. *Phytopat. Medit.* **2020**, *59*(2), 377-384. 493
494
495
27. Tapiero, H., Townsend, D. M., Tew, K. D. The role of carotenoids in the prevention of human pathologies. *Biomed. Pharmacother.* **2004**, *58*(2), 100-110. 496
497
28. Chalker-Scott, L., Fuchigami, L. H. The role of phenolic compounds in plant stress responses. In *Low temperature stress physiology in crops*; Li, P.H., Ed.; CRC press: Boca Rato, Florida, 1989; pp. 67-80. 498
499
29. Vetrani, C., Costabile, G., Vitale, M., Giacco, R. (Poly) phenols and cardiovascular diseases: Looking in to move forward. *JFF* **2020**, *71*, 104013. 500
501

30. Brunner, K., Zeilinger, S., Ciliento, R., Woo, S. L., Lorito, M., Kubicek, C. P., Mach, R. L. Improvement of the fungal biocontrol agent *Trichoderma atroviride* to enhance both antagonism and induction of plant systemic disease resistance. *AEM* **2005**, *71*(7), 3959-3965. 502
503
504
31. Colla, G., Roupheal, Y., Di Mattia, E., El-Nakhel, C., Cardarelli, M. Co-inoculation of *Glomus intraradices* and *Trichoderma atroviride* acts as a biostimulant to promote growth, yield and nutrient uptake of vegetable crops. *J. Sci. Food Agric.* **2015**, *95*(8), 1706-1715. 505
506
507
32. Pascual, J.A. The Use of Arbuscular Mycorrhizal Fungi in Combination with *Trichoderma* spp. in Sustainable Agriculture. In *Bioformulations: for Sustainable Agriculture*. Arora N., Mehnaz S., Balestrini R., Eds.; Springer: New Delhi, India; pp. 137-146. 508
509
33. Alonso-Ramírez, A., Poveda, J., Martín, I., Hermosa, R., Monte, E., Nicolás, C. Salicylic acid prevents *Trichoderma harzianum* from entering the vascular system of roots. *Mol. Plant Pathol.* **2014**, *15*(8), 823-831. 510
511
34. Mendoza-Mendoza, A., Zaid, R., Lawry, R., Hermosa, R., Monte, E., Horwitz, B. A., Mukherjee, P. K. Molecular dialogues between *Trichoderma* and roots: role of the fungal secretome. *Fungal Biol. Rev.* **2018**, *32*(2), 62-85. 512
513
35. Poveda, J., Hermosa, R., Monte, E., Nicolás, C. (2019). *Trichoderma harzianum* favours the access of arbuscular mycorrhizal fungi to non-host Brassicaceae roots and increases plant productivity. *Sci. Rep.* **2019**, *9*(1), 1-11. 514
515
36. Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M. I., Ashraf, M., Ahmed, N., Zhang, L. Role of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress tolerance. *Front. Plant Sci.* **2019**, *10*, 1068. 516
517
37. Savolainen, T., Kytöviita, MM. Mycorrhizal symbiosis changes host nitrogen source use. *Plant Soil* **2022**, *471*, 643-654. 518
38. Yang, H., Fang, C., Li, Y., Wu, Y., Fransson, P., Rillig, M. C., Weih, M. Temporal complementarity between roots and mycorrhizal fungi drives wheat nitrogen use efficiency. *New Phyt.* **2022**, *236*(3), 1168-1181. 519
520
39. Lin, C., Wang, Y., Liu, M., Li, Q., Xiao, W., & Song, X. Effects of nitrogen deposition and phosphorus addition on arbuscular mycorrhizal fungi of Chinese fir (*Cunninghamia lanceolata*). *Sci. Rep.* **2020**, *10*(1), 1-8. 521
522
40. Emery, S. M., Bell-Dereske, L., Stahlheber, K. A., Gross, K. L. Arbuscular mycorrhizal fungal community responses to drought and nitrogen fertilization in switchgrass stands. *Appl. Soil Ecol.* **2022**, *169*, 104218. 523
524
41. Muneer, M. A., Tarin, M. W. K., Chen, X., Afridi, M. S., Iqbal, A., Munir, M. Z., Ji, B. Differential response of mycorrhizal fungi linked with two dominant plant species of temperate grassland under varying levels of N-addition. *Appl. Soil Ecol.* **2022**, *170*, 104272. 525
526
527
42. Yang, H., Fang, C., Li, Y., Wu, Y., Fransson, P., Rillig, M. C., Weih, M. Temporal complementarity between roots and mycorrhizal fungi drives wheat nitrogen use efficiency. *New Phytol.* **2022**, *236*(3), 1168-1181. 528
529
43. Wang, X. X., Wang, X., Sun, Y., Cheng, Y., Liu, S., Chen, X., Kuyper, T. W. Arbuscular mycorrhizal fungi negatively affect nitrogen acquisition and grain yield of maize in a N deficient soil. *Front. Microb.* **2018**, *9*, 418. 530
531
44. Chen, J., Li, J., Yang, Y., Wang, Y., Zhang, Y., & Wang, P. Effects of Conventional and Organic Agriculture on Soil Arbuscular Mycorrhizal Fungal Community in Low-Quality Farmland. *Front. Microb.* **2022**, *13*. 532
533
45. Hoeksema, J. D., Chaudhary, V. B., Gehring, C. A., Johnson, N. C., Karst, J., Koide, R. A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecol. Lett.* **2010**, *13*, 394-407. 534
535
46. Zaller, J. G., Frank, T., and Drapela, T. Soil sand content can alter effects of different taxa of mycorrhizal fungi on plant biomass production of grassland species. *Eur. J. Soil Biol.* **2011**, *47*, 175-181. 536
537
47. Shen, K., He, Y., Xu, X., Umer, M., Liu, X., Xia, T., Yan, J. Effects of AMF on plant nutrition and growth depend on substrate gravel content and patchiness in the karst species *Bidens pilosa* L. *Front. Plant Sci.*, **2022**, *13*. 538
539
540

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content. 541
542
543