

On the Berger conjecture for manifolds all of whose geodesics are closed

Marco Radeschi¹ · Burkhard Wilking¹

Received: 8 March 2016 / Accepted: 20 June 2017
© Springer-Verlag GmbH Germany 2017

Abstract We define a Besse manifold as a Riemannian manifold (M, g) all of whose geodesics are closed. A conjecture of Berger states that all prime geodesics have the same length for any simply connected Besse manifold. We firstly show that the energy function on the free loop space of a simply connected Besse manifold is a perfect Morse–Bott function with respect to a suitable cohomology. Secondly we explain when the negative bundles along the critical manifolds are orientable. These two general results, then lead to a solution of Berger’s conjecture when the underlying manifold is a sphere of dimension at least four.

Mathematics Subject Classification 53C22 · 58E10

Contents

1	Free loop space, and Morse–Bott theory	
1.1	Critical submanifolds	
2	Orientability of the negative bundle	
2.1	Idea behind the proof	
2.2	The plan	
2.3	Algebraic data	

✉ Marco Radeschi
mradesch@nd.edu

Burkhard Wilking
wilking@uni-muenster.de

¹ University of Münster, Einsteinstraße 62, 48149 Münster, Germany

2.4	Paths of closed geodesics
2.5	The variation
2.6	Making the Poincaré map generic
2.7	The modified negative bundle
2.8	Local orientability of pseudo vector bundles
2.9	Local orientability of $N \oplus \mathcal{E}$
2.10	If A_S is not nullhomotopic
3	Free loop spaces of cohomology CROSSes
3.1	Integral cohomology CROSSes
4	Index gap
5	Perfectness of the energy functional
5.1	When the negative bundles are all orientable
5.2	The general case
6	The proof of the Main Theorem
	Appendix A. Small subsets of $\mathrm{Sp}(n-1, \omega)$
	References

Riemannian manifolds with all geodesics closed have been object of study since the beginning of the XX century, when the first nontrivial examples were produced by Tannery and Zoll. To this day the famous book of Besse [3] describes the state of art to a large extent with a few but notable exceptions. We define a Besse metric g on a manifold M , as a Riemannian metric all of whose geodesics are closed and a Besse manifold as a manifold endowed with a Besse metric.

The “trivial” examples of Besse manifolds are the compact rank one symmetric spaces (also called CROSSes), with their canonical metrics: round spheres \mathbb{S}^n , complex and quaternionic projective spaces $\mathbb{C}\mathbb{P}^n$, $\mathbb{H}\mathbb{P}^n$ with their Fubini-Study metric, and the Cayley projective plane $Ca\mathbb{P}^2$. To this day, these are also the only manifolds that are known to admit a Besse metric. Moreover, on the projective spaces the canonical metric is the only known Besse metric. On the other hand, in the case of spheres many other Besse metrics have been discovered by Zoll, Berger, Funk, and Weinstein.

Given a Besse manifold, a theorem of Wadsley states that all the prime geodesics (i.e. which are not iterates of a shorter one) have a common period L . Using this result, a combination of theorems by Bott–Samelson [3, Thm 7.37] and McCleary [10, Cor. A] proves that any simply connected Besse manifold M has the integral cohomology ring of a CROSS, the so called *model* of M .

A conjecture of Berger states that on a simply connected Besse manifold all the prime geodesics have the *same* length L . This strengthens Wadsley’s result in the simply connected case. If all prime geodesics have the same length one knows, for example, that the volume of M is an integer multiple of the volume of a round sphere of radius $L/2\pi$ [15].

This conjecture was proved for 2 spheres by Gromoll and Grove [7]. Partial results in dimension three have been obtained by Olsen [11]. On the other hand, the conjecture of Berger does not hold for irreversible Finsler metrics.

Counterexamples come from certain so-called *Katok metrics* on CROSSes, as pointed out for example in [19].

These counterexamples suggest that reversibility should come as a crucial ingredient in the proof of the Berger’s Conjecture.

The main goal of this paper is to prove the conjecture for all topological spheres of dimension at least 4.

Theorem A *Let M be a topological sphere of dimension ≥ 4 . Then, for every Besse metric on M , all prime geodesics have the same length.*

To prove Theorem A, we study the free loop space $\Lambda M = \mathcal{H}^1(\mathbb{S}^1, M)$ (the Sobolev space of maps $\mathbb{S}^1 \rightarrow M$) of any Besse manifold. In particular, we study the energy functional $E : \Lambda M \rightarrow \mathbb{R}$, which associates to any free loop $\gamma : \mathbb{S}^1 \rightarrow M$ the energy

$$E(\gamma) = \int_{\mathbb{S}^1} \|\gamma'(t)\|^2 dt.$$

Basically, the proof of Theorem A is by contradiction: if not all prime geodesics in M have the same length, then the set C_{min} in ΛM consisting of nonconstant geodesics of shortest length (which corresponds to the set of critical points of the energy E , with lowest nonzero critical value) must have the integral (co)homology of a sphere. On the other hand, since the metric is Riemannian and, in particular, reversible (in the sense that if $\gamma(t)$ is a closed geodesic, then $\gamma(-t)$ is also a geodesic, of the same length as $\gamma(t)$), then the energy functional, and thus the set C_{min} , is invariant under the natural action of $O(2)$ acting by reparametrization. In particular, C_{min} admits a free $\mathbb{Z}_2 \times \mathbb{Z}_2 \subset O(2)$ action, which is known not to be possible if C_{min} is an integral homology sphere.

To prove that the set C_{min} has the required homology, we combine the simplicity of the topology of the the free loop space of M with Morse–Bott theory. To that end we prove two general results which hold on all simply connected Besse manifolds. The second named author proved in [17] that E is a Morse–Bott function. In particular, the critical points of E , which precisely correspond to the closed geodesics in M , form smooth “critical” submanifolds. Moreover, along every critical submanifold C the negative eigenspaces of the Hessian of E give rise to a “negative bundle” $\mathbf{N} \rightarrow C$, of finite rank. In order to apply Morse–Bott theory, it is important to make sure that these negative bundles are orientable.

For the following theorem, notice that a closed curve $c : \mathbb{S}^1 \rightarrow C$ in a critical manifold C , $s \mapsto c_s$, induces a curve $\alpha : \mathbb{S}^1 \rightarrow M$, $\alpha(s) = c_s(0)$, and a bundle α^*TM over \mathbb{S}^1 . Choosing a framing of TM along α induces a trivialization of α^*TM . The map $A : \mathbb{S}^1 \rightarrow SO(n)$ sending s to the holonomy

map $A_s \in \mathrm{SO}(T_{c_s(0)}M) \simeq \mathrm{SO}(n)$ along c_s may depend on the choice of the framing, but its (free) homotopy type does not.

Theorem B *Let M be an orientable Besse manifold, and $C \subseteq \Lambda M$ a critical submanifold of the energy functional. Then for any curve $c : \mathbb{S}^1 \rightarrow C$ the negative bundle over c is orientable if and only if the map $A : \mathbb{S}^1 \rightarrow \mathrm{SO}(n)$ along c is nullhomotopic in $\mathrm{SO}(n)$.*

The proof of Theorem B allows in a way to use the Poincaré map to define a natural orientation on the negative bundle. One might hope to use similar ideas in other contexts, e.g. to define (under suitable assumptions) a natural orientation on the unstable manifold along a periodic Reeb orbit. If M is a spin manifold, the holonomy $A : C \rightarrow \mathrm{SO}(n)$ always lifts to a map $\tilde{A} : C \rightarrow \mathrm{Spin}(n)$, and in particular A is always contractible. It follows in particular:

Corollary C *Let M be an orientable spin Besse manifold. Then for every critical submanifold $C \subseteq \Lambda M$ for the energy functional, the negative bundle is orientable.*

Almost all Besse manifolds are spin: in fact, this is the case if the model of M is not given by $\mathbb{C}\mathbb{P}^{2m}$. Since M has the same integral cohomology as its model this follows either from $H^2(M; \mathbb{Z}_2) = 0$ or in case of the model $\mathbb{C}\mathbb{P}^{2m+1}$ from the fact that then M is homotopy equivalent to its model and the homotopy invariance of Stiefel-Whitney classes $w_2(M) = w_2(\mathbb{C}\mathbb{P}^{2m+1}) = 0$.

Recall that $\mathrm{O}(2)$ acts on ΛM by reparametrization, and the energy functional is $\mathrm{O}(2)$ -invariant. In particular, we can use E as an \mathbb{S}^1 -equivariant Morse–Bott function, with $\mathbb{S}^1 = \mathrm{SO}(2) \subseteq \mathrm{O}(2)$. Let $M \rightarrow \Lambda M$ denote the embedding sending $p \in M$ to the constant curve $\gamma \equiv p$, and denote the image of such embedding again by M .

Theorem D *Let M be a simply connected Besse manifold. Then the energy functional is a perfect Morse–Bott function for the rational, \mathbb{S}^1 -equivariant cohomology of the pair $(\Lambda M, M)$.*

The proof of the perfectness of the energy functional uses three main ingredients.

- (1) *Index parity*: On an orientable Besse manifold M the index of every closed geodesic has the same parity as $\dim M + 1$.
- (2) *Lacunarity*: If a manifold M has the integral cohomology of a CROSS then, up to inverting a finite number of primes, the cohomology $H_{\mathbb{S}^1}^q(\Lambda M, M; \mathbb{Z})$ is zero whenever q has the same parity as $\dim M$.
- (3) *Index gap*: If c is a closed geodesic on a simply connected Besse manifold M of length iL/q , where L is the common period of all geodesics and $i, q \in \mathbb{N}$, then, roughly speaking, the differences $\mathrm{ind} c^q - \mathrm{ind} c^{q'}$, $q' \neq q$, are bounded away from zero by some constant independent from c .

The index parity was proved by the second-named author in [17]. The Lacunarity and the Index gap are proved in the present paper.

The paper is structured as follows. Section 2 contains the proof of Theorem B. Several technical results are proved in the appendix. Sections 3 and 4 contain the precise statements and proofs of the Lacunarity and the Index gap, respectively. Finally, in Sect. 5 Theorem D is proved, while Sect. 6 contains the proof of Theorem A.

1 Free loop space, and Morse–Bott theory

Let M be a Besse manifold. As defined above a Besse manifold is a Riemannian manifold all of whose geodesics are closed. Given $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$, we define the *free loop space of M* , to be the Hilbert manifold $\Lambda M = \mathcal{H}^1(\mathbb{S}^1, M)$ of maps $c : \mathbb{S}^1 \rightarrow M$ of class \mathcal{H}^1 . The *energy functional* $E : \Lambda M \rightarrow \mathbb{R}$ is defined as

$$E(c) = \int_{\mathbb{S}^1} \|c'(t)\|^2 dt.$$

As pointed out in [17], the energy functional is a Morse–Bott function, and any critical set K of energy e consists of all closed geodesics of length $\ell = \sqrt{e}$.

From now on, we will use the following notation:

- We denote by $0 = e_0 \leq e_1 \leq \dots$ the sequence of critical values of E (or *critical energies*). Each critical value appears as many times as the number of connected components of the corresponding critical manifold.
- For every critical energy e_i , we denote by $K_i \subseteq E^{-1}(e_i)$ the corresponding critical manifold, and $\Lambda^i = E^{-1}([0, e_i + \epsilon))$ the corresponding sub-level set. Since each critical value appears with multiplicity, each critical manifold K_i is connected.
- Along a critical manifold K_i , the number of negative eigenvalues of $\text{Hess}(E)|_{\nu(K_i)}$, counted with multiplicity, is constant, as otherwise there would exist a point in K_i where the Hessian has some nontrivial kernel—contradicting the fact that E is a Morse–Bott function. We can thus define the *index* of K_i , $\text{ind}(K_i)$, to be this number.
- For every critical manifold K_i , denote by $\mathbf{N}_i \rightarrow K_i$ the subbundle of $\nu(K_i)$ consisting of the negative eigenspaces of $H_i = \text{Hess}(E)|_{\nu(K)}$. We call \mathbf{N}_i the *negative bundle* of K_i .

For $i = 0$, the critical set K_0 consists of point curves and will be from now on identified with M . For each $i > 0$, let $\mathbf{N}_i^{\leq 1} = \{v \in \mathbf{N}_i \mid \|v\| \leq 1\}$, for some choice of norm on \mathbf{N}_i . By Morse–Bott theory, each sublevel set Λ^k is homotopy equivalent to the complex obtained by successfully attaching the unit disk bundles $\mathbf{N}_i^{\leq 1}$, $i = 1, \dots, k$, to $M = K_0$.

If one were to know the relative cohomology of the pairs $(\Lambda^i, \Lambda^{i-1})$, it would be possible to reconstruct the cohomology groups $H^*(\Lambda^i; \mathbb{Z})$ iteratively, using the long exact sequence of the pair $(\Lambda^i, \Lambda^{i-1})$, and in the limit one would reconstruct the cohomology of ΛM . By excision, the relative homology group of the pair $(\Lambda^i, \Lambda^{i-1})$ is equal to the relative homology group of $(\mathbb{N}_i^{\leq 1}, \partial\mathbb{N}_i^{\leq 1})$. If $\mathbb{N}_i \rightarrow K_i$ is orientable, then by the Thom isomorphism the cohomology of $(\mathbb{N}_i^{\leq 1}, \partial\mathbb{N}_i^{\leq 1})$ equals the shifted cohomology of K_i :

$$H^k(\Lambda^i, \Lambda^{i-1}; \mathbb{Z}) \simeq H^k(\mathbb{N}_i^{\leq 1}, \partial\mathbb{N}_i^{\leq 1}; \mathbb{Z}) \simeq H^{k-\text{ind}(K_i)}(K_i; \mathbb{Z}).$$

Notice that the group \mathbb{S}^1 acts on ΛM by reparametrization, and the energy functional is \mathbb{S}^1 -invariant. In particular, the critical manifolds and the sublevel sets are all \mathbb{S}^1 invariant, and there is a natural \mathbb{S}^1 -action on the negative bundles such that the maps $\mathbb{N}_i \rightarrow K_i$ are all \mathbb{S}^1 -equivariant. Moreover, by choosing an \mathbb{S}^1 -invariant metric on \mathbb{N}_i , the inclusion $(\mathbb{N}_i^{\leq 1}, \partial\mathbb{N}_i^{\leq 1}) \rightarrow (\Lambda^i, \Lambda^{i-1})$ is also \mathbb{S}^1 -equivariant. In particular, using \mathbb{S}^1 -equivariant cohomology we have that whenever $\mathbb{N}_i \rightarrow K_i$ is orientable, there is an isomorphism

$$H_{\mathbb{S}^1}^k(\Lambda^i, \Lambda^{i-1}; \mathbb{Z}) \simeq H_{\mathbb{S}^1}^{k-\text{ind}(K_i)}(K_i; \mathbb{Z}).$$

1.1 Critical submanifolds

Given a Besse manifold M , by Wadsley Theorem [14] the prime geodesics of M have a common period L , thus the geodesic flow induces a \mathbb{S}^1 -action on the unit tangent bundle T^1M of M . Moreover, if a critical set $K \subseteq E^{-1}(\ell^2)$ contains prime geodesics, then the common length ℓ of the geodesics in K equals $\ell = L/k$ for some integer k , and the map

$$K_i \rightarrow T^1M, \quad c \mapsto (c(0), c'(0)/\ell)$$

defines a diffeomorphism of K_i into the subset of T^1M fixed by $\mathbb{Z}_k \subseteq \mathbb{S}^1$. This diffeomorphism is \mathbb{S}^1 -equivariant, where \mathbb{S}^1 acts on K_i by reparametrization, and on T^1M by the geodesic flow. In particular if the length of the geodesics in K is a multiple of L , then K is diffeomorphic to T^1M .

In the rest of the paper we will use the following notation:

- We let $\{C_1, \dots, C_k\}$ denote the set of critical manifolds containing prime geodesics.
- We call *regular* any geodesic whose length is a multiple of L . Similarly, we say that a critical set is *regular* if it contains regular geodesics.
- Given a closed geodesic $c : [0, 1] \rightarrow M$, we denote by $c^q : [0, 1] \rightarrow M$, $c^q(t) = c(qt)$, the q -iterate of c .

Any critical set of positive energy can be written as $C^q = \{c^q \mid c \in C\}$ for some integer q and some C in $\{C_1, \dots, C_k\}$.

2 Orientability of the negative bundle

Let M be a Besse manifold. The goal of this section is to prove Theorem B. The techniques used in this sections are completely independent from the remaining sections, and therefore can be read independently or used as a black-box.

2.1 Idea behind the proof

In [17] the second-named author proved that the the Poincaré map of a closed geodesic determines the parity of its index. In a way, we will prove here the orientability of the negative bundle by showing that the Poincaré map of a closed geodesic c can help define a natural orientation on the negative space at c .

More precisely, consider a “formal closed geodesic”, modelled by a map $R \in \text{Map}([0, \pi], \text{Sym}(n - 1, \mathbb{R}))$ (the curvature operator along the geodesic) and a map $A \in \text{SO}(n - 1)$ (the holonomy along the geodesic). Given this data it is possible to define an “index form” H for (R, A) , modelling the Hessian of the energy functional, and a “negative space” N defined as the sum of the negative eigenspaces of H , which models the space of negative directions in the free loop space (see Sect. 2.3).

Choose a generic path $(R_\tau, A_\tau)_{\tau \in [0,1]}$ of data from (R, A) to the fixed data (Id, Id) . This path induces a family of index forms H_τ , and a corresponding family of negative spaces N_τ . The idea is that, by fixing an orientation of N_1 , we want to induce an orientation on $N_0 = N$ along the path. This can be easily done if H_τ does not change index along the path, in which case the collection $\{N_\tau\}_{\tau \in [0,1]}$ defines a vector bundle on $[0, 1]$. In general however H_τ does change index, and at the transition points H_τ has nontrivial kernel, as some eigenvalue of H_τ is changing sign.

For each τ it also makes sense to define a Poincaré map P_τ of the “formal geodesic” modelled by (R_τ, A_τ) , and one can consider the space

$$\mathcal{E}_\tau = \bigoplus_{\lambda \in (0,1)} E_\lambda(P_\tau),$$

with $E_\lambda(P_\tau)$ the eigenspace of P_τ of eigenvalue λ . Just like in the geometric setting, there is a natural identification between the kernel of H_τ (given by “periodic Jacobi fields”) and the eigenspace $E_1(P_\tau)$. Moreover, it was proved in [17] that the dimension of $\mathcal{E}_\tau \oplus N_\tau$ has constant parity at generic points. If P_τ is generic enough, it has eigenvalue 1 of geometric multiplicity at most 1,

and therefore at every transition point both the index of H_τ (which equals the dimension of \mathbf{N}_τ) and the number of real eigenvalues of P in $(0, 1)$ (which equals the dimension of \mathcal{E}_τ) only change by 1.

In other words: at any transition point, the form H_τ will develop a one-dimensional kernel V_H , the Poincaré map P_τ will develop a one-dimensional fixed space V_P , such that there is a natural identification $V_H \simeq V_P$. Thus, at transition points, the sum $\mathbf{N}_\tau \oplus \mathcal{E}_\tau$ either gains or loses two one-dimensional subspaces that can be canonically identified. Since the sum of two identical subspaces carries a natural orientation, there is an obvious way to induce an orientation from $\mathbf{N}_{\tau-\epsilon} \oplus \mathcal{E}_{\tau-\epsilon}$ to $\mathbf{N}_{\tau+\epsilon} \oplus \mathcal{E}_{\tau+\epsilon}$. In our situation, we have $\mathcal{E}_0 = 0 = \mathcal{E}_1$ and therefore the orientation on $\mathbf{N}_1 \oplus \mathcal{E}_1 = \mathbf{N}_1$ induces naturally an orientation on $\mathbf{N}_0 \oplus \mathcal{E}_0 = \mathbf{N}$.

Of course one needs to prove that the induced orientation on \mathbf{N} does not depend on the path chosen, and therefore one needs a 2-dimensional variation, in which case more problems arise because the dimension of \mathcal{E}_τ can jump for other reasons (two real eigenvalues of P_τ could “collide and disappear”, for example), which should be taken into account.

2.2 The plan

We sketch here the main steps of the proof: in Sect. 2.3 we define certain algebraic data $(R_x, A_x)_{x \in \mathfrak{M}}$ parametrized by a manifold \mathfrak{M} , which formally model geometric structures around a closed geodesic. Moreover we show how one can construct, out of these data, a map $\mathbf{N} \rightarrow \mathfrak{M}$ which resembles a vector bundle (we call these *pseudo vector bundles*).

In Sect. 2.4 we consider a Besse manifold M and a loop $c_s : \mathbb{S}^1 \rightarrow \Lambda M$ in a critical manifold for the energy functional. We then explain how to associate an algebraic data set $(R_s, A_s)_{s \in \mathbb{S}^1}$ to c_s , and that proving the orientability of the negative bundle along c_s is equivalent to proving the orientability of the pseudo vector bundle $\mathbf{N} \rightarrow \mathbb{S}^1$ induced by the algebraic data.

At first we assume that $A : \mathbb{S}^1 \rightarrow \mathrm{SO}(n-1)$ is nullhomotopic, leaving the other case to the last Sect. 2.10. In Sects. 2.5 and 2.6 we produce a one-parameter deformation

$$(R_{s,\tau}, A_{s,\tau})_{(s,\tau) \in \mathbb{S}^1 \times [0,1]}$$

of algebraic data, which is the original one for $\tau = 0$ and has trivial negative bundle for $\tau = 1$. This variation induces a pseudo vector bundle $\mathbf{N} \rightarrow \mathbb{S}^1 \times [0, 1]$, whose restriction to $\mathbb{S}^1 \times \{0\}$ is the original one, and whose restriction to $\mathbb{S}^1 \times \{1\}$ is orientable since the bundle is trivial.

In Sect. 2.7 we define a “modified” pseudo vector bundle $\mathbf{N} \oplus \mathcal{E} \rightarrow \mathbb{S}^1 \times [0, 1]$, whose restriction to $\mathbb{S}^1 \times \{0\}$ and $\mathbb{S}^1 \times \{1\}$ coincides with \mathbf{N} .

In Sect. 2.9 we prove that a notion of orientability can be defined for $\mathbf{N} \oplus \mathcal{E}$, in such a way that $(\mathbf{N} \oplus \mathcal{E})|_{\mathbb{S}^1 \times \{0\}}$ is orientable if and only if $(\mathbf{N} \oplus \mathcal{E})|_{\mathbb{S}^1 \times \{1\}}$ is orientable.

Finally, in Sect. 2.10, we discuss the case in which $A : \mathbb{S}^1 \rightarrow \text{SO}(n - 1)$ is not nullhomotopic. Using the results of the previous sections, we prove that in this case the negative bundle is not orientable.

2.3 Algebraic data

Let M be a Riemannian manifold and $c : [0, 2\pi] \rightarrow M$ a closed geodesic. Parallel translation along c allows to identify the spaces $c'(t)^\perp$ with $V = c'(0)^\perp$, and it defines a map $A \in \text{O}(V)$ defined by $A(e(0)) = e(2\pi)$ for every parallel normal vector field $e(t)$ along c . Moreover, the curvature operator of M defines a map $R : [0, 2\pi] \rightarrow \text{Sym}^2(V)$ by

$$\langle R(t) \cdot e_1(0), e_2(0) \rangle = \langle R(e_1(t), c'(t))c'(t), e_2(t) \rangle$$

for every parallel normal vector fields $e_1(t), e_2(t)$ along c .

We say that the data (R, A) model a “formal geodesics”, because out of such data one can formally recover a number of objects usually related to real geodesics.

In fact, given the data of an Euclidean vector space $V \simeq \mathbb{R}^{n-1}$, a piecewise continuous map $R \in \text{Map}([0, 2\pi], \text{Sym}^2(n - 1))$ and a $A \in \text{O}(n - 1)$, we can define:

- The space $\mathfrak{X} = \{X : [0, 2\pi] \rightarrow V \mid X(2\pi) = A \cdot X(0)\}$ of *periodic vector fields*.
- The space $\mathfrak{J} = \{J : [0, 2\pi] \rightarrow V \mid J''(t) + R(t) \cdot J(t) = 0\}$ of *Jacobi fields*.
- The *Poincaré map* $P : V \oplus V \rightarrow V \oplus V$ which sends (u, v) to $(A^{-1} \cdot J(2\pi), A^{-1} \cdot J'(2\pi))$ where J is the unique Jacobi field with $J(0) = u, J'(0) = v$. This map preserves the symplectic form $\omega((u_1, v_1), (u_2, v_2)) = \langle u_1, v_2 \rangle - \langle u_2, v_1 \rangle$ and thus $P \in \text{Sp}(n - 1, \omega)$.
- The *index operator* $H : \mathfrak{X} \times \mathfrak{X} \rightarrow \mathbb{R}$ defined as

$$H(X, Y) = \int_0^{2\pi} \langle X'(t), Y'(t) \rangle - \langle R(t)(X(t)), Y(t) \rangle dt$$

The index operator is symmetric, and by standard arguments it can be checked that the sum $\mathbf{N} \subseteq \mathfrak{X}$ of the negative eigenspaces of H is finite dimensional. We call this sum the *negative space* of the pair (R, A) .

Given a manifold \mathfrak{M} and maps

$$A : \mathfrak{M} \rightarrow \text{SO}(n - 1), \quad R : \mathfrak{M} \rightarrow \text{Map}([0, 2\pi], \text{Sym}^2(n - 1))$$

then for every $x \in \mathfrak{M}$ the data (R_x, A_x) determine, in particular, a Poincaré map $P_x \in \text{Sp}(n - 1, \omega)$ an index form H_x and a negative space $N_x \in \mathfrak{X}$.

This determines a map $P : \mathfrak{M} \rightarrow \text{Sp}(n - 1, \omega)$ and a space $N = \coprod_{x \in \mathfrak{M}} N_x$ with a projection $N \rightarrow \mathfrak{M}$ sending N_x to x . The space N has a natural topology, induced by the inclusion $N \subseteq \mathfrak{M} \times \text{Map}([0, 2\pi], V)$, and $N \rightarrow \mathfrak{M}$ has the structure of a fiberwise vector space.

In general, however, the map $N \rightarrow \mathfrak{M}$ is not a vector bundle, since the dimension of the fibers might change from point to point.

Remark 2.1 Notice that, given data $(R_x, A_x)_{x \in \mathfrak{M}}$, there is always an isomorphism $\ker H_x \rightarrow E_1(P_x)$ ($E_1(P_x)$ is the eigenspace of P_x with eigenvalue 1). In fact, a vector field X in $\ker H_x$ is a periodic Jacobi field, in the sense that $X''(t) + R_x(t) \cdot X(t) = 0$ and $(X(2\pi), X'(2\pi)) = (A \cdot X(0), A \cdot X'(0))$. In particular, the vector $(X(0), X'(0))$ is a fixed vector for P_x and therefore the map $X \mapsto (X(0), X'(0))$ defines the isomorphism $\ker H_x \rightarrow E_1(P_x)$.

In particular, when dimension of the fibers of $N \rightarrow \mathfrak{M}$ changes, at the transition point the kernel of H_x must be nontrivial and therefore P_x must have eigenvalue 1. This will be very important later on.

2.4 Paths of closed geodesics

Let $C \subseteq \Lambda M$ be a critical submanifold of the energy functional E and let $c : \mathbb{S}^1 \rightarrow C, s \mapsto c_s$ be a closed curve. Each c_s defines a unit speed closed geodesic which, possibly after scaling, can be parametrized as $c_s : [0, 2\pi] \rightarrow M$.

Along $\alpha(s) = c_s(0)$, the bundle $c'_s(0)^\perp$ over \mathbb{S}^1 is orientable, and every orientable vector bundle over \mathbb{S}^1 is trivializable. Thus, let $\{e_1(s), \dots, e_{n-1}(s)\}$ be an orthonormal frame of $c'_s(0)^\perp$ such that $e_i(0) = e_i(1), i = 1, \dots, n - 1$. Moreover, let $e_i(s, t)$ denote the parallel transport of $e_i(s)$ along c_s . For each $s \in [0, 1]$, let $A_s \in \text{SO}(n - 1)$ denote the holonomy map along c_s , i.e. $e_i(s, 2\pi) = A_s \cdot e_i(s, 0)$. Using the frame $\{e_1(s, t), \dots, e_{n-1}(s, t)\}$ we identify each space $c'_s(t)^\perp$ with $V = c'_0(0)^\perp \simeq \mathbb{R}^{n-1}$.

We can see A_s and $R_s(t) = R(\cdot, \dot{c}_s(t))\dot{c}_s(t)$ as maps

$$A : \mathbb{S}^1 \rightarrow \text{SO}(n - 1)$$

$$R : \mathbb{S}^1 \rightarrow \text{Map}([0, 2\pi], \text{Sym}^2(n - 1))$$

and by the previous section, we can associate to the data $(R_s, A_s)_{s \in \mathbb{S}^1}$ a Poincaré map $P : \mathbb{S}^1 \rightarrow \text{Sp}(n - 1, \omega)$ and a negative bundle $N \rightarrow \mathbb{S}^1$. In

this case the negative bundle is indeed a vector bundle, which coincides with the usual definition of negative bundle in Morse–Bott theory. In particular, the goal of this section is to prove that $\mathbf{N} \rightarrow \mathbb{S}^1$ is orientable.

2.5 The variation

As explained in Sect. 2.1 from now on we assume to have algebraic data (R_s, A_s) , $s \in \mathbb{S}^1$, relative to the geometric setup, such that $A : \mathbb{S}^1 \rightarrow \text{SO}(n - 1)$ is nullhomotopic. The goal of this section is to produce a deformation $(R_{(s,\tau)}, A_{(s,\tau)})$, $(s, \tau) \in \mathbb{S}^1 \times [0, 1]$, such that $(R_{(s,0)}, A_{(s,0)})$ is related to the data of our geometric setup, $(R_{(s,1)}, A_{(s,1)})$ has trivial negative bundle, and in such a way that we can somehow “keep track” of the negative bundle along the deformation. In this section we emphasize the properties of the variation near $\tau = 0$ and $\tau = 1$, while in the next sections we concentrate on the interior of $\mathbb{S}^1 \times [0, 1]$.

Before defining the deformation, we slightly modify the initial data. We replace the t -interval $[0, 2\pi]$ with a longer interval $[0, 6\pi]$, and define $A_{(s,0)} = A_s$, and

$$R_{(s,0)}(t) = \begin{cases} R_s(t) & t \in [0, 2\pi] \\ 4 \cdot Id & t \in (2\pi, 4\pi] \\ Id & t \in (4\pi, 6\pi] \end{cases}$$

In general, given real numbers $0 < L_1 \dots < L_k$ and curvature operators $R_i \in \text{Map}((L_{i-1}, L_i], \text{Sym}^2(n - 1))$, $i = 1, \dots, k$ we define the concatenation $R_1 \star \dots \star R_k \in \text{Map}((0, L_k], \text{Sym}^2(n - 1))$ to be the operator whose restriction to $(L_{i-1}, L_i]$ is R_i for every $i = 1, \dots, k$. In our case can then write

$$R_{(s,0)} = R_s \star (4 \text{ Id}) \star \text{Id}.$$

Notice that R does not need to be continuous in t , but only piecewise continuous.

Recall

Lemma 2.2 *The negative bundle of $(R_{(s,0)}, A_{(s,0)})$ is a vector bundle, and it is orientable if and only if the negative bundle of (R_s, A_s) is orientable.*

Proof For any $s \in \mathbb{S}^1$, the negative space $\mathbf{N}_{(s,0)}$ of $(R_{(s,0)}, A_{(s,0)})$ is the (finite dimensional) sum of the negative eigenspaces of the bilinear map $H_s : \mathfrak{X}_s \times \mathfrak{X}_s \rightarrow \mathbb{R}$ given by

$$H_s(X, Y) = \int_0^{6\pi} \langle X'(t), Y'(t) \rangle - \langle R_{(s,0)}(t) \cdot X(t), Y(t) \rangle dt$$

where $\mathfrak{X}_s = \{X : [0, 6\pi] \rightarrow V \mid X(6\pi) = A_{(s,0)} \cdot X(0)\}$.

The kernel of H_s is given by the periodic Jacobi fields on $[0, 6\pi]$ (i.e. vector fields $J : [0, 6\pi] \rightarrow V$ such that $J''(t) + R_{(s,0)}(t) \cdot J(t) = 0$, $J(6\pi) = A_{(s,0)} \cdot J(0)$ and $J'(6\pi) = A_{(s,0)} \cdot J'(0)$). Therefore the nullity of H_s corresponds to the multiplicity of the eigenvalue 1 of the Poincare map $P_{(s,0)}$ of $(R_{(s,0)}, A_{(s,0)})$ which, from the discussion above, in turn coincides with the Poincare map of (R_s, A_s) . In particular the dimension of the kernel of H_s is constant, and the same holds for the index of H_s (i.e., the sum of the negative space $N_{(s,0)}$).

Using Equation (1.4) of [2], it follows that for every $s \in \mathbb{S}^1$, the difference between the dimension of the negative space $N_{(s,0)}$ of $(R_{(s,0)}, A_{(s,0)})$ and the dimension of the negative space N_s of (R_s, A_s) , equals the difference between the number of conjugate points (i.e. zeroes of Jacobi fields with $J(0) = 0$ counted with multiplicity) on the interval $[0, 6\pi]$ and the number of conjugate points on the interval $[0, 2\pi]$. This equals the number of conjugate points on the interval $[2\pi, 6\pi]$, which is $6(n - 1)$ (recall that on $(2\pi, 4\pi]$ and $(4\pi, 6\pi]$ the curvature tensor is that of a sphere of constant curvature).

For any $s \in \mathbb{S}^1$, extend each vector field X in N_s (which is defined only on $[0, 2\pi]$) to a vector field \tilde{X} on $[0, 6\pi]$, by adding on the interval $(2\pi, 6\pi]$ a $R_{(s,0)}$ -Jacobi field for in such a way that \tilde{X} remains C^2 . Since all such Jacobi fields are periodic on $[2\pi, 6\pi]$, any such \tilde{X} is periodic and H_s is defined on these extensions. Furthermore, we define N' (independent of s) as the sum of all nonpositive eigenspace of the bilinear form H_s restricted to the space $\{X \in \mathfrak{X}_s \mid X_{[0,2\pi]} = 0\}$. Clearly the dimension of N' is the number of conjugate points on the interval $[2\pi, 6\pi]$, which again equals $6(n - 1)$.

The form H_s is nonnegative definite on the space $N_s \oplus N'$. Finally, the space $N_s \oplus N'$ does not contain any periodic Jacobi fields. Thus for every s , the orthogonal projection of $N_s \oplus N'$ to the negative space $N_{(s,0)}$ of H_s is an isomorphism.

Thus we see that the negative bundle $N_{(s,0)}$ of $(R_{(s,0)}, A_{(s,0)})$ is isomorphic to the sum of the negative bundle N_s , plus a trivial bundle N' . In particular, $N_{(s,0)}$ is orientable if and only if N_s is orientable. □

In particular, in the proof of Theorem B we can switch our attention to the orientability of the negative bundle of $(R_{(s,0)}, A_{(s,0)})$.

Once again, we assume now that $A_{(s,0)}$, as a loop in $SO(n - 1)$, is nullhomotopic. Notice that this is always the case if the manifold is spin (for example, when $M = \mathbb{S}^n$ or $\mathbb{H}\mathbb{P}^n$). In this case, let

$$H_{(s,\tau)} : \mathbb{S}^1 \times [0, 1] \rightarrow SO(n - 1)$$

be a homotopy with $H_{(s,\epsilon)} = A_{(s,0)}$ and $H_{(s,1)} = \text{Id}$.

Given a function $\varphi : [0, 1] \rightarrow \mathbb{R}$ such that

$$\begin{cases} \varphi(\tau) \equiv 0 \text{ around } \tau = 0 \\ \varphi(\tau) \equiv 1 \text{ around } \tau = 1 \\ \varphi'(\tau) \geq 0 \ \forall \tau \in (0, 1) \end{cases}$$

we now define the variation $(R_{(s,\tau)}, A_{(s,\tau)})$ by

$$\begin{aligned} R_{(s,\tau)} &= ((1 - \varphi(\tau))R_s + \varphi(\tau)\text{Id}) \star ((4 - 3\tau) \cdot \text{Id}) \star \tilde{R}_{(s,\tau)} \\ A_{(s,\tau)} &= H_{(\varphi(\tau),s)} \end{aligned} \tag{2.1}$$

The operator $\tilde{R}_{(s,\tau)}$ is a “generic” operator, sufficiently close to the constant map Id, which we will define in the next section, and whose goal is to make the Poincaré map generic. Before that, however, we will focus on the first two components of $R_{(s,\tau)}$. Namely, we consider now

$$\hat{R}_{(s,\tau)} = ((1 - \varphi(\tau))R_s + \varphi(\tau)\text{Id}) \star ((4 - 3\tau) \cdot \text{Id}) \tag{2.2}$$

Let us set the notation and call

$$\mathcal{C} = \mathbb{S}^1 \times [0, 1]$$

the space parametrised by s and τ . The variation $(\hat{R}_{(s,\tau)}, A_{(s,\tau)})$ can be seen as parametrised data $(\hat{R}_x, A_x)_{x \in \mathcal{C}}$, as defined in Sect. 2.3. In particular, we have an associated Poincaré map $\hat{P} : \mathcal{C} \rightarrow \text{Sp}(n - 1, \omega)$.

For reasons that will be clearer later, we want to have some control over the real eigenvalues of \hat{P} . At the moment we do not have such control, although in the following lemma we show that, sufficiently close to the boundary of \mathcal{C} , the map \hat{P} has no real eigenvalues.

Lemma 2.3 *There is a neighbourhood U of $\partial\mathcal{C}$ such that the Poincaré map \hat{P} of $(\hat{R}_x, A_x)_{x \in \mathcal{C}}$ does not have any real eigenvalue on $U \setminus \partial\mathcal{C}$.*

Proof It is enough to show that for every $s \in \mathbb{S}^1$, $P_{(s,\tau)}$ does not have eigenvalues in $(0, 1)$ in some neighbourhoods of $\tau = 0$ and $\tau = 1$. For $\tau = 0$, $\hat{P}_{(s,0)} = P_s$ is the Poincaré map of a closed geodesic in our geometric situation, and we know that $P_s^k = Id$ for some k . In particular the eigenvalues of P_s are roots of unity. Since the Poincaré map depends continuously on τ , its eigenvalues vary continuously as well and thus we only have to check that the eigenvalue 1 of P_s disappears for any $\tau > 0$ small.

Recall that we constructed the variation $(\hat{R}_{(s,\tau)}, A_{(s,\tau)})$ so that for small values of τ , $A_{(s,\tau)} = A_s$ and $\hat{R}_{(s,\tau)}$ only changes in the t -interval $[2\pi, 4\pi]$,

thus it is not hard to prove that, with respect to the canonical basis of $V \oplus V = \mathbb{R}^{2(n-1)}$, $\hat{P}_{(s,\tau)}$ can be written as

$$\hat{P}_{(s,\tau)} = \Delta A_s^{-1} \cdot (\cos \theta(\tau) \text{Id} + \sin \theta(\tau) J) \cdot \Delta A_s \cdot P_s$$

where $J = \begin{pmatrix} & -Id_V \\ Id_V & \end{pmatrix}$, $\Delta A_s = \text{diag}(A_s, A_s)$ and $\theta(\tau) = 2\pi \sqrt{4 - 3\tau}$.

Since we are only interested in the eigenvalues of $\hat{P}_{(s,\tau)}$ we can, up to conjugation of $\hat{P}_{(s,\tau)}$ and P_s with ΔA_s , reduce ourselves to the case

$$\hat{P}_{(s,\tau)} = (\cos \theta(\tau) \text{Id} + \sin \theta(\tau) J) \cdot P_s$$

We claim that $\hat{P}_{(s,\tau)}$ cannot have real eigenvalues for small τ . Suppose in fact that $\hat{P}_{(s,\tau)}$ has an eigenvector v_τ with real eigenvalue λ_τ , and v_τ tends to a fixed point of P_s for $\tau \rightarrow 0^+$. Around $\tau = 0$, the map $\tau \mapsto v_\tau$ is smooth. Differentiating the equation $\hat{P}_{(s,\tau)} v_\tau = \lambda_\tau v_\tau$ and taking the limit as $\tau \rightarrow 0$ we obtain

$$\theta'(0) \cdot J v_0 + P_s v'_0 = \lambda'_0 v_0 + v'_0.$$

By evaluating the two sides of the equation using the symplectic form $\omega(\cdot, v_0)$ we get

$$\theta'(0) \cdot \omega(J v_0, v_0) + \omega(P_s v'_0, v_0) = \omega(v'_0, v_0).$$

Since v_0 is a fixed point for P_s , we get $\omega(P_s v'_0, v_0) = \omega(v'_0, v_0)$ and the equation becomes $\theta'(0) \cdot \omega(J v_0, v_0) = 0$, which is not possible since $\theta'(0) \neq 0$ and $\omega(J v_0, v_0) \neq 0$.

The case around $\tau = 1$ can be handled in a similar fashion. □

The goal of the next section is to produce the last piece of curvature operator

$$\tilde{R} : \mathcal{C} \rightarrow \text{Map}((4\pi, 6\pi], \text{Sym}^2(n - 1)),$$

in the variation $R_{(s,\tau)}$ as in Eq. 2.1. The curvature operator will be arbitrarily close to the constant map $\equiv \text{Id}$ (and in fact equal to the constant map in a neighbourhood of $\partial\mathcal{C}$), such that the Poincaré map of $(R_x = \hat{R}_x \star \tilde{R}_x, A_x)_{x \in \mathcal{C}}$, as defined in Eq. 2.1, is controlled.

2.6 Making the Poincaré map generic

By [17, Lemma 3.8] there are curvature operators

$$R_0, R_1, \dots, R_h : [0, 2\pi] \rightarrow \text{Sym}^2(n - 1),$$

with $h = \dim \text{Sp}(n - 1, \omega)$, such that the map

$$\begin{aligned} (-1, 1)^h &\longrightarrow \text{Sp}(n - 1, \mathbb{R}) \\ (a_1, \dots, a_h) &\longmapsto \text{Poincaré map of } (R_0 + a_1 R_1 + \dots + a_h R_h, \text{Id}) \end{aligned}$$

is a diffeomorphism between $(-1, 1)^h$ and a neighbourhood U of the identity in $\text{Sp}(n - 1, \omega)$. In particular, given a map $\tilde{P} : \mathcal{C} \rightarrow U$ there is a map

$$\tilde{R} : \mathcal{C} \rightarrow \text{Map}([0, 2\pi], \text{Sym}^2(n - 1))$$

such that \tilde{P} is the Poincaré map of (\tilde{R}, Id) .

It is easy to see in general that that if P_1, P_2, P_3 are the Poincaré maps of $(R_1, A_1), (R_2, A_2)$ and $(R_1 \star R_2, A_3)$ respectively, then

$$(\Delta A_1 \cdot P_1) \cdot (\Delta A_2 \cdot P_2) = \Delta A_3 \cdot P_3$$

where $\Delta A = \text{diag}(A, A) \in \text{Sp}(n - 1, \omega)$.

In our case, given \hat{P}_x the Poincaré map of the pair $(\hat{R}_x, A_x)_{x \in \mathcal{C}}$ defined in Eq. (2.2), then the Poincaré map of $(R_x \star \tilde{R}_x, A_x)$ is

$$\Delta A_x^{-1} \cdot \tilde{P}_x \cdot \Delta A_x \cdot \hat{P}_x$$

In particular, for any map $P : \mathcal{C} \rightarrow \text{Sp}(n - 1, \omega)$ sufficiently close to \hat{P} , such that $P|_{\partial \mathcal{C}} = \hat{P}|_{\partial \mathcal{C}}$, we can find an operator $\tilde{R} : \mathcal{C} \rightarrow \text{Map}([0, 2\pi], \text{Sym}^2(n - 1))$ sufficiently close to the identity, such that $\tilde{R}|_{\partial \mathcal{C}} \equiv \text{Id}$ and $(\hat{R}_x \star \tilde{R}_x, A_x)_{x \in \mathcal{C}}$ has Poincaré map P .

The goal of this section is to prove that for a “generic” choice of \tilde{R} , the Poincaré map $P : \mathcal{C} \rightarrow \text{Sp}(n - 1, \omega)$ of $(\hat{R}_x \star \tilde{R}_x, A_x)_{x \in \mathcal{C}}$ has the following properties:

- For every $x \in \mathcal{C}$, the positive real eigenvalues of P_x have geometric multiplicity 1.
- The set of points $x \in \mathcal{C}$, whose Poincaré map P_x has eigenvalue 1, is a smooth subvariety.
- The function $\bar{\chi} : \mathcal{C} \times \mathbb{R}_+ \rightarrow \mathbb{R}$, $\bar{\chi}(x, \lambda) = \det(P_x - \lambda \text{Id})$ does not have critical value 0 in the interior of \mathcal{C} .

To this end, consider the following subsets of $\mathbf{Sp}(n - 1, \omega) \times \mathbb{R}_+$:

$$\begin{aligned} \mathcal{G} &= \{(Q, \lambda) \in \mathbf{Sp}(n - 1, \omega) \times \mathbb{R}_+ \mid \dim \ker(Q - \lambda \text{Id}) \leq 1\} \\ \mathcal{G}_1 &= \{(Q, \lambda) \in \mathbf{Sp}(n - 1, \omega) \times \mathbb{R}_+ \mid \dim \ker(Q - \lambda \text{Id}) = 1\} \\ \mathcal{G}_0 &= \{(Q, 1) \in \mathcal{G}_1\} = \mathcal{G}_1 \cap (\mathbf{Sp}(n - 1, \omega) \times \{1\}) \end{aligned}$$

Clearly \mathcal{G} is open in $\mathbf{Sp}(n - 1, \omega) \times \mathbb{R}_+$. We show in Proposition A.4 and Lemma A.6 that \mathcal{G}_1 is a smooth hypersurface of \mathcal{G} and \mathcal{G}_0 is a smooth hypersurface of \mathcal{G}_1 .

Lemma 2.4 *In any neighbourhood of $\hat{P} : \mathcal{C} \rightarrow \mathbf{Sp}(n - 1, \omega)$ there exists a $P : \mathcal{C} \rightarrow \mathbf{Sp}(n - 1, \omega)$ such that the image of*

$$(P \times \text{Id})|_{\text{int}(\mathcal{C}) \times \mathbb{R}_+} : \text{int}(\mathcal{C}) \times \mathbb{R}_+ \rightarrow \mathbf{Sp}(n - 1, \omega) \times \mathbb{R}_+$$

(int(C) denotes the interior of C) is contained in \mathcal{G} , and it intersects \mathcal{G}_0 and \mathcal{G}_1 transversely.

Proof Let $\mathbf{Sp}_1(n - 1, \omega)$ denote the set of symplectic matrices whose positive real eigenvalues have geometric multiplicity 1, and let $\mathbf{Sp}_0(n - 1, \omega) \subseteq \mathbf{Sp}_1(n - 1, \omega)$ denote the subset of matrices with eigenvalue 1. By Proposition A.3 $\mathbf{Sp}_1(n - 1, \omega)$ is open, and its complement has codimension ≥ 3 . Moreover, $\mathbf{Sp}_0(n - 1, \omega)$ is a smooth algebraic subvariety, and it has codimension at least 1 in $\mathbf{Sp}_1(n - 1, \omega)$.

Since $\dim \mathcal{C} = 2$ it is possible to find a $P : \mathcal{C} \rightarrow \mathbf{Sp}_1(n - 1, \omega)$, close to \hat{P} and with $P|_{\partial \mathcal{C}} = \hat{P}|_{\partial \mathcal{C}}$, such that $P|_{\text{int}(\mathcal{C})}$ intersects $\mathbf{Sp}_0(n - 1, \omega)$ transversely. Moreover, we can do it in such a way that $P|_{\text{int}(\mathcal{C})}$ is an embedding (because $n \geq 3$) and $P(\text{int}(\mathcal{C}))$ is disjoint from $P(\partial \mathcal{C})$. By construction, the image of $(P \times \text{Id})|_{\text{int}(\mathcal{C}) \times \mathbb{R}_+} : \text{int}(\mathcal{C}) \times \mathbb{R}_+ \rightarrow \mathbf{Sp}(n - 1, \omega) \times \mathbb{R}_+$ lies in \mathcal{G} and it intersects \mathcal{G}_0 transversely.

Let $\chi : \mathcal{G} \rightarrow \mathbb{R}$ denote the function $\chi(Q, \lambda) = \det(Q - \lambda \text{Id})$. This function is a submersion by Proposition A.4. Clearly $\mathcal{G}_1 = \chi^{-1}(0)$ and $P \times \text{Id}$ fails to meet \mathcal{G}_0 transversely if and only if $(P \times \text{Id})^* \chi : \mathcal{C} \times \mathbb{R}_+ \rightarrow \mathbb{R}$ has some critical point of value 0. Suppose then that $(P \times \text{Id})^* \chi$ has critical value 0. By Proposition A.5, there exists a vector field V in $\mathbf{Sp}_1(n - 1, \omega)$ such that $d_{(Q,\lambda)} \chi(V) > 0$ for every (Q, λ) in \mathcal{G} . We fix a nonnegative function f supported on a neighbourhood of the image of P , which is 0 on $P(\partial \mathcal{C})$ and let Φ_t be the flow of fV for some time t . If t is small enough, then $(P \circ \Phi_t) \times \text{Id}$ does not have critical point 0 and we obtain the result, up to replacing P with $P \circ \Phi_t$. \square

We will from now on consider a variation

$$\left(R_x = \hat{R}_x \star \tilde{R}_x, A_x \right)_{x \in \mathcal{C}}$$

where \hat{R} , A are defined in Eq. (2.2), and \tilde{R} is defined in such a way that the Poincaré map P of $(R_x, A_x)_{x \in \mathcal{C}}$ satisfies the conditions of Lemma 2.4.

2.7 The modified negative bundle

Recall from Sect. 2.3 that to the data $(R_x, A_x)_{x \in \mathcal{C}}$ there is an associated negative bundle $\mathbf{N} \rightarrow \mathcal{C}$, which has the structure of a fiberwise vector space, but it is not a vector bundle in general since the dimension of the fibers might (and does, in general) change from point to point. Nevertheless, whenever the restriction of \mathbf{N} to a subset of $U \subseteq \mathcal{C}$ has constant rank then $\mathbf{N}|_U \rightarrow U$ is a vector bundle in the usual sense. We make this precise with the following definition.

Definition 2.5 A pseudo vector bundle consists of topological spaces E, B , a continuous surjection $\pi : E \rightarrow B$, a section $0 : B \rightarrow E$ (the zero section), a map $a : E \times_B E \rightarrow E$ (addition) over B and, for every $\lambda \in \mathbb{R}$, an operation $\lambda \cdot : E \rightarrow E$ (scalar multiplication) over B , satisfying the usual axioms of vector bundles. Moreover, we require that there exists an open dense set B_{reg} of B (the regular part) and an open cover $\{U_i\}_{i \in I}$ of B_{reg} , such that $E|_{U_i} \simeq U_i \times \mathbb{R}^{n_i}$ (where n_i might depend on U_i).

By abuse of language, we will sometimes denote a pseudo vector bundle by $\pi : E \rightarrow B$, or simply by E when B and π are understood. It is clear that \mathbf{N} admits the maps in the definition of pseudo vector bundle.

The following lemma shows that there exists an open dense set $\mathcal{C}_{reg} \subseteq \mathcal{C}$ over which \mathbf{N} is a vector bundle, thus proving that $\mathbf{N} \rightarrow \mathcal{C}$ is a pseudo vector bundle.

Lemma 2.6 Let $x \in \mathcal{C}$. The following hold:

- (1) The index form H_x has nontrivial kernel if and only if the Poincaré map P_x has a fixed vector. Moreover, there is an isomorphism between the eigenspace of P with eigenvalue 1, $E_1(P_x)$, and $\ker(H_x)$.
- (2) If $\ker H_x = 0$ for some $x \in \mathcal{C}$, then $\mathbf{N} \rightarrow \mathcal{C}$ is a vector bundle around x . It thus makes sense to define $\mathcal{C}_{reg} = \{x \in \mathcal{C} \mid \ker H_x = 0\}$.
- (3) \mathcal{C}_{reg} is open and dense in \mathcal{C} .

Proof (1) If $X \in \ker H_x$ then by, the first variation formula for the energy function, X is a periodic Jacobi field and

$$P_x(X(0), X'(0)) = (X(6\pi), X'(6\pi)) = (X(0), X'(0)).$$

In particular, there is an isomorphism $\bar{\phi}_x : \ker H_x \rightarrow E_1(P_x)$, $\bar{\phi}(X) = (X(0), X'(0))$.

(2) Since H_x is a symmetric map it has real eigenvalues and, since they change continuously with x , it follows that the number of negative eigenvalues

of H_x remains constant if H_x has empty kernel. Therefore, if $\ker H_x = 0$ then \mathbf{N} is a vector bundle around x .

(3) By the previous points, the complement of \mathcal{C}_{reg} consists of the points $x \in \mathcal{C}$ such that P_x has eigenvalue 1, which by construction is closed and has codimension at least 1 in \mathcal{C} . \square

Definition 2.7 A pseudo vector bundle $\pi : E \rightarrow B$ is *locally orientable* if there exists a \mathbb{Z}_2 cover $O \rightarrow B$ such that, for every component B_i of B_{reg} , the restriction $O|_{B_i}$ is isomorphic to the orientation bundle of $E|_{B_i} \rightarrow B_i$.

By the construction of the variation (R_x, A_x) there is some $\epsilon > 0$ small enough, such that the restriction of \mathbf{N} to the curves

$$\gamma_0(s) = (\epsilon, s) \quad \gamma_1(s) = (1 - \epsilon, s)$$

has constant rank and therefore $\mathbf{N}|_{\gamma_i}, i = 0, 1$, is a vector bundle. If we could prove that \mathbf{N} is locally orientable, it would follow that $\mathbf{N}|_{\gamma_0}$ is orientable if and only if $\mathbf{N}|_{\gamma_1}$ is orientable. Since $\mathbf{N}|_{\gamma_0}$ is isomorphic to the sum of the geometric negative bundle with a trivial bundle, while $\mathbf{N}|_{\gamma_1}$ is trivial, this would prove the orientability of the geometric negative bundle.

With this goal in mind, we introduce a second pseudo-vector bundle $\mathcal{E} \rightarrow \mathcal{C}$, $\mathcal{E} \subseteq \mathcal{C} \times (V \oplus V)$, whose fiber at $x \in \mathcal{C}$ is the vector space consisting of the eigenspaces of P_x , with eigenvalues in $(0, 1)$. It makes sense to define a fiberwise direct sum

$$\mathbf{N} \oplus \mathcal{E} \rightarrow \mathcal{C}$$

which we call *modified negative bundle* and this is the bundle that we will consider from now on.

By Lemma 2.3, $\mathcal{E} = 0$ around $\partial\mathcal{C}$ and thus we are not changing anything along γ_1 and γ_2 . In particular, we can prove the local orientability of $\mathbf{N} \oplus \mathcal{E}$ instead of \mathbf{N} , and we will still obtain that the geometric negative bundle is orientable. The next two sections are devoted to proving the local orientability of $\mathbf{N} \oplus \mathcal{E}$.

2.8 Local orientability of pseudo vector bundles

In this section we show some classes of pseudo vector bundles that are always locally orientable.

We start by remarking that local orientability of a pseudo vector bundle $\pi : E \rightarrow B$ can be proved by showing that there is an open cover $\{U_i\}$ of B such that $E|_{U_i}$ is locally orientable, and such that the corresponding orientation bundles $O(U_i)$ agree on intersections, in the sense that for every $U_{ij} = U_i \cap U_j$

there are bundle maps $O(U_i)|_{U_{ij}} \rightarrow O(U_j)|_{U_{ij}}$ satisfying the usual cocycle conditions.

Let M, N be manifolds of the same dimension, let $E \rightarrow N$ be a vector bundle, and let $f : N \rightarrow M$ a closed map with finite fibers. For each $p \in M$, let $F_p = \bigoplus_{q \in f^{-1}(p)} E_q$. Define f_*E as the space $f_*E = \coprod_{p \in M} F_p$ with projection $f_*\pi : F \rightarrow M$ sending F_p to p . By Sard’s theorem, the set $M_{reg} \subseteq M$ of regular points for f is open and dense. Around each point $x \in M_{reg}$ there is a neighbourhood U with $f^{-1}(U) = \coprod U_i$, such that $f|_{U_i} : U_i \rightarrow U$ is a diffeomorphism, and there is a canonical bijection $\phi_U : f_*E|_U \rightarrow \bigoplus_i E|_{U_i}$. The space f_*E , endowed with the roughest topology that makes the maps $f_*\pi$ and ϕ_U continuous, is clearly a pseudo vector bundle, which we call the *push-forward* of E via f .

Lemma 2.8 *Given a pseudo vector bundle $E \rightarrow M$ and a vector bundle $F \rightarrow M$, $E \oplus F \rightarrow M$ is locally orientable if and only if $E \rightarrow M$ is locally orientable.*

Proof If $O_E \rightarrow M, O_F \rightarrow M$ are the orientation bundles of E, F respectively, then the orientation bundle of $E \oplus F$ exists and it is given by $O_{E \oplus F} := (O_E \times O_F)/\mathbb{Z}_2$, where \mathbb{Z}_2 acts diagonally. Therefore, $E \oplus F$ is locally orientable. On the other hand, if $O_{E \oplus F} \rightarrow M, O_F \rightarrow M$ are the orientation bundles of $E \oplus F$ and E respectively, then $O_{E \oplus F} \otimes O_F$ is the orientation bundle of E . □

Proposition 2.9 *Let M, N be manifolds without boundary of the same dimension, and let $f : N \rightarrow M$ be a closed map with finite fibers. Then for every vector bundle $E \rightarrow N$ the push forward $f_*E \rightarrow M$ is locally orientable.*

Proof We will define a pre-sheaf on \mathcal{O} on N such that $\mathcal{O}(U) = \mathbb{Z}_2$ for any U small enough, and whose restriction to any open set in N_{reg} is isomorphic to the sheaf of local sections of the orientation bundle. Then the étale space $O = \coprod_{p \in N} O_p, O_p = \varinjlim_{U \ni p} \mathcal{O}(U)$, together with the projection $O \rightarrow N$ sending O_p to p , will be the \mathbb{Z}_2 -cover we need to prove the result.

For any $p \in M, f^{-1}(p)$ is a discrete set $\{q_1, \dots, q_r\}$. Given a small neighbourhood U of $p, f^{-1}(U)$ is a disjoint union of neighbourhoods U_i of $q_i, i = 1, \dots, r$. Defining $f_i = f|_{U_i} : U_i \rightarrow U$, the preimage of f_i has almost everywhere constant parity, and we define $\epsilon_i = 0$ if given parity is even, and $\epsilon_i = 1$ if it is odd.

Let $b = (b_1, \dots, b_r)$ be an r -tuple where each b_i is a local basis of sections of $E|_{U_i}$, and let \mathcal{B} denote the set of such r -tuples. Given two r -tuples b, b' , there exists an r -tuple (J_1, \dots, J_r) where each J_i is a local section of $GL(E|_{U_i})$ taking b_i to b'_i . We can finally define $\mathcal{O}(U)$ to be the set of maps $\theta : \mathcal{B} \rightarrow \{\pm 1\}$

such that for any b, b' in \mathcal{B} ,

$$\theta(b) = \theta(b') \cdot \prod_{i=1}^r \operatorname{sgn}(\det J_i)^{\epsilon_i}.$$

It is easy to see that if U is a small neighbourhood of a point in M_{reg} then $f^{-1}(U)$ is a disjoint union of open sets U_1, \dots, U_r such that $f_i : U_i \rightarrow U$ is a homeomorphism. In particular $\epsilon_i = 1$ for all $i = 1, \dots, r$, and $\mathcal{O}(U)$ coincides with the set of sections of the orientation bundle of $f_*E|_U$. \square

2.9 Local orientability of $\mathbf{N} \oplus \mathcal{E}$

In this section we prove that the modified vector bundle is locally orientable, by showing that it locally takes the form $E_1 \oplus f_*E_2$ for some vector bundle $E_1 \rightarrow \mathcal{C}$ and some push forward $f_*E_2 \rightarrow \mathcal{C}$ with respect to some function $f : \bar{\mathcal{C}} \rightarrow \mathcal{C}$. In order to do this, we must first construct the space $\bar{\mathcal{C}}$, which will be defined by gluing two spaces \mathcal{C}_P and \mathcal{C}_H along their (diffeomorphic) boundaries.

Consider the subset $\mathcal{C}_P \subseteq \mathcal{C} \times (0, 1]$ defined as

$$\mathcal{C}_P = \{(x, \lambda) \in \mathcal{C} \times (0, 1] \mid \lambda \text{ is an eigenvalue of } P_x\}$$

and let $f_P : \mathcal{C}_P \rightarrow \mathcal{C}$ denote the obvious projection.

Proposition 2.10 *The following hold:*

- (1) *The set \mathcal{C}_P is a submanifold with boundary $\partial \mathcal{C}_P = \mathcal{C}_P \cap \mathcal{C} \times \{1\}$.*
- (2) *There is a line bundle $E_P \rightarrow \mathcal{C}_P$, such that $\mathcal{E} = f_{P*}E_P|_{\operatorname{Int}(\mathcal{C}_P)}$, where $\operatorname{Int}(\mathcal{C}_P)$ denotes the interior of \mathcal{C}_P .*

Proof (1) As in Sect. 2.6, let $\mathcal{G}_1 \subseteq \operatorname{Sp}(n - 1, \mathbb{R}) \times \mathbb{R}_+$ denote the subset of couples (Q, λ) where $\dim \ker(Q - \lambda \operatorname{Id}) = 1$, and let \mathcal{G}_0 denote the subset of couples $(Q, 1)$ in \mathcal{G}_1 . By the construction of $(R_x, A_x)_{x \in \mathcal{C}}$ (cf. Lemma 2.4) the image of

$$P \times \operatorname{Id} : \mathcal{C} \times \mathbb{R}_+ \longrightarrow \operatorname{Sp}(n - 1, \mathbb{R}) \times \mathbb{R}_+$$

intersects \mathcal{G}_1 and \mathcal{G}_0 transversely. In particular $(P \times \operatorname{Id})^{-1}(\mathcal{G}_1)$ is a smooth hypersurface, $(P \times \operatorname{Id})^{-1}(\mathcal{G}_0)$ is a smooth hypersurface in $(P \times \operatorname{Id})^{-1}(\mathcal{G}_1)$ dividing it in two components, and it is easy to see that \mathcal{C}_P is one such component. In particular, \mathcal{C}_P is a smooth manifold with boundary $\partial \mathcal{C}_P = (P \times \operatorname{Id})^{-1}(\mathcal{G}_0)$.

(2) By construction, for every $x \in \mathcal{C}$, every positive real eigenvalue of P_x has geometric multiplicity 1 (cf. Lemma 2.4). In particular, the space $E \subseteq$

$\mathcal{C}_P \times (V \oplus V)$ given by

$$E_P = \{((x, \lambda), v) \in \mathcal{C}_P \times (V \oplus V) \mid P_x v = \lambda v\}$$

with the obvious projection $E_P \rightarrow \mathcal{C}_P$ is a line bundle, and by definition $\mathcal{E} = f_{P*}(E_P)|_{\text{Int}(\mathcal{C}_P)}$. □

By Lemma 2.6, the set

$$\Sigma = f_P(\partial \mathcal{C}_P)$$

coincides with the set of points x such that $\ker H_x \neq 0$ and, therefore, the set of points around which \mathbf{N} is not a vector bundle.

Since $\ker H_x$ is isomorphic to the eigenspace of P_x of eigenvalue 1, and this is 1 dimensional, it follows that $\dim \ker H_x = 1$ for every $x \in \Sigma$. Since Σ is compact, it is possible to find a neighbourhood U_Σ of Σ and some $\epsilon > 0$ such that H_x has at most one eigenvalue in $(-\epsilon, \epsilon)$ for every x in U_Σ . Consider now

$$\psi : U_\Sigma \times (-\epsilon, \epsilon) \rightarrow \mathbb{R} \quad \psi(x, \lambda) = \det(H_x - \lambda I)$$

and define $\mathcal{C}_H = (U_\Sigma \times (-\epsilon, 0]) \cap \psi^{-1}(0)$. Equivalently, \mathcal{C}_H is the set of pairs (x, λ) such that λ is a nonpositive eigenvalue of H_x . Let $f_H : \mathcal{C}_H \rightarrow U_\Sigma$ denote the obvious projection, and let $E_H \rightarrow \mathcal{C}_H$ be the line bundle whose fiber at (x, λ) is the (one dimensional by definition) eigenspace of H_x with eigenvalue λ .

Proposition 2.11 *The following hold:*

- (1) *There is a diffeomorphism $\phi : \partial \mathcal{C}_P \rightarrow \mathcal{C}_H \cap (\mathcal{C} \times \{0\})$ such that $f_H \circ \phi = f_P$.*
- (2) *\mathcal{C}_H is a smooth submanifold, with boundary $\partial \mathcal{C}_H = \mathcal{C}_H \cap \mathcal{C} \times \{0\}$.*
- (3) *There is an isomorphism of line bundles*

$$\bar{\phi} : E_P|_{\partial \mathcal{C}_P} \longrightarrow E_H|_{\partial \mathcal{C}_H}$$

over ϕ .

Proof (1) The set $\mathcal{C}_H \cap (\mathcal{C} \times \{0\})$ consists of points $(x, 0)$ where $x \in U$ and $\ker H_x \neq 0$. By Lemma 2.6 the map $\phi : \partial \mathcal{C}_P \rightarrow \mathcal{C}_H \cap (\mathcal{C} \times \{0\})$ sending $(x, 1)$ to $(x, 0)$ is a diffeomorphism.

(2) We first prove that 0 is a regular value of $\psi : U_\Sigma \times (-\epsilon, \epsilon) \rightarrow \mathbb{R}$. Given $(x, \lambda) \in \psi^{-1}(0)$, λ is an eigenvalue of H_x . By construction λ has geometric multiplicity 1 and, since H_x is symmetric, it has algebraic multiplicity 1 as

well. Therefore, by letting $\frac{\partial}{\partial t}$ denote the vector at (x, λ) tangent to $(-\epsilon, \epsilon)$, we have

$$d_{(x,\lambda)}\psi\left(\frac{\partial}{\partial t}\right) = \frac{d}{dt}\Big|_{t=\lambda} \det(H_x - tI) \neq 0.$$

Then $\psi^{-1}(0)$ is a smooth submanifold of $U \times (-\epsilon, \epsilon)$. By the previous point, the subset $(\mathcal{C} \times \{0\}) \cap \psi^{-1}(0) = \mathcal{C}_H \cap (\mathcal{C} \times \{0\})$ is a smooth submanifold of $\psi^{-1}(0)$ which divides it into two components, one of which is \mathcal{C}_H .

(3) For any point $(x, 1) \in \partial \mathcal{C}_P$, we have a map $\bar{\phi}_{(x,1)} : E_P|_{(x,1)} \rightarrow E_H|_{(x,0)}$ sending the fixed point $(v, w) \in E_1(P_x) = E_P|_{(x,1)}$ of P_x to the unique Jacobi field $J \in \ker H_x = E_H|_{(x,0)}$ with $J(0) = v, J'(0) = w$. □

By Proposition 2.11, it makes sense to define:

- The manifold $\bar{\mathcal{C}} = \mathcal{C}_P \sqcup_{\phi} \mathcal{C}_H$, without boundary.
- The map $f = f_P \sqcup_{\phi} f_E : \bar{\mathcal{C}} \rightarrow U$. It is easy to check that this map is closed.
- The line bundle $E = E_P \sqcup_{\bar{\phi}} E_H \rightarrow \bar{\mathcal{C}}$.

Proposition 2.12 *The modified negative bundle $\mathbf{N} \oplus \mathcal{E}$ is locally orientable.*

Proof It is enough to argue locally around a point $p \in \mathcal{C}$. If p does not belong to Σ , then we can find a neighbourhood U disjoint from Σ . Therefore, $\mathbf{N}|_U$ is a vector bundle, and $f_P : f_P^{-1}(U) \rightarrow U$ is a proper map with finite fibers, with $\mathcal{E}|_U = f_*E_P$. By Proposition 2.9 and Lemma 2.8, $(\mathbf{N} \oplus \mathcal{E})|_U$ is locally orientable.

If p lies in Σ , we can take a neighbourhood U of p that is contained in U_{Σ} . Let $F \rightarrow U$ denote the subbundle of \mathbf{N} whose fiber at $x \in U$ is the sum of the negative eigenspaces of H_x with eigenvalue $\leq -\epsilon$. By the construction of U , F is a vector bundle and

$$(\mathbf{N} \oplus \mathcal{E})|_U = F \oplus f_*E|_{f^{-1}(U)}.$$

Again by Proposition 2.9 and Lemma 2.8, $\mathbf{N} \oplus \mathcal{E}|_U$ is locally orientable. □

2.10 If A_s is not nullhomotopic

We just finished proving that given data $(R_s, A_s)_{s \in \mathbb{S}^1}$ with $A : \mathbb{S}^1 \rightarrow \text{SO}(n - 1)$ nullhomotopic, the corresponding negative bundle is orientable. We now complete the proof of Theorem B by showing that \mathbf{N} is not orientable if $A : \mathbb{S}^1 \rightarrow \text{SO}(n - 1)$ is not nullhomotopic. In fact, in this case we can replace the

data V, R_s, A_s with

$$\begin{aligned} V^+ &= V \oplus \mathbb{R}^2 = \mathbb{R}^{n+1} \\ R_s^+ &= \text{diag}(R_s, Q_s) \\ A_s^+ &= \text{diag}(A_s, \text{Rot}_{-s}) \end{aligned}$$

where $\text{Rot}_s \in \text{SO}(2)$ denotes rotation by s and $Q_s \equiv Q : [0, 2\pi] \rightarrow \text{Sym}^2(2)$ is an operator (independent of $s \in \mathbb{S}^1$) with Poincaré map equal to

$$B = \left(\begin{array}{c|c} & \begin{array}{c} -1 \\ 1 \end{array} \\ \hline \begin{array}{c} 1 \\ -1 \end{array} & \end{array} \right)$$

One can check that the Poincaré map P_s of $(Q_s, \text{Rot}_{-s})_{s \in \mathbb{S}^1}$ is given by the product $\text{diag}(\text{Rot}_s, \text{Rot}_s) \cdot B$. This family of Poincaré maps has constant eigenvalues $\pm i$, and in particular no eigenvalue 1. By Remark 2.1, the index of $(Q, \text{Rot}_{-s})_{s \in \mathbb{S}^1}$ remains constant. Therefore the negative bundle $\mathbf{N}' \rightarrow \mathbb{S}^1$ of the data $(Q, \text{Rot}_{-s})_{s \in \mathbb{S}^1}$ is indeed a vector bundle and so is the negative bundle $\mathbf{N}^+ = \mathbf{N} \oplus \mathbf{N}'$ of the data $(R_s^+, A_s^+)_{s \in \mathbb{S}^1}$.

Since A_s^+ now defines a contractible loop in $\text{SO}(n + 1)$, by the result above \mathbf{N}^+ is orientable. However, we claim that \mathbf{N}' is not orientable, from which it follows that \mathbf{N} is not orientable either. In fact, for every s let U_s denote the space of vector fields

$$U_s = \left\{ J \mid \exists t_0 \in (0, 2\pi) \text{ s.t. } J(0) = J(t_0) = 0, J|_{[0, t_0]} \text{ } Q_s\text{-Jacobi, } J|_{[t_0, 2\pi]} \equiv 0 \right\}$$

In this case, since Q_s does not depend on $s \in \mathbb{S}^1$, neither does U_s , which then describes a trivial bundle over \mathbb{S}^1 . Moreover, let W_s denote a maximal space of Q_s -Jacobi fields $J : [0, 2\pi] \rightarrow \mathbb{R}^2$ such that

$$J(0) = \text{Rot}_s(J(2\pi)), \langle \text{Rot}_s(J'(2\pi)) - J'(0), J(0) \rangle < 0.$$

Using Equations (1.4') of [2] it is easy to show that \mathbf{N}'_s is isomorphic to $U \oplus W_s$. In this case, the space W_s can be chosen to be the (1-dimensional) space of spanned by the Jacobi field with initial conditions

$$\begin{aligned} J(0) &= (\sin(-s/2), \cos(-s/2)) \\ J'(0) &= (\cos(-s/2), \sin(-s/2)). \end{aligned}$$

In particular W_s forms a non-orientable vector bundle and, since U_s is a trivial vector bundle, $\mathbf{N}' \simeq U \oplus W$ is not orientable.

3 Free loop spaces of cohomology CROSSes

The goal of this section is to prove that for every manifold M with the integral cohomology ring of a CROSS,

$$H_{\mathbb{S}^1}^i(\Lambda M, M; R) = 0 \quad \forall i \equiv \dim M \pmod{2},$$

where $R = \mathcal{P}^{-1}\mathbb{Z}$ is an extension \mathbb{Z} where we allow to divide by the primes in a finite set \mathcal{P} . We prove this by first showing it for CROSSes, using the energy function $E : \Lambda M \rightarrow \mathbb{R}$ of their canonical metric as an \mathbb{S}^1 -equivariant Morse–Bott function, and then proving the result in general.

Most computations here are certainly not new. The integral homology of the tangent bundle T^1M and the homology $H_*(\Lambda M, \mathbb{Z})$ of CROSSes was computed by Ziller in [18]. In [8], Hingston used Equivariant Morse theory to compute the rational, relative, \mathbb{S}^1 -equivariant cohomology $H_{\mathbb{S}^1}^*(\Lambda M, M, \mathbb{Q})$ for CROSSes. Moreover, the authors were informed by a referee that Schwarz computed in [12] the rational, \mathbb{S}^1 -equivariant cohomology of $\Lambda \mathbb{S}^n$ (the article is in Russian, but an account of this result can be found for example in [13], Theorem 2).

Since the canonical metrics on CROSSes are Besse metrics and all geodesics have the same period π , the critical energies are $e_k = \frac{1}{2}k^2\pi^2$ for $k \geq 0$.

The critical set C_0 consists of constant curves, and thus it is homeomorphic to M . The set C_1 consists of all the simple geodesics in M and, for $k > 1$, the set $C^k = E^{-1}(e_k)$ consists of the k -iterates of the geodesics in C^1 . Since all the geodesics of M are closed of the same length, for every $k \geq 1$ and every unit tangent vector v the geodesic $c_v = \exp(ktv)$, $t \in [0, 2\pi]$ is a geodesic in C^k , and the map sending v to c_v is a homeomorphism $T^1M \rightarrow C^k$.

Let i_k denote the index of C^k . For every $a \in \mathbb{R}$ we let $\Lambda^k \subseteq \Lambda M$ denote

$$\Lambda^k = E^{-1}([0, e_k + \epsilon)) \simeq E^{-1}([0, e_{k+1} - \epsilon))$$

for some ϵ sufficiently small.

Since the negative bundle of every critical submanifold is orientable, we have

$$H_{\mathbb{S}^1}^*(\Lambda^0; \mathbb{Z}) = H_{\mathbb{S}^1}^*(M; \mathbb{Z}), \quad (3.1)$$

$$H_{\mathbb{S}^1}^*(\Lambda^k, \Lambda^{k-1}; \mathbb{Z}) = H_{\mathbb{S}^1}^{*-i_k}(C^k; \mathbb{Z}). \quad (3.2)$$

The action of \mathbb{S}^1 on ΛM reduces to a trivial action on C^0 , a free action on C^1 and, for $k > 1$, an almost free action on C^k with ineffective kernel \mathbb{Z}_k .

Moreover, if $k > 1$ then the action of $\mathbb{S}^1/\mathbb{Z}_k$ on C^k becomes free. In particular,

$$H_{\mathbb{S}^1}^*(C^0; \mathbb{Z}) = H^*(M \times B\mathbb{S}^1; \mathbb{Z}) \tag{3.3}$$

$$H_{\mathbb{S}^1}^*(C^1; \mathbb{Z}) = H^*(T^1M/\mathbb{S}^1; \mathbb{Z}) \tag{3.4}$$

$$H_{\mathbb{S}^1}^*(C^k; \mathbb{Z}) = H^*(T^1M/\mathbb{S}^1 \times B\mathbb{Z}_k; \mathbb{Z}), \quad \forall k > 1. \tag{3.5}$$

Proposition 3.1 *Given M a CROSS, then the group $H^*(T^1M/\mathbb{S}^1; \mathbb{Z})$ is isomorphic to $H^*(N; \mathbb{Z})$ where N is given in the following Table:*

M	\mathbb{S}^{2m}	\mathbb{S}^{2m+1}	$\mathbb{C}P^m$	$\mathbb{H}P^m$	$Ca\mathbb{P}^2$
N	$\mathbb{C}P^{2m-1}$	$\mathbb{S}^{2m} \times \mathbb{C}P^m$	$\mathbb{C}P^{m-1} \times \mathbb{C}P^m$	$\mathbb{H}P^{m-1} \times \mathbb{C}P^{2m+1}$	$\mathbb{S}^8 \times \mathbb{C}P^{11}$

Proof In the Serre spectral sequence of $\mathbb{S}^{n-1} \rightarrow T^1M \rightarrow M$ the only nonzero differential is the transgression map $\mathbb{Z} = H^{n-1}(\mathbb{S}^{n-1}; \mathbb{Z}) \rightarrow H^n(M; \mathbb{Z}) = \mathbb{Z}$ which is the multiplication by $\chi(M)$. It is thus easy to compute the integral cohomology groups of T^1M (only the nontrivial groups are mentioned):

$$M = \mathbb{S}^n, n \text{ even} \quad H^q(T^1M; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = 0, 2n - 1 \\ \mathbb{Z}_2 & q = n \end{cases}$$

$$M = \mathbb{S}^n, n \text{ odd} \quad H^q(T^1M; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = 0, n - 1, n, 2n - 1 \end{cases}$$

$$M = \mathbb{C}P^m, n = 2m \quad H^q(T^1M; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = 2j \quad j = 0, \dots, m - 1 \\ & q = 2(m + j) + 1 \\ \mathbb{Z}_{n+1} & q = 2m \end{cases}$$

$$M = \mathbb{H}P^m, n = 4m \quad H^q(T^1M; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = 4j \quad j = 0, \dots, m - 1 \\ & q = 4(m + j) + 3 \\ \mathbb{Z}_{n+1} & q = 4m \end{cases}$$

$$M = Ca\mathbb{P}^2, n = 16 \quad H^q(T^1M; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = 0, 8, 23, 31 \\ \mathbb{Z}_3 & q = 16 \end{cases}$$

The result follows by analysing the Gysin sequence of the principal bundle $\mathbb{S}^1 \rightarrow T^1M \rightarrow (T^1M)/\mathbb{S}^1$. We show the explicit computations for the case of $M = \mathbb{H}P^m$.

It is enough to show that for $q \leq \dim(T^1\mathbb{H}P^m/\mathbb{S}^1)/2 = 4m - 1$, the cohomology of $T^1\mathbb{H}P^m/\mathbb{S}^1$ is

$$H^q(T^1\mathbb{H}P^m/\mathbb{S}^1; \mathbb{Z}) = \begin{cases} \mathbb{Z}^{j+1} & q = 4j, 4j + 2 \\ 0 & \text{otherwise} \end{cases}$$

which coincides with the cohomology of $\mathbb{H}\mathbb{P}^{m-1} \times \mathbb{C}\mathbb{P}^{2m+1}$ in that range. Since $T^1\mathbb{H}\mathbb{P}^m/\mathbb{S}^1$ and $\mathbb{H}\mathbb{P}^m \times \mathbb{C}\mathbb{P}^{2m+1}$ have the same dimension and satisfy Poincaré duality, the isomorphism in cohomology follows for all q .

Recall that, given a principal bundle $\mathbb{S}^1 \rightarrow E \rightarrow B$, the Gysin sequence reads

$$\dots \rightarrow H^{q-1}(B; \mathbb{Z}) \xrightarrow{\cup e} H^{q+1}(B; \mathbb{Z}) \rightarrow H^{q+1}(E; \mathbb{Z}) \rightarrow \dots \quad (3.6)$$

where $e \in H^2(B; \mathbb{Z})$ is the Euler class of the bundle.

For the sake of notation, we will denote $E = T^1\mathbb{H}\mathbb{P}^m$ and $B = T^1\mathbb{H}\mathbb{P}^m/\mathbb{S}^1$ for the rest of the proof. Notice that for If $q \leq 4m - 1$, $H^q(E; \mathbb{Z}) = 0$ unless $q = 4j$, and by (5.1) it follows that $H^{4q}(B; \mathbb{Z}) = H^{4q+2}(B; \mathbb{Z})$ and $H^{4q+1}(B; \mathbb{Z}) = H^{4q+3}(B; \mathbb{Z})$. In particular, $H^0(B; \mathbb{Z}) = H^2(B; \mathbb{Z}) = \mathbb{Z}$. Again from (5.1), $0 \rightarrow H^1(B; \mathbb{Z}) \rightarrow H^1(E; \mathbb{Z}) = 0$ and therefore $H^1(B; \mathbb{Z}) = H^3(B; \mathbb{Z}) = 0$, thus proving the claim for $q < 4$.

Suppose by induction that the claim is true for $k < 4j$, for $j < m$ (if it were $j = m$ we would be done). Again by (5.1) we have

$$0 = H^{4j-1}(B; \mathbb{Z}) \rightarrow H^{4j+1}(B; \mathbb{Z}) \rightarrow H^{4j+1}(E; \mathbb{Z}) = 0$$

and thus $H^{4j+1}(B; \mathbb{Z}) = H^{4j+3}(B; \mathbb{Z}) = 0$. Moreover,

$$0 \rightarrow H^{4j-2}(B; \mathbb{Z}) \xrightarrow{\cup e} H^{4j}(B; \mathbb{Z}) \rightarrow H^{4j}(E; \mathbb{Z}) \rightarrow 0$$

Since $j < m$, then $H^{4j}(E; \mathbb{Z}) = \mathbb{Z}$ and therefore $H^{4j}(B; \mathbb{Z}) = H^{4j-2}(B; \mathbb{Z}) \oplus \mathbb{Z}$. Since $H^{4j+2}(B; \mathbb{Z}) = H^{4j}(B; \mathbb{Z})$, this proves the induction step. □

Corollary 3.2 *Let M be a CROSS of dimension n . The relative equivariant cohomology $H_{\mathbb{S}^1}^*(\Delta M, M; \mathbb{Z})$ satisfies*

$$\begin{cases} H_{\mathbb{S}^1}^{odd}(\Delta M, M; \mathbb{Z}) = 0 & \text{if } n \text{ is odd} \\ H_{\mathbb{S}^1}^{ev}(\Delta M, M; \mathbb{Z}) = 0 & \text{if } n \text{ is even} \end{cases}$$

Proof It follows from Proposition 3.1 that $H^{odd}(T^1M/\mathbb{S}^1; \mathbb{Z}) = 0$ for all cases. Recall that for any $k > 0$, $H_{\mathbb{S}^1}^*(C^k; \mathbb{Z}) = H^*(T^1M/\mathbb{S}^1 \times B\mathbb{Z}_k; \mathbb{Z})$ and thus, since $H^{odd}(B\mathbb{Z}_k; \mathbb{Z}) = 0$ as well, it follows that $H_{\mathbb{S}^1}^{odd}(C^k; \mathbb{Z}) = 0$ for all $k > 0$. Moreover, by [17] the index i_k is even if and only if n is odd for any $k = 0$. By Eq. (3.2), it follows that for every $k > 0$,

$$\begin{cases} H_{\mathbb{S}^1}^{odd}(\Lambda^k, \Lambda^{k-1}; \mathbb{Z}) = 0 & \text{if } n \text{ is odd} \\ H_{\mathbb{S}^1}^{ev}(\Lambda^k, \Lambda^{k-1}; \mathbb{Z}) = 0 & \text{if } n \text{ is even} \end{cases} \quad (3.7)$$

Using the long exact sequence in cohomology

$$\dots \rightarrow H_{\mathbb{S}^1}^q(\Lambda^{k_1}, M; \mathbb{Z}) \rightarrow H_{\mathbb{S}^1}^q(\Lambda^{k_2}, M; \mathbb{Z}) \rightarrow H_{\mathbb{S}^1}^{q+1}(\Lambda^{k_1}, \Lambda^{k_2}; \mathbb{Z}) \rightarrow \dots$$

for any $k_1 > k_2$, we obtain by induction that for every k

$$\begin{cases} H_{\mathbb{S}^1}^{odd}(\Lambda^k, M; \mathbb{Z}) = 0 & \text{if } n \text{ is odd} \\ H_{\mathbb{S}^1}^{even}(\Lambda^k, M; \mathbb{Z}) = 0 & \text{if } n \text{ is even} \end{cases}$$

Taking the direct limit as $k \rightarrow \infty$ we obtain the result. □

Remark 3.3 By the Universal Coefficient Theorem, Corollary 3.2 also holds with coefficients in $R = \mathcal{P}^{-1}\mathbb{Z}$.

3.1 Integral cohomology CROSSes

Let now M be a manifold whose integral cohomology is that of a CROSS. The goal of this section is to prove the following generalization of Corollary 3.2.

Proposition 3.4 *Let M be a compact manifold whose rational cohomology ring is isomorphic to that of a CROSS M' . Then there is a ring isomorphism*

$$\phi : H_{\mathbb{S}^1}^*(\Lambda M, M; R) \rightarrow H_{\mathbb{S}^1}^*(\Lambda M', M'; R),$$

where $R = \mathcal{P}^{-1}\mathbb{Z}$ for a suitably chosen finite collection of primes \mathcal{P} .

Proof Since M has the rational cohomology of a CROSS M' then it is *formal*, i.e. its rational homotopy type only depends on the rational cohomology ring (cf. for example [1, Cor. 2.7.9]). Therefore there is a space M_0 and maps $M \rightarrow M_0 \leftarrow M'$ that induce isomorphisms in rational cohomology. Since the cohomology groups of M and M' are finitely generated, it follows that there is a finite set of primes \mathcal{P} and a space $M_{\mathcal{P}}$ (called *localisation* of M at \mathcal{P}) with maps $M \rightarrow M_{\mathcal{P}} \leftarrow M'$, which induce isomorphism in cohomology with coefficients in the localised ring $R = \mathcal{P}^{-1}\mathbb{Z}$ (cf. for example [9, Cor. 5.4(c)]). By Corollary 4.4 of [6] there is a homotopy commutative diagram

$$\begin{array}{ccccc} \Lambda M & \xrightarrow{\bar{\varphi}} & \Lambda M_{\mathcal{P}} & \xleftarrow{\bar{\varphi}'} & \Lambda M' \\ \downarrow & & \downarrow & & \downarrow \\ M & \xrightarrow{\varphi} & M_{\mathcal{P}} & \xleftarrow{\varphi'} & M' \end{array}$$

with horizontal arrows inducing isomorphism in $H^*(\cdot; R)$, where the map $\bar{\varphi} : [\mathbb{S}^1, M] \rightarrow \Lambda M_{\mathcal{P}} = [\mathbb{S}^1, M_{\mathcal{P}}]$ takes a curve c to $\varphi \circ c$, and similarly

for $\bar{\varphi}'$. In particular, $\bar{\varphi}$ and $\bar{\varphi}'$ are \mathbb{S}^1 equivariant, and therefore they induce isomorphisms in relative equivariant cohomology

$$H_{\mathbb{S}^1}^*(\Lambda M, M; R) \xleftarrow{\bar{\varphi}^*} H_{\mathbb{S}^1}^*(\Lambda M_{\mathcal{P}}, M_{\mathcal{P}}; R) \xrightarrow{\bar{\varphi}'^*} H_{\mathbb{S}^1}^*(\Lambda M', M'; R)$$

The composition $\phi = \bar{\varphi}'^* \circ (\bar{\varphi}^*)^{-1}$ is the isomorphism we wanted. □

Corollary 3.5 *Suppose the manifold M is a rational cohomology CROSS of dimension n . Then there is a finite set \mathcal{P} of primes, such that the relative equivariant cohomology $H_{\mathbb{S}^1}^*(\Lambda M, M; R)$, $R = \mathcal{P}^{-1}\mathbb{Z}$, satisfies*

$$\begin{cases} H_{\mathbb{S}^1}^{odd}(\Lambda M, M; R) = 0 & \text{if } n \text{ is odd} \\ H_{\mathbb{S}^1}^{ev}(\Lambda M, M; R) = 0 & \text{if } n \text{ is even} \end{cases}$$

Proof Suppose for sake of simplicity that n is odd, the case n even follows in the same way. By Proposition 3.4, it is enough to check that the theorem holds for M a CROSS. In this case, by Corollary 3.2 we have that $H_{\mathbb{S}^1}^{odd}(\Lambda M, M; \mathbb{Z}) = 0$ and, by the Universal Coefficient Theorem, we have that $H_{odd}((\Lambda M)_{\mathbb{S}^1}, M_{\mathbb{S}^1}; \mathbb{Z})$ is torsion and $H_{even}((\Lambda M)_{\mathbb{S}^1}, M_{\mathbb{S}^1}; \mathbb{Z})$ is free. Therefore, $\text{Hom}(H_{odd}((\Lambda M)_{\mathbb{S}^1}, M_{\mathbb{S}^1}; \mathbb{Z}), R) = 0$ since R is torsion free, and $\text{Ext}(H_{even}((\Lambda M)_{\mathbb{S}^1}, M_{\mathbb{S}^1}; \mathbb{Z}), R) = 0$. Again by the Universal Coefficient Theorem, $H_{\mathbb{S}^1}^{odd}(\Lambda M, M; R) = 0$. □

4 Index gap

Let (M, g) be a Besse manifold. For a geodesic $c : \mathbb{S}^1 \rightarrow M^n$ let c^q denote the q -iterate of c . Recall that by Wadsley theorem, there is a number L such that every prime geodesic has length equal to L/k for some integer k .

Definition 4.1 A closed geodesic c in M is called *regular* if its length is a multiple of L . Moreover, a critical set $C \subseteq \Lambda M$ for the energy functional is called *regular* if it contains regular geodesics.

For every primitive closed geodesic c there is some q such that c^q is regular. It follows in particular that any regular set C , containing geodesics of length, say, kL , is homeomorphic to the unit tangent bundle T^1M , via the map $T^1M \rightarrow C$ sending (p, v) to $c(t) = \exp_p(tkv)$.

Recall that the index $\text{ind}(c)$ of a closed geodesic is the index of the Hessian of the Energy functional $E : \Lambda M \rightarrow \mathbb{R}$, at c . Along a critical manifold C the index of the hessian remains constant, so sometimes we will also refer to the index $\text{ind}(C)$. Similarly, the *extended index* is given by $\text{ind}_0(c) = \text{ind}(c) + \text{null}(c)$, where $\text{null}(c)$ is the dimension of the kernel of the Hessian of E at c . Notice that $\text{null}(c)$ equals the number of periodic Jacobi fields along c , which

equals the dimension of the subspace of $V \oplus V$, $V = \langle c'(0) \rangle^\perp$, fixed by the Poincaré map of c .

Proposition 4.2 (Index Gap) *Let $c : \mathbb{S}^1 \rightarrow M^n$ be a geodesic such that c^q is regular. Then, for any l , $0 < l < q$:*

$$\text{ind}(c^{q+l}) = \text{ind}(c^q) + \text{ind}(c^l) + (n - 1) \tag{4.1a}$$

$$\text{ind}_0(c^q) = \text{ind}_0(c^{q-l}) + \text{ind}(c^l) + (n - 1) \tag{4.1b}$$

Proof Recall, for example from [2], that the index and extended index of a geodesic $c : [0, 2\pi] \rightarrow M$, with Poincaré map P can be computed as

$$\text{ind}(c) = \text{ind}_\Omega(c) + (\text{ind} + \dim \ker) \tilde{H} - \dim \ker(P - \text{Id})$$

$$\text{ind}_0(c) = \text{ind}_\Omega(c) + (\text{ind} + \dim \ker) \tilde{H}$$

where $\text{ind}_\Omega c$ is the number of conjugate points of $c(0)$ along c , and \tilde{H} is the concavity form defined on $(P - \text{Id})^{-1}(0 \oplus V)$ as

$$\tilde{H}(X, Y) = -\omega((P - \text{Id})X, Y).$$

As the summand $(\text{ind} + \dim \ker) \tilde{H} - \ker(P - \text{Id})$ only depends on P , we will call this $\text{ind}_P(c)$.

To prove Eq. (4.1a) it is enough to prove that

$$\text{ind}_\Omega(c^{q+l}) = \text{ind}_\Omega(c^q) + \text{ind}_\Omega(c^l) + (n - 1)$$

$$\text{ind}_P(c^{l+q}) = \text{ind}_P(c^q) + \text{ind}_P(c^l).$$

The first equation holds because for every conjugate point t_0 of c^{q+l} , we have $t_0 \in (0, q)$, $t_0 = q$, or $t_0 \in (q, q + l)$. By definition there are exactly $\text{ind}_\Omega(c^q)$ many conjugate points of the first type. Since every Jacobi field J with $J(0) = 0$ also satisfies $J(q) = 0$, we have in particular that $t_0 = q$ has multiplicity $n - 1$. Finally, since every Jacobi field is periodic on $[0, q]$, a Jacobi field J on $[0, q + l]$ satisfies $J(0) = J(t_0) = 0$ for some $t_0 \in (q, q + l)$ if and only if $K(t) = J|_{[q, q+l]}(t - q)$ satisfies $K(0) = K(t_0 - t) = 0$, thus there are exactly $\text{ind}_\Omega(c^l)$ many conjugate points of the last type.

For the second equation, just notice that since c^q has Poincaré map Id , it follows that $\text{ind}_P(c^q) = 0$, and since c^l and c^{q+l} have the same Poincaré map, it follows that $\text{ind}_P(c^l) = \text{ind}_P(c^{q+l})$.

To prove Eq. (4.1b), we first prove a couple of easy lemmas. □

Lemma 4.3 *Let \mathcal{L} be a Lagrangian subspace of $(V \oplus V, \omega)$ and let K be a symplectic subspace. Then*

$$\dim(\mathcal{L} \cap K^\perp) - \dim(\mathcal{L} \cap K) + \dim K = \dim V.$$

where $K^\perp = \{x \in V \oplus V \mid \omega(x, K) = 0\}$.

Proof We compute

$$\begin{aligned} \dim(\mathcal{L} \cap K^\perp) &= \dim(V \oplus V) - \dim(\mathcal{L} \cap K^\perp)^\perp \\ &= 2 \dim(V) - \dim(\mathcal{L}^\perp + K) \\ &= 2 \dim(V) - \dim(\mathcal{L}^\perp) - \dim(K) + \dim(\mathcal{L} \cap K) \\ &= \dim(V) - \dim(K) + \dim(\mathcal{L} \cap K). \end{aligned}$$

□

Lemma 4.4 *Let $K = \ker(P - \text{Id}) \subseteq V \oplus V$, where $P \in \text{U}(n - 1) \subseteq \text{Sp}(n - 1, \mathbb{R})$. Then for any subspace \mathcal{L} of $V \oplus V$,*

$$\dim(P - \text{Id})^{-1}(\mathcal{L}) = \dim K + \dim(\mathcal{L} \cap K^\perp).$$

Proof Notice first that K is the (generalised) eigenspace of P with eigenvalue 1, and therefore it is a symplectic subspace of $V \oplus V$ (cf. for example [2], p. 220-222). Because P lies in the maximal compact subgroup of $\text{Sp}(n - 1, \mathbb{R})$, it is possible to write $P = \begin{pmatrix} \text{Id}_K & \\ & P|_{K^\perp} \end{pmatrix}$. In particular, $\text{Im}(P - \text{Id}) \subseteq K^\perp$, K^\perp is $(P - \text{Id})$ -invariant, and the restriction $(P - \text{Id})|_{K^\perp}$ is invertible. Therefore

$$\begin{aligned} \dim(P - \text{Id})^{-1}(\mathcal{L}) &= \dim(P - \text{Id})^{-1}(\mathcal{L} \cap K^\perp) \\ &= \dim(K) + \dim(P - \text{Id})|_{K^\perp}^{-1}(\mathcal{L} \cap K^\perp) \\ &= \dim(K) + \dim(\mathcal{L} \cap K^\perp). \end{aligned}$$

□

We can now prove Eq. (4.1b). Let P be the Poincaré map of c , so that c^l and c^{q-l} have Poincaré maps P^l and P^{q-l} , respectively, and $P^q = \text{Id}$. Let us denote the Lagrangian subspace $0 \oplus V$ with \mathcal{L} , and the symplectic space $\ker(P^l - \text{Id}) = \ker(P^{q-l} - \text{Id})$ with K . By definition, $\dim K = \text{null}(c^l) = \text{null}(c^{q-l})$.

Since P satisfies $P^q = \text{Id}$, we have $\text{ind}_P(c^q) = 0$, $\text{null}(c^q) = 2(n - 1)$ and therefore $\text{ind}_0(c^q) = \text{ind}_\Omega(c^q) + 2(n - 1)$. Equation (4.1b) can be then

simplified as

$$\text{ind}_0(c^{q-l}) + \text{ind}(c^l) = \text{ind}_\Omega(c^q) + (n - 1)$$

To prove the equation above it is enough to prove that

$$\begin{aligned} \text{ind}_\Omega(c^{q-l}) + \text{ind}_\Omega(c^l) &= \text{ind}_\Omega(c^q) - \mu(q - l) \\ \text{ind}_P(c^{q-l}) + \text{ind}_P(c^q) + \text{null}(c^{q-l}) &= (n - 1) + \mu(q - l) \end{aligned}$$

where $\mu(q - l)$ denotes the number multiplicity of $q - l$ as a conjugate point of $c(0)$.

The first equation holds because for every conjugate point t_0 of c^q , we have either $t_0 \in (0, q - l)$, $t_0 = q - l$, or $t_0 \in (q - l, q)$, and by definition there are exactly $\text{ind}_\Omega(c^{q-l})$ many conjugate points of the first type, and $\text{ind}_\Omega(c^l)$ many conjugate points of the last type.

For the second equation notice that, since the Poincaré maps of c^l and c^{q-l} are inverses of each other, their concavity forms \tilde{H}_1 and \tilde{H}_2 (defined on the same space $(P^l - \text{Id})^{-1}(\mathcal{L}) = (P^{q-l} - \text{Id})^{-1}(\mathcal{L})$) satisfy the relation $\tilde{H}_1 = -\tilde{H}_2$. Therefore

$$(\text{ind} + \dim \ker) \tilde{H}_1 + (\text{ind} + \dim \ker) \tilde{H}_2 = \dim(P^l - \text{Id})^{-1}(\mathcal{L}) + \dim \ker \tilde{H}_2.$$

By Lemma 4.4, $\dim(P^l - \text{Id})^{-1}(\mathcal{L}) = \dim(K) + \dim(\mathcal{L} \cap K^\perp)$, and the equation after (1.3) in [2] gives

$$\dim \ker \tilde{H}_2 = \dim K + \mu(q - l) - \dim(\mathcal{L} \cap K).$$

Putting these equations together, we compute $\text{ind}_P(c^{q-l}) + \text{ind}_P(c^q) + \text{null}(c^{q-l})$ as

$$\begin{aligned} &(\text{ind} + \dim \ker) \tilde{H}_1 + (\text{ind} + \dim \ker) \tilde{H}_2 - \text{null}(c^l) \\ &= \dim(P^l - \text{Id})^{-1}(\mathcal{L}) + \dim \ker \tilde{H}_2 - \dim K \\ &= \mu(q - l) + \dim(K) + \dim(\mathcal{L} \cap K^\perp) - \dim(\mathcal{L} \cap K) \\ &= \mu(q - l) + (n - 1) \end{aligned}$$

where in the last equality we used Lemma 4.3. Thus Eq. (4.1b) holds. \square

Given a Besse manifold M , let $i(M)$ denote the minimal index of a critical set for the energy functional in ΛM . It is easy to see that $i(M)$ is the lowest degree q such that $H_{\mathbb{S}^1}^q(\Lambda M, M; \mathbb{Q}) \neq 0$. In fact, letting e be the smallest critical energy with index $i(M)$, clearly $q = i(M)$ would be the lowest degree such that $H_{\mathbb{S}^1}^q(\Lambda^{e+\epsilon}, M; \mathbb{Q}) \neq 0$. However, by the index parity for closed geodesics [17] the index of every critical set has the same parity

of $i(M)$. In particular, the index any further critical energy e_k would either be $= i(M)$, or it would be $\geq i(M) + 2$. In either case, we would have $H_{\mathbb{S}^1}^{i(M)}(\Lambda^{e_k+\epsilon}, M; \mathbb{Q}) \neq 0$ and moreover this would be the first nonzero degree. In the limit, $H_{\mathbb{S}^1}^{i(M)}(\Lambda M, M; \mathbb{Q})$ is the first nonzero homology group.

We thus have the following values for CROSSes:

$$i(\mathbb{S}^n) = n - 1, \quad i(\mathbb{C}P^n) = 1, \quad i(\mathbb{H}P^n) = 3, \quad i(CaP^2) = 7.$$

In general, since M is a simply connected rational cohomology CROSS then by Proposition 3.4 $i(M)$ only depends on the CROSS M is modelled on.

The following is a straightforward consequence of Proposition 4.2.

Corollary 4.5 *Given a rational cohomology CROSS M and a geodesic c such that c^q is regular, we have*

$$\text{ind}(c^k) \geq \text{ind}(c^q) + (n - 1) + i(M) \quad \text{if } k > q \quad (4.2)$$

$$\text{ind}_0(c^k) \leq \text{ind}(c^q) + (n - 1) - i(M) \quad \text{if } k < q. \quad (4.3)$$

Moreover, the inequality in (4.2) (resp. the inequality in (4.3)) is strict unless $k = q + 1$ (resp. $k = q - 1$) and $\text{ind}(c) = i(M)$.

5 Perfectness of the energy functional

The goal of this section is to prove Theorem D, that is, that for every simply connected Besse manifold M , the energy functional $E : \Lambda M \rightarrow \mathbb{R}$ is perfect with respect to the \mathbb{S}^1 equivariant, rational cohomology of $(\Lambda M, M)$.

In Sect. 5.1 we prove Theorem D in the special case in which all negative bundles are orientable. As pointed out in Corollary C, this is the case of spin manifolds, like manifolds with the integral cohomology ring of spheres, quaternionic projective spaces, or the Cayley plane. Finally, in 5.2 we prove Theorem D in the general case.

5.1 When the negative bundles are all orientable

Let M be a Besse manifold, and let $C_1, \dots, C_k \subseteq \Lambda M$ be the critical sets of E containing prime geodesics. For every $C \in \{C_1, \dots, C_k\}$, and every $q \in \mathbb{Z}$ let C^q denote the critical set consisting of q -iterates of geodesics in C . Clearly, every critical set of E is of the form C^q for some $q \in \mathbb{Z}$ and $C \in \{C_1, \dots, C_k\}$.

The core result of the section is Proposition 5.4, where we prove that the rational, \mathbb{S}^1 -equivariant cohomology of every critical set is concentrated in even degrees. This fact, together with the index parity result of the index in [17], will allow us to prove Theorem D by the lacunarity principle.

Before that, however, we first need to analyse the structure of p -torsion in the cohomology of the critical sets, for big primes p .

Remark 5.1 For the rest of this section, we will denote by R the localisation ring $R = \mathcal{P}^{-1}\mathbb{Z}$ where \mathcal{P} is a finite collection of primes such that

- every prime dividing the order of some element in $H^*(C; \mathbb{Z})$ or $H_{\mathbb{S}^1}^*(C; \mathbb{Z})$, $C \in \{C_1, \dots, C_k\}$ is contained in \mathcal{P} .
- Proposition 3.4 and Corollary 3.5 hold for $R = \mathcal{P}^{-1}\mathbb{Z}$.
- For every prime geodesics of length L/k , all the prime divisors of k are contained in \mathcal{P} . Equivalently, for any $p \notin \mathcal{P}$ and any critical set K , the set K^p is also a critical set.

For this to make sense, one must first make sure that there are only finitely many primes that divide the orders of the elements of $H^*(C; \mathbb{Z})$ and $H_{\mathbb{S}^1}^*(C; \mathbb{Z})$. However, this is clearly true for $H^*(C; \mathbb{Z})$ because it is finitely generated and in particular contains finitely many torsion elements. As for $H_{\mathbb{S}^1}^*(C; \mathbb{Z}) = H^*(C_{\mathbb{S}^1}; \mathbb{Z})$, from the Gysin sequence of the \mathbb{S}^1 -bundle $C \rightarrow C_{\mathbb{S}^1}$ we obtain that there is an isomorphism $H_{\mathbb{S}^1}^i(C; \mathbb{Z}) \rightarrow H_{\mathbb{S}^1}^{i+2}(C; \mathbb{Z})$ for any $i > \dim(C)$ and therefore the torsion of $H_{\mathbb{S}^1}^*(C; \mathbb{Z})$ is the same as the torsion of $H_{\mathbb{S}^1}^{\leq \dim(C)}(C; \mathbb{Z})$, which is finite.

We recall the following basic fact

Lemma 5.2 *For any critical energy e_r with critical manifold K_r , the group $H_{\mathbb{S}^1}^i(\Lambda^r, \Lambda^{r-1}; \mathbb{Z})$ can contain torsion free elements only in degrees $i \in \{\text{ind}(K_r), \dots, \text{ind}_0(K_r)\}$.*

Proof This is equivalent to showing that $H_{\mathbb{S}^1}^i(\Lambda^r, \Lambda^{r-1}; \mathbb{Q}) = 0$ for $i \notin \{\text{ind}(K_r), \dots, \text{ind}_0(K_r)\}$. Recall that $H_{\mathbb{S}^1}^i(\Lambda^r, \Lambda^{r-1}; \mathbb{Q}) = H_{\mathbb{S}^1}^{i-\text{ind}(K_r)}(K_r; \mathbb{Q})$. Moreover, since \mathbb{S}^1 acts almost freely on K_r , the quotient K_r/\mathbb{S}^1 is an orbifold of dimension $\dim(K_r) - 1 = \text{null}(K_r)$, and therefore $H_{\mathbb{S}^1}^{i-\text{ind}(K_r)}(K_r; \mathbb{Q}) = H^{i-\text{ind}(K_r)}(K_r/\mathbb{S}^1; \mathbb{Q})$. This is clearly 0 for $i \notin \{0, \dots, \text{null}(K_r)\}$ and this proves the result. □

Proposition 5.3 *Let K be a critical set of the energy functional, and let $p \notin \mathcal{P}$ be a prime. Then:*

- (1) *If K is a critical manifold with $H_{\mathbb{S}^1}^{\text{odd}}(K; \mathbb{Q}) = 0$, then any critical manifold of the form K^q has no p -torsion in $H_{\mathbb{S}^1}^{\text{odd}}(K^q; \mathbb{Z})$.*
- (2) *If K is a critical manifold with $H_{\mathbb{S}^1}^{\text{odd}}(K; \mathbb{Q}) \neq 0$, then $H_{\mathbb{S}^1}^{2h+1}(K^p; \mathbb{Z})$ contains p -torsion for every $2h + 1 \geq \dim(K) - 1$.*

Proof Recall that for every coefficient ring, $H_{\mathbb{S}^1}^*(K^q) = H^*(K_{\mathbb{S}^1}^q)$ with

$$K_{\mathbb{S}^1}^q \simeq K \times_{\mathbb{S}^1} E\mathbb{S}^1$$

where \mathbb{S}^1 acts on $K \times ES^1$ by $z \cdot (x, a) = (z^q \cdot x, z \cdot a)$. We can rewrite this slightly differently but homotopy equivalent as

$$K_{\mathbb{S}^1}^q \simeq K \times_{\mathbb{S}^1} ES^1 \times ES^1$$

with $z \cdot (x, a, b) = (z^q x, z^q a, z b)$. Since this \mathbb{S}^1 -action extends naturally to the free 2-torus action

$$(z_1, z_2) \cdot (x, a, b) = (z_1 x, z_1 a, z_2 b), \quad (z_1, z_2) \in T^2 = \mathbb{S}^1 \times \mathbb{S}^1,$$

we see that $K_{\mathbb{S}^1}^q = (K \times ES^1 \times ES^1)/\mathbb{S}^1$ is an \mathbb{S}^1 -bundle over $K_{\mathbb{S}^1} \times BS^1 = (K \times ES^1 \times ES^1)/T^2$.

The Euler class of this bundle $\xi : K_{\mathbb{S}^1}^q \rightarrow K_{\mathbb{S}^1} \times BS^1$ is given by

$$\bar{e} = e \otimes 1 - q(1 \otimes c) \in H^2(K_{\mathbb{S}^1} \times BS^1; R) \subset H_{\mathbb{S}^1}^*(K; R) \otimes H^*(BS^1; R).$$

where c is the generator of $H^2(BS^1; R)$ and e is the Euler class of the bundle $K \rightarrow K_{\mathbb{S}^1}$. The Gysin sequence for the bundle $K_{\mathbb{S}^1}^q \rightarrow K_{\mathbb{S}^1} \times BS^1$ reads

$$H^{2k+1}(K_{\mathbb{S}^1} \times BS^1; R) \xrightarrow{\xi^*} H^{2k+1}(K_{\mathbb{S}^1}^q; R) \xrightarrow{\xi_!} H^{2k}(K_{\mathbb{S}^1} \times BS^1; R) \tag{5.1}$$

If $H^{2k+1}(K_{\mathbb{S}^1}^q; R)$ had some p -torsion element x , it would lie in the kernel of $\xi_!$ because $H^*(K_{\mathbb{S}^1} \times BS^1; R)$ does not have any p -torsion by definition of \mathcal{P} . Then it would be $x = \xi^*(y)$ for some $y \in H^{2k+1}(K_{\mathbb{S}^1} \times BS^1; R)$. By the choice of p , y cannot be p -torsion and thus it must be torsion free, which implies that $H^{odd}(K_{\mathbb{S}^1}; \mathbb{Q}) \neq 0$. This proves the first point.

Suppose now that $H_{\mathbb{S}^1}^{odd}(K; \mathbb{Q}) \neq 0$, and let $x \in H^{h_0}(C_{\mathbb{S}^1}; R)$, h_0 odd, be a torsion-free element not divisible by p , such that $x \cup e = 0$. Such an x exists because $H^{odd}(K_{\mathbb{S}^1}; \mathbb{Q})$ is nonzero and, since the \mathbb{S}^1 -action on K is almost free, the cohomological dimension of $H^*(K_{\mathbb{S}^1}; \mathbb{Q})$ is at most $\dim(K) - 1$. The Gysin sequence of ξ reads

$$H^{h_0+2m-2}(K_{\mathbb{S}^1} \times BS^1; R) \xrightarrow{\cup \bar{e}} H^{h_0+2m}(K_{\mathbb{S}^1} \times BS^1; R) \xrightarrow{\xi^*} H^{h_0+2m}(K_{\mathbb{S}^1}^p; R)$$

The map $\cup \bar{e}$ is easily seen to be injective. Combining with the fact that the primitive element $(x \otimes c^{m-1})$ is mapped to $p \cdot (x \otimes c^m)$, we deduce that p -torsion can be found in the image of ξ^* . \square

Recall from Sect. 1 that we denote by K_1, K_2, \dots the list of the critical sets of E of positive energy, in increasing order $e_1 < e_2 < \dots$. For every

$k \geq 0$, let i_k denote the index of K_k . Moreover, let us define the sublevel sets $\Lambda^k = E^{-1}([0, e_k + \epsilon]) \subseteq \Lambda M$ for some $\epsilon > 0$ small enough.

Proposition 5.4 *For every critical set K of the energy functional, one has $H_{\mathbb{S}^1}^{odd}(K; \mathbb{Q}) = 0$.*

Proof We choose a large prime p and consider the ring

$$S := \mathbb{Z}[\{1/q \mid q \text{ prime } q \neq p\}]$$

all cohomology groups in this proof will be with respect to coefficients in S . Equivalently one can work with integral coefficients and ignore all torsion except for p -torsion.

We argue by contradiction and consider the smallest length l for which the set $C = C_l$ of geodesics of length l has nontrivial rational cohomology in some odd degree. We let C_1, \dots, C_{i-1} be the critical manifolds of smaller length containing primitive geodesics and $C_{i+1}, \dots, C_m \cong T^1M$ the ones of larger length containing primitive geodesics. Choose $\epsilon > 0$ such that there are no closed geodesics of length $l' \in ([l - \epsilon, l + \epsilon] \setminus \{l\})$.

We may assume that p is so large that we can find positive integers $u_1 < u_2$ such that $p(l - \epsilon) < u_1L < pl < u_2L < p(l + \epsilon)$, where L is the common period of all unit speed geodesics.

Recall that any critical manifold is given by iterating the geodesics in $\{C_1, \dots, C_m\}$, that is $K_h = (C_{j(h)})^{l(h)}$. Let r be the index corresponding to the critical manifold $C_i^p = K_r$. Furthermore let $r_1 < r$ and $r_2 > r$ denote the index corresponding to length u_1L and u_2L , respectively.

By construction K_h does not contain a p -times iterated geodesic if $h = r_1, \dots, r - 1$ or $h = r + 1, \dots, r_2$. In the following we denote by $[x]$ the Gauß bracket of a real number x .

Step 1. Suppose $r_1 \leq h \leq r - 1$. Then the inclusion $\Lambda^h \rightarrow \Lambda M$ induces an epimorphism $H_{\mathbb{S}^1}^i(\Lambda M, M) \rightarrow H_{\mathbb{S}^1}^i(\Lambda^h, M)$ for $i \geq \text{ind}(K_{r_2}) + n - 2$.

Let $e \in H^2(\Lambda M \times_{\mathbb{S}^1} E\mathbb{S}^1)$ denote the Euler class of the \mathbb{S}^1 -bundle $\Lambda M \rightarrow \Lambda M \times_{\mathbb{S}^1} E\mathbb{S}^1$.

We first consider the Λ^{r_1-1} . By the index gap Lemma we have $\text{ind}_0(K_h) \leq \text{ind}(K_{r_1}) + (n - 1)$, $h = 1, \dots, r_1 - 1$. Hence $H^i(\Lambda^{r_1-1}, M) = 0$ for $i \geq \text{ind}(K_{r_1}) + n$. Using the Gysin sequence of the \mathbb{S}^1 -bundle we see that $\cup_e: H_{\mathbb{S}^1}^i(\Lambda^{r_1-1}, M) \rightarrow H_{\mathbb{S}^1}^{i+2}(\Lambda^{r_1-1}, M)$ is an epimorphism for $i \geq \text{ind}(K_{r_1}) + 2[(n - 1)/2]$.

Furthermore, for each $h < r_1$ we know that either $H^{odd}(K_h) = 0$ or K_h does not contain a p -iterated geodesic. In either case, by Lemma 5.2 and Proposition 5.3 one has $H_{\mathbb{S}^1}^i(\Lambda^h, \Lambda^{h-1}) = H_{\mathbb{S}^1}^{i-\text{ind}(K_h)}(K_h) = 0$ if $i \equiv n \pmod 2$ with $i \geq \text{ind}(K_{r_1}) + 2[(n - 1)/2]$. Let $i_1 = \text{ind}(K_{r_1})$. From the exact sequence of the triple $(\Lambda^{r_1}, \Lambda^{r_1-1}, M)$ we then obtain, for each $i \equiv n + 1$

mod 2, the diagram

$$\begin{array}{ccccccc}
 H_{\mathbb{S}^1}^{i+2-i_1}(K_{r_1}) & \rightarrow & H_{\mathbb{S}^1}^{i+2}(\Lambda^{r_1}, M) & \xrightarrow{i^*} & H_{\mathbb{S}^1}^{i+2}(\Lambda^{r_1-1}, M) & \xrightarrow{d} & 0 \\
 \uparrow & & \uparrow & & \uparrow & & \\
 H_{\mathbb{S}^1}^{i-i_1}(K_{r_1}) & \rightarrow & H_{\mathbb{S}^1}^i(\Lambda^{r_1}, M) & \xrightarrow{i^*} & H_{\mathbb{S}^1}^i(\Lambda^{r_1-1}, M) & \xrightarrow{d} & 0
 \end{array}$$

where the vertical arrows are given by cupping with e and the horizontal sequences are exact. As explained the last vertical map is an epimorphism for $i \geq i_1 + 2[(n - 1)/2]$. Since $K_{r_1} = T^1M$ is a regular level the first vertical map is an epimorphism for $i \geq i_1 + 2[(n - 1)/2]$ as well. By the Four Lemma this implies that the middle map is an epimorphism.

The Index Gap Lemma implies for any critical manifold K_j with $j > r_1$ that $\text{ind}(K_j) \geq j_1 = i_1 + 2[(n - 1)/2] + 2$. Hence the map $H^{j_1}(\Lambda M, M) \rightarrow H^{j_1}(\Lambda^{r_1}, M)$ is an isomorphism. By the previous discussion we deduce that $H_{\mathbb{S}^1}^j(\Lambda M, M) \rightarrow H_{\mathbb{S}^1}^j(\Lambda^{r_1}, M)$ is surjective for each $j \geq j_1$.

Similarly, the natural map $H_{\mathbb{S}^1}^j(\Lambda^h, M) \rightarrow H_{\mathbb{S}^1}^j(\Lambda^{r_1}, M)$ is an isomorphism for $j \geq \text{ind}(K_{r_2}) + n - 2$ and $h = r_1, \dots, r - 1$, since $K_{r_1+1}, \dots, K_{r-1}$ do not contain p -iterated geodesics. Thus the claim of Step 1 follows.

Step 2. $H_{\mathbb{S}^1}^j(\Lambda^r M, M) \neq 0$ for all $j \geq \text{ind}(K_{r_2}) + n - 2$ with $j \equiv n \pmod 2$.

By Proposition 5.3, $H_{\mathbb{S}^1}^j(K_r) \neq 0$ for all odd $j \geq \text{dim}(K_r)$. Using the exact sequence of the triple $(\Lambda^r, \Lambda^{r-1}, M)$

$$H_{\mathbb{S}^1}^{j-1}(\Lambda^r, M) \xrightarrow{\iota^*} H_{\mathbb{S}^1}^{j-1}(\Lambda^{r-1}, M) \xrightarrow{d} H_{\mathbb{S}^1}^{j-ir}(K_r) \longrightarrow H_{\mathbb{S}^1}^j(\Lambda^r, M) \rightarrow$$

The map ι^* is a factor of $H_{\mathbb{S}^1}^{j-1}(\Lambda, M) \rightarrow H_{\mathbb{S}^1}^{j-1}(\Lambda^{r-1}, M)$, which is surjective by Step 1, and therefore so is ι^* . Hence the last map in the sequence above is injective and the result follows.

Step 3. $H_{\mathbb{S}^1}^j(\Lambda^{r_2}, M) \neq 0$ for all $j \geq \text{ind}(K_{r_2}) + 2[(n - 1)/2] + 1$ with $j \equiv n \pmod 2$.

The critical manifolds K_{r+1}, \dots, K_{r_2} do not contain p -iterated geodesics. By the index gap Lemma $H_{\mathbb{S}^1}^j(\Lambda^h, \Lambda^{h-1}) = 0$ for all $j \geq \text{ind}(K_{r_2}) + 2[(n - 1)/2] + 1$ and $h = r + 1, \dots, r_2 - 1$. This readily implies $H^j(\Lambda^{r_2-1}, M) \neq 0$ for all $j \geq \text{ind}(K_{r_2}) + 2[(n - 1)/2] + 1$. The critical manifold $K_{r_2} \cong T^1M$ is regular and $H_{\mathbb{S}^1}^j(\Lambda^{r_2}, \Lambda^{r_2-1}) = 0$ if $j \equiv n \pmod 2$ while $H_{\mathbb{S}^1}^j(\Lambda^{r_2}, \Lambda^{r_2-1})$ is torsion free if $j \equiv n + 1 \pmod 2$. Clearly the result follows.

Finally, the following step provides a contradiction to Corollary 3.5.

Step 4. $H_{\mathbb{S}^1}^{j_0}(\Lambda M, M) \neq 0$ for $j_0 = \text{ind}(K_{r_2}) + 2[(n - 1)/2] + 1$.

By the Index Gap Lemma all indices of critical manifold of energy $> e(K_{r_2})$ have indices $> j_0$. Furthermore the relative cohomology groups

$H^{j_0+1}(\Lambda^h, \Lambda^{h-1})$ are torsion free for $h > r_2$ while $H_{\mathbb{S}^1}^{j_0}(\Lambda^{r_2}, M)$ consists of nontrivial p -torsion. Thus the map $H_{\mathbb{S}^1}^{j_0}(\Lambda M, M) \rightarrow H_{\mathbb{S}^1}^{j_0}(\Lambda^{r_2}, M)$ is surjective. □

Corollary 5.5 *Let M be a Besse manifold. Then the energy function $E : \Lambda M \rightarrow \mathbb{R}$ is rationally \mathbb{S}^1 -equivariantly perfect, relatively to $M = \Lambda^0 \subseteq \Lambda M$ when all the negative bundles of all the critical sets of E are orientable.*

Proof We will prove this for M even dimensional, the other case follows in the same way. It is enough to prove that for every i , the map

$$H_{\mathbb{S}^1}^{*-ind(K_i)}(K_i; \mathbb{Q}) \simeq H_{\mathbb{S}^1}^*(\Lambda^i, \Lambda^{i-1}; \mathbb{Q}) \rightarrow H_{\mathbb{S}^1}^*(\Lambda^i; \mathbb{Q})$$

is injective. We prove this, together with the statement that $H_{\mathbb{S}^1}^{ev}(\Lambda^i, M; \mathbb{Q}) = 0$, by induction on i .

For $i = 0$ there is nothing to prove, so suppose that the both statements hold for $i - 1$. By the long exact sequence of $(\Lambda^i, \Lambda^{i-1}, M)$ we have

$$\begin{aligned} H_{\mathbb{S}^1}^{2m-ind(K_i)}(K_i; \mathbb{Q}) &\rightarrow H_{\mathbb{S}^1}^{2m}(\Lambda^i, M; \mathbb{Q}) \rightarrow H_{\mathbb{S}^1}^{2m}(\Lambda^{i-1}, M; \mathbb{Q}) \rightarrow \\ &\rightarrow H_{\mathbb{S}^1}^{2m+1-ind(K_i)}(K_i; \mathbb{Q}) \rightarrow H_{\mathbb{S}^1}^{2m+1}(\Lambda^i, M; \mathbb{Q}) \rightarrow \dots \end{aligned}$$

Since M is even dimensional, $ind(K)$ is odd, thus $H_{\mathbb{S}^1}^{2m-ind(K_i)}(K_i; \mathbb{Q}) = 0$ by Proposition 5.4. Moreover, by the induction hypothesis $H_{\mathbb{S}^1}^{2m}(\Lambda^{i-1}, M; \mathbb{Q}) = 0$, which gives

$$\begin{aligned} H_{\mathbb{S}^1}^{2m}(\Lambda^i, M; \mathbb{Q}) &= 0 \\ 0 &\rightarrow H_{\mathbb{S}^1}^{2m+1-ind(K_i)}(K_i; \mathbb{Q}) \rightarrow H_{\mathbb{S}^1}^{2m+1}(\Lambda^i, M; \mathbb{Q}), \end{aligned}$$

thus proving the induction step. □

5.2 The general case

We now remove the assumption that the negative bundles $N \rightarrow K$ are all orientable. By Corollary C, this can only happen if the manifold M has the integral cohomology of $\mathbb{C}P^{2n}$. In particular, M has even dimension, and therefore:

For the rest of the section, we will assume that the manifold M is even dimensional. In particular, by [17], $ind(K)$ is odd for every critical set K of the energy functional.

Let K be a critical manifold with non orientable negative bundle $N \rightarrow K$. In this case, we denote by $\delta : \hat{K} \rightarrow K$ the double cover such that N pulls back to an orientable bundle \hat{N} over \hat{K} . By Theorem B, \hat{K} can be realized as the

quotient \tilde{K}/H where \hat{K} is the universal cover of K , and $H \subseteq \pi_1(K)$ is the kernel of the homomorphism $A_* : \pi_1(K) \rightarrow \pi_1(\text{SO}(n - 1)) \simeq \mathbb{Z}_2$ induced by the holonomy map $A : K \rightarrow \text{SO}(n - 1)$. Notice that, given an orbit γ for the \mathbb{S}^1 -action on K , $A(\gamma(t))$ is constant. Hence, $A_*([\gamma]) = 1$ and therefore the \mathbb{S}^1 -action on $N \rightarrow K$ lifts to $\hat{N} \rightarrow \hat{K}$.

Lemma 5.6 *Let $K, K' \subseteq \Lambda M$ be critical sets of the energy functional, such that $K' = K^q$ for some q . Then*

- (1) *If $N \rightarrow K$ is orientable, then $N' \rightarrow K'$ is orientable as well.*
- (2) *If $N \rightarrow K$ is non-orientable, then $N' \rightarrow K'$ is orientable if and only if q is even.*
- (3) *If both $N \rightarrow K$ and $N' \rightarrow K'$ are not orientable, the diffeomorphism $f : K \rightarrow K'$ sending c to c^q lifts to a \mathbb{Z}_2 -equivariant diffeomorphism $\hat{K} \rightarrow \hat{K}'$.*

Proof The map $K \rightarrow K'$ sending $c \rightarrow c^q$ allows us to identify K and K' . To prove 1) and 2), it is enough to observe that when K has holonomy map A and $K' = K^q$ then, under the identification $K \sim K'$, the holonomy map $A' : K' \rightarrow \text{SO}(n - 1)$ equals A^q and in particular $A'_* = qA_*$.

To prove 3), it is sufficient to further notice that when $N \rightarrow K$ and $N' \rightarrow K'$ are both non orientable, in particular q is odd, and therefore the map $f_* : \pi_1(K) \rightarrow \pi_1(K')$ sends the kernel of A_* isomorphically to the kernel of A'_* . □

Let K be a critical manifold with non orientable negative bundle, and let \hat{K} be the 2-fold cover defined above. The \mathbb{Z}_2 -action on \hat{K} induces a \mathbb{Z}_2 -action on $H^*(\hat{K})$. Letting g denote the generator of \mathbb{Z}_2 , we define

$$H^*(\hat{K})^{-\mathbb{Z}_2} = \{x \in H^*(\hat{K}) \mid g \cdot x = -x\}$$

Proposition 5.7 *Let K_i be a critical manifold for the energy functional, with non-orientable negative bundle. Then if R is a ring where 2 is invertible, we have*

$$H^*(\Lambda^i, \Lambda^{i-1}; R) \simeq H^{*-\text{ind}(K_i)}(\hat{K}_i; R)^{-\mathbb{Z}_2}$$

and the same holds for \mathbb{S}^1 equivariant cohomology.

Proof By excision, $H^*(\Lambda^i, \Lambda^{i-1}; R) \simeq H^*(N_i, \partial N_i; R)$. Let $\hat{\eta} : \hat{N}_i \rightarrow \hat{K}_i$ denote the lift of $\eta : N_i \rightarrow K_i$. Then $(\hat{N}_i, \partial \hat{N}_i) \rightarrow (N_i, \partial N_i)$ is a \mathbb{Z}_2 -cover as well and, since 2 is invertible in R , by [5, Thm. 2.4] it induces an isomorphism

$$H^*(N_i, \partial N_i; R) \simeq H^*(\hat{N}_i, \partial \hat{N}_i; R)^{\mathbb{Z}_2}.$$

Moreover, since \hat{N}_i is orientable, by the Thom isomorphism there is a class $\hat{\tau} \in H^{\text{ind}(K_i)}(\hat{N}_i, \partial\hat{N}_i; R)$ such that the map

$$T : H^{k-\text{ind}(K_i)}(\hat{K}_i; R) \rightarrow H^k(\hat{N}_i, \partial\hat{N}_i; R), \quad \alpha \mapsto \hat{\eta}^*(\alpha) \cup \hat{\tau}$$

induces an isomorphism of groups for every $q > 0$. By construction, the Thom class $\hat{\tau}$ satisfies $g \cdot \hat{\tau} = -\hat{\tau}$, and thus

$$\begin{aligned} g \cdot T(\alpha) &= g \cdot (\hat{\eta}^*(\alpha) \cup \hat{\tau}) \\ &= (g \cdot \hat{\eta}^*(\alpha)) \cup (g \cdot \hat{\tau}) \\ &= -\hat{\eta}^*(g \cdot \alpha) \cup \hat{\tau} = -T(g \cdot \alpha) \end{aligned}$$

Therefore, T sends $H^{q-\text{ind}(K_i)}(\hat{K}_i; R)^{-\mathbb{Z}_2}$ isomorphically into $H^q(\hat{N}_i, \partial\hat{N}_i; R)^{\mathbb{Z}_2}$. Therefore

$$H^k(N_i, \partial N_i; R) \simeq H^k(\hat{N}_i, \partial\hat{N}_i; R)^{\mathbb{Z}_2} \simeq H^{q-\text{ind}(K_i)}(\hat{K}_i; R)^{-\mathbb{Z}_2}.$$

Because all the maps involved are \mathbb{S}^1 -equivariant, and all the properties used hold for \mathbb{S}^1 -equivariant cohomology as well, the result follows for \mathbb{S}^1 -equivariant cohomology as well:

$$H_{\mathbb{S}^1}^k(N_i, \partial N_i; R) \simeq H_{\mathbb{S}^1}^{q-\text{ind}(K_i)}(\hat{K}_i; R)^{-\mathbb{Z}_2}.$$

□

We are now ready to modify the proof of Theorem D in the previous section, in the case of non orientable bundles. This time, since we do not have the Thom isomorphism at hand, we want to use the relative cohomology $H_{\mathbb{S}^1}^*(\Lambda^i, \Lambda^{i-1}; \mathbb{Q})$ instead of $H_{\mathbb{S}^1}^*(K_i; \mathbb{Q})$, and prove by contradiction that it satisfies

$$H_{\mathbb{S}^1}^{ev}(\Lambda^i, \Lambda^{i-1}; \mathbb{Q}) = 0. \tag{5.2}$$

Supposing that this is not the case, then among the pairs which do not satisfy (5.2), we focus on the one whose corresponding critical set C has minimal energy. Just as in the previous section, we provide the contradiction by showing show that for some prime p big enough, the pair $(\Lambda^i, \Lambda^{i-1})$ corresponding to C^p introduces some p -torsion element on $H_{\mathbb{S}^1}^{ev}(\Lambda M, M; R)$ which cannot be removed, contradicting Corollary 3.5 according to which $H_{\mathbb{S}^1}^{ev}(\Lambda M, M; R) = 0$.

The following is the equivalent of Proposition 5.3.

Proposition 5.8 *Let e_i be a critical energy with critical manifold K_i , and let p be a big enough prime. Then:*

- (1) If $H_{\mathbb{S}^1}^{ev}(\Lambda^i, \Lambda^{i-1}; \mathbb{Q}) = 0$ then for any critical energy e_j with critical set $K_j = K_i^q$, q odd, the group $H^{ev}(\Lambda^j, \Lambda^{j-1}; R)$ contains no p -torsion.
- (2) If $H_{\mathbb{S}^1}^{ev}(\Lambda^i, \Lambda^{i-1}; \mathbb{Q}) \neq 0$, then for the critical energy e_j with critical set $K_j = K_i^p$, the group $H^{2h}(\Lambda^j, \Lambda^{j-1}; R)$ contains p -torsion for every $2h \geq \text{ind}_0(K_i)$.

Proof When $N_i \rightarrow K_i$ is orientable, the result follows directly from Proposition 5.3 and the Thom isomorphism.

When $N_i \rightarrow K_i$ is non-orientable, then we can repeat the same constructions in Proposition 5.3 to the \mathbb{Z}_2 -coverings \hat{K}_i, \hat{K}_i^q of K_i and K_i^p , and obtain a \mathbb{Z}_2 equivariant \mathbb{S}^1 -bundle

$$\hat{\xi} : (\hat{K}_i^q)_{\mathbb{S}^1} \simeq \hat{K}_i \times_{\mathbb{S}^1} B\mathbb{Z}_q \rightarrow (\hat{K}_i)_{\mathbb{S}^1} \times B\mathbb{S}^1. \tag{5.3}$$

whose \mathbb{Z}_2 -quotient is the bundle ξ defined in Proposition 5.3. The Gysin sequence of $\hat{\xi}$ is a \mathbb{Z}_2 -equivariant long exact sequence

$$H^{2k+1} \left((\hat{K}_i)_{\mathbb{S}^1} \times B\mathbb{S}^1; R \right) \xrightarrow{\hat{\xi}^*} H_{\mathbb{S}^1}^{2k+1}(\hat{K}_i^q; R) \xrightarrow{\hat{\xi}_!} H^{2k} \left((\hat{K}_i)_{\mathbb{S}^1} \times B\mathbb{S}^1; R \right)$$

equivalent to (5.1). Arguing in the same way as in Proposition 5.3 and taking the $-\mathbb{Z}_2$ -invariant part, we can see that $H_{\mathbb{S}^1}^{odd}(\hat{K}_i^q; R)^{-\mathbb{Z}_2}$ (which equals $H_{\mathbb{S}^1}^{ev}(\Lambda^j, \Lambda^{j-1}; R)$) cannot have p -torsion unless $H_{\mathbb{S}^1}^{odd}(\hat{K}_i; \mathbb{Q})^{-\mathbb{Z}_2}$ ($= H_{\mathbb{S}^1}^{ev}(\Lambda^i, \Lambda^{i-1}; \mathbb{Q})$) is nonzero.

On the other hand, if $H_{\mathbb{S}^1}^{odd}(\hat{K}_i; \mathbb{Q})^{-\mathbb{Z}_2} \neq 0$ then we can find some torsion-free element $x \in H_{\mathbb{S}^1}^{odd}(\hat{K}_i; R)^{-\mathbb{Z}_2}$ not divisible by p and such that $x \cup \hat{e} = 0$, where $\hat{e} \in H_{\mathbb{S}^1}^2(\hat{K}_i; R)$ is the Euler class of $\hat{K}_i \rightarrow \hat{K}_{\mathbb{S}^1}$. Then, again as in Proposition 5.3, one can prove that for every $k > 0$, the element $\hat{\xi}^*(x \otimes c^k) \in H_{\mathbb{S}^1}^{odd}(\hat{K}_i^p; R)^{-\mathbb{Z}_2}$ is a non trivial p -torsion element. \square

The new version of Proposition 5.4 is the following:

Proposition 5.9 *For every critical energy e_i , one has $H_{\mathbb{S}^1}^{ev}(\Lambda^i, \Lambda^{i-1}; \mathbb{Q}) = 0$.*

The proof is, for a large part, the same as the one of Proposition 5.4: we will give a sketch of the proof of Proposition 5.9, by focusing on the parts that differ from Proposition 5.4.

Proof As in Proposition 5.4, we choose a large prime and consider the localization $S = \mathbb{Z}[\frac{1}{q} \mid q \text{ prime} \neq p]$ as coefficient ring. We argue by contradiction and consider the smallest length l in correspondence to which one has $H_{\mathbb{S}^1}^{ev}(\Lambda^i, \Lambda^{i-1}; \mathbb{Q}) \neq 0$. From Proposition 5.7 and Proposition 5.8, one can see that the critical manifold K_i has one of the following forms:

- $K_i = C_j$ for some $C \in \{C_1, \dots, C_m\}$ with $N \rightarrow C$ orientable and $H_{\mathbb{S}^1}^{odd}(C; \mathbb{Q}) \neq 0$.
- $K_i = C_j$ for some $C \in \{C_1, \dots, C_m\}$ with $N \rightarrow C$ non orientable and $H_{\mathbb{S}^1}^{odd}(\hat{C}; \mathbb{Q})^{-\mathbb{Z}_2} \neq 0$.
- $K_i = C_j^2$ for some $C \in \{C_1, \dots, C_m\}$ with $N \rightarrow C$ non orientable, $H_{\mathbb{S}^1}^{odd}(\hat{C}; \mathbb{Q})^{-\mathbb{Z}_2} = 0$ and $H_{\mathbb{S}^1}^{odd}(C; \mathbb{Q}) \neq 0$.

Consider the set \mathcal{S} of critical manifolds K_j of the form $K_j = C, C \in \{C_1, \dots, C_m\}$, or $K_j = C^2$ with $N \rightarrow C$ non orientable, and pick an ϵ small enough that there are no critical sets in \mathcal{S} of length $l' \in [l - \epsilon, l + \epsilon] \setminus \{l\}$. By choosing p large enough, we can assume that there are integers $u_1 < u_2$ such that $p(l - \epsilon) < u_1L < pl < u_2L < p(l + \epsilon)$.

Let now K_r denote the critical set $(K_i)^p$, and K_{r_1}, K_{r_2} the critical manifolds of length u_1L and u_2L , respectively. By Proposition 5.8, it follows that K_h does not contain p -iterates for $h = r_1, \dots, r - 1$ and therefore $H_{\mathbb{S}^1}^{2j}(\Lambda^h, \Lambda^{h-1}) = 0$ for every $2j \geq \text{ind}(K_h) + (n - 1)$. Using this, the steps of Proposition 5.4 follow identically. □

6 The proof of the Main Theorem

Let $\mathbb{S}^n, n > 3$, be a topological sphere endowed with a Besse metric, and let L be the common period of the geodesics. We want to prove by contradiction that all geodesics have the same length L or, equivalently, that the geodesic flow $\mathbb{S}^1 \curvearrowright T^1\mathbb{S}^n$ acts freely. If not, there are critical sets of the energy functional $E : \Lambda\mathbb{S}^n \rightarrow \mathbb{R}$ which consist of geodesics of length $L/m, m \in \mathbb{Z}$, and they can be identified with the fixed point set, in $T^1\mathbb{S}^n$, of the subgroup $\mathbb{Z}_m \subseteq \mathbb{S}^1$. Since these sets have even codimension in $T^1\mathbb{S}^n$, in particular every critical set of positive energy has odd dimension.

By Proposition 4.2, the critical set C of lowest index consists of geodesics of length L/m for some integer m and, by the discussion above, it must be

$$\dim C \leq 2n - 3.$$

We now divide the discussion into two cases, according to whether n is even or odd.

If n is even, the integral cohomology and rational \mathbb{S}^1 -equivariant cohomology groups of $(\Lambda\mathbb{S}^n, \mathbb{S}^n)$ are the following

$$H^q(\Lambda\mathbb{S}^n, \mathbb{S}^n; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = (2k - 1)(n - 1), k \geq 1 \\ \mathbb{Z} & q = (2k + 1)(n - 1) + 1, k \geq 1 \\ \mathbb{Z}_2 & q = 2k(n - 1) + 1, k \geq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$H_{\mathbb{S}^1}^q(\Lambda\mathbb{S}^n, \mathbb{S}^n; \mathbb{Q}) = \begin{cases} \mathbb{Q} & q \geq (n - 1) \text{ odd, } q \neq 3(n - 1), 5(n - 1), \dots \\ \mathbb{Q}^2 & q = 3(n - 1), 5(n - 1), \dots \\ 0 & \text{otherwise} \end{cases}$$

By the perfectness of the energy functional, C consists of one component only, it is the unique critical set with minimal index, and $\text{ind}(C) = n - 1$. Moreover, the $\mathbb{S}^1/\mathbb{Z}_m$ -action on C is free, otherwise there would be some closed geodesics $c \notin C$ such that $c^k \in C$ for some k . If this were the case, we claim that this would imply $\text{ind}(c) \leq \text{ind}(c^k) = \text{ind}(C)$, contradicting the minimality of C . To prove the claim, recall that by Bott iteration formula (cf. for example [2] Theorem 2.1) for any closed geodesic c there exists a non-negative function I defined on the unit circle in \mathbb{C} , such that

$$\text{ind}(c^m) = \sum_{z^m=1} I(z).$$

From this formula, it is clear that $\text{ind}(c^k) \geq \text{ind}(c)$ for every closed geodesic c and every non-negative integer k .

In particular, $O(2)/\mathbb{Z}_m \simeq O(2)$ acts freely on C .

The quotient (manifold) C/\mathbb{S}^1 is embedded in $T^1\mathbb{S}^n/\mathbb{S}^1$, which is a symplectic orbifold (cf. [16]). It is easy to see that the symplectic form on $T^1\mathbb{S}^n/\mathbb{S}^1$ restricts to a symplectic form on C/\mathbb{S}^1 , and therefore C/\mathbb{S}^1 is a symplectic manifold. In particular,

$$\dim H_{\mathbb{S}^1}^{2q}(C; \mathbb{Q}) = \dim H^{2q}(C/\mathbb{S}^1; \mathbb{Q}) \geq 1 \tag{6.1}$$

for every $2q \leq \dim(C/\mathbb{S}^1)$. However, by the perfectness of the energy functional in rational equivariant cohomology, for any $q \leq 2n - 3$ we have

$$\begin{aligned} \dim H_{\mathbb{S}^1}^q(C; \mathbb{Q}) &= \dim H_{\mathbb{S}^1}^{q+(n-1)}(\Lambda^{e+\epsilon}, \Lambda^{e-\epsilon}; \mathbb{Q}) \\ &\leq \dim H_{\mathbb{S}^1}^{q+(n-1)}(\Lambda\mathbb{S}^n, \mathbb{S}^n; \mathbb{Q}) = \begin{cases} 1 & q \text{ even} \\ 0 & q \text{ odd} \end{cases} \end{aligned} \tag{6.2}$$

From inequalities (6.1) and (6.2) it follows that for any $q \leq \dim(C/\mathbb{S}^1) = \dim(C) - 1$,

$$H_{\mathbb{S}^1}^q(C; \mathbb{Q}) = H_{\mathbb{S}^1}^{q+(n-1)}(\Lambda^{e+\epsilon}, \Lambda^{e-\epsilon}; \mathbb{Q}) = H_{\mathbb{S}^1}^{q+(n-1)}(\Lambda\mathbb{S}^n, \mathbb{S}^n; \mathbb{Q})$$

where $e = E(C)$. Again by perfectness of the energy functional, it follows that every critical set different from C cannot contribute to the rational equivariant cohomology in degrees $\leq (n - 1) + \dim(C) - 1$, and in particular the index of every critical set different from C must be $\geq (n - 1) + \dim(C)$.

The index of the critical sets, however, does not depend on the cohomology we are using. In particular, if we now switch to regular integral cohomology, we still have that the only contribution to the cohomology $H^q(\Lambda S^n, S^n; \mathbb{Z})$ in degrees $q \leq (n - 1) + \dim(C) - 1$ is given by $H^{q-(n-1)}(C; \mathbb{Z})$ and therefore

$$H^q(C; \mathbb{Z}) = H^{q+(n-1)}(\Lambda S^n, S^n; \mathbb{Z}) \quad \forall q = 0, \dots, \dim(C) - 1.$$

In particular, $H^q(C; \mathbb{Z}) = 0$ for every $q = 1, \dots, m_0 = \min\{\dim(C) - 1, n - 1\}$. For $n \geq 4$, we have $m_0 \geq \frac{1}{2} \dim(C) + 1$ and therefore, by Poincaré duality, C is an integral cohomology sphere. However, $\mathbb{Z}_2 \times \mathbb{Z}_2 \subseteq O(2)$ acts freely on C and this contradicts the well-known result, that a finite abelian group acting freely on an integral cohomology sphere must be cyclic (cf. [5] Theorem 8.1).

If n is odd, the integral cohomology and the rational S^1 -equivariant cohomology of $(\Lambda S^n, S^n)$ are as follows:

$$H^q(\Lambda S^n, S^n; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = k(n - 1) \text{ or } q = (k + 1)(n - 1) + 1, \quad k \geq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$H_{S^1}^q(\Lambda S^n, S^n; \mathbb{Q}) = \begin{cases} \mathbb{Q} & q \geq (n - 1) \text{ even, } q \neq 2(n - 1), 3(n - 1), \dots \\ \mathbb{Q}^2 & q = 2(n - 1), 3(n - 1), \dots \\ 0 & \text{otherwise} \end{cases}$$

As in the previous case $\text{ind}(C) = n - 1$, C is the unique critical set with minimal index, and $O(2)/\mathbb{Z}_m \simeq O(2)$ acts freely on C . Moreover, C/S^1 is a symplectic manifold and

$$\dim H^{2q}(C/S^1; \mathbb{Q}) = \dim H_{S^1}^{2q}(C; \mathbb{Q}) \geq 1 \quad \forall q \leq \frac{1}{2} \dim(C/S^1). \quad (6.3)$$

By the perfectness of the energy functional in rational equivariant cohomology, for any $q \leq n - 2$ we have

$$\begin{aligned} \dim H_{S^1}^q(C; \mathbb{Q}) &= \dim H_{S^1}^{q+(n-1)}(\Lambda^{e+\epsilon}, \Lambda^{e-\epsilon}; \mathbb{Q}) \\ &\leq \dim H_{S^1}^{q+(n-1)}(\Lambda S^n, S^n; \mathbb{Q}) = \begin{cases} 1 & q \text{ even} \\ 0 & q \text{ odd} \end{cases} \end{aligned} \quad (6.4)$$

For any $q \leq m_0 = \min\{\dim(C/S^1), n - 2\}$ we then have

$$H_{S^1}^q(C; \mathbb{Q}) = H_{S^1}^{q+(n-1)}(\Lambda^{e+\epsilon}, \Lambda^{e-\epsilon}; \mathbb{Q}) = H_{S^1}^{q+(n-1)}(\Lambda S^n, S^n; \mathbb{Q}) \quad (6.5)$$

For $n \geq 3$ we have $m_0 \geq \frac{1}{2} \dim(C/S^1)$ and, by Poincaré duality, we obtain that $H^*(C/S^1; \mathbb{Q}) = H^*(\mathbb{C}P^{\dim(C/S^1)/2}; \mathbb{Q})$.

We claim that $\dim(C/S^1) \leq n - 1$. If this was not the case, the critical set C' with second lowest index, would have index $2(n - 1)$. Call e' its energy. If $\dim C' = 1$, then C' consists of at least two geodesics, because for any geodesic $\gamma(t)$ in C' , the inverted geodesic $\gamma(-t)$ belongs to C' as well. Therefore $H_{S^1}^0(C'; \mathbb{Q}) = H_{S^1}^{2(n-1)}(\Lambda^{e'+\epsilon}, \Lambda^{e'-\epsilon}; \mathbb{Q})$ would have dimension ≥ 2 , hence by perfectness

$$\begin{aligned} 2 &= \dim H_{S^1}^{2(n-1)}(\Lambda S^n, S^n; \mathbb{Q}) \\ &\geq H_{S^1}^{2(n-1)}(\Lambda^{e+\epsilon}, \Lambda^{e-\epsilon}; \mathbb{Q}) + H_{S^1}^{2(n-1)}(\Lambda^{e'+\epsilon}, \Lambda^{e'-\epsilon}; \mathbb{Q}) \geq 3 \end{aligned}$$

which gives a contradiction. If $\dim C'/S^1 \geq 1$, the quotient C'/S^1 would be a symplectic orbifold just as C and hence it would satisfy

$$\dim H^2(C'/S^1; \mathbb{Q}) = \dim H_{S^1}^{2n}(\Lambda^{e'+\epsilon}, \Lambda^{e'-\epsilon}; \mathbb{Q}) \geq 1,$$

but again by perfectness we would then have

$$\begin{aligned} 1 &= \dim H_{S^1}^{2n}(\Lambda S^n, S^n; \mathbb{Q}) \\ &\geq \dim H_{S^1}^{2n}(\Lambda^{e+\epsilon}, \Lambda^{e-\epsilon}; \mathbb{Q}) + \dim H_{S^1}^{2n}(\Lambda^{e'+\epsilon}, \Lambda^{e'-\epsilon}; \mathbb{Q}) \geq 2 \end{aligned}$$

which would provide a contradiction as well.

Therefore $\dim(C/S^1) \leq n - 1$ and, by (6.5), every other critical set has index $\geq (n - 1) + \dim(C) - 1$. Then shifting our attention to integral cohomology, the only contribution to $H^q(\Lambda S^n, S^n; \mathbb{Z})$ in degrees $q \leq (n - 1) + \dim(C) - 1$ is given by $H^{q-(n-1)}(C; \mathbb{Z})$. In particular,

$$H^q(C; \mathbb{Z}) = H^{q+(n-1)}(\Lambda S^n, S^n; \mathbb{Z}) = \begin{cases} \mathbb{Z} & q = 0 \\ 0 & q = 1, \dots, \dim(C) - 1 \end{cases}$$

For $n \geq 3$, this covers more than half of the cohomology of C and therefore C is an integral cohomology sphere. As in the previous case, the fact that $\mathbb{Z}_2 \oplus \mathbb{Z}_2 \subseteq O(2)/\mathbb{Z}_m$ acts freely on C provides a contradiction.

Appendix A. Small subsets of $\mathbf{Sp}(n - 1, \omega)$

Let $(\mathbb{R}^{2(n-1)}, \omega)$ be the symplectic vector space, and let $\mathbf{Sp}(n - 1, \omega) = \{P \in \text{GL}(2(n - 1), \mathbb{R}) \mid \omega(Ax, Ay) = \omega(x, y)\}$ be the real symplectic group. In this appendix we focus on the subspaces of symplectic matrices with real eigenvalues of higher geometric multiplicity.

Lemma A.1 *Let $P \in \mathbf{Sp}(n - 1, \omega)$ and let λ be a positive real eigenvalue of P of algebraic multiplicity a .*

- (a) *If $\lambda \neq 1$ then up to conjugation with an element of $\mathbf{Sp}(n - 1, \omega)$, the matrix P can be written as $P = \text{diag}(\lambda U^{tr}, (\lambda U)^{-1}, R)$ where $U \in \text{GL}(a, \mathbb{R})$ is unipotent, $R \in \mathbf{Sp}(n - a - 1, \omega)$, and U^{tr} denotes the transpose of U .*
- (b) *If $\lambda = 1$ then up to conjugation with some element of $\mathbf{Sp}(n - 1, \omega)$, the matrix P can be written as $P = \text{diag}(U, R)$ where $U \in \mathbf{Sp}(a, \omega)$ is unipotent and $R \in \mathbf{Sp}(n - a - 1, \omega)$.*

Proof By the so-called refined Jordan decomposition, there are commuting matrices $P_s, P_u \in \mathbf{Sp}(n - 1, \omega)$ such that P_s is diagonalizable over \mathbb{C} , P_u is unipotent, and $P = P_s P_u$. In particular, P_s has the same eigenvalues of P with the same algebraic multiplicities.

Let E_1 denote the direct sum of the eigenspaces of P_s of eigenvalues λ and λ^{-1} , and E_2 the sum of the other eigenspaces. Since E_1 and E_2 are symplectic subspaces (see for example [2]) then, up to conjugation with a symplectic matrix, we can write $P_s = \text{diag}(\lambda \text{Id}_a, \lambda^{-1} \text{Id}_a, R_1)$ (if $\lambda \neq 1$) or $P_s = \text{diag}(\text{Id}_{2a}, R_1)$ (if $\lambda = 1$) for some $R_1 \in \mathbf{Sp}(n - a - 1, \omega)$.

Since P_u commutes with P_s , it can be written either as $P_u = \text{diag}(U^{tr}, U^{-1}, R_2)$ for some $U \in \text{GL}(a, \mathbb{R})$ unipotent and $R_2 \in \mathbf{Sp}(n - a - 1, \omega)$ (if $\lambda \neq 1$), or $P_u = \text{diag}(U, R_2)$ for some $U \in \mathbf{Sp}(a, \omega)$ unipotent and $R_2 \in \mathbf{Sp}(n - a - 1, \omega)$ (if $\lambda = 1$).

Since $P = P_u P_s$ we have proved the lemma. □

Given an algebraic group $G \subseteq \text{GL}(N, \mathbb{R})$, recall that a torus $T \subseteq G$ is a connected, abelian subgroup whose elements are diagonalizable over \mathbb{C} . Every algebraic group admits at least one torus of maximal dimension, called *maximal torus*, which is unique up to conjugation by an element of G , and the *rank* of G , denoted $\text{rk } G$, is defined as the dimension of a maximal torus of G . We will be mostly concerned with $G = \mathbf{Sp}(N, \omega)$, in which case $\text{rk } G = N$. The following Lemma is a consequence of well-known results, but we could not find any reference in the literature.

Lemma A.2 *Given an algebraic group $G \subseteq \text{GL}(N, \mathbb{R})$, the set G_u of unipotent elements in G is an affine variety of codimension equal to the rank of G .*

Proof The set G_u is invariant under the action of G by conjugation. Fixing B a Borel (i.e., maximal connected solvable) subgroup of G , let B_u denote the subset of unipotent elements in B . Every G -orbit meets B at least once by [4, 11.10], and therefore the map $G \times B_u \rightarrow G_u$ sending (g, A) to gAg^{-1} is surjective. The normalizer $N(B_u) = \{g \in G \mid gB_u g^{-1} \subseteq B_u\}$ coincides with B by a theorem of Chevalley [4, 11.16] and therefore $\dim B_u + \dim G -$

Clearly a matrix $P = \text{diag}(P_1, R)$ in Σ is uniquely determined by $R \in \text{Sp}(n' - 1, \omega)$, $n' = n - n_1 - n_2$, and therefore $\dim \Sigma = \dim \text{Sp}(n' - 1, \omega)$.

We now compute $N(\Sigma)$. Suppose that $n_1 \leq n_2$, and let $\mathcal{A} \subseteq \text{GL}(n_1 + n_2, \mathbb{R})$ be the set of matrices such that

$$A^{\text{tr}} = \left(\begin{array}{ccc|ccc} a_1 & \cdots & a_{n_1} & 0 & b_1 & \cdots & b_{n_1} \\ & & \vdots & \vdots & & \ddots & \vdots \\ & & a_1 & 0 & & & b_1 \\ \hline c_1 & \cdots & c_{n_1} & d_1 & d_2 & \cdots & d_{n_2} \\ & & \vdots & & d_1 & \ddots & \vdots \\ & & c_1 & & & \ddots & d_2 \\ 0 & \cdots & 0 & & & & d_1 \end{array} \right)$$

Clearly $\dim \mathcal{A} = 3n_1 + n_2 \geq 4$. For any matrix $A \in \mathcal{A}$ and $B \in \text{Sp}(n' - 1, \omega)$, the matrix $\text{diag}(A^{\text{tr}}, A^{-1}, B)$ lies in $N(\Sigma)$. In particular, $\dim N(\Sigma) \geq \dim \mathcal{A} + \text{Sp}(n' - 1, \omega)$ and therefore $\dim \mathcal{M} \leq \dim \text{Sp}(n - 1, \omega) - \dim \mathcal{A} \leq \text{Sp}(n - 1, \omega) - 4$.

If $\lambda = 1$ then any $P \in \mathcal{M}_\lambda$ can be written, in a suitable symplectic basis, as $\text{diag}(U, R)$ where $U \in \text{Sp}(n_0, \omega)$, for some n_0 , is unipotent with at least two linearly independent eigenvectors, and $R \in \text{Sp}(n - n_0 - 1, \omega)$. We now define Σ to be the set of matrices that, under the same fixed basis, can be written as $\text{diag}(U', R')$ for some $R' \in \text{Sp}(n' - 1, \omega)$ and some unipotent matrix $U' \in \text{Sp}(n_0, \omega)$. If we let $\text{Sp}(n_0, \omega)_u$ denote the set of unipotent matrices in $\text{Sp}(n_0, \omega)$, we have

$$\dim \Sigma = \dim \text{Sp}(n_0, \omega)_u + \dim \text{Sp}(n - n_0 - 1, \omega).$$

where $\text{Sp}(n_0, \omega)_u$ denote the unipotent matrices in $\text{Sp}(n_0, \omega)$. The normalizer $N(\Sigma)$ contains the matrices of the form $\text{diag}(P_1, R')$ with $P_1 \in \text{Sp}(n_0, \omega)$ and $R' \in \text{Sp}(n - n_0 - 1, \omega)$, thus

$$\dim N(\Sigma) \geq \dim \text{Sp}(n_0, \omega) + \dim \text{Sp}(n - n_0 - 1, \omega).$$

Once again $\text{Sp}(n - 1, \omega)$ acts on \mathcal{M}_λ and by Lemma A.1 every orbit meets Σ . In particular, \mathcal{M}_λ is contained in the space spanned by the orbits of Σ , and $\dim \mathcal{M}_\lambda \leq \dim \text{Sp}(n - 1, \omega) - (\dim N(\Sigma) - \dim \Sigma)$. By the computation above and Lemma A.2, we have

$$\begin{aligned} \dim \mathcal{M}_\lambda &\leq \dim \text{Sp}(n - 1, \omega) - \text{rk } \text{Sp}(n_0, \omega) \\ &= \dim \text{Sp}(n - 1, \omega) - n_0. \end{aligned}$$

The codimension of \mathcal{M}_λ is then ≥ 3 , unless possibly when $n_0 = 1$ or 2 .

If $n_0 = 1$, then every matrix in \mathcal{M}_λ can be written, under some basis, as

$$\text{diag}(\text{Id}_2, R), \quad R \in \text{Sp}(n - 2, \omega) \tag{A.2}$$

Fixing a basis and letting Σ denote the space of matrices that, in the fixed basis, can be written as in (A.2), we have that $\dim \Sigma = \dim \text{Sp}(n - 2, \omega)$ and the normalizer $N(\Sigma)$ contains $\text{Sp}(1, \omega) \times \text{Sp}(n - 2, \omega)$. Therefore, $\dim N(\Sigma) - \dim \Sigma \geq 3$. Using Eq. (A.1), we obtain $\dim \mathcal{M}_\lambda \leq \dim \text{Sp}(n - 1, \omega) - 3$.

If $n_0 = 2$, then every matrix in \mathcal{M}_λ can be written, under some basis, as

$$U = \text{diag}(U_1, U_2) \quad U_i = \begin{pmatrix} 1 & \\ \sigma_i & 1 \end{pmatrix}, \sigma_i \in \{-1, 0, 1\}. \tag{A.3}$$

Fixing a basis and letting Σ denote the space of matrices that, in the fixed basis, can be written as in (A.3), we have $\dim \Sigma = \dim \text{Sp}(n - 3, \omega)$. If for example $\sigma_1 = \sigma_2 = 1$ then $N(\Sigma)$ contains all the matrices of the form $\text{diag}(P_1, R)$ where $R \in \text{Sp}(n - 3, \omega)$ is any matrix, and $P_1 \in \text{Sp}(2, \omega)$ has the form

$$P_2 = \left(\begin{array}{cc|cc} \cos \theta & a & -\sin \theta & b \\ & \cos \theta & & -\sin \theta \\ \sin \theta & c & \cos \theta & d \\ & \sin \theta & & \cos \theta \end{array} \right),$$

where a, b, c, d satisfy the linear equation $(a + b) \cos \theta = (c - d) \sin \theta$. Therefore $\dim N(\Sigma) \geq \dim \text{Sp}(n - 3, \omega) + 4$ and, by Eq. (A.1), we obtain $\dim \mathcal{M}_\lambda \leq \dim \text{Sp}(n - 1, \omega) - 4$. The same computations can be checked for the other values of σ_1 and σ_2 . □

Let $\mathcal{G}, \mathcal{G}_1$ denote the subspaces of $\text{Sp}(n - 1, \omega) \times \mathbb{R}_+$ given by

$$\begin{aligned} \mathcal{G} &= \{(P, \lambda) \in \text{Sp}(n - 1, \omega) \times \mathbb{R}_+ \mid \dim \ker(P - \lambda \text{Id}) \leq 1\} \\ \mathcal{G}_1 &= \{(P, \lambda) \in \text{Sp}(n - 1, \omega) \times \mathbb{R}_+ \mid \dim \ker(P - \lambda \text{Id}) = 1\} \end{aligned}$$

Clearly \mathcal{G} is open and dense in $\text{Sp}(n - 1, \omega) \times \mathbb{R}_+$, $\mathcal{G}_1 \subseteq \mathcal{G}_0$ and, by construction, we have $\text{Sp}_1(n - 1, \omega) \times \mathbb{R}_+ \subseteq \mathcal{G}$.

Proposition A.4 *The map*

$$\begin{aligned} \chi : \mathcal{G} &\longrightarrow \mathbb{R} \\ (P, \mu) &\longmapsto \det(P - \mu \text{Id}) \end{aligned}$$

is a submersion in a neighbourhood of \mathcal{G}_1 .

Proof We are going to prove a stronger statement. In fact, we prove that for any $(S, \lambda) \in \mathcal{G}_0$ we can find a vector $v_{(S,\lambda)} \in T_S \mathbf{Sp}(n-1, \omega)$ such that $d_{(S,\lambda)} \chi(v_{(S,\lambda)}) > 0$.

Let a denote the algebraic multiplicity of λ in S . By Lemma A.1, there is a symplectic basis such that S can be written as $S = \text{diag}(S_1, S_2)$ where $S_1 \in \mathbf{Sp}(a, \omega)$ only contains the eigenvalues λ, λ^{-1} and $S_2 \in \mathbf{Sp}(n-1-a, \omega)$ has eigenvalues different from λ and λ^{-1} .

If $\lambda \neq 1$ then by Lemma A.1 we can write $S_1 = \text{diag}(\lambda U^{\text{tr}}, (\lambda U)^{-1})$, where

$$\lambda U^{\text{tr}} = \begin{pmatrix} \lambda & 1 & & \\ & \lambda & \ddots & \\ & & \ddots & 1 \\ & & & \lambda \end{pmatrix}.$$

If $\lambda > 1$ let $v = \text{diag}(-E_{a,1}, E_{1,a}) \in \mathfrak{sp}(a, \omega)$ otherwise let $v = (-1)^a \text{diag}(-E_{a,1}, E_{1,a})$. In either case, let $v' = \text{diag}(v, 0) \in \mathfrak{sp}(n-1, \omega)$ and $v_{(S,\lambda)} = L_{S*} v' \in T_S \mathbf{Sp}(n-1, \omega)$, one can compute

$$d_{(S,\lambda)} \chi(v_{(S,\lambda)}) = \lambda |\lambda - \lambda^{-1}|^a > 0.$$

If $\lambda = 1$, then S_1 can be written in the following block form

$$S_1 = \begin{pmatrix} U^{-1} & \\ & U^{\text{tr}T} U^{\text{tr}} \end{pmatrix}$$

where T is a symmetric matrix, and

$$U^{\text{tr}} = \begin{pmatrix} 1 & \cdots & 1 \\ & \ddots & \vdots \\ & & 1 \end{pmatrix}, \quad U^{-1} = \begin{pmatrix} 1 & & \\ -1 & \ddots & \\ & & -1 & 1 \end{pmatrix}$$

In order for S_1 to have geometric multiplicity 1, it must be $(UT)_{a,a} = c \neq 0$. Without loss of generality we can suppose that the sign of c is $(-1)^{a-1}$. Define

$$v = \begin{pmatrix} 0 & | & E_{1,1} \\ \hline 0 & | & 0 \end{pmatrix} \in \mathfrak{sp}(a, \omega).$$

Letting $v' = \text{diag}(v, 0) \in \mathfrak{sp}(n - 1, \omega)$ and $v_{(S,\lambda)} = L_{S*}v' \in T_S\mathbf{Sp}(n - 1, \omega)$ we can easily compute that

$$d_{(S,\lambda)}\chi(v_{(S,\lambda)}) = (-1)^{a-1}c > 0.$$

□

From Proposition A.4 we obtain the following stronger, more global result.

Proposition A.5 *There exists a vector field V on $\mathbf{Sp}_1(n - 1, \omega)$ such that for every $S \in \mathbf{Sp}_1(n - 1, \omega)$ and every real eigenvalue λ of S , $d_{(S,\lambda)}\chi(V) > 0$.*

Proof Given $S \in \mathbf{Sp}_1(n - 1, \omega)$ and $\lambda \in (0, 1]$ a real eigenvalue of S , let $v_{(S,\lambda)} \in T_S\mathbf{Sp}_1(n - 1, \omega)$ be the vector constructed in the previous proposition, so that $d_{(S,\lambda)}\chi(v_{(S,\lambda)}) > 0$. It can be easily checked that, at the point (S, λ^{-1}) , the differential of χ is

$$d_{(S,\lambda^{-1})}\chi(v_{(S,\lambda)}) = -\lambda^{1-2a}(\lambda - \lambda^{-1})^a > 0$$

and moreover for any other eigenvalue μ of S different from λ and λ^{-1} , one has $d_{(S,\mu)}\chi(v_{(S,\lambda)}) = 0$. In particular, letting $v_S = \sum_{\lambda} v_{(S,\lambda)}$, where the sum is taken over all the real eigenvalues of S in $(0, 1]$, the v_S satisfies

$$d_{(S,\lambda)}\chi(v_S) > 0$$

for every real eigenvalue λ of S . By continuity, we can find a neighbourhood U_S of S and an extension V_S of v_S such that for every $S' \in U_S$ and λ' real eigenvalue of S' , we have $d_{(S',\lambda')}\chi(V_S) > 0$.

The open sets $\{U_S\}_{S \in \mathbf{Sp}_1(n-1,\omega)}$ form an open cover of $\mathbf{Sp}_1(n - 1, \omega)$. Choosing a countable subcover $\{U_{S_i}\}_i$ with a subordinate partition of unity $\{\lambda_i\}_i$, the vector field

$$V = \sum_i \lambda_i V_{S_i}$$

has the required properties. □

Proposition A.4 implies that \mathcal{G}_1 is a smooth hypersurface in \mathcal{G} . Consider now the projection $\pi : \mathcal{G}_0 \rightarrow \mathbb{R}$, sending (Q, λ) to λ , and let $\mathcal{G}_0 = \pi^{-1}(1)$.

Lemma A.6 *The map $\pi : \mathcal{G}_1 \rightarrow \mathbb{R}$ is a submersion around \mathcal{G}_0 .*

Proof It is enough to find, for every point $(Q, 1) \in \mathcal{G}_1$, a vector $v \in T_{(Q,1)}\mathcal{G}_1$ such that $d_{(Q,1)}\pi(v) \neq 0$. Fixing $(Q, 1)$, we know in particular that 1 is an eigenvalue of Q and therefore Q can be written, in some basis, as $Q = \text{diag}(P, R)$

where $P \in \text{Sp}(a, \omega)$, $R \in \text{Sp}(n - 1 - a, \omega)$ and moreover

$$P = \begin{pmatrix} U^{-1} & 0 \\ U^{\text{tr}}T & U^{\text{tr}} \end{pmatrix}$$

with T symmetric and U unipotent such that

$$U^{\text{tr}} = \begin{pmatrix} 1 & \cdots & 1 \\ & \ddots & \vdots \\ & & 1 \end{pmatrix}.$$

For some t small, let

$$P(t) = \begin{pmatrix} e^{-t}U^{-1} & 0 \\ e^tU^{\text{tr}}T & e^tU^{\text{tr}} \end{pmatrix}, \quad Q(t) = \text{diag}(P(t), R).$$

Then the path $(Q(t), e^{-t})$ lies in \mathcal{G}_0 for small t , and $\pi(Q(t), e^{-t}) = e^{-t}$. In particular, letting

$$v = \left. \frac{d}{dt} \right|_{t=0} (Q(t), e^{-t})$$

we obtain $d_{(Q,1)}\pi(v) \neq 0$ thus proving the Lemma. □

The following Corollary is straightforward

Corollary A.7 *The subset $\pi^{-1}((0, 1]) \subseteq \mathcal{G}_1$ is a smooth manifold, with boundary \mathcal{G}_0 .*

References

1. Allday, C., Puppe, V.: Cohomological Methods in Transformation Groups. Cambridge Studies in Advanced Mathematics, vol. 32. Cambridge University Press, Cambridge (1993)
2. Ballmann, W., Thorbergsson, G., Ziller, W.: Closed geodesics on positively curved manifolds. *Ann. Math.* **116**(2), 213–247 (1982)
3. Besse, A.L.: Manifolds all of Whose Geodesics are Closed. *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 93 (1978)
4. Borel, A.: Linear Algebraic Groups. Second Enlarged Edition. Springer, New York (1991)
5. Bredon, G.E.: Introduction to Compact Transformation Groups, vol. 46 (Pure and Applied Mathematics). Elsevier, New York (1972)
6. Cornea, O., Lupton, G., Oprea, J., Tanré, D.: Lusternik-Schnirelmann Category. *Mathematical Surveys and Monographs*, vol. 103 (2003)
7. Gromoll, D., Grove, K.: On metrics on S^2 all of whose geodesics are closed. *Invent. Math.* **65**, 175–177 (1981)
8. Hingston, N.: Equivariant Morse theory and closed geodesics. *J. Differ. Geom.* **19**, 85–116 (1984)

9. Hilton, P.J., Mislin, G., Roitberg, J.: Localization of Nilpotent Groups and Spaces, *Notas de Matematica*. North-Holland, Amsterdam (1975)
10. McCleary, J.: On the mod- p Betti numbers of loop spaces. *Invent. Math.* **87**(3), 643–654 (1987)
11. Olsen, J.: Three-dimensional manifolds all of whose geodesics are closed. *Ann. Glob. Anal. Geom.* **37**(2), 173–184 (2010)
12. Schwarz, A.S.: The homologies of spaces of closed curves. *Trudy Moskov. Mat. Obšč.* **9**, 3–44 (1960)
13. Taimanov, I.: The type numbers of closed geodesics. *Regul. Chaot. Dyn.* **15**(1), 84–100 (2010)
14. Wadsley, A.W.: Geodesic foliations by circles. *J. Differ. Geom.* **10**(4), 541–549 (1975)
15. Weinstein, A.: On the volume of manifolds all of whose geodesics are closed. *J. Differ. Geom.* **9**(4), 513–517 (1974)
16. Weinstein, A.: Symplectic V-manifolds, periodic orbits of Hamiltonian systems, and the volume of certain Riemannian manifolds. *Commun. Pure Appl. Math.* **30**, 265–271 (1977)
17. Wilking, B.: Index parity of closed geodesics and rigidity of Hopf fibrations. *Invent. Math.* **144**(2), 281–295 (2001)
18. Ziller, W.: The free loop space of globally symmetric spaces. *Invent. Math.* **41**, 1–22 (1977)
19. Ziller, W.: Geometry of the Katok examples. *Ergod. Theory Dyn. Syst.* **3**(1), 135–157 (1983)